

Connectivity Bounds in Graphs and Beyond

Erdős Problem 915, from Proof to Search

Louis Vandenbruwaene

Supervisor: Prof. Stijn Cambie
KU Leuven

Thesis presented in
fulfillment of the requirements
for the degree of Master of Science
in Mathematics

Academic year 2026–2027

Open Access Disclaimer

This master's thesis is a student work submitted as part of an academic programme at KU Leuven. The work is made available for examination and archiving in accordance with the applicable KU Leuven regulations and open-access policy for master's theses.

© Copyright by KU Leuven

Without written permission from the supervisors and the author, it is forbidden to reproduce or adapt in any form or by any means any part of this publication. Requests for obtaining the right to reproduce or utilize parts of this publication should be addressed to KU Leuven, Faculty of Science, Celestijnenlaan 200H - box 2100, 3001 Leuven (Heverlee), Telephone +32 16 32 14 01.

Written permission from the supervisor is also required to use the methods, products, schematics, and programs described in this work for industrial or commercial use, and for submitting this publication in scientific contests.

Foreword

This thesis studies how many edges or arcs a network can contain when no pair of vertices is allowed to have too many independent routes between them. The starting point is Erdős Problem 915, an extremal question about edge-disjoint paths in simple graphs. The work is told as a journey. It begins with the variants whose answers can be proved cleanly by hand, watches those proofs grow longer and more delicate as the model changes, and at the point where a single argument no longer suffices it turns to a unified program: one representation for every variant, an exact connectivity checker built on maximum flow (a standard routine for finding the most independent routes between two points), a certificate engine that turns finite checks into proofs, and a temperature-guided search that hunts for extremal graphs in the space of all graphs.

I am grateful to my supervisor, Prof. Stijn Cambie, and my daily advisor for their guidance, corrections, and mathematical feedback throughout the project. I also thank the people who discussed early versions of the arguments, checked examples, or helped clarify the presentation. Any remaining mistakes are my own.

Contribution Statement

The author was responsible for the literature study, the formulation of the variants, the design and implementation of the unified program, the construction and verification of the examples, the writing of the proofs included in the thesis, and the final $\text{\LaTeX}$ implementation. Results from the literature are cited explicitly, in particular the theorems of Mader, Leonard, Sørensen-Thomassen, Menger, Gomory-Hu, Baranyai, the hypergraph connectivity results of Bérczi, Chandrasekaran, Király and Kulkarni, and the probabilistic connectivity results used for random graphs.

The connectivity checker computes local connectivity exactly through maximum flow. The certificate engine yields a genuine upper-bound proof for a fixed size whenever it reports an optimal solution with zero gap over a finite domain. The temperature-guided search is a discovery tool: it produces constructions and therefore lower bounds, proposes conjectures, and rediscovers known extremal graphs, but it never certifies an upper bound on its own. Every general statement in the thesis rests on a hand proof or a cited theorem. Computational searches are supporting evidence, not substitutes for proofs. AI language models, including Claude Code, ChatGPT, and Gemini, were used as tools for literature search assistance, $\text{\LaTeX}$ editing, proofreading, and exploring proof drafts. Every mathematical statement and proof included in the thesis was checked by the author, who takes full responsibility for the content.

The following new results are contributed by this thesis. The hypergraph vertex problem is solved completely at $m = 2$ and $m = 3$ for all uniformities r and all n (Theorems A.34 and A.37): the vertex-disjoint and edge-disjoint extremal values coincide, equalling $\lfloor (m-1)(n-1)/(r-1) \rfloor$. The $m = 3$ proof rests on a new rank bound for incidence graphs, derived via Tutte's triconnected decomposition (Lemma A.36). For the directed multigraph problem at $m = 3$, an unconditional attachment lemma (Lemma A.26) and an odd-step theorem (Theorem A.28, Remark A.29) reduce the entire problem for all n to two finite machine-checkable statements about multigraphs on seven vertices, without requiring the unproved minimum-degree conjecture.

Short Summary

Erdős Problem 915, posed by Bollobás and Erdős [4], asks for the maximum number of edges in a graph on n vertices such that no two vertices are connected by m edge-disjoint paths. In modern notation this asks for

$$\ell_m(n) = \max\{|E(G)| : |V(G)| = n, \lambda_G^{\max} \leq m - 1\}.$$

Mader proved that, for simple graphs and $n \geq m \geq 2$,

$$\ell_m(n) = \left\lfloor \frac{m(n-1)}{2} \right\rfloor.$$

This thesis revisits this theorem and studies the corresponding extremal problem under three independent changes: the graph model (simple graphs, multigraphs, and hypergraphs), the path-separation notion (edge-disjoint or internally vertex-disjoint), and the presence or absence of directions. Sampled random graphs $G(n, p)$ run alongside throughout as a lens on the typical case rather than as a separate model.

The thesis is organised as a journey from hand proofs to computation. The early variants are settled rigorously: the multigraph value $L_m(n) = (m-1)(n-1)$, the random-graph threshold $p^* = m/n$, the equality of the edge and vertex problems for $m \leq 4$, and the first divergence at $m = 5$, where Sørensen and Thomassen prove $k_5(n) = \lfloor 8n/3 \rfloor - 3$. The directed case forces the turn: the value $\ell_2^{\text{dir}}(n) = \max(2(n-1), \lfloor n^2/4 \rfloor)$ is proved by a long induction, and for $m \geq 3$ the hand proof gives way to a conjecture supported by an explicit counterexample. The centerpiece is then a unified program. One multiplicity-matrix representation models almost all twelve variants, an exact maximum-flow checker measures local connectivity, a cut-counting optimisation certifies upper bounds for small sizes, and a simulated-annealing search guided by edge sensitivity discovers extremal graphs. The program rediscovers the double star, the hub, the one-directional bipartite digraph, and the augmented-bipartite construction as extremal graphs, certifies the directed multigraph values $L_m^{\text{dir}}(n) = 2(n-1)(m-1)$ for $n \leq 6$, and frames the remaining open problems on the directed frontier.

The interplay then pays back in the other direction. The hypergraph vertex problem is settled completely at $m = 2$ and $m = 3$: $k_m^{(r)}(n) = \lfloor (m-1)(n-1)/(r-1) \rfloor$ for every uniformity r , the $m = 3$ case through a new rank bound on incidence graphs proved by way of Tutte's triconnected decomposition. On the directed side, an unconditional attachment lemma and a conditional odd-step theorem reduce the entire directed multigraph problem at $m = 3$, value and extremal characterisation for all n , to two finite, machine-checkable statements about multigraphs on seven vertices, and directed hypergraphs receive their first bounds, quadratic in n .

Abbreviations and Symbols

GH Gomory-Hu
w.h.p. with high probability

$G = (V, E)$	graph with vertex set V and edge set E
$D = (V, A)$	directed graph with vertex set V and arc set A
n	number of vertices
m	forbidden-connectivity parameter, with all local connectivities at most $m - 1$
$\lambda_G(u, v)$	maximum number of edge-disjoint u - v paths
$\lambda_G^{\max}$	maximum local edge-connectivity over all vertex pairs
$\kappa_G(u, v)$	maximum number of internally vertex-disjoint u - v paths
$\kappa_G^{\max}$	maximum local vertex-connectivity over all vertex pairs
$\ell_m(n)$	simple undirected edge-disjoint extremal function
$L_m(n)$	undirected multigraph edge-disjoint extremal function
$k_m(n)$	simple undirected vertex-disjoint extremal function
$\ell_m^{\text{dir}}(n)$	simple directed arc-disjoint extremal function
$L_m^{\text{dir}}(n)$	directed multigraph arc-disjoint extremal function
$k_m^{\text{dir}}(n)$	simple directed vertex-disjoint extremal function
$\mu(u, v)$	multiplicity of an edge or arc between u and v
$\sigma(e)$	edge sensitivity $\lambda^{\max}(G) - \lambda^{\max}(G - e)$
$G(n, p)$	binomial random graph

Contents

Foreword	II
Contribution Statement	III
Short Summary	IV
Abbreviations and Symbols	V
1 The Problem and Its Base Cases	1
1.1 Graphs and routes	1
1.2 Erdős Problem 915	2
1.3 Three axes of variation	3
1.4 Cases with short proofs	4
1.5 The first divergence: vertex-disjoint paths	5
1.6 The directed case and the quadratic phenomenon	6
1.7 The failure of direct proof for $m \geq 3$	9
1.8 The hypergraph variant	11
1.9 Random graphs as intuition	12
1.10 Where this work sits	14
2 Certifying Bounds by Machine	16
2.1 The solver	17
2.2 A single representation for all variants	17
2.3 The connectivity checker built on maximum flow	18
2.4 Checking every graph of a fixed size	22
2.5 Reaching further: a pruned search for the directed base cases	22
2.6 Proving every m at once	23
2.7 What the solver may claim	27
2.8 Reproducibility	27

3	Discovering Bounds by Search	28
3.1	The wall: how enumeration explodes	28
3.2	The temperature search	29
3.3	Temperature and edge sensitivity	31
3.4	Keeping the checker cheap inside the search	32
3.5	Rediscovering the known extremisers	34
3.6	The trichotomy: proved, certified, conjectured	35
4	Synthesis, Results, and Open Problems	42
4.1	The directed frontier and the single open lemma	42
4.2	Three principal phenomena	44
4.3	Summary of the twelve variants	48
4.4	Contributions and limitations of the program	48
4.5	Open problems	53
A	Full Proofs and Computational Transcripts	55
A.1	Menger, Gomory–Hu, and the degree bound	55
A.2	Mader’s theorem in full	56
A.3	The directed arc value at $m = 2$	59
A.4	Structural propositions on the directed frontier	61
A.5	The directed multigraph problem at large n	62
A.6	The hypergraph bounds	67
A.7	The hypergraph vertex problem	69
A.8	The random threshold	73
A.9	Computational transcripts	74
A.10	How the program checks itself	75
B	Supplementary Figures	77
B.1	Search traces across the variants	77
B.2	The variant landscape at $m = 6$ and in three dimensions	80

Chapter 1

The Problem and Its Base Cases

A network is useful not only because it connects, but because it connects in more than one way. A single road between two towns is a liability, and a second independent road is insurance. The mathematical content of that intuition is the number of routes between two points that share no segment, and the question this thesis circles is how many connections a network can hold before some pair of points is forced to have too many such routes. This chapter builds the question from nothing, settles the variants whose proofs are short enough to read in an afternoon, and then follows the same question along three axes until the proofs stretch past breaking. By the end the case for a machine has made itself.

1.1 Graphs and routes

To state the question precisely we need only a few ideas. A *graph* G is a finite set of points, called *vertices*, together with a set of links, called *edges*, each joining two of the vertices. One pictures the vertices as dots and the edges as lines between them.




Figure 1.1. A simple graph on five vertices: K_5 with the two edges $\{0, 4\}$ and $\{1, 3\}$ removed, leaving eight of the original ten edges. The five dots are the vertices and the lines between them are the edges. Since at most one line joins any given pair and no edge loops back to its own endpoint, this is a simple graph.

A graph is the bare skeleton of a network: the vertices are the towns or the devices, and the edges are the roads or the links between them. We write n for the number of vertices, and the *degree* of a vertex is the number of edges meeting it. Unless we say otherwise our graphs are *simple*, meaning that at most one edge joins any given pair of vertices and its endpoints are distinct.

A *path* from a vertex u to a vertex v is a sequence of distinct vertices that starts at u , ends at v , and joins each consecutive pair by an edge. It is the formal version of a route through the network. Two paths between the same endpoints are *edge-disjoint* if no edge belongs to both, so that the

loss of a single edge can break at most one of them. The largest number of pairwise edge-disjoint paths between u and v is their *local edge-connectivity* $\lambda_G(u, v)$, the count of genuinely independent routes the network offers to that pair. The pair with the most independent routes sets the ceiling for the whole graph:

$$\lambda_G^{\max} = \max_{u \neq v} \lambda_G(u, v),$$

the largest local edge-connectivity anywhere in G . A theorem of Menger [15], recalled in [Chapter A](#), gives this quantity a second face: $\lambda_G(u, v)$ also equals the least number of edges one must remove to cut every route from u to v . Many independent routes between a pair is exactly the same thing as a costly cut between them.

1.2 Erdős Problem 915

The constraint at the heart of this thesis is a ceiling on connectivity. Requiring $\lambda_G^{\max} \leq m - 1$ says that no pair of vertices is joined by as many as m edge-disjoint paths, or equivalently that every pair can be severed by removing fewer than m edges. The question is how many edges a graph can hold while staying under that ceiling.

Question 1.1 (Erdős Problem 915 [5]). How many edges can a graph on n vertices have if no two of its vertices are joined by m edge-disjoint paths?

Writing the answer as an extremal function,

$$\ell_m(n) = \max\{ |E(G)| : |V(G)| = n, \lambda_G^{\max} \leq m - 1 \},$$

the problem asks for $\ell_m(n)$, the most edges possible under the ceiling. Mader settled the simple-graph case completely.

Theorem 1.2 (Mader [12]). *For all $n \geq m \geq 2$,*

$$\ell_m(n) = \left\lfloor \frac{m(n-1)}{2} \right\rfloor.$$

An elegant proof by way of the Gomory–Hu tree, found independently of Mader, is given in full in [Chapter A](#), together with a characterisation of the graphs that attain equality ([Theorem A.11](#)). We single out the one identity that makes it work, because the same idea reappears when the program of [Chapter 2](#) stores connectivities as a tree (graph without cycles): a Gomory–Hu tree of G is a weighted tree on the same vertex set in which $\lambda_G(u, v)$ equals the lightest edge on the tree path from u to v . All $\binom{n}{2}$ local connectivities are therefore encoded by $n - 1$ numbers, and $\lambda_G^{\max}$ is simply the heaviest edge of the tree.

The whole argument ([Chapter A](#)) is one act of double counting on that tree. Each of the $n - 1$ tree edges stands for a minimum cut of G , and an original edge of G is charged once for every tree edge lying between its endpoints, so summing the tree-edge weights counts every edge of G .

at least once and, crucially, counts most of them at least twice. The only edges charged a single time are those joining tree-adjacent vertices, and a *simple* graph has at most $n - 1$ of those, one per tree edge. Every other edge is charged twice. Since each tree weight is a local connectivity and so at most $m - 1$, the two counts meet at $2|E(G)| - (n - 1) \leq (m - 1)(n - 1)$, which rearranges to the floor. The factor of two, and with it the gap between Mader's $\lfloor m(n - 1)/2 \rfloor$ and the multigraph value $(m - 1)(n - 1)$ proved below, is exactly the dividend of simplicity, the one place the argument uses that no pair carries a parallel edge.

1.3 Three axes of variation

Mader's theorem is the floor we build on, and everything afterwards is a variation on it. There are three independent ways to change the rules, and they generate the landscape this thesis traverses.

The first axis is the *model*, drawn in Figure 1.2. A simple graph allows at most one edge between two vertices. A *multigraph* allows parallel edges, recorded by a multiplicity $\mu(u, v)$, so a single pair can already carry several independent routes. A *hypergraph* lets an edge join more than two vertices at once: a hyperedge is a set of vertices rather than a pair, and a route may pass through it between any two of the vertices it contains. These three models are the three rows of every comparison figure in this thesis.

The second axis is the *direction*: arcs may be one-way, and then a network can be lopsided in a manner no undirected graph can imitate. Although it might seem a generalization of the undirected setting, they are distinct in complexity and methodology: the arcs $u \rightarrow v$ and $v \rightarrow u$ offer a round trip while the edge $u \rightarrow v$ doesn't. This distinction fades when we no longer distinguish paths by their edges, as in our next choice of interpretation.

The third axis is the *separation*: routes may be required to avoid sharing an edge, as above, or the stricter *internally vertex-disjoint*, sharing no intermediate vertex, which gives the local vertex-connectivity $\kappa_G(u, v)$ and its maximum $\kappa_G^{\max}$. Whitney's inequality $\kappa_G(u, v) \leq \lambda_G(u, v)$ [20]




Figure 1.2. The first axis of variation, the *model*. In a *simple* graph each pair of vertices is joined by at most one edge. A *multigraph* permits parallel edges, here a triple and a double, so the multiplicity $\mu(u, v)$ counts how many copies join a pair. A *hypergraph* replaces pairwise edges by hyperedges: the violet region spans three vertices and the blue region spans four, and they share a single vertex where the two colours blend, illustrating hyperedges of different sizes that may overlap. When all hyperedges have the same size r we call it an r -uniform hypergraph. The same connectivity question, how many independent routes a pair can be forced to share, is asked of all three.

records that avoiding shared vertices is at least as demanding as avoiding shared edges, and [Figure 1.3](#) shows the gap on a graph where two routes are edge-disjoint yet forced through one common vertex.




Figure 1.3. The third axis, the *separation*. The two s – t routes drawn here, one orange and one blue, share no edge, so they count as two edge-disjoint paths and $\lambda(s, t) = 2$. Both pass through the single vertex w , however, so they are *not* internally vertex-disjoint, and since every s – t route must cross w we have $\kappa(s, t) = 1$. This is Whitney’s inequality $\kappa \leq \lambda$ made concrete: the gap between the two separations is exactly what the vertex-disjoint problem measures, and the rest of the chapter shows it stays shut until $m = 5$.

Three models, two directions, two separations: twelve versions of one question. The thesis is the attempt to answer as many of them as honest mathematics allows, and to build a machine for the rest. Some answers are short and we prove them here. Some answers are long, and their length is the reason [Chapter 2](#) exists. The rest of this chapter collects the variants whose proofs are short enough to belong in the body, so that the reader meets the easy slope before the climb.

1.4 Cases with short proofs

Allowing parallel edges removes the parity subtlety that produces the floor in Mader’s theorem, and the answer becomes exactly linear. Write $L_m(n)$ for the multigraph analogue of $\ell_m(n)$.

Theorem 1.3. *For multigraphs on n vertices, $L_m(n) = (m - 1)(n - 1)$.*

Proof. For the lower bound, take a spanning tree of K_n , that is, a connected cycle-free subgraph spanning all n vertices, which necessarily has $n - 1$ edges, and give every tree edge multiplicity $m - 1$. The result has $(m - 1)(n - 1)$ edges. Any two vertices are joined by a unique path in the tree, of some length $k \geq 1$. The minimum edge cut between them is the lightest tree edge on that path, of capacity $m - 1$, so $\lambda(u, v) = m - 1$ and $\lambda^{\max} = m - 1$.

For the upper bound, suppose $\lambda_G^{\max} \leq m - 1$ and build a Gomory–Hu tree T of G ([Theorem A.2](#)). By that theorem, each tree edge e corresponds to a minimum cut of G whose capacity $c(e) = \lambda_G(u, v) \leq m - 1$ for the pair (u, v) it represents. Every original edge $\{x, y\}$ of G crosses the cut corresponding to the minimum-weight tree edge on the tree path from x to y , so summing cut capacities over the $n - 1$ tree edges counts each original edge at least once. Hence $|E(G)| \leq \sum_{e \in T} c(e) \leq (m - 1)(n - 1)$. $\square$

The vertex-disjoint question is just as quick at the first nontrivial value. Forbidding two internally vertex-disjoint paths is a strong constraint: it forbids cycles.

Proposition 1.4. *For simple graphs, $k_2(n) = n - 1$, attained exactly by trees.*

Proof. If G contains a cycle, then any two vertices on that cycle are joined by the two arcs of the cycle, which are internally vertex-disjoint, so $\kappa_G^{\max} \geq 2$. Hence $\kappa_G^{\max} \leq 1$ forces G to be acyclic, and an acyclic graph on n vertices has at most $n - 1$ edges, with equality for trees. A tree indeed has $\kappa^{\max} = 1$, since removing the single intermediate vertex on any path disconnects its endpoints. $\square$

Here $k_m(n)$ denotes the vertex-disjoint extremal function, the counterpart of $\ell_m(n)$ for $\kappa^{\max}$. We have just seen $k_2(n) = n - 1 = \ell_2(n)$. The agreement of the edge and vertex problems is no accident at small m , and the moment it breaks is the first place the proofs begin to strain.

1.5 The first divergence: vertex-disjoint paths

Replacing edge-disjoint by internally vertex-disjoint routes changes nothing at first. Leonard proved that the edge and vertex problems give the same answer while m stays small.

Theorem 1.5 (Leonard [11]). *For simple graphs and $m \leq 4$,*

$$k_m(n) = \ell_m(n) = \left\lfloor \frac{m(n-1)}{2} \right\rfloor.$$

For $m \leq 4$ the extremal graphs for the two problems coincide: Whitney's inequality $\kappa \leq \lambda$ is tight where it matters and the harder vertex constraint costs nothing (Figure 1.4, the three overlapping blue curves). One is tempted to expect this forever. It fails at the very next value, and the failure is sharp.

Theorem 1.6 (Sørensen-Thomassen [18]). *For simple graphs and all $n \geq 10$,*

$$k_5(n) = \left\lfloor \frac{8n}{3} \right\rfloor - 3 > \left\lfloor \frac{5(n-1)}{2} \right\rfloor = \ell_5(n).$$

The threshold $n \geq 10$ is where the inequality first holds with strict positivity: at $n = 10$ the vertex formula gives 23 and the edge formula gives 22, while at $n = 7$ they tie at 15. For $n \leq 9$ one must verify the formula directly against the structural classification Sørensen-Thomassen supply; the asymptotic formula equals or overestimates $k_5(n)$ for small values, so the gap between k_5 and ℓ_5 is not yet guaranteed by the formula alone.

At $m = 5$ the vertex problem genuinely diverges from the edge problem, visible in Figure 1.4 as the shaded gap that opens between the dashed blue and solid orange curves. The reason is structural. Leonard's method leans on the extremal graphs being assembled from cliques glued along shared cliques. A k -tree is the graph built by starting from a complete graph K_{k+1} and repeatedly adding a new vertex adjacent to exactly the vertices of some k -clique already present. These chordal scaffolds are exactly the building blocks the method uses. Through $m = 4$ those scaffolds are forced, and counting their edges gives the bound. At $m = 5$ they are no longer the unique extremal shape. Additional configurations appear, and the clean count breaks. Sørensen

and Thomassen recover the exact value by a delicate structural induction, which we cite rather than reproduce. For $m \geq 6$ even that much is unknown: the vertex-disjoint problem has stood open since 1974, and the extremal structure is genuinely not understood. It is the one direction this thesis does not attempt to force, and [Chapter 4](#) explains why.




Figure 1.4. The edge and vertex extremal functions for $m = 2, 3, 4$ (blue shades, single curve per m since they agree) and $m = 5$ (dashed blue for edge, solid orange for vertex). Through $m = 4$ the two problems give identical answers, and at $m = 5$ the vertex formula $k_5(n) = \lfloor 8n/3 \rfloor - 3$ separates above the edge formula $\ell_5(n) = \lfloor 5(n-1)/2 \rfloor$, with the gap shaded.

1.6 The directed case and the quadratic phenomenon

Direction breaks a symmetry that undirected graphs never had. In an undirected graph the number of edges is tied to the connectivity through every vertex's degree. In a digraph a vertex can have many arcs in and none out, and a whole graph can lean entirely one way. Send every arc from one half of the vertices to the other and nothing comes back: between any two vertices there is at most one route, yet the number of arcs is quadratic.

Construction 1.7 (One-directional bipartite digraph). Split V into parts A and B with $|A| = \lfloor n/2 \rfloor$ and $|B| = \lceil n/2 \rceil$, and include every arc from A to B . All arcs run from A to B , and none within A , within B , or from B back to A . Since B has no outgoing arcs, a vertex in A cannot reach any other vertex of A , and two vertices of B cannot communicate at all. For any $a \in A$ and $b \in B$ the only directed path from a to b is the single arc $a \rightarrow b$, so $\lambda(a, b) = 1$ and $\lambda^{\max} = 1$. The arc count is $|A| \cdot |B| = \lfloor n^2/4 \rfloor$.

[Figure 1.5](#) draws the construction. Every arrowhead lands in B and none leaves it, so the picture is a wall of one-way traffic: from any a on the left there is exactly one way to reach any b on the right, the single arc between them, and there is no way at all to travel between two vertices on the same side. The arc count $|A| \cdot |B|$ is quadratic, yet no pair carries more than one route. This is the lopsidedness direction buys, and it has no undirected analogue. It competes with a

second construction built around a single centre.




Figure 1.5. The one-directional complete bipartite digraph $B_{k,k}$ of [Construction 1.7](#), drawn here at $n = 8$ with $|A| = |B| = 4$. Every one of the $|A| \cdot |B| = \lfloor n^2/4 \rfloor$ arcs runs left to right, from A to B . There are no arcs inside A , inside B , or back from B to A . Because B has no outgoing arcs and A no incoming ones, the only route from any a_i to any b_j is the single arc $a_i \rightarrow b_j$, so every pair has local edge-connectivity 1 while the arc count grows quadratically.

In a digraph each vertex has two distinct degree counts. The *out-degree* $d^+(v)$ is the number of arcs leaving v , and the *in-degree* $d^-(v)$ is the number entering v . A vertex with high out-degree fans many arcs outward; one with high in-degree receives many. Keeping both counts balanced is the structural idea behind the next construction.

Construction 1.8 (Directed hub). For $n \geq m \geq 2$, take a hub vertex h and label the remaining $N = n - 1$ vertices $0, 1, \dots, N - 1$. Add the $2N$ hub arcs $h \rightarrow i$ and $i \rightarrow h$ for every i , and on the non-hub vertices add the circulant arcs $i \rightarrow i + j \pmod{N}$ for $j = 1, \dots, m - 2$. The total is $2N + (m - 2)N = m(n - 1)$ arcs. Every non-hub vertex has $d^+ = d^- = 1 + (m - 2) = m - 1$, and any ordered pair of vertices contains a non-hub endpoint, so the directed degree bound ([Lemma A.3](#)) gives $\lambda^{\max} \leq m - 1$.

At $m = 2$ this construction is easiest to picture: the circulant layer is empty and what remains is the *bidirected star* of [Figure 1.6](#), a hub joined to each spoke by a matched pair of opposite arcs. Every spoke can reach every other only by going in to the hub and back out, two hops, so no pair is ever joined by two independent routes, and the arc total is exactly $2(n - 1)$. For $m \geq 3$ not all $m(n - 1)$ arcs can touch the hub, since a simple digraph has at most $2(n - 1)$ arcs incident to one vertex: the surplus lives in the circulant layers among the spokes, tuned so that no degree exceeds the budget. The two constructions compete on different scales, linear against quadratic, and which one wins depends on n . At $m = 2$ the competition resolves exactly.

Theorem 1.9 (Directed arc value at $m = 2$). *For simple digraphs,*

$$\ell_2^{\text{dir}}(n) = \max\left(2(n - 1), \left\lfloor \frac{n^2}{4} \right\rfloor\right).$$

The hub construction leads for $n < 7$, the one-directional bipartite construction leads for $n > 7$,




Figure 1.6. The directed hub at $n = 6$ for two values of m . *Left* ($m = 2$): the bidirected star. The hub h is joined to each spoke by one arc in each direction, giving $2(n-1) = 10$ arcs. No circulant layer is added; all spoke communication routes through h , so $\lambda^{\max} = 1$. *Right* ($m = 3$): a single circulant layer is added among the five spokes. The blue arcs are the same hub-spoke pairs as on the left. The orange arcs are that one circulant layer, the cycle $0 \rightarrow 1 \rightarrow 2 \rightarrow 3 \rightarrow 4 \rightarrow 0$, so each spoke has $d^+ = d^- = m-1 = 2$. The total is $m(n-1) = 15$ arcs and $\lambda^{\max} = 2$. For $m \geq 3$ the spokes are no longer isolated from one another, and tracing all the routes becomes the argument that fills the rest of the chapter.

and the two tie at $n = 7$, where both reach 12 arcs.

The lower bound is the better of the two constructions. The upper bound is where the trouble starts.

Idea of the upper bound at $m = 2$. The value $M(n) = \max(2(n-1), \lfloor n^2/4 \rfloor)$ is a contest between a linear term and a quadratic one, and which leads is pure arithmetic: $2(n-1) \geq \lfloor n^2/4 \rfloor$ precisely for $n \leq 7$, with equality at $n = 7$, where both give 12. So M follows the bidirected star up to the crossover and the bipartite digraph past it, and $n = 7$ is the single value where the two constructions tie. The proof that no digraph beats $M(n)$ is an induction on n (Chapter A). Delete a vertex of *minimum* total degree: since the degrees sum to twice the arc count, an averaging argument bounds that degree by $2a/n$, where a is the number of arcs, and the remaining digraph on $n-1$ vertices still respects the ceiling, so the inductive bound applies to it. Feeding the two facts together gives $a \leq \frac{n}{n-2} \lfloor (n-1)^2/4 \rfloor$, and a short parity check shows the right-hand side never exceeds $\lfloor n^2/4 \rfloor$ once $n \geq 8$: for even n it lands exactly on the bipartite value, and for odd n a sub-unit slack is swallowed by the fact that a is an integer. The averaging closes the step with no separate hub case, which is why the difficulty is not depth but bulk. Nothing in the argument is deep. It is a careful walk through two parity regimes, with the base cases $n \leq 7$ settled by exhaustive computer search rather than by hand. That experience of a correct but long and almost entirely mechanical case-check is exactly what motivates the chapters that follow. One genuinely wants to hand the case-checking to a machine.

The vertex-disjoint version of the same question needs no separate machinery at $m = 2$. Vertex-disjoint routes are at most as numerous as arc-disjoint ones, and the two constructions above keep $\kappa^{\max} = 1$ as readily as $\lambda^{\max} = 1$, so the value transfers exactly.

Theorem 1.10 (Directed vertex value at $m = 2$). *For simple digraphs,*

$$k_2^{\text{dir}}(n) = \max\left(2(n-1), \left\lfloor \frac{n^2}{4} \right\rfloor\right) = \ell_2^{\text{dir}}(n).$$

1.7 The failure of direct proof for $m \geq 3$

For $m \geq 3$ the induction no longer closes, and the natural guess is wrong. The natural initial conjecture was that the two constructions above continued to be the whole story, giving $\max(m(n-1), \lfloor n^2/4 \rfloor)$. A single graph on ten vertices refutes it.

Remark 1.11 (A thirty-arc counterexample). Take $A = \{a_1, \dots, a_5\}$ and $B = \{b_1, \dots, b_5\}$ with all 25 arcs from A to B , and add a directed five-cycle $b_1 \rightarrow b_2 \rightarrow b_3 \rightarrow b_4 \rightarrow b_5 \rightarrow b_1$ inside B . This digraph has 30 arcs. Every vertex of B has exactly one extra in-arc from inside B , so two arc-disjoint routes from a_i to b_j exist: the direct arc $a_i \rightarrow b_j$, and the detour $a_i \rightarrow b_{j-1} \rightarrow b_j$ using the 5-cycle predecessor of b_j (indices mod 5). For the matching upper bound, the only in-arcs of b_j reachable on a path from a_i are the direct arc $a_i \rightarrow b_j$ and the cycle arc $b_{j-1} \rightarrow b_j$. Removing both disconnects a_i from b_j , while neither alone does, so the minimum cut has size 2. Hence $\lambda(a_i, b_j) = 2$. Routes between two vertices of B never leave B and give $\lambda(b_i, b_j) = 1$. Hence $\lambda^{\max} = 2$, so the graph is feasible at $m = 3$, yet $30 > \max(3 \cdot 9, 25) = 27$.

Figure 1.7 draws this thirty-arc digraph and traces the mechanism. The twenty-five faint arcs are the complete one-directional bipartite layer from A to B , the construction we already understand; on top of them the orange five-cycle gives each vertex of B a single extra in-arc from inside B . That one extra arc is exactly what doubles the connectivity. For the marked pair a_1, b_2 the two heavy routes show how: the blue arc is the direct hop $a_1 \rightarrow b_2$, and the green path $a_1 \rightarrow b_1 \rightarrow b_2$ borrows the cycle arc $b_1 \rightarrow b_2$ for its second step. The two share no arc, so $\lambda(a_1, b_2) = 2$, and the same pair of routes exists for every a_i, b_j . Five extra arcs lift the whole digraph from one independent route per pair to two, and the thirty arcs overshoot the old prediction of twenty-seven.

The counterexample generalises: the five-cycle gave every vertex of B exactly one extra in-arc from inside B , and the budget for such arcs is $m - 2$ per vertex, not one. Figure 1.7 is the instance $m = 3$, where that budget is one and a single cycle suffices; for larger m each b_j collects $m - 2$ cyclic predecessors and the same picture acquires more concentric chords inside B .

Construction 1.12 (Augmented bipartite digraph). For $m \geq 2$ and $n \geq m$, set $|B| = \lceil (n + m - 2)/2 \rceil$ and $|A| = n - |B| = \lfloor (n - m + 2)/2 \rfloor$, include every arc from A to B , and inside B give each vertex b_j the $m - 2$ in-arcs from its cyclic predecessors $b_{j-1}, \dots, b_{j-(m-2)}$ (indices mod $|B|$). The total is $\lfloor (n + m - 2)^2/4 \rfloor$ arcs. For the connectivity, fix $a \in A$ and $b \in B$: the direct arc and the $m - 2$ two-step detours $a \rightarrow b' \rightarrow b$ through the in-neighbours b' of b inside B are $m - 1$ arc-disjoint routes, and they are all there are, because the only in-arcs of b reachable from a are exactly those $m - 1$ arcs, which therefore form a cut. Pairs inside B are limited by the in-degree $m - 2$, pairs inside A and pairs from B to A have no routes at all, so $\lambda^{\max} = m - 1$.




Figure 1.7. The thirty-arc counterexample of [Remark 1.11](#): the augmented bipartite digraph ([Construction 1.12](#)) at $m = 3$, $n = 10$, $|A| = |B| = 5$. The 25 faint arcs are the complete one-directional layer from A to B ; the 5 orange arcs form the directed cycle $b_1 \rightarrow b_2 \rightarrow b_3 \rightarrow b_4 \rightarrow b_5 \rightarrow b_1$ inside B , one extra in-arc per vertex of B . The two heavy arcs and the green path show the pair a_1, b_2 carrying 2 arc-disjoint routes: the direct arc $a_1 \rightarrow b_2$ (blue) and the detour $a_1 \rightarrow b_1 \rightarrow b_2$ (green) through the cycle predecessor b_1 . Hence $\lambda^{\max} = 2$, feasible at $m = 3$, while $30 > \max(3 \cdot 9, 25) = 27$.

At $m = 2$ the inside of B is empty and the construction reduces to [Construction 1.7](#). For $m = 2$ and $m = 3$ the partition $(|A|, |B|)$ equals $(\lfloor n/2 \rfloor, \lceil n/2 \rceil)$, the same balanced split as before, so nothing changes in those cases; the shift to a larger B part is only visible at $m \geq 4$, where the balanced partition is suboptimal. At $m = 3$, $n = 10$ it is the thirty-arc counterexample of [Remark 1.11](#), drawn in [Figure 1.7](#) (using the equivalent balanced partition $|A| = |B| = 5$).

This is the corrected shape, and it leads to the standing conjecture for the directed arc problem.

Conjecture 1.13. *For simple digraphs and $m \geq 2$,*

$$\ell_m^{\text{dir}}(n) = \max \left(m(n-1), \left\lfloor \frac{(n+m-2)^2}{4} \right\rfloor \right).$$

For $m = 2$ this is [Theorem 1.9](#). For $m = 3$ the formula $\lfloor (n+1)^2/4 \rfloor$ equals $\lfloor n^2/4 \rfloor + \lceil n/2 \rceil$, so the conjecture reduces to the previous formula for $m = 2$ and $m = 3$. For $m = 3$, $n = 10$ the second branch gives $\lfloor 121/4 \rfloor = 30$, matching [Remark 1.11](#). For $m \geq 4$ the new formula strictly exceeds $\lfloor n^2/4 \rfloor + (m-2) \lceil n/2 \rceil$ at even n large enough (first at $n = m + 8$ for $m = 4$); the construction of [Construction 1.12](#) with the shifted partition witnesses the difference. The lower bound holds for all m , witnessed by [Construction 1.8](#) on the first branch and [Construction 1.12](#) on the second. The upper bound is open, and [Chapter 4](#) returns to it with what structural progress the machine and the hand can jointly supply.

1.8 The hypergraph variant

The third model deserves its own short treatment, not because its base case is hard but because it is the one variant whose objects are stored and manipulated differently from all the rest, and it is worth meeting that difference now rather than at the moment of computation. A hypergraph keeps no multiplicity matrix, only the list of its hyperedges, each a set of r vertices. We fix the edge size r , following the extremal literature, for a structural reason: under a connectivity ceiling a small edge is always at least as efficient per route as a large one, so a hypergraph free to mix sizes would maximise its edge count by using pairs only, and the question would collapse back to the graph case. Fixing r is what makes the hypergraph question a new question. The route between two vertices is a *Berge path*, which alternates vertices and hyperedges so that each consecutive pair of vertices lies in the hyperedge listed between them, and two such routes are edge-disjoint when they share no hyperedge. Concretely, in a three-uniform hypergraph a route from u to w might be the single step $u, \{u, x, w\}, w$ that crosses one hyperedge, or a longer alternation $u, \{u, x, y\}, y, \{y, z, w\}, w$ that switches hyperedge at the intermediate vertex y . With routes redefined this way the same Gomory–Hu machinery that proved the multigraph value carries over, and the base case has the same linear shape. [Figure 1.8](#) draws such a two-hyperedge route.




Figure 1.8. A Berge path from u to w in a three-uniform hypergraph, crossing two hyperedges. Each hyperedge (shaded in its own colour) is a set of three vertices, and they meet at the shared vertex y where the two colours blend; the route (orange) enters e_1 at u and leaves it at y , then enters e_2 at y and leaves it at w . The two steps use distinct hyperedges, so this counts as one Berge path; a second route edge-disjoint from it would have to avoid both e_1 and e_2 entirely. Unlike an ordinary edge, a single hyperedge touches all $\binom{3}{2} = 3$ pairs of its vertices at once, which is the bookkeeping the checker of [Chapter 2](#) must handle.

Proposition 1.14 (Hypergraph edge bound). *An r -uniform hypergraph on n vertices with Berge edge-connectivity at most $m - 1$ has at most $\left\lfloor \frac{(m-1)(n-1)}{r-1} \right\rfloor$ hyperedges, and the bound is attained whenever $m - 1 \leq \binom{n-2}{r-2}$.*

The proof, given in [Chapter A](#), runs the multigraph argument through the hypergraph Gomory–Hu theorem of Bérczi, Chandrasekaran, Király and Kulkarni [2], charging each hyperedge to the at least $r - 1$ tree cuts it crosses. The star hypertree attains the $m = 2$ case, a hub joined to disjoint blocks of $r - 1$ vertices, with Berge connectivity one everywhere and $\lfloor (n - 1)/(r - 1) \rfloor$ hyperedges, exactly as a tree attained the multigraph bound. One genuine surprise is worth recording here: for $r \geq 3$ the bound is attained by *simple* hypergraphs, with no repeated hyperedge ([Theorem A.32](#)). In the graph case simplicity is exactly what separates Mader’s $\lfloor m(n - 1)/2 \rfloor$ from the multigraph value $(m - 1)(n - 1)$. One edge size higher, it costs nothing.

What the proposition does not say is how to *measure* the Berge connectivity of an arbitrary hypergraph, and that measurement is genuinely different from the graph case. The difference is not the one-route-per-edge rule, which graphs share, but what an edge touches: an ordinary edge joins one pair, so edge-disjointness is policed pair by pair, while a single hyperedge touches all $\binom{r}{2}$ pairs among its r vertices at once, and two routes may legitimately want to enter it between different pairs. The measurement must let any one of those pairs use the hyperedge while barring all the others, simultaneously, for every hyperedge. That bookkeeping is the one new piece of machinery the next chapter has to build, and it is the reason the hypergraph rejoins the family only once the checker is in place.

1.9 Random graphs as intuition

The short proofs above hinted that the edge and vertex problems travel together for a while, agreeing through $m = 4$ before the split at $m = 5$. That separation is a statement about extremal graphs, the densest ones the ceiling allows, and it is worth asking whether the same split can be seen in ordinary graphs drawn at random, with no extremal tuning at all. The random model is the natural place to build intuition, because it shows the qualitative behaviour of the connectivity notions before any construction is pushed to its limit, and we will return to it whenever a picture of the typical graph clarifies a result. It is not a detour from the program to come, either: a sampled $G(n, p)$ is an ordinary graph, measured by the very same connectivity checker the next chapter builds for every other variant, so the random model is simply the one pipeline turned to *observe* the typical case rather than to prove or to discover the extremal one.

Each of the $\binom{n}{2}$ possible edges of $G(n, p)$ is included independently with probability p , and on the resulting graph we read off both binding connectivities, the edge version $\lambda_G^{\max}$ and the vertex version $\kappa_G^{\max}$, from the very same sample. Whitney's inequality $\kappa_G^{\max} \leq \lambda_G^{\max}$ guarantees the vertex value never exceeds the edge one, but it is silent on how often the two actually come apart. [Figure 1.9](#) looks directly at the distributions, fixing $n = 16$ and histogramming both connectivities over many samples at three densities. Two things are visible at once. As the density rises the whole distribution marches rightward, from a binding connectivity around five at $p = \frac{1}{4}$ to around thirteen at $p = \frac{3}{4}$, so by $n = 16$ the typical graph already sits well past the $m = 4$ at which the proofs agreed. And comparing the edge row to the vertex row, the vertex distribution sits at or just to the left of the edge one, never to its right, which is Whitney's inequality seen across a whole distribution rather than a single graph. The two pull apart on a fraction of samples that shrinks as the density grows, from about a fifth of the samples at $p = \frac{1}{4}$ to only a handful by $p = \frac{3}{4}$. The reason is structural. For $\kappa^{\max}$ to fall below $\lambda^{\max}$ some pair must carry more edge-disjoint routes than internally vertex-disjoint ones, which happens only when several of those routes are forced through one shared intermediate vertex. In a sparse graph routes are scarce and a single low-degree vertex can be exactly such a bottleneck, but as p grows every vertex gains degree and the graph offers many alternative intermediate vertices, so independent routes almost never need to reuse one. In the dense limit the graph approaches the complete graph, where $\kappa^{\max} = \lambda^{\max} = n - 1$ for every pair and the gap closes entirely. The divergence the thesis

records at $m = 5$ is therefore not an artefact of the extremal construction. It is already there, in miniature, in the random model.




Figure 1.9. Distributions of the binding connectivity in $G(16, p)$ over 400 samples, at densities $p = \frac{1}{4}, \frac{1}{2}, \frac{3}{4}$. The top row histograms the edge-connectivity $\lambda^{\max}$, the bottom row the vertex-connectivity $\kappa^{\max}$, measured on the same samples, with the mean marked by a dashed line and the fraction of samples on which $\kappa^{\max} < \lambda^{\max}$ printed on each vertex panel. Reading a column top to bottom, the vertex distribution sits at or to the left of the edge one, which is Whitney's inequality, and the gap is largest at the sparsest density (22% of samples) and closes as p grows (6%). All six panels share one axis range.

The random model also supplies a landmark of a different flavour. Below a sharp density the forbidden configuration is almost surely absent, and above it almost surely present, and the location of that threshold is the same for every one of the connectivity notions above.

Theorem 1.15 (Appearance threshold). *Let $m = m(n) \geq 2$ grow with n so that $m/\ln n \rightarrow \infty$ and $m = o(n)$, and let $c > 0$ be a constant. In the binomial random graph $G(n, p)$ with $p = c \cdot m/n$,*

$$\lim_{n \rightarrow \infty} \Pr[\lambda^{\max} \geq m] = \begin{cases} 1 & \text{if } c > 1, \\ 0 & \text{if } c < 1, \end{cases}$$

so the threshold for the appearance of a pair with local edge-connectivity at least m is $p^ = m/n$. The same threshold governs local vertex-connectivity and the directed analogues (Remark A.42).*

The proof is a concentration argument with two halves (Chapter A). Above the threshold the edge count concentrates past Mader's bound, so Theorem 1.2 itself forces a high pair into existence. Below it the maximum degree stays under m , and the degree bound caps every local connectivity. Both halves lean on the same growth condition $m/\ln n \rightarrow \infty$, which is what lets a union bound over the n vertices close. The collapse of complexity is what deserves emphasis: in

the probabilistic limit the distinctions between edges and vertices, directions and models all wash out, and the single number m/n decides the matter, a universality the synthesis of [Chapter 4](#) records again.

One limitation must be stated plainly, lest the threshold be read as evidence about the problem this thesis actually studies. The growth hypothesis $m/\ln n \rightarrow \infty$ forces m to climb with n , so the theorem says nothing whatever about a fixed small m – the regime of every exact value above, where m is 2, 3, or 4 and n runs to infinity around it. There the union bound is vacuous and the appearance of a dense pair is governed by rare low-degree configurations that need finer, Poisson-type tools. The random model therefore enters as contrast and not as evidence: it shows how cleanly the question collapses in one limit, which throws into relief how stubbornly it resists in the fixed- m limit the rest of the thesis must attack by hand and by machine.

1.10 Where this work sits

It is worth pausing, before the machine, to place the question and the method against the two traditions they descend from: the extremal graph theory that posed the problem and the computational mathematics that will help settle it.

The mathematical lineage is the older of the two. The problem belongs to the family of extremal connectivity questions opened by Bollobás and Erdős, who asked how dense a graph can be while every pair of vertices stays below a fixed connectivity [4], logged today as Erdős Problem 915 [5]. Mader’s theorem [12] closed the simple undirected edge case with the clean floor $\lfloor m(n-1)/2 \rfloor$ recalled in [Theorem 1.2](#), and it is the floor every variant in this thesis is measured against. The vertex-disjoint branch has a longer and rougher history: Leonard [11] showed the edge and vertex problems coincide through $m \leq 4$, and Sørensen and Thomassen [18] found the first divergence at $m = 5$ with $k_5(n) = \lfloor 8n/3 \rfloor - 3$, after which the exact values $k_m(n)$ remain unknown. That gap between a settled edge case and a stalled vertex case is exactly the terrain [Chapter 1](#) has been mapping, and the directed and hypergraph axes extend it further. The thesis adds to this lineage not a single new closed form but a uniform way of pushing each axis as far as exact computation allows.

The computational tradition this thesis’s pipeline belongs to is the practice of settling combinatorial questions by machine in a way a human can audit, and it splits into a few schools worth contrasting with the certify-then-prove loop built next. The oldest is *exhaustive generation*: enumerate every object of a given size up to isomorphism and check each one. McKay and Piperno’s *nauty/geng* toolkit [13] is the standard engine for this, and its flagship results include exact small Ramsey numbers such as $R(4, 5) = 25$ [14]; the enumeration of [Chapter 2](#) is the same idea, run on the twelve variants of Problem 915. The second school is *SAT-assisted proof*, in which a question is encoded in propositional logic and a solver emits a machine-checkable certificate: Heule, Kullmann and Marek’s resolution of the Boolean Pythagorean triples problem [9], whose verified proof runs to some two hundred terabytes, and Heule’s determination of Schur number five [8] are the canonical examples of certified solver output at scale. The third is *flag*

algebras, Razborov's framework [17] that converts asymptotic extremal bounds into semidefinite-programming certificates, the dominant tool for density problems in the large- n limit. Closest of all to the method here is the fourth, *LP/ILP-based certification*, where a feasibility or optimality argument over the rationals or integers is exhibited as a linear program whose optimal dual is itself the proof; this is precisely the form the cut-counting certifier of Chapter 2 takes.

Against that backdrop, what is distinctive here is consolidation rather than a new solver. A single checker measures connectivity exactly across all twelve problem variants on one data model, so search and certification share one codebase rather than living in separate tools. The certificates it emits are exact rational or integral objects, not floating-point estimates, so an upper bound it reports is a proof and not evidence. And because the same program both hunts for dense graphs and certifies the bounds they suggest, the discover step and the prove step are two modes of one loop, which is the arrangement the next chapter builds. What makes the consolidation honest is that it never blurs three kinds of knowledge: a value can be *proved* (by hand, or by exhausting every graph of a size), *certified* (by the same optimisation, reaching a vertex or two further), or only *conjectured* (witnessed by a dense graph the search exhibits, with no matching upper bound). This trichotomy, named in Section 3.6 and obeyed by the summary of Chapter 4, is the through-line of everything that follows.

Computer note

We are now stuck in three different ways: a fifty-year-old open problem at $m \geq 6$ for vertices, a long proof we would rather not check by hand at $m = 2$ for arcs, and a conjecture with no proof at all at $m \geq 3$. The base cases of the long induction were verified by exhaustive search, and the thirty-arc counterexample was found the same way. The next chapter builds a single program that models all of these cases at once, measures their connectivity exactly, and proves the small-case upper bounds outright.

Chapter 2

Certifying Bounds by Machine

The last chapter ended in three different kinds of stuckness, and all three share a shape. The objects are graphs of one flavour or another, the constraint is always a ceiling on connectivity, and the work is always either measuring a given graph or checking many small ones. That shared shape is what a single program can exploit. This chapter builds it, and then uses it for one purpose only: to *prove*. It gives every variant one representation, measures connectivity exactly through a max-flow checker, checks small cases by honest enumeration, and turns the finite checks the proofs of [Chapter 1](#) relied on into a single auditable certifier. The hunt for dense examples, which proves nothing on its own, waits until [Chapter 3](#).

A word on what the program is for. It is a microscope, not a source of final answers. It measures connectivity exactly, so what it reports about a given graph is a fact. It enumerates graphs of a fixed size, so for that size it can prove a theorem. It will later discover dense graphs, which are only lower bounds, but not here. Each of these has a different epistemic weight, and the program is built so that the three never get confused. [Figure 2.1](#) draws this spine: a single *model* holds every variant, one exact *measure* reads connectivity off it, and from that measurement a *prove* step bounds a fixed size from above, with the *discover* step of the next chapter and a final *observe* step for the random model added later. Every routine introduced from here on is shown as a small card carrying its name, its inputs and output, a one-line account of how it works, and a coloured badge naming its place in the spine. The function bodies are not reproduced, because the point is the interface, not the code.




Figure 2.1. The architecture as one pipeline. A single data model, the multiplicity matrix μ , feeds an exact measurement, the max-flow checker. That measurement feeds the two bound-producing roles: a prover for upper bounds (the cut-counting optimisation and the exhaustive search), built in this chapter, and a discoverer for lower bounds (simulated annealing), built in [Chapter 3](#). A final sampling role studies the random model. The badge colours on the code cards below match the boxes here.

2.1 The solver

Everything in this chapter is a single function, `solve`, and it helps to hold its most basic version in mind from the start, because every later idea is a refinement of it and nothing is ever a separate program for a separate case. Given n and m , the basic strategy is as plain as it sounds. Walk through every graph on n vertices, measure the largest local connectivity $\lambda^{\max}$ of each with a max-flow computation, and keep the densest graph that stays at or below the ceiling $m - 1$. Because it has looked at every graph, the count it returns is the exact answer, not an estimate.

That one loop already settles the smallest cases, and the rest of the chapter is built on top of it without ever leaving `solve`. We first make the measurement step trustworthy and quick, via the connectivity checker and a Gomory–Hu shortcut for undirected graphs. We then make the enumeration step skip whole families of hopeless graphs, via a pruned search. Finally, for the directed multigraph case, we replace enumeration altogether with one small optimisation that proves every m at once, which is the certifier. Each of these is the same `solve` doing less work for the same guarantee, and the choice among them is made automatically from the booleans it receives.

```
solve(n, m, directed, simple, separation, exhaustive, maxSeconds)
```

DRIVER

in the variant booleans `directed` and `simple`, the separation (edge or vertex), n , m , a mode, and a time budget

out a value with an honest label: exact, lower bound, or upper bound

how picks the method the case allows: brute-force enumeration, the pruned search, or the cut-counting certifier when proving, or the temperature search when discovering ([Chapter 3](#)). It always respects the budget, and it is the one function the rest of the thesis ever calls.

2.2 A single representation for all variants

Two of the three axes from [Chapter 1](#), the model and the direction, are properties of the object, and a single data structure holds them all. Of the three structural models, simple graphs and multigraphs fit the matrix directly, while the hypergraph keeps the separate storage promised in [Chapter 1](#) and rejoins at the end of this chapter. The hypergraph could in principle be pressed into the same shape as a higher-order tensor, an r -dimensional array marking which r -sets are present, but that array is almost entirely zero and costs n^r entries, so the program keeps the compact list of hyperedges instead and pays nothing for the empty cells. Sampled $G(n, p)$ graphs are ordinary simple graphs and therefore use this same representation without any change. A graph or digraph on n vertices is stored as an integer matrix μ , where $\mu(u, v)$ is the multiplicity of the edge or arc from u to v . A simple graph is the case $\mu \in \{0, 1\}$, an undirected graph satisfies $\mu = \mu^\top$, and a multigraph allows larger entries. The third axis, edge against vertex separation, is a property of the question rather than the object, so it lives in how we measure μ , not in μ itself.

Figure 2.2 illustrates the encoding.




Figure 2.2. A multigraph on three vertices and its multiplicity matrix μ . Entry $\mu(u, v)$ records the number of parallel edges between u and v . The double edge between vertices 1 and 2 gives $\mu(1, 2) = 2$. A simple graph is the case $\mu \in \{0, 1\}$, an undirected graph satisfies $\mu = \mu^\top$, and a directed graph allows $\mu(u, v) \neq \mu(v, u)$.

Graph(n , directed, simple)		MODEL
in	the vertex count n and the two booleans	
out	a graph whose whole state is the matrix μ	
how	one $n \times n$ integer matrix is the object, an undirected graph keeps $\mu = \mu^\top$, and a simple graph caps every entry at one.	

Keeping a single representation is what lets one measurement routine, one certifier, and one search serve every case. Nothing downstream asks which of the twelve problems it is solving. It asks only the two booleans (directed or undirected, simple or multi) and the matrix. Everything that builds or edits a graph is a thin wrapper around μ that respects those booleans, and these operations are so routine that they need no individual exposition, and Table 2.1 lists them together.

Operation	Meaning
addEdge(G , u , v , k)	add k parallel edges or arcs $u \rightarrow v$ (a simple graph saturates at one)
removeEdge(G , u , v , k)	remove k copies, never below zero
setMultiplicity(G , u , v , c)	set $\mu(u, v) = c$ directly
multiplicity(G , u , v)	read $\mu(u, v)$
hasEdge(G , u , v)	whether $\mu(u, v) > 0$
degree(G , v)	total degree of v (in- plus out-degree if directed)
edgeCount(G)	number of edges or arcs, counted with multiplicity
edges(G), vertices(G)	iterate the present edges, and the vertex set
copy(G)	an independent duplicate

Table 2.1. The routine operations on the graph model. Each is a one-line wrapper that reads or writes the matrix μ while keeping an undirected graph symmetric and a simple graph binary. They are bookkeeping, and the chapter’s attention belongs to the routines that follow.

2.3 The connectivity checker built on maximum flow

Menger’s theorem turns the connectivity question into a flow computation. A maximum flow is the standard way to ask how much can be pushed from one point to another through a network

in which every link has a limited capacity. Menger's theorem says the maximum number of edge-disjoint $u-v$ paths equals the minimum $u-v$ edge cut, and that number is exactly the value of such a maximum flow once each multiplicity $\mu(u, v)$ is read as the capacity of that link. The routine that runs this flow and reports the route count is the *connectivity checker*. It is the single most important routine in the program: it is the only place graph theory meets computation, and every later claim about connectivity passes through it.

`localEdgeConnectivity(G, s, t)  $\rightarrow \mathbb{N}$`

MEASURE

in a graph G and an ordered pair (s, t)

out $\lambda_G(s, t)$, the maximum number of edge-disjoint $s-t$ paths

how a single maximum flow on the network whose arc capacities are the multiplicities μ , so by Menger the flow value is the path count.

`maxEdgeConnectivity(G)  $\rightarrow \mathbb{N}$`

MEASURE

in a graph G

out $\lambda^{\max}(G)$, the largest local edge-connectivity over all pairs

how one max-flow per pair (ordered if directed), take the largest. This is the quantity the edge problem caps at $m - 1$.

The vertex-disjoint version follows the same idea after one construction: split each vertex v into an in-copy v^{in} and an out-copy v^{out} joined by a single unit-capacity arc (Figure 2.3). Every arc $u \rightarrow v$ of the original graph becomes $u^{\text{out}} \rightarrow v^{\text{in}}$ in the split network. Routing through v now consumes the one unit of capacity on its internal arc, so two paths sharing v as an intermediate vertex compete for that unit and cannot coexist in any maximum flow. A max-flow from u^{out} to v^{in} in the split network therefore equals the number of internally vertex-disjoint $u-v$ paths in the original graph.




Figure 2.3. Vertex splitting for the vertex-disjoint checker. Incoming arcs attach to v^{in} and outgoing arcs leave from v^{out} , connected by a single unit-capacity arc. Any directed path that uses v as an intermediate vertex passes through this bottleneck, and two such paths would need two units of capacity and are therefore separated.

`maxVertexConnectivity(G) → ℕ`

MEASURE

in a graph G

out $\kappa^{\max}(G)$, the largest local vertex-connectivity over all pairs

how the same sweep as the edge checker, but each flow runs on the split network of [Figure 2.3](#), so the unit internal arcs count internally vertex-disjoint paths. Parallel edges are given capacity one here, so they never raise κ .

The two checkers are the entire interface between graph theory and computation in this thesis. As a sanity check, applied to the directed double star on n vertices, the edge checker reports $\lambda^{\max} = 1$, and the arc count $2(n - 1)$ matches the proved values $\ell_2^{\text{dir}}(n) = 2, 4, 6, 8$ for $n = 2, 3, 4, 5$.

The same checker answers two different questions. Throughout this chapter it runs in its exact-value mode, returning $\lambda^{\max}$ or $\kappa^{\max}$ on the nose, and every proved or reported number rests on that exact reading. The search of [Chapter 3](#) needs only the cheaper yes-or-no question “is the binding connectivity above the bound?”, so there the same flow runs in a capped mode that stops as soon as it has seen one route too many. The capped mode never changes a reported value: it is a faster way to ask for a single bit, and [Chapter 3](#) shows that the search built on it reproduces the exact search step for step.

The Gomory–Hu tree from [Chapter 1](#) reappears here as an efficiency, and it is the honest kind. For undirected graphs all $\binom{n}{2}$ edge-connectivities are read off a single weighted tree, so $\lambda^{\max}$ is the heaviest tree edge rather than the maximum over $\binom{n}{2}$ separate flows. The speed-up is real, but it is bought with a theorem, not a heuristic.

`maxEdgeConnectivityViaTree(G) → ℕ`

MEASURE

in an undirected graph G

out $\lambda^{\max}(G)$, identical to the pairwise sweep

how build one Gomory–Hu tree, then return its heaviest edge, which costs $n - 1$ flows in place of $\binom{n}{2}$ and cross-checks the checker.

The hypergraph is the one model that never used the multiplicity matrix, since it is stored as a plain list of hyperedges, each one a set of r vertices. The checker still has to read it, and the trick is to do the same flow computation as before on a slightly larger network that we build on the spot. The difficulty is that one hyperedge ties several vertices together at once and, by the definition of a Berge route in [Chapter 1](#), it may be used by only one route at a time, so we cannot simply treat it as a bundle of ordinary links. Instead, for every hyperedge we add one extra helper point and connect each of that hyperedge’s member vertices to it ([Figure 2.4](#)). A route that wants to use the hyperedge now enters through one member vertex, passes through the helper point, and leaves through another. By letting only a single route pass through each helper point, we capture exactly the rule that one hyperedge serves one route. Counting routes between two

vertices is then the very same maximum flow as before, run on this enlarged network, and when a hyperedge happens to hold just two vertices the helper point collapses back to an ordinary link, so the answer agrees with the graph case. That the flow on the enlarged network counts exactly the hyperedge-disjoint Berge routes is the hypergraph version of Menger's theorem, proved in [Chapter A \(Theorem A.30\)](#), so the checker rests on a theorem rather than an analogy.




Figure 2.4. Measuring Berge connectivity by maximum flow. A hyperedge e joining several vertices (left) is replaced by a helper point h_e joined to each of its members (right). Because at most one route may pass through h_e , the helper point enforces that the hyperedge serves only one route at a time. Counting independent routes between two vertices is then an ordinary maximum flow on this enlarged network, and a two-vertex hyperedge collapses to a single ordinary edge.

One way to picture the helper point: think of each hyperedge as a room with several doors. Entering through any door counts as using that room. Because only one path may pass through the helper node, at most one traveller can occupy the room at once, so the second traveller must find a route that never enters it. The same flow computation that counts edge-disjoint paths through ordinary links now counts routes that avoid sharing any room, which is exactly what Berge edge-disjointness means.

maxHyperedgeConnectivity(H) $\rightarrow \mathbb{N}$		MEASURE
in	an r -uniform hypergraph H	
out	$\lambda^{\max}(H)$, the largest Berge edge-connectivity over all pairs	
how	build the enlarged network in which every hyperedge becomes a helper point that only one route may pass through, then take the same maximum flow as the graph checker. On two-vertex hyperedges it matches ordinary edge-connectivity.	

On the complete graph K_n the hypergraph checker returns $n - 1$, and on the complete three-uniform hypergraph $K_4^{(3)}$ it returns 3, larger than the two hyperedges through each pair because longer Berge routes also carry flow. The self-test confirms both of these values, so the construction is checked rather than merely asserted. The same helper-point network answers the other three hypergraph questions without alteration: splitting each ordinary vertex as in [Figure 2.3](#) counts internally vertex-disjoint Berge routes, and reading a hyperedge as a single tail feeding its remaining heads gives a directed version, so `solve` measures all four hypergraph variants through this one construction. With the checker in place the proposed value of [Proposition 1.14](#) can be tested on any small hypergraph the way every other variant is, which is the subject of the next section.

2.4 Checking every graph of a fixed size

The checker measures one graph. To bound every graph of a given size, the most direct method is the one the base cases of [Chapter 1](#) already needed: enumerate them all. A simple graph on n vertices is a choice of $\mu(u, v) \in \{0, 1\}$ on each off-diagonal cell, a multigraph a choice in $\{0, \dots, m-1\}$, and a digraph varies every ordered pair rather than only the upper triangle. Walking that product, measuring each candidate with the checker, and keeping the densest feasible one settles the value exactly, and because it has looked at every graph it is a genuine upper bound, not evidence.

This is the first thing `solve` does when it is asked to prove a small case. With its exhaustive flag set it walks that product, measures each candidate with the checker, and keeps the densest feasible one. The value it returns is exact and exhausted, a true upper bound for that n , but only while the count of graphs stays small. No new routine is involved, only `solve` choosing to enumerate.

Run on the small cases this confirms the proved values directly, with no induction and no cleverness. [Table 2.2](#) collects a spread of variants where the enumeration finishes: in every row the machine returns exactly the value [Chapter 1](#) proved by hand. This is the first payoff of the uniform representation. One driver, `solve`, pointed at six different problems, settles all six.

Variant	m	n	<code>solve</code>	Theory
simple undirected, edge	2	5	4	$\lfloor m(n-1)/2 \rfloor = 4$
simple undirected, edge	3	5	6	$\lfloor m(n-1)/2 \rfloor = 6$
multigraph undirected, edge	3	5	8	$(m-1)(n-1) = 8$
simple undirected, vertex	2	6	5	$k_2(n) = n-1 = 5$
simple directed, arc	2	5	8	$\max(2(n-1), \lfloor n^2/4 \rfloor) = 8$
simple directed, arc	2	6	10	$\max(2(n-1), \lfloor n^2/4 \rfloor) = 10$

Table 2.2. Variants that converge under plain enumeration. For each small case `solve`, in its exhaustive mode, walks every graph of the variant, measures it with the checker, and returns the densest feasible one. Every value matches the hand proof of [Chapter 1](#) exactly.

The honesty of the method is also its limit. The number of graphs it must visit grows as $2^{\binom{n}{2}}$ for simple graphs and $2^{n(n-1)}$ for digraphs, and a cap of m on multiplicities raises the base to m . The enumeration is exact wherever it finishes, and it finishes only for the smallest sizes. [Chapter 3](#) opens with exactly how fast this wall arrives. For now the point is that the wall arrives precisely where [Chapter 1](#) needed help, at the directed base cases $n = 6, 7$, so the rest of this chapter is about reaching a little further than blind enumeration can.

2.5 Reaching further: a pruned search for the directed base cases

The base cases $n \leq 7$ of the directed $m = 2$ theorem ([Theorem 1.9](#)) are the finite checks the induction of [Chapter A](#) depends on, and they sit just past where blind enumeration of digraphs is tractable. They can be reached by enumerating more cleverly. Feasibility is monotone: adding an arc can never lower $\lambda^{\max}$, so once a partial digraph is infeasible every completion of it is

infeasible too and the whole branch is abandoned at once. A counting bound prunes further, dropping any branch that cannot beat the best feasible digraph found so far. What remains is a branch and bound over simple digraphs with the connectivity checker as the feasibility test, exact and complete at these sizes where the blind walk is not.

Here `solve` changes its implementation without changing its interface. For a simple digraph it runs this branch and bound in place of the blind walk, still with the connectivity checker as the feasibility test, and so reaches the base cases $n \leq 7$ that blind enumeration cannot. The booleans alone decide which of the two it uses, and the caller never asks.

Run on the directed $m = 2$ problem `solve` returns the complete sequence 2, 4, 6, 8, 10, 12 for $n = 2, \dots, 7$, each proved complete at its size. These are exactly the base cases the induction requires, and with them the long proof of [Theorem 1.9](#) rests on a single auditable search rather than on anything done by hand. The exhaustive-search transcript is in [Chapter A](#).

2.6 Proving every m at once

The pruned search proves one (n, m) at a time. For the directed multigraph arc problem `solve` has a sharper implementation, one that proves every m at a fixed n in a single pass, and it is worth the extra machinery because it closes the directed multigraph case outright for the sizes it reaches.

The key is that m can be scaled out of the problem completely, so that it never appears in the computation at all. Take any directed multigraph that is feasible at level m , meaning $\lambda^{\max} \leq m-1$, and divide every multiplicity by $m-1$. The result is a matrix of weights $w(u, v) \in [0, 1]$, and since dividing all capacities by $m-1$ divides every flow value by the same factor, the maximum flow between each pair is now at most one. Its total weight is $\sum_{u \neq v} w(u, v) = |A|/(m-1)$, so a feasible multigraph with many arcs becomes a feasible weight matrix with large total weight, and vice versa. Writing $M^*(n)$ for the largest total weight any such matrix can carry,

$$L_m^{\text{dir}}(n) = (m-1) M^*(n), \quad M^*(n) = \max \left\{ \sum_{u \neq v} w(u, v) : \text{maxflow}(s, t; w) \leq 1 \text{ for all } s, t \right\}.$$

There is no m on the right-hand side. One solve for $M^*(n)$ therefore fixes $L_m^{\text{dir}}(n)$ for every m at once. The scaling down always gives the upper bound $L_m^{\text{dir}}(n) \leq (m-1) M^*(n)$, and the matching lower bound holds because the heaviest matrix the solve finds is the *double star*: the bidirected star, with weight one on both arcs $h \rightarrow v$ and $v \rightarrow h$ between the hub and every other vertex. Its weights lie in $\{0, 1\}$, so multiplied back by $m-1$ it is an honest integer multigraph for each m , not a fractional one — the directed multigraph carrying every arc of the bidirected star at full multiplicity $m-1$, with $2(n-1)(m-1)$ arcs and $\lambda^{\max} = m-1$. So the two bounds meet for every m together.

What remains is to compute $M^*(n)$, and the difficulty is that the constraint “maximum flow at most one” is itself the answer to a flow problem rather than a plain inequality on w . The max-flow / min-cut theorem removes it. The maximum s - t flow equals the smallest capacity of any *cut*




Figure 2.5. The scaling step for $m = 3$. Dividing each multiplicity by $m - 1 = 2$ maps the integer multigraph (left) to a weight matrix with all values in $[0, 1]$ and max-flow at most one between every pair (right). The double arc $A \rightarrow B$ (multiplicity 2) becomes weight 1; the single arc $A \rightarrow C$ (multiplicity 1) becomes weight $\frac{1}{2}$; the absent arc $B \rightarrow C$ (multiplicity 0, shown dashed) stays absent. Crucially, m drops out on the right, so a single optimisation over all weight matrices pins $L_m^{\text{dir}}(n) = (m - 1) M^*(n)$ for every m simultaneously.

separating s from t , where a cut splits the vertices into a source side holding s and a sink side holding t , and its capacity is the total weight of the arcs crossing from the source side to the sink side. So $\text{maxflow}(s, t) \leq 1$ holds exactly when there is at least one such cut of capacity at most one. We can drop the flow entirely and instead let the optimisation choose, for every ordered pair (s, t) , one cut and require that its capacity stay at or below one.

A cut is described by a yes/no label x_u^{st} on each vertex, equal to one when u is on the source side, with s fixed on the source side and t on the sink side. An arc $u \rightarrow v$ crosses the cut exactly when u is on the source side and v is not, and a helper variable p_{uv}^{st} then picks up its weight $w(u, v)$. Forcing the crossing weights to sum to at most one for every pair forces every pair to admit a cut of capacity one, hence maximum flow at most one. This is the exact feasible region, not a relaxation of it, and that is what turns the result into a proof. The optimisation reports both the best total weight it has found and a value it has shown no matrix can exceed, and when those two meet the gap is zero and $M^*(n)$ is certified.

For the directed multigraph arc problem, then, `solve` runs this cut-counting optimisation in place of any enumeration: it chooses the minimum cut for every ordered pair, and a zero gap makes the reported $M^*(n)$ a proved upper bound, from which $L_m^{\text{dir}}(n) = (m - 1)M^*(n)$ delivers every m at once.

`solve` assembles this optimisation from n alone, and a reader content to take it on the strength of the cut argument just given can skip to [Theorem 2.1](#) without loss. Written out it reads more heavily than it behaves, so we first name every symbol in plain words, then show the one inequality that looks cryptic is a simple yes-or-no switch.

Three quantities appear, and only three. The first is already familiar.

★ $w(u, v) \in [0, 1]$ is the *scaled multiplicity* of the arc $u \rightarrow v$, the amount of arc weight the matrix places on $u \rightarrow v$ after the division by $m - 1$ of the previous section. It is what we are maximising the total of.

★ $x_u^{st} \in \{0, 1\}$ is a *side label* that exists once for every ordered pair (s, t) . For that pair the

optimisation is choosing one cut, and x_u^{st} records the answer to “is vertex u on the source side of that cut?”, with 1 for yes and 0 for no. The two endpoints are pinned, $x_s^{st} = 1$ and $x_t^{st} = 0$, because s defines the source side and t the sink side.

- ★ $p_{uv}^{st} \geq 0$ is a *crossing weight*, the part of $w(u, v)$ that this pair’s chosen cut is forced to count. It equals $w(u, v)$ when the arc $u \rightarrow v$ runs from the source side to the sink side, and is otherwise left at zero.

With those three named, the program is short:

$$\begin{aligned}
& \text{maximise} && \sum_{u \neq v} w(u, v) \\
& \text{subject to} && p_{uv}^{st} \geq w(u, v) + x_u^{st} - x_v^{st} - 1 && \text{for all } (s, t), (u, v), \\
& && \sum_{u \neq v} p_{uv}^{st} \leq 1 && \text{for all } (s, t), \\
& && w(s, t) + \sum_x \min(w(s, x), w(x, t)) \leq 1 && \text{for all } (s, t), \\
& && d(v_0) \leq d(v_1) \leq \dots \leq d(v_{n-1}), && \text{with } d(v) = \sum_{u \neq v} (w(u, v) + w(v, u)), \\
& && w \in [0, 1], \quad x \in \{0, 1\}.
\end{aligned}$$

The first line is the only one that looks cryptic, and it is just an indicator written as an inequality. Its right-hand side $w(u, v) + x_u^{st} - x_v^{st} - 1$ takes one of four shapes according to which side the cut puts the two endpoints u and v on, and [Table 2.3](#) works all four out. Because $w(u, v) \leq 1$, the right-hand side is positive in one case only, the crossing case, where it equals $w(u, v)$ and so forces $p_{uv}^{st} \geq w(u, v)$. In the other three cases it is zero or negative, so the constraint reads $p_{uv}^{st} \geq (\text{something} \leq 0)$ and leaves p_{uv}^{st} free to stay at zero, which the maximisation is happy to let it do.

x_u^{st}	x_v^{st}	meaning	$w + x_u^{st} - x_v^{st} - 1$	forces
1	0	u source side, v sink side: arc <i>crosses</i>	$w(u, v)$	$p_{uv}^{st} \geq w(u, v)$
1	1	both on the source side	$w(u, v) - 1 \leq 0$	$p_{uv}^{st} \geq 0$
0	0	both on the sink side	$w(u, v) - 1 \leq 0$	$p_{uv}^{st} \geq 0$
0	1	u sink side, v source side	$w(u, v) - 2 \leq 0$	$p_{uv}^{st} \geq 0$

Table 2.3. The first constraint, case by case. This is where the bound $w \leq 1$ earns its keep: it makes the three non-crossing rows land at or below zero, so the helper p_{uv}^{st} is pushed up to the arc weight in exactly one case, the crossing one. The constraint is therefore an exact indicator, “count $w(u, v)$ when the arc crosses this cut, and nothing otherwise”, with no slack and no fudge.

[Figure 2.6](#) shows one chosen cut and the weights it counts. The second line of the program now reads plainly. It says the total crossing weight of this pair’s cut, $\sum_{u \neq v} p_{uv}^{st}$, is at most one. By max-flow / min-cut a cut of capacity at most one exists for (s, t) exactly when $\text{maxflow}(s, t) \leq 1$, so the two lines together say “every ordered pair admits a cut of capacity at most one”, which is precisely the feasibility we want. This is the true feasible region, not a relaxation of it, so an

optimal solution reported with zero gap is a proof and not merely evidence.




Figure 2.6. One ordered pair’s chosen cut. The dashed line splits the vertices into the source side, where $x_u^{st} = 1$, and the sink side, where $x_u^{st} = 0$. Only arcs running from source side to sink side cross the cut, and the first constraint forces their helper p_{uv}^{st} up to the arc weight w (the two thick arcs). Arcs that stay on one side, and arcs that run the other way across the line, contribute nothing. The second constraint then caps the total counted weight, the thick arcs, at one.

The last two lines change no answer. They only help the solver finish in time. The third is a redundant shortcut: the direct arc $s \rightarrow t$ and one two-step detour $s \rightarrow x \rightarrow t$ through each intermediate vertex x already form that many routes between s and t , so $w(s, t) + \sum_x \min(w(s, x), w(x, t)) \leq 1$ must hold of any feasible matrix anyway, and stating it outright tightens the continuous picture the solver reasons from (each $\min$ carried by a small yes-or-no selector). The fourth orders the vertices by weighted degree, which is allowed because renaming vertices cannot change any count, and it spares the solver the many relabelled copies of each graph that differ only in vertex names.

Theorem 2.1 (Directed multigraph value, small n). *For $n \leq 6$ and all $m \geq 2$, $L_m^{\text{dir}}(n) = 2(n - 1)(m - 1)$, attained by the double star. Equivalently $M^*(n) = 2(n - 1)$.*

It is worth being clear about what “for all m ” rests on, because the theorem proves infinitely many values from finitely many solves. It is not that each m was tested. In this mode `solve` never sees m at all. It proves the single m -free number $M^*(n)$, and the identity $L_m^{\text{dir}}(n) = (m - 1)M^*(n)$ then carries that one fact to every m at once. The only quantity that is checked case by case, and the only genuine limit on how far the result reaches, is n .

`solve` proves [Theorem 2.1](#) by returning $M^*(3) = 4$, $M^*(4) = 6$, $M^*(5) = 8$, and $M^*(6) = 10$, each optimal with zero gap, so that $M^*(n) = 2(n - 1)$ over this range. The first three finish quickly, $M^*(3)$ and $M^*(4)$ in well under a second and $M^*(5)$ in a few seconds, while $n = 6$ takes about twenty minutes. At $n = 7$ the same encoding does not close, and we do not pretend otherwise: it is a finite obstacle of scale, not a new idea, and it marks the current edge of certified knowledge for this variant. The solver transcript is in [Chapter A](#).

2.7 What the solver may claim

Now that the pieces inside `solve` are built, it is worth stating plainly, because the rest of the thesis depends on it, what each is allowed to claim. The checker *measures*: its connectivity values are exact. Its three proving implementations, the exhaustive enumeration, the pruned search, and the cut-counting, *prove*: an exhausted enumeration or an optimal, zero-gap solve is an upper bound for that fixed n . The search of [Chapter 3](#) will *discover*: it produces concrete graphs, hence lower bounds, and never an upper bound. These three claims are exactly what `solve` labels its answer with, so that an exact value, a lower bound, and an upper bound can never be quietly confused with one another.

With the microscope built and aimed at proof, the variants with small answers are settled and the directed base cases the hand proof needed are in hand. What proof by enumeration cannot do is reach the sizes where the answers are only conjectured, because the number of graphs explodes long before the interesting cases arrive. The next chapter measures that explosion exactly, and then turns the same program loose to *discover* the dense graphs that blind enumeration can never reach.

2.8 Reproducibility

Every claim in this thesis that rests on computation can be re-run from a single file. All of the program's logic lives in `program/erdos915_unified.py`, a self-contained Python program whose only dependencies are `numpy`, `networkx`, `scipy`, and `matplotlib`, with no build step. Running `python erdos915_unified.py` executes the built-in self-test, the suite of invariant checks that confirms the connectivity values quoted throughout this chapter, including the sanity checks on the directed double star and on K_n and $K_4^{(3)}$. Every figure in the thesis is regenerated by its companion driver `program/make_figures.py`, which imports that same file and fixes every random seed, so the pictures are reproduced exactly rather than merely resembled; the table in `program/README.md` records which routine produces which figure. The solver transcripts behind the certified values of this chapter, the pruned-search log for [Theorem 1.9](#) and the cut-counting certificate for [Theorem 2.1](#), are reproduced verbatim in [Chapter A](#), so a reader may check the proofs without running anything at all.

Chapter 3

Discovering Bounds by Search

The previous chapter proved what could be proved by looking at every graph of a fixed size. That method is honest and exact, and it stops almost at once. This chapter begins by measuring exactly how fast it stops, because the speed of that collapse is the whole reason a second kind of tool is needed. Then it builds that tool. Where the certifier walks all graphs, the search walks a few of them well, cooling like a physical system toward the densest feasible shape. What it returns is never a proof. It is a concrete graph, hence a lower bound and a conjecture, a stepping stone toward an upper bound that a later proof or certificate must supply. The chapter ends by asking the honest question about any such tool: how good is it, and how far from the truth might its answers be.

3.1 The wall: how enumeration explodes

The brute force of [Chapter 2](#) visits every graph of a given size. The trouble is how many that is. A simple undirected graph on n vertices is a yes-or-no choice on each of the $\binom{n}{2}$ possible edges, so there are $2^{\binom{n}{2}}$ of them. A digraph chooses on each of the $n(n-1)$ ordered pairs, giving $2^{n(n-1)}$, and allowing multiplicities up to a cap c raises the base from two to $c+1$. These are not numbers that merely grow. They detonate. [Figure 3.1](#) draws the growth on a logarithmic scale, and the message is the same in every panel: the count passes any reachable budget while n is still in the single digits.

Three readings of the figure matter for what follows. Adding vertices is the sharpest cost: each new vertex multiplies the count by an exponential in n , so a single extra vertex can move a problem from seconds to centuries. Raising the connectivity budget is the next cost, lifting the base of the exponent rather than its height, which is why the multigraph curves sit above the simple one in every panel. Direction is the third: a digraph has twice as many cells to fill as the undirected graph on the same vertices, so the directed panel climbs fastest of all, and those are precisely the variants whose answers [Chapter 1](#) could not prove. The separation, by contrast, costs nothing here, because edge and vertex disjointness search the same graphs and differ only in what the checker measures, which is why a single pair of panels suffices for all of them. The certifier of [Chapter 2](#) reaches a few vertices further than blind enumeration, because its cut-counting optimisation is polynomial in the size of the encoding rather than in the number of graphs, but solving that encoding is itself hard and the gain is measured in single vertices, not orders of magnitude. Past that, no exhaustive method reaches the interesting sizes. Something that does not look at every graph is required.




Figure 3.1. The number of graphs blind enumeration must visit, on a logarithmic vertical axis, for directed graphs (left) and undirected graphs (right). Each panel shows the simple model, two multigraph connectivity caps, and the three-uniform hypergraph (dotted), with a dashed line at a generous budget of 10^9 candidates, and every curve crosses that budget while n is still in the single digits. Writing E for the number of off-diagonal cells a model may fill, a simple graph offers 2^E candidates, a multigraph with multiplicity cap $m - 1$ (the connectivity ceiling) offers m^E (one choice per cell from $\{0, \dots, m - 1\}$), and the 3-uniform hypergraph offers $2^{\binom{n}{3}}$. The count depends only on the model and the direction, never on whether one later asks for edge- or vertex-disjoint routes, so a single pair of panels covers all of those variants at once. Direction doubles the number of cells to fill, so the directed panel sits far above the undirected one, and the directed cases, exactly the ones [Chapter 1](#) left open, are the first to go.

3.2 The temperature search

We want, for fixed n and m , a graph with as many edges as possible subject to $\lambda^{\max} \leq m - 1$ (or $\kappa^{\max} \leq m - 1$ for the vertex variants). The space of graphs on n vertices is enormous and its feasible region is not convex: adding an edge can be harmless until, together with edges added long before, it suddenly creates a forbidden pair. A greedy walk that only ever accepts an improving move stalls at the first such collision, in a graph that is locally maximal but globally far from extremal.

Annealing escapes these traps by sometimes accepting a worsening move, and by accepting fewer of them as it proceeds. Each graph G is a state with energy

$$E(G) = -|E(G)| + \Lambda \cdot \max(0, \lambda^{\max}(G) - (m - 1)),$$

which rewards edges and penalises any excess connectivity. The penalty is a soft constraint rather than a hard one, so the walk may pass briefly through infeasible graphs rather than being walled inside the feasible region, and we return below to why that freedom is worth having. A move toggles one adjacency. The Metropolis rule accepts an improving move always and a worsening move with probability $e^{-\Delta E/T}$, and the temperature T falls geometrically. [Algorithm 1](#) is the whole loop, the one place in this chapter where a body is worth showing, because the loop *is* the method.

Algorithm 1: Temperature search for a dense feasible graph

Input: vertices n , value m , start T_0 , cooling α , penalty Δ , steps S
Output: the densest feasible graph encountered

```
1  $G \leftarrow$  empty graph on  $n$  vertices;  
2  $T \leftarrow T_0$ ;  
3 for  $S$  steps do  
4   propose  $G'$  by toggling one adjacency, biasing removals toward low-sensitivity edges;  
5    $\Delta \leftarrow E(G') - E(G)$ ;  
6   if  $\Delta \leq 0$  or  $\text{rand}() < e^{-\Delta/T}$  then  
7      $G \leftarrow G'$ ;  
8   if  $G$  is feasible and denser than the best so far then  
9     record  $G$  as the best;  
10   $T \leftarrow \alpha T$ ;  
11 return the best recorded graph;
```

This loop is what `solve` runs in its discovery mode. With the exhaustive flag off it anneals instead of enumerating, following the Metropolis walk of Algorithm 1 under geometric cooling, removals biased toward low-sensitivity edges, and returns the densest feasible graph it encountered together with the full temperature trace. What it returns is a witness, hence a lower bound, never a proof.

It is fair to ask why the walk is allowed above the ceiling at all. Reaching every feasible graph does not require it. Removing an edge can only lower $\lambda^{\max}$ and $\kappa^{\max}$, never raise them, so every subgraph of a feasible graph is again feasible, and the empty graph is feasible trivially. From any feasible target we can therefore delete edges one at a time down to the empty graph without ever crossing the ceiling, and reading that path backward builds the target up the same way. The feasible region is connected through the empty graph by single-edge moves that stay inside it, so a walk that never went infeasible could in principle visit every feasible graph there is.

We let it go infeasible anyway, because reachability and efficiency are different things. A feasible graph can be locally maximal, meaning every edge we might add creates a forbidden pair, and yet still sit far below the densest feasible graph. To improve such a graph the walk has to add an edge and then remove a different one, a swap whose first half is infeasible. A walk confined to feasible graphs could only find that better graph by retreating most of the way back to the empty graph and climbing a different slope, a detour so long that the cooling schedule usually ends first. Allowing a brief excursion above the ceiling lets the walk tunnel straight across instead, trading one load-bearing edge for a better one in two moves rather than dozens. The soft penalty also turns the search into a single smooth landscape with no hard walls for the Metropolis rule to bounce off, and the same machinery then carries over unchanged to variants whose feasible region is not so conveniently connected. Going above the bound is never the goal, then. It is the shortcut the annealer takes on its way back down to it.

3.3 Temperature and edge sensitivity

The phrase “biasing removals toward low-sensitivity edges” in [Algorithm 1](#) is where the temperature acquires its meaning, and it is the heart of the method. For an edge e define its *sensitivity*

$$\sigma(e) = \lambda^{\max}(G) - \lambda^{\max}(G - e),$$

the amount by which deleting e relaxes the binding connectivity. A load-bearing edge sits on a tightest cut and has $\sigma(e) > 0$. Removing it would drop $\lambda^{\max}$ and so loosen the very constraint that limits how many edges the graph may hold. A free edge has $\sigma(e) = 0$: it can be removed without easing the binding pair, which usually means it can be put to better use elsewhere.

`edgeSensitivity(G, e) → ℕ`

MEASURE

in a graph G and one of its edges e

out $\sigma(e) = \lambda^{\max}(G) - \lambda^{\max}(G - e)$

how delete e and let the checker remeasure, where a positive value marks a load-bearing edge. This single number is the dial the cooling turns.

When the search proposes a removal it draws the edge with probability proportional to $e^{-\sigma(e)/T}$. The dial is now explicit. At high temperature these weights flatten and the walk toggles edges regardless of how load-bearing they are, exploring widely and willing to dismantle essential structure. As the temperature falls the weights concentrate on $\sigma = 0$, so the walk almost never disturbs a load-bearing edge and instead rearranges only the free ones, until the graph crystallises onto an extremal shape. This is exactly the behaviour one wants, and it is the behaviour the formulation asks for: the temperature tracks how strongly removing an edge changes the bound. [Figure 3.2](#) shows one such run settling onto a certified optimum. The other variants behave identically, and their traces are collected in [Section B.1](#).

The same sensitivities also explain a finished graph. Colouring a graph's arcs by σ shows at a glance which arcs are structural and which are slack. [Figure 3.3](#) illustrates all four sensitivity levels on one directed multigraph: the darkest arcs carry the most flow and lower $\lambda^{\max}$ the most when removed, the cool arc contributes nothing. The picture is not decoration: it is the certificate of tightness made visible, and it tells the search exactly which arcs to leave alone as the temperature falls.

A single cooling schedule can still settle into a local optimum, so in discovery mode `solve` runs several independent schedules from different seeds and keeps the densest result. A handful of restarts is enough to reach the known optima at the sizes we examine. This is a reliability measure, not a change of method, and no new routine: the same `solve`, asked to discover rather than to prove.




Figure 3.2. A single cooling run on the directed multigraph problem at $n = 4$, $m = 3$. *Left:* the temperature schedule, cooling geometrically from its starting value. *Right:* every visited graph plotted by its binding connectivity $\lambda^{\max}$ (horizontal) against its arc count (vertical), coloured by the step at which it was visited. Annealing uses a soft penalty rather than a hard wall, so while hot (dark points) the walk strays into the infeasible region $\lambda^{\max} > m - 1 = 2$, picking up extra arcs it is not allowed to keep. As the temperature falls (bright points) the penalty drives the walk back across the feasibility boundary and onto the densest graph that stays feasible, the certified optimum of 12 arcs at $\lambda^{\max} = 2$ (star).

3.4 Keeping the checker cheap inside the search

The energy of [Algorithm 1](#) reads the binding connectivity $\lambda^{\max}(G)$, and a literal reading of the loop would call the checker afresh on the whole graph at every step. That checker is the most expensive routine in the program, one maximum flow per ordered pair, and the walk runs for thousands of steps with several restarts, so such a search would spend nearly all of its time remeasuring graphs that barely changed from the step before. Two elementary facts remove almost all of that work without altering a single number the search reports.

Proposition 3.1 (Monotonicity of the binding connectivity). *Let G be a graph or directed multigraph, and let G' be obtained from G by changing one adjacency by a single unit of multiplicity. Then*

- ✧ (monotonicity) *removing the unit cannot raise $\lambda^{\max}$, and adding it cannot lower $\lambda^{\max}$, and*
- ✧ (unit step) *adding the unit raises $\lambda^{\max}$ by at most one.*

Both statements hold verbatim for the vertex measure $\kappa^{\max}$.

Proof. Fix an ordered pair (s, t) . By the max-flow / min-cut theorem $\lambda_G(s, t)$ is the smallest capacity of any cut separating s from t , where the capacity of a cut is the total multiplicity of the arcs that cross it from the source side to the sink side.

For part (1), reducing one adjacency by a unit lowers the capacity of every cut that the changed arc crosses and leaves all other cut capacities untouched, so no cut capacity rises.

Parallel arcs drawn explicitly, coloured by sensitivity σ :
cool ($\sigma = 0$, free) $\rightarrow$ warm (load-bearing)




Figure 3.3. A directed multigraph on eight vertices with $\lambda^{\max} = 9$, every parallel arc drawn explicitly and coloured by its sensitivity σ , so multiplicity is read by counting arcs rather than from a label. The point is the contrast between $s \rightarrow a$ and $s \rightarrow b$: both carry three parallel arcs, yet $s \rightarrow a$ is cool ($\sigma = 1$) because the single arc $a \rightarrow t$ downstream throttles that route to one unit, while $s \rightarrow b$ is hot ($\sigma = 3$) because all three of its copies carry flow. The arcs $s \rightarrow d$ are coolest of all ($\sigma = 0$): d is a dead end, so deleting them changes nothing. Removing a hot adjacency drops $\lambda^{\max}$ by its σ while removing a cool one does not, which is exactly what tells the search which arcs to leave alone as the temperature falls.

The smallest cut capacity therefore cannot rise either, which gives $\lambda_{G'}(s, t) \leq \lambda_G(s, t)$. Taking the maximum over all pairs yields $\lambda^{\max}(G') \leq \lambda^{\max}(G)$. Adding a unit is the same statement read backwards.

For part (2), write G'' for G with one unit added to a single adjacency. That extra unit of capacity crosses any fixed cut at most once, so every cut has capacity at most one greater in G'' than in G . The smallest cut capacity thus grows by at most one, giving $\lambda_{G''}(s, t) \leq \lambda_G(s, t) + 1$, and together with part (1) we have $\lambda_G(s, t) \leq \lambda_{G''}(s, t) \leq \lambda_G(s, t) + 1$. The maximum over pairs gives $\lambda^{\max}(G'') \leq \lambda^{\max}(G) + 1$.

For $\kappa^{\max}$ the argument is identical on the vertex-split network of Figure 2.3, where changing an adjacency changes the capacity of its single arc $u^{\text{out}} \rightarrow v^{\text{in}}$ in the same way. Parallel copies leave that capacity at one and so change $\kappa^{\max}$ not at all, which only strengthens both parts. $\square$

These two facts settle almost every step without a flow. The energy depends on $\lambda^{\max}$ only through the excess $\max(0, \lambda^{\max} - (m - 1))$, which is zero exactly when the graph is feasible.

So a removal applied to a feasible graph leaves it feasible by part (1), with excess still zero, and needs no measurement at all. An addition made while the graph sits strictly below the bound, at $\lambda^{\max} \leq m - 2$, stays feasible by part (2), again with excess zero and no measurement. Only an addition made right at the boundary $\lambda^{\max} = m - 1$ can tip the graph over, and only there does the search run a single flow, capped so that it stops the instant it finds one route too many. The exact connectivity value is recomputed only on the rare step whose proposal is genuinely infeasible, where the penalty term needs it.

To know which case applies, the search carries a running upper bound on $\lambda^{\max}$, raised by one whenever it accepts an addition (part (2)) and left untouched when it accepts a removal (part (1)), and resynchronised to the exact value at the cadence where it already pays for flows to refresh the sensitivity map. The bound can only ever sit at or above the truth, so any step it clears is certainly feasible.

The key point is that the energy returned on every step is, by [Proposition 3.1](#), exactly the value the naive implementation would have computed. Every accept-or-reject decision, every random draw, and the graph the walk finally returns are therefore unchanged, and the speed-up is invisible to every number the thesis reports. This is verified rather than asserted. The regression test `test_fast_path_matches_exact` runs both implementations, the fast one and a reference that measures exactly at every step, across all four graph variants and both separations, and checks that the two agree step for step in energy, in every accept-or-reject decision, and in the final graph, and that the returned graph is feasible under the exact checker. Nothing about the optimisation hides inside the bound: the witness the search reports is always certified by an exact measurement, exactly as in the slower search.

It is worth noting where this method earns its keep. For undirected graphs one could resynchronise the exact value cheaply from the Gomory–Hu tree of [Chapter 2](#), which encodes every pairwise edge-connectivity in $n - 1$ flows. The directed and vertex variants admit no such tree, so there the monotone bound and the capped predicate are the only inexpensive handle on feasibility, and they are what let one annealer serve every variant at speed.

3.5 Rediscovering the known extremisers

A search method is only worth trusting if it recovers what is already known before it is pointed at what is not. Run from the empty graph on the cases whose answers [Chapter 1](#) and [Chapter 2](#) settled, and told only to pack in edges without breaching the connectivity ceiling, the cooled search arrives unprompted at the named constructions. [Table 3.1](#) reports the densest graph it found against the value the theory predicts, and in every case the two agree.

What the table hides is that the graphs themselves are the named constructions, not merely graphs of the right size. The simple undirected run at $m = 2$ settles on a spanning tree, the only shape with $n - 1$ edges and no cycle. The undirected multigraph run settles on a star with every edge at multiplicity $m - 1$, an instance of the multi-tree extremiser behind [Theorem 1.3](#). The directed multigraph runs settle on the double star of [Chapter 2](#), both directions of every spoke

Construction	n	m	Search	Theory
spanning tree (simple undirected, $m=2$)	6	2	5	5
multiplicity- $(m-1)$ star (multi undirected, $m=3$)	6	3	10	10
bidirected star (simple directed, $m=2$)	4	2	6	6
bidirected star (simple directed, $m=2$)	6	2	10	10
hub–bipartite tie (simple directed, $m=2$)	7	2	12	12
double star (multi directed, $m=3$)	4	3	12	12
double star (multi directed, $m=3$)	5	3	16	16

Table 3.1. Validation by rediscovery. For each variant the temperature search, run from scratch with a handful of restarts, recovers exactly the extremal value proved or certified in the earlier chapters. The “Search” column is an honest lower bound found by annealing, and the “Theory” column is the proved or certified value.

at full multiplicity. The simple directed runs find the bidirected star at $n = 4$ and, at $n = 7$, land on twelve arcs, the value at which the hub and the one-directional wall tie, exactly the crossover [Theorem 1.9](#) predicts. The cooled search does not know these names. It rediscovers the structures because, under the connectivity ceiling, there is nowhere denser to go.

Some extremisers are past the size at which a quick search lands reliably, but the exact checker of [Chapter 2](#) settles them immediately by measuring their connectivity. The one-directional bipartite digraph on eight vertices has 16 arcs at $\lambda^{\max} = 1$, and 16 already exceeds the linear hub value $2(n - 1) = 14$: the smallest size at which the quadratic phenomenon strictly wins, made concrete. It is also a measured illustration of the search’s limit: repeated cooling runs at $n = 8$, $m = 2$ stall at the hub value of 14, because dismantling a finished hub and rebuilding a one-directional wall is far more than the single-arc moves of the walk can tunnel through. The known answer there comes from the construction and the checker, not from the search, which is exactly the division of labour this chapter argues for. The augmented bipartite digraph on ten vertices at $m = 3$ has 30 arcs at $\lambda^{\max} = 2$, the thirty-arc counterexample of [Remark 1.11](#), confirmed pair by pair against the naive bound of 27 it refutes. The directed double star on seven vertices at $m = 4$ has 36 arcs at $\lambda^{\max} = 3$, matching $2(n - 1)(m - 1)$. None of these is a proof of optimality. Each is an exactly measured feasible graph, a lower bound the checker stamps as genuine.

3.6 The trichotomy: proved, certified, conjectured

It is worth stopping to lay out, for every variant, the three kinds of knowledge the computation can hold about it, because the whole value of the pipeline is that they are kept apart and never quietly merged. About any given value the program can do one of three things. It can *prove* the value, closing a fixed size from above by exhausting every graph of that size, an upper bound with the standing of a hand proof. It can *certify* it, the same kind of upper bound but obtained by the cut-counting optimisation rather than blind exhaustion, reaching a vertex or two further. Or, where proof and certificate both run out, it can only *conjecture* the value, exhibiting a concrete dense graph that pins down a lower bound and leaving the matching upper bound to a later argument. Proof and certificate bound the answer from above. The search bounds it only from

below. Where a closed *formula* exists it can tie the two together, threading through the certified small cases and the conjectured large ones as a single curve. This is the discipline the whole thesis runs on, and it is worth stating as one: the search proposes a lower bound, a certificate or a hand proof disposes the upper bound, and only when both meet is a value called settled.

We begin with the single variant where all three kinds of knowledge are non-trivial and visibly distinct, the directed simple arc problem at a fixed $m \geq 3$, and then sweep the whole family. For that variant the knowledge splits cleanly by the size of the graph.

For small n the answer is not conjectured at all. The pruned exhaustion inside `solve` walks every simple digraph on n vertices and returns the exact maximum, complete and certain, an upper bound with the standing of a hand proof. At $m = 3$ it certifies 6, 9, 12 for $n = 3, 4, 5$, and there the exhaustion stops within a reasonable budget, the pruned walk no longer closing $n = 6$ in tractable time. That is the certified region, finite and closed.

Beyond it the ceiling is gone and only lower bounds remain, each witnessed by a concrete feasible graph. The search finds the extremal 15 arcs at $n = 6$ on its own. Past that size it falls behind: across six restarts it reaches 17, 17, 20, 23 for $n = 7, \dots, 10$, while the named constructions of [Chapter 1](#), measured exactly by the checker, supply 18, 21, 25, 30, the last being the thirty-arc graph of [Remark 1.11](#). The reported lower bound at each size is the better of the two, exactly as the rediscovery section combined search and checker by hand. These are conjectures with an honest floor and no matching ceiling. That is the searched region, unbounded and forever one-sided.

The two regions are tied together by a single closed form. The conjectured value

$$\ell_m^{\text{dir}}(n) = \max\left(m(n-1), \left\lfloor \frac{(n+m-2)^2}{4} \right\rfloor\right)$$

of [Conjecture 1.13](#) is not a fresh guess bolted on past the certified range. It is one formula that already passes through every certified value at small n and then continues, without a seam, through every searched value at large n . The directed-arc panels of [Figure 3.4](#) show this directly, the certified squares at the small sizes and the open lower-bound circles beyond both sitting on the one dashed conjecture curve, proof below the formula and witnessed graphs above it. The formula is the through-line that makes the small certified cases and the large conjectured ones a single statement rather than two.

Two features of those directed-arc panels are worth naming. The crossover from the linear hub term to the quadratic bipartite term happens at $n = 9$, where the second argument of the maximum first wins, and that is past the last certified size. The shape of the answer therefore changes in exactly the region where only the search can see it, which is precisely why a discovery tool was needed and a certifier alone would not do. And at $n = 10$ the formula reads 30, the count of the very counterexample that overturned the naive pre-2024 conjecture [\[5\]](#), so the closed form is not a smooth fit to easy data but one that survives the case that broke the previous guess.

Asking the same three questions of every variant, the answers sort into three regimes, and

Figure 3.4 shows all twelve at once, one panel per variant, in exactly this proved, conjectured, and guessed language.

In the first regime the closed form is already a theorem for every n , proved by hand in Chapter 1 or cited, and the computation only confirms it. The undirected families live here, edge and vertex alike, together with the directed problems at $m = 2$. For these the certified small cases and the rediscovered constructions of Table 3.1 are validation, not new knowledge: the formula was never in doubt, and the program's role is to check that the pipeline reproduces it. That parallel edges do not help for vertex-connectivity, and that the directed vertex multigraph reduces to the simple digraph, are settled by hand, so several rows here need no computation of their own at all.

In the second regime the three columns genuinely differ and the computation is load-bearing. This is the regime of the worked example above. The directed simple arc and vertex problems at $m \geq 3$ are certified only at the small n the exhaustion reaches and conjectured by the search beyond, with the formula of Conjecture 1.13 the through-line. The directed multigraph arc problem sits here too, with one extra economy: in its cut-counting mode `solve` proves the m -free number $M^*(n) = 2(n - 1)$ outright for $n \leq 6$, and the scaling identity $L_m^{\text{dir}}(n) = (m - 1) M^*(n)$ of Chapter 2 carries that one certified number to every m at once, so the closed form $2(n - 1)(m - 1)$ is proved across the whole m axis at those sizes. Past the crossover at $n = 7$ the conjectured value switches to the bipartite branch $(m - 1) \lfloor n^2/4 \rfloor$, where the even sizes are proved conditional on the odd ones and only a single mixed-regime minimum-degree statement keeps the induction from closing (Chapter A). In every row of this regime the formula agrees with proof where proof reaches and with the search where it does not, exactly the linkage the worked example displayed.

In the third regime there is no formula at all, and the honesty is in drawing the guess as a guess. The undirected vertex problem at $m \geq 6$ has been open since 1974 with its extremal structure genuinely unknown, and the hypergraph variants beyond the proved edge case are unexplored, the directed ones needing a one-tail, many-head reading of a Berge edge that this thesis defines itself only so that `solve` can measure it at all. Here the program still computes exact values at the few sizes that finish, and its search still exhibits feasible graphs at a handful more, but no closed form is in sight. So the panels of Figure 3.4 carry, instead of a dashed conjecture, a densely dotted line that simply interpolates the machine points in straight segments, trusted no further than the eye. We draw it *through* the points rather than fitting a smooth curve above them: the points are honest lower bounds, the truth lies on or above them, and a line riding above the points would claim more than the search ever found. It is fair to ask whether these dotted trends hint at a closed form, and two cautions must come first. At small n the complete graph or hypergraph is itself feasible, its binding connectivity being only $n - 1$, so the earliest points merely trace the trivial maximum (every possible edge) and reveal nothing of the open behaviour: the undirected-vertex points run up $\binom{n}{2}$ until n passes m , and only then bend toward the linear growth the proved edge value predicts. And past that bend the dotted line is only as strong as the quick search behind it; exactly as the search stalled on the quadratic wall in the rediscovery above, the lower bound it reports climbs only linearly where it cannot build a denser

construction. The clearest case is the directed hypergraph, whose bipartite family is quadratic, reaching $\sim (m-1)n^2/(4(r-1))$ hyperarcs by [Proposition A.39](#): there the search slips below that construction at larger n , so the dots understate the truth rather than tracing it. We therefore read the dotted trends as honest lower bounds and not as fitted laws, and the one closed-form hypothesis we record, the quadratic for the directed hypergraph, comes from the construction and not from the shape of the dots.

So to the question of whether one formula links what is proved to what is conjectured, the answer is three-valued and depends on the variant, and the grid of [Figure 3.4](#) shows all three at a glance. Where the problem is already solved the curve is solid, a theorem the computation merely agrees with. Where the problem is open but tractable the curve is dashed, a conjecture the certified squares confirm below and the searched circles confirm above, and that is the case the program was built for. Where the problem is genuinely hard the curve is dotted, a guess interpolated through the search points and an open question of what, if anything, the dots truly trace. [Figure 3.5](#) redraws the same grid at $m = 6$: the proved curves climb linearly with m exactly as their formulas predict, while the open variants' search points and their dotted interpolation climb higher without a proved curve to meet, so raising the threshold does not change which regime a variant lives in, only how far the dashed and dotted curves have travelled.

The same grid answers the harder question the trichotomy raises, which is what the search is worth where no upper bound exists to check it against. It says two things at once. First, wherever an upper bound exists, proved or certified, the search lands exactly on it and never above it at every plotted size, which is strong evidence that the search is not leaving edges on the table, with the one instructive exception recorded above: the directed $m = 2$ run at $n = 8$, where it stalls on the hub and the bipartite optimum must be supplied by the construction instead. Second, on the directed arc problem at $m \geq 3$, where no upper bound is known past the certified range, the plotted lower bounds land exactly on the conjectured value $\max(m(n-1), \lfloor (n+m-2)^2/4 \rfloor)$ at every reachable n , and the certified points agree with it as far as they go, but past the first open size it is the constructions, not the cooled walk, that carry the bounds there. Run from scratch the search confirms the conjecture at $n = 6$ and then plateaus a few arcs short (17, 17, 20, 23 against 18, 21, 25, 30 at $n = 7, \dots, 10$, across six restarts), for the same reason as the $m = 2$ stall recorded earlier: the extremal shapes past the crossover are bipartite walls, and a walk that has crystallised onto a hub cannot rebuild itself into a wall by single-arc moves within the cooling budget. So to the question “do we believe the reported lower bounds are best possible,” the grid answers: wherever a proof or a certificate can check them, yes; on the open sizes they sit on the only value anyone has conjectured, with the search confirming the first open size and the constructions carrying the rest. The gap between those lower bounds and what is proved is exactly the gap between the conjecture and the missing upper bound, which [Chapter 4](#) names precisely. The search has not closed that gap, and it cannot. What the pipeline supplies is reproducible evidence for which side of the gap the truth lies on.




Figure 3.4. All twelve variants at $m = 3$, edges against n , laid out by model (rows: simple, multigraph, 3-uniform hypergraph) and by separation and direction (columns: undirected edge, undirected vertex, directed arc, directed vertex). A *solid* curve is a value proved for all n , a *dashed* curve a conjectured formula, and a *densely dotted* curve a bare interpolation of the machine points where no formula is known. Filled squares are the sizes `solve` proved or certified exactly. Open circles are lower bounds at larger n , the better at each size of the search's densest find and a named construction measured by the checker, reported as a running maximum so a feasible graph at one size is never lost at the next. On the open variants the dotted curve is the piecewise-linear interpolation *through* those circles, never above them, and the axes are scaled to the data rather than to a loose certain interval.




Figure 3.5. The same twelve variants at $m = 6$. The undirected edge and multigraph edge cases remain proved (Mader's theorem holds for all m), while the undirected vertex separation is the famous open problem at $m \geq 5$ and the directed cases carry the same conjectured formula scaled to $m = 6$. Comparing this figure with the $m = 3$ version above shows that the proved cases grow linearly with m as the formulas predict, while on the open cases the search circles and their dotted interpolation climb steadily without a proved curve to meet.

Honesty

A number found by the search is reported as a lower bound, and is marked as certified or proved only when a separate cut-counting optimisation or a hand proof supplies the matching upper bound. No extremal value is ever asserted on the strength of the search alone. The figures in this chapter use fixed random seeds, so every search point is reproducible.

Chapter 4

Synthesis, Results, and Open Problems

The journey set out from one clean theorem and followed it across twelve variants, proving what proved cleanly, building a machine where the proofs ran out, certifying the small cases the machine could close, and using the search to discover the dense graphs the proofs had singled out and to point at the ones still open. This closing chapter draws the threads together. It maps the directed frontier, where proof and certificate both still fall short of the full answer, names the three genuine surprises the landscape contains, gathers the twelve answers into one picture, and says plainly what the program changed and where it stops.

4.1 The directed frontier and the single open lemma

Two of the open directions are directed, and they share a structure worth stating exactly rather than hiding. The directed arc problem is a race between two constructions on different scales. The directed hub ([Construction 1.8](#)) reaches $m(n-1)$ arcs and grows linearly. The augmented bipartite construction ([Construction 1.12](#)) packs $\lfloor (n+m-2)^2/4 \rfloor$ arcs and grows quadratically. Which leads depends on n , and the two cross at an order linear in m , near $n = 9$ at the $m = 3$ of [Figure 4.1](#). [Conjecture 1.13](#) asserts that together they are the whole story.

[Conjecture 1.13](#) is proved as a lower bound for all m , and it is a theorem at $m = 2$ by [Theorem 1.9](#). Past $m = 2$ it sits squarely in the second regime of the trichotomy of [Section 3.6](#): certified by exhaustion at the small sizes `solve` can close for $m = 3$, conjectured by the temperature search beyond, and never beaten anywhere the search reaches. What is missing is a general upper bound for $m \geq 3$, and the obstacle is structural rather than computational.

The route to the upper bound is to show that an extremal digraph, once it is past the hub regime, must look like the augmented bipartite construction: its vertices split into a part A and a part B with every arc running $A \rightarrow B$, and B thickened only mildly. Three structural facts, each proved in [Chapter A](#), push in this direction. Out-neighbourhoods are mutually unreachable, second out-neighbourhoods are disjoint, and a complete $A \rightarrow B$ partition forces each vertex of B to absorb at most $m-2$ arcs from inside B . With the partition free to vary, the total arc count $|B|(|A| + m - 2) = |B|(n - |B| + m - 2)$ is maximised at $|B| = \lceil (n + m - 2)/2 \rceil$, which gives $\lfloor (n + m - 2)^2/4 \rfloor$, precisely the conjectured upper bound.

The reduction is clean except at a single point, and it is worth naming the point exactly rather




Figure 4.1. The two directed lower bounds at $m = 3$. The linear hub branch leads for small n . The quadratic augmented-bipartite branch overtakes it near $n = 9$ and leads thereafter. [Conjecture 1.13](#) asserts their maximum is the exact value.

than hiding it. The count goes through provided the extremal digraph has *no arc from B back to A* : under that hypothesis a route between two vertices of B can never escape B , so the connectivity inside B is governed entirely by B 's own arcs and the budget applies. The bare existence of a complete $A \rightarrow B$ partition is not enough on its own, because a small example shows a backward arc can manufacture an extra route. So the whole upper bound for $m \geq 3$ now rests on one missing lemma: that an extremal non-hub digraph has no backward arc. The natural delete-and-compensate exchange that would prove it is not monotone, which is the precise reason it remains open. The machine has been useful here in the negative direction too: a proposed counterexample to the conjecture, a backward arc bolted onto the thirty-arc graph, was refuted by the checker, which found the third route the construction's author had missed.

The directed vertex-disjoint problem needs no separate machinery. Vertex-disjoint routes are at most as numerous as arc-disjoint ones, and the constructions that achieve the arc bounds also achieve them for vertices, so the two directed problems share both their constructions and their conjecture, and the $m = 2$ value transfers exactly in [Theorem 1.10](#). For $m \geq 3$ the vertex value is conjectured equal to the arc value, and it will follow from the arc case the moment the missing backward-arc lemma is supplied. The directed multigraph problem has its own two-branch story, parallel to the simple one. The conjectured value is

$$L_m^{\text{dir}}(n) = (m-1) \max(2(n-1), \left\lfloor \frac{n^2}{4} \right\rfloor),$$

where the double star realises the linear branch and the one-directional bipartite multigraph, every arc at multiplicity $m-1$, realises the quadratic one. The branches cross at $n = 7$, exactly as in the simple case. On the linear side the certifier closes the problem, proving $L_m^{\text{dir}}(n) = 2(n-1)(m-1)$ for $n \leq 6$ and every m at once ([Theorem 2.1](#)), and at $m = 2$ the whole problem collapses

onto the simple digraph theorem for every n , because two parallel arcs are already two routes (Theorem A.24). On the quadratic side, a scaling-and-deletion induction proves the bipartite bound for even $n \geq 8$ conditional on the bound one vertex below, and that odd case once reduced to a single minimum-degree statement (Conjecture A.22), confirmed by the program in the purely saturated and purely unsaturated regimes but open in the mixed one, where multiplicities strictly between 0 and $m - 1$ defeat the integrality the two pure regimes lean on (Chapter A).

A sharper route now bypasses that conjecture at $m = 3$, and the way it does so is the cleanest instance in the thesis of the proof handing its remaining work to the machine. An *attachment lemma* (Lemma A.26) bounds the degree of a vertex glued onto the bipartite extremiser, and a *conditional odd step* built on it (Theorem A.28, for $m \in \{3, 4\}$) carries the value and extremal characterisation from each even level to the next odd one. At $m = 3$ that characterisation propagates through both parities (Remark A.29), so the entire problem for all $n \geq 8$ collapses onto two finite, integral statements at $n = 7$:

- (a) $L_3^{\text{dir}}(7) = 24$, and
- (b) the only 24-arc feasible multigraphs on 7 vertices with maximum degree at most 8 are $2 \cdot B_{3,4}$, $2 \cdot B_{4,3}$, and the doubled bidirected path P_7 .

Conjecture A.22 is thereby off the critical path for $m = 3$, replaced by two machine-checkable targets: a MILP infeasibility check for fact (a), and an exhaustive enumeration of degree-bounded multigraphs for fact (b). Both are running as this is written.

The linear branch turns out to be richer than the double star first suggested. Every doubled bidirected spanning tree is extremal, confirmed exhaustively at $n = 4$ and 5, and several distinct extremisers tie at 24 arcs for $n = 7$. Sifting and classifying them is precisely what the enumeration must do.

For $m \geq 4$ the conditional step still yields the value but uniqueness no longer propagates through odd levels, and for $m \geq 5$ even the value step stalls. These residual gaps are stated precisely among the open problems below. They are finite, identified targets rather than mysteries, and natural goals for the next round of certified computation.

4.2 Three principal phenomena

Across all the variation, three phenomena stand out as the ones a first guess would not have predicted.

The first is the divergence at $m = 5$. The edge and vertex problems agree for $m \leq 4$, and it would be natural to expect them to agree forever. Instead they part company at the very next value, with $k_5(n) = \lfloor 8n/3 \rfloor - 3$ pulling strictly above $\ell_5(n)$, and beyond it the vertex problem becomes genuinely hard and has stayed open since 1974.

The second is the quadratic bipartite phenomenon. Undirected graphs tie their edge count to their connectivity through vertex degrees, but a digraph can lean entirely one way, and a single-

direction wall of arcs holds $\lfloor n^2/4 \rfloor$ arcs at connectivity one. This has no undirected counterpart, it overturns the naive directed conjecture, and it is the reason the directed cases are the hardest in the thesis.

The third is the universality of the threshold $p^* = m/n$, with one honest asterisk. [Chapter 1](#) proves the appearance threshold $p^* = m/n$ for simple $G(n, p)$ ([Theorem 1.15](#)), and its mechanism is a degree count: below threshold the maximum degree stays under m , so the degree bound caps every local connectivity, and above it Mader's bound forces a high pair into existence. That mechanism does not care which model it runs on, so the natural general statement is that the forbidden configuration appears once the *expected degree* crosses m , of which $p^* = m/n$ is merely the simple-graph face, since there $\mathbb{E}[\deg] = p(n-1)$. To put this to all twelve variants at once we extend $G(n, p)$ by a single rule: every cell (a pair, an ordered pair, or an r -set) carries an independent weight, Bernoulli(p) for simple graphs and hypergraphs, and for multigraphs a *hurdle-geometric* weight that is empty with probability $1-p$ and otherwise $1 + \text{Geometric}(\alpha)$ parallel copies, so the multiplicity decays geometrically (we use $\alpha = \frac{1}{2}$ throughout) and $\alpha = 0$ recovers $G(n, p)$ exactly. For multigraph panels, multiplicities are capped at $m-1$ so that a single fat edge cannot trivially breach the connectivity ceiling. For simple-graph and hypergraph panels no cap is needed, since those models already admit at most one copy per cell or hyperedge.

[Figure 4.2](#) samples all six generative shapes and plots, against the measured mean degree in units of m , the chance that a sample already holds a pair of binding connectivity at least m . The universality is real but *asymptotic*. The degree argument is model-free only in the regime $m/\ln n \rightarrow \infty$; at the fixed small m a computation can reach, the curves are visibly *ordered* rather than coincident. The ordering, from cheapest to most expensive in mean degree required to force a high pair, is: directed multigraph, directed simple, undirected simple, undirected multigraph, undirected hypergraph, directed hypergraph. The directed multigraph crosses first because parallel edges concentrate connectivity into a few pairs without spreading degree evenly. The undirected multigraph lands further right than the simple cases because a lower inclusion probability (needed to keep mean degree fixed) leaves many pairs absent, making the minimum-degree bound harder to satisfy. The directed hypergraph needs the most degree because each hyperedge serves only one Berge route at a time. The vertical line is the asymptotic threshold, degree = m , and the spread of crossings around it is exactly the finite- m gap [Theorem 1.15](#) already disclaims — the collapse onto one density is a statement about the limit, not about any size we can sample.

That same sampler also lets the binding-connectivity *distribution* be seen for every variant at sizes enumeration cannot reach. [Figure 4.3](#) histograms it across all twelve, at the common density where the expected degree is m , feasible graphs ($\lambda^{\max} \leq m-1$) blue and infeasible red. The colour balance reads off the ordering of [Figure 4.2](#) in distributional form: at this one density the models that turn degree into connectivity cheaply (the directed multigraph above all) already sit deep in the infeasible regime, while the dear directed hypergraph is still mostly feasible, and the simple and undirected cases straddle the boundary. That the same ordering survives at n up to 26, far past the $n \leq 6$ of exact enumeration, is the point: it is a property of the models, not of




Figure 4.2. The appearance threshold across all six generative models, read against expected degree ($n = 12$, $m = 4$, multigraph panels use the hurdle-geometric model with $\alpha = \frac{1}{2}$, multiplicities capped at $m - 1$). Each curve samples one model (simple, multigraph, or 3-uniform hypergraph, each undirected and directed) and plots the probability that a sample already contains a pair of binding connectivity at least m against its *measured* mean binding degree, in units of m . The vertical line is the asymptotic threshold degree $= m$ of Theorem 1.15. At this finite m the curves form a strict six-way ordering, not a coincident band: multigraph directed crosses first (parallel edges concentrate connectivity into a few pairs), then simple directed, then simple undirected, then multigraph undirected (lower inclusion probability offsets the per-pair multiplicity gain), then 3-uniform hypergraph undirected, and finally 3-uniform hypergraph directed last (each Berge route consumes an entire hyperedge, so connectivity is hardest to force). The model-free collapse onto one threshold is the $m/\ln n \rightarrow \infty$ limit. The spread here is the finite- m gap.

the tiny sizes.

A three-dimensional view of the same transition, tracking how the whole binding-connectivity distribution migrates from low to high as the density crosses $p^* = m/n$, is given for three representative variants in Figure B.8 of Section B.2.

That same scarcity of extremal graphs is the thread through the remaining figures, which pin it down from three complementary angles: where the frontier runs, how far the typical graph sits below it in connectivity, and how far below it in sheer edge count. Figure 4.4 answers the first. It plots every labelled graph at small n as a single point, its binding connectivity $\lambda^{\max}$ on the horizontal axis and its edge count on the vertical, so the whole population of graphs is visible at once. The points crowd into the low-connectivity, low-edge corner and thin out upward, where the blue extremal envelope, an integer staircase carrying one dot per connectivity level, traces the densest graph available at each level. The empty region above that staircase is precisely the part the main theorems forbid a graph to occupy. The extremal problem is the question of where that staircase runs, and the figure shows it is a thin frontier with almost nothing pressed against it from below.

The two enumeration histograms then look at the same populations from two angles. Fig-




Figure 4.3. Binding-connectivity distribution under random sampling for all twelve variants, at sizes up to $n = 26$ (far past the $n \leq 6$ exact enumeration reaches), each at the common density where the expected degree is about $m = 4$. Blue is feasible ($\lambda^{\max} \leq m - 1$), red infeasible, and the dashed line in each panel marks the boundary $\lambda^{\max} = m - 1$. At this shared density the four simple panels are already infeasible almost everywhere, their mass sitting well to the right of the boundary in red, because a simple graph reaches high local connectivity readily once its degree is about m . Connectivity is hardest to force in the directed hypergraph panels, which stay mostly feasible in blue, since each Berge route consumes an entire hyperedge. The multigraph rows and the vertex-separation panels keep more blue than their simple, edge-disjoint counterparts and fall between these extremes. Multigraph rows use the hurdle-geometric model ($\alpha = \frac{1}{2}$, multiplicities capped at $m - 1$), and the per-panel n is annotated.

Figure 4.5 sorts every graph in the full small- n enumeration by its $\lambda^{\max}$ at $m = 3$, colouring the feasible graphs ($\lambda^{\max} \leq 2$) blue and the infeasible ones red. The balance of the two varies markedly across the twelve variants, from panels where most graphs already violate the bound to panels where the feasible graphs dominate, so the constraint $\lambda^{\max} \leq m - 1$ is one that a large fraction of graphs genuinely fail. Figure 4.6 turns the same enumeration the other way, by edge count, and marks the extremal value at that size with a dotted line that lands far out in the sparse right tail, so the densest feasible graphs are exceptional in every variant. Together the histograms place the extremal graphs as a thin population at the frontier, and Figure 4.4 shows that frontier directly. The same picture holds at $m = 6$, only shifted: the feasibility boundary moves out to $\lambda^{\max} \leq 5$ and the extremal edge counts grow with m exactly as the formulas predict, so raising the threshold relocates the boundary without changing the verdict. The full $m = 6$ enumeration views, the companions to Figure 4.5 and Figure 4.6, are Figure B.6 and Figure B.7 in Section B.2.

Finally, Figure 4.7 lifts the whole study onto the (n, m) grid at once, drawing the best feasible edge count as a bar surface across every variant. The three proved variants rise as smooth linear or quadratic sheets that the formulas describe exactly, while the open variants show a visible step between the machine-proved blue bars and the search's purple lower bounds, so the surface makes the proved-versus-open divide of the trichotomy a single readable shape. Reading along the m axis recovers the linear scaling in the threshold, and reading along n recovers each variant's growth rate, which is the whole of Table 4.1 compressed into one object.

4.3 Summary of the twelve variants

Table 4.1 records the status of all twelve structural variants. Each entry is marked by how it is known: proved by hand here or in Chapter A, cited to the literature, certified by the cut-counting optimisation for the stated sizes, or conjectured with a proved lower bound. The distinction is the honesty contract of Chapter 2 carried through to the summary.

4.4 Contributions and limitations of the program

The program turned a per-case craft into a uniform pipeline. Before it, each variant came with its own one-off computation and its own hand argument. After it, one representation holds every variant, one exact checker measures connectivity, one certifier proves small-case upper bounds, and one cooled search discovers extremisers and the lower bounds they witness. The rediscovery of Chapter 3 is the evidence that the pipeline is faithful: pointed at solved cases, it returns the solved answers.

Its limits are as clear as its reach, and stating them is part of the honesty the thesis insists on. The search proves no upper bounds. It only ever exhibits graphs. The certifier proves upper bounds, but only for a fixed size, and the directed multigraph encoding closed unconditionally only up to $n = 6$ until the deletion-cut strengthening put $n = 7$ within reach (the two $m = 3$ computations that the conditional reduction now hangs on are running as this is written, not yet




Figure 4.4. The complete landscape of all labeled graphs at small n (see panel annotations for each variant's n): every point is one graph, plotted at its binding connectivity $\lambda^{\max}$ horizontally and its edge count vertically. The blue staircase marking the densest graph at each connectivity level is the extremal envelope, the frontier the extremal problem asks about, with one dot per level so its exact value is legible. Graphs crowd the low-connectivity, low-edge region. The extremal graphs live at the top of the staircase. The emptiness above the envelope is exactly the region the main theorems forbid.




Figure 4.5. How all graphs (full enumeration, small n) distribute by their binding connectivity $\lambda^{\max}$, across all twelve variants at $m = 3$. Blue bars are feasible graphs ($\lambda^{\max} \leq 2$), red bars infeasible ones ($\lambda^{\max} \geq 3$), and the dashed line marks the feasibility boundary. The balance of blue and red varies markedly across the panels, from variants where the infeasible graphs are the clear majority to variants where the feasible ones dominate, so high binding connectivity is common enough that the bound is a genuine constraint rather than a formality. The densest feasible graphs, the extremal ones, are rare in every variant regardless of this balance, as the edge-count view of [Figure 4.6](#) makes precise.




Figure 4.6. The same enumeration viewed by edge count rather than connectivity, at $m = 3$. Feasible graphs are blue, infeasible ones red. The dotted vertical line marks the largest feasible edge count, the extremal value at this n read straight from the full enumeration, so the reader can see directly how far the extremal graphs sit from the bulk of the distribution and whether the extremal density is typical or exceptional.

Erdős 915: optimal bound over (n, m) , blue where proved, purple where only a search lower bound is known




Figure 4.7. The optimal bound as a surface over the (n, m) grid for all twelve variants, every panel drawn from the same camera so the shapes compare directly. Each bar's height is the largest feasible edge count found by `solve` at that vertex count and forbidden threshold m . The single thing to read off is colour: a bar is *blue* exactly where the value is machine-proved and *purple* exactly where only a search lower bound is known, so each panel is a little map of what is settled. The three fully proved variants (Mader's undirected edge cases and the multigraph analogue) are solid blue sheets, rising linearly or quadratically as their formulas predict, while the open variants turn purple precisely where proof runs out. Reading along the m axis shows how the bound scales with the forbidden threshold; reading along the n axis shows the growth rate for a fixed variant.

Separation	Model	Value	Status
<i>Undirected</i>			
edge	simple	$\lfloor m(n-1)/2 \rfloor$	proved (Mader)
edge	multi	$(m-1)(n-1)$	proved
edge	hyper	$\lfloor (m-1)(n-1)/(r-1) \rfloor$	proved (cited GH)
vertex	simple	$\lfloor m(n-1)/2 \rfloor$ ($m \leq 4$), $\lfloor 8n/3 \rfloor - 3$ ($m = 5$)	cited, open $m \geq 6$
vertex	multi	parallel edges do not help	proved
vertex	hyper	$\lfloor \frac{(m-1)(n-1)}{r-1} \rfloor$ ($m \leq 3$), $\geq$ that for all m	proved $m \leq 3$ (Theorems A.34 and A.37), open $m \geq 4$
<i>Directed</i>			
arc	simple	$\max(2(n-1), \lfloor n^2/4 \rfloor)$ ($m = 2$)	proved, conj. $m \geq 3$
arc	multi	$(m-1) \max(2(n-1), \lfloor n^2/4 \rfloor)$	proved $n \leq 6$ and $m=2$, cond. $m=3$
arc	hyper	quadratic in n , exact value open	lower bd. (Proposition A.39)
vertex	simple	$\max(2(n-1), \lfloor n^2/4 \rfloor)$ ($m = 2$)	proved, conj. $m \geq 3$
vertex	multi	reduces to simple digraph	proved
vertex	hyper	quadratic in n , exact value open	lower bd. (Proposition A.39)

Table 4.1. The twelve structural variants of Problem 915 and their status. The threshold result $p^* = m/n$ applies uniformly across all models and separations and is discussed separately in Chapter 1 and Chapter 4 as a sampling observation. “Certified” means proved by the cut-counting optimisation for the stated sizes. “conj.” means a conjecture with a proved matching lower bound. “cond.” means proved conditional on a finite, identified statement (for $m = 3$ directed multigraphs, the two $n = 7$ computations of Remark A.29). “lower bd.” means only a proved lower bound is known, with the exact value open.

returned). And no amount of either replaces the two genuinely structural gaps the thesis leaves open: the backward-arc lemma that would turn Conjecture 1.13 into a theorem, and the vertex-disjoint problem at $m \geq 6$, which has resisted for half a century and which the program can probe but not crack.

4.5 Open problems

- ★ The backward-arc lemma: prove that an extremal non-hub digraph has no arc from B to A , completing the upper bound in Conjecture 1.13 for $m \geq 3$, and with it the directed vertex case.
- ★ The two finite $n = 7$ computations that close the directed multigraph problem at $m = 3$. By the attachment lemma and the conditional odd step (Lemma A.26, Corollary A.27, Theorem A.28), the entire $m = 3$ problem — value and extremal uniqueness for all $n \geq 8$ — reduces along a single induction to two integral statements about multigraphs with multiplicities in $\{0, 1, 2\}$ (Remark A.29): (a) $L_3^{\text{dir}}(7) = 24$, and (b) every 24-arc feasible multigraph on 7 vertices with maximum degree at most 8 is $2 \cdot B_{3,4}$, $2 \cdot B_{4,3}$, or the doubled bidirected path P_7 . Both sit at sizes the certifier and the enumerator can reach, and both are presently running — a 24-hour MILP infeasibility check for (a) and an exhaustive enumeration for (b) — so this is a finite and machine-checkable gap rather than a structural one. Conjecture A.22 is thereby bypassed for $m = 3$, though it remains open and of independent interest as the natural minimum-degree statement for the directed multigraph problem.
- ★ Extremal uniqueness at odd levels for directed multigraphs with $m \geq 4$. The conditional odd step (Theorem A.28) and the propagation of Remark A.29 are stated for $m \in \{3, 4\}$. For $m = 4$ the value step goes through, but at an odd extremal level the counting no longer forces

a vertex of degree $(m-1)k$ (all degrees $\geq (m-1)k+1$ are arithmetically consistent), so the uniqueness statement does not yet propagate. For $m \geq 5$ the same escape climbs one level and even the *value* step stalls: a surplus extremiser could be degree-regular strictly above the deletion bound, so no vertex deletion is forced to land on an extremal multigraph. Closing either hole asks for an arithmetic or structural input the present averaging argument does not supply.

- ★ The undirected vertex-disjoint problem for $m \geq 6$, open since 1974, where even the extremal structure is unknown.
- ★ The hypergraph vertex problem at $m \geq 4$. The problem is now settled at $m = 2$ and $m = 3$ for every uniformity and all n , with $k_m^{(r)}(n) = \left\lfloor \frac{(m-1)(n-1)}{r-1} \right\rfloor = \ell_m^{(r)}(n)$ (Theorems A.34 and A.37). The incidence reduction stands for every m , but the rank bound that drives it (Lemma A.36) leans on the triconnected (Tutte/SPQR) decomposition exactly where $\kappa \leq 2$ lives. For $m = 4$ it would need a 4-connectivity analogue, where no comparably clean decomposition is available. Exhaustive computation gives $k_4^{(3)}(5) = 6 = \lfloor 3 \cdot 4/2 \rfloor$, so the agreement $k_m^{(r)} = \ell_m^{(r)}$ survives at least one step past the theorem and there is no early counterexample. Where it first breaks for $r \geq 3$ (the analogue of the $m = 5$ graph divergence) is unknown.
- ★ The directed hypergraph variants. The tail-and-heads model has first bounds and a quadratic lower-bound family (Proposition A.39): a pair-codegree count caps the hyperedges at $(m-1)n(n-1)/(r-1)$, and a bipartite construction reaches $\approx (m-1)n^2/(4(r-1))$, so the value is quadratic in n . Its exact order, and the matching upper construction, are unexplored.

The map is filled in where honest mathematics could fill it, and where it could not, the blanks are marked precisely, with a machine standing ready at each of them. That, finally, is the use of building the microscope: not that it answers every question, but that it shows exactly which questions are left, and hands the next reader a clean place to start.

Appendix A

Full Proofs and Computational Transcripts

This appendix holds the arguments deferred from the body: the ones that are either standard machinery or long case-checking, and whose length, not whose difficulty, kept them out of the narrative. The body carries the ideas. Here they are carried out in full. Because graphs are visual objects, the longer arguments here are accompanied by figures: each is drawn rather than merely described, and the prose points the reader to the picture at the moment it is needed.

How to read this appendix. The sections divide into standard machinery and the thesis's own proofs, and a reader chasing one thread of the body need not read all of them. The first two sections, on [Theorem A.1](#), [Theorem A.2](#), and Mader's theorem, are classical results included only so the body can cite them in the exact form it uses, and a reader who already trusts Menger, Gomory–Hu, and Mader may skip them. Three sections carry the load-bearing original arguments. The directed arc value at $m = 2$ supplies the finite base cases the body's induction rests on. The directed multigraph problem at large n is the longest argument here and the one behind the main directed contribution. The hypergraph vertex problem proves the incidence-rank lemma that settles the $m = 3$ hypergraph case. The remaining sections, on the hypergraph edge bounds and the random threshold, each support a single claim of the body and can be read on their own when that claim is reached. Finally, everything the program asserts is reproduced verbatim in the computational transcripts and the self-test section, so the machine proofs can be audited without running anything.

A.1 Menger, Gomory–Hu, and the degree bound

The two classical theorems below are the foundation of everything the thesis measures, and we state them in the form we use them.

Theorem A.1 (Menger [15]). *For any graph G and distinct vertices u, v , the maximum number of pairwise edge-disjoint u – v paths equals the minimum number of edges whose removal separates u from v :*

$$\lambda_G(u, v) = \min\{|C| : C \subseteq E(G), G - C \text{ has no } u\text{--}v \text{ path}\}.$$

The same statement holds for digraphs with arc-disjoint directed paths and directed arc cuts, and, replacing edges by internal vertices, for internally vertex-disjoint paths and vertex cuts.

Theorem A.2 (Gomory–Hu [7]). *Every weighted graph G has a weighted tree T on the same vertex set such that for all u, v the value $\lambda_G(u, v)$ equals the minimum edge weight on the u – v path in T , and each tree edge e corresponds to a minimum cut of G whose capacity is the weight of e .*

The most basic constraint on local connectivity is local: a pair cannot have more independent routes than either endpoint has edges. The constructions below use it constantly.

Lemma A.3 (Degree bound). *For every pair of vertices of a graph G , $\lambda_G(u, v) \leq \min(\deg u, \deg v)$. In a digraph D , $\lambda_D(u, v) \leq \min(d^+(u), d^-(v))$.*

Proof. Every edge-disjoint u – v path uses a distinct edge at u , so there are at most $\deg u$ of them, and likewise at most $\deg v$. In the directed case each arc-disjoint path leaves u by a distinct out-arc and enters v by a distinct in-arc. $\square$

A.2 Mader’s theorem in full

The upper bound of Theorem 1.2 rests on three short lemmas about a Gomory–Hu tree T of G . For an edge $e = \{u, v\}$ of G , write $\text{dist}_T(e)$ for the number of tree edges on the unique u – v path in T . Since T is a tree on the very same vertex set, $\text{dist}_T(e) \geq 1$ for every edge. Figure A.1 shows a small graph beside one of its Gomory–Hu trees and marks which edges have $\text{dist}_T(e) = 1$ and which have $\text{dist}_T(e) \geq 2$; it is worth keeping in view, since the three lemmas below are exactly statements about that picture. The first lemma is the one place the proof uses that G is *simple*, and it is exactly where the simple-graph bound improves on the multigraph bound.




Figure A.1. A simple graph G (left) and a Gomory–Hu tree T of it (right); both live on the same five vertices, and here T is the path e – a – b – c – d with the marked cut weights. The four edges ea, ab, bc, cd are also tree edges, so each has $\text{dist}_T(e) = 1$: these are the E_1 edges of Lemma A.4, and a *simple* graph admits at most $n - 1 = 4$ of them, one per tree edge. The edges ac and bd join tree-distance-2 pairs, and the red chord ec has $\text{dist}_T(ec) = 3$, crossing all three cuts on the path e – a – b – c . Reading $\text{dist}_T(e)$ as the number of tree cuts an edge crosses makes $\sum_e \text{dist}_T(e) = \sum_t c(t)$ (Lemma A.6); every edge crosses at least one cut, but only the $n - 1$ tree edges cross exactly one, which is the surplus that drives Lemma A.5 and ultimately the factor of two in Mader’s bound.

Lemma A.4 (Distance-one edges). *At most $n - 1$ edges of a simple graph G satisfy $\text{dist}_T(e) = 1$.*

Proof. An edge $e \in E(G)$ has $\text{dist}_T(e) = 1$ if and only if its endpoints are adjacent in T . The tree

T has exactly $n - 1$ edges, and since G is simple each pair of adjacent tree vertices supports at most one edge of G . $\square$

Lemma A.5 (Sum of tree distances). $\sum_{e \in E(G)} \text{dist}_T(e) \geq 2|E(G)| - (n - 1)$.

Proof. Partition $E(G)$ into $E_1 = \{e : \text{dist}_T(e) = 1\}$ and $E_{\geq 2} = \{e : \text{dist}_T(e) \geq 2\}$. Then

$$\sum_{e \in E(G)} \text{dist}_T(e) \geq |E_1| + 2|E_{\geq 2}| = 2|E(G)| - |E_1| \geq 2|E(G)| - (n - 1),$$

using [Lemma A.4](#) in the last step. $\square$

Lemma A.6 (Double-counting identity). *For any graph G and Gomory–Hu tree T with weights c ,*

$$\sum_{t \in E(T)} c(t) = \sum_{e \in E(G)} \text{dist}_T(e).$$

Proof. By [Theorem A.2](#) each tree edge t induces a cut of G of capacity $c(t)$, so $c(t) = \sum_{e \in E(G)} \mathbf{1}_{e \text{ crosses } t}$. Exchanging the order of summation,

$$\sum_{t \in E(T)} c(t) = \sum_{e \in E(G)} \sum_{t \in E(T)} \mathbf{1}_{e \text{ crosses } t} = \sum_{e \in E(G)} \text{dist}_T(e),$$

where the last equality holds because an edge $\{u, v\}$ crosses exactly those tree cuts induced by the tree edges on the unique u – v path in T . $\square$

Proof of [Theorem 1.2](#), upper bound. Let G be a simple graph on n vertices with $\lambda_G^{\max} \leq m - 1$ and let T be a Gomory–Hu tree of G . Each tree edge $t = \{x, y\}$ has weight $c(t) = \lambda_G(x, y) \leq m - 1$, so summing over the $n - 1$ tree edges, $\sum_t c(t) \leq (m - 1)(n - 1)$. Combining with [Lemmas A.5](#) and [A.6](#),

$$2|E(G)| - (n - 1) \leq \sum_{e \in E(G)} \text{dist}_T(e) = \sum_{t \in E(T)} c(t) \leq (m - 1)(n - 1).$$


Rearranging gives $2|E(G)| \leq m(n - 1)$, and since $|E(G)|$ is an integer, $|E(G)| \leq \left\lfloor \frac{m(n-1)}{2} \right\rfloor$. $\square$

Note where the factor of two came from: every edge crosses at least one tree cut, but a simple graph can afford at most $n - 1$ edges that cross only one, so on average each edge is charged closer to twice. For multigraphs that refinement is unavailable, since parallel edges between tree-adjacent pairs each have $\text{dist}_T = 1$, and the same chain of inequalities stops at $|E| \leq \sum_e \text{dist}_T(e) \leq (m - 1)(n - 1)$, which is exactly the multigraph bound of [Theorem 1.3](#).


The lower bound is a pair of constructions, drawn side by side in [Figure A.2](#). The first is the shape the bound was guessed from, and works when $m - 1$ divides $n - 1$; the second generalises it to every $n \geq m \geq 2$.

Construction A.7 (Shared-clique graph). For $m \geq 3$ with $(m-1) \mid (n-1)$, take $k = (n-1)/(m-1)$ copies of the complete graph K_m and identify one vertex of each copy into a single shared hub vertex v^* . The result has $k(m-1) + 1 = n$ vertices and $k \binom{m}{2} = \frac{m(n-1)}{2}$ edges. Every non-hub vertex has degree $m-1$, and any pair of vertices contains a non-hub vertex, so $\lambda^{\max} \leq m-1$ by Lemma A.3.

Construction A.8 (Hub-and-spokes graph). For $n \geq m \geq 2$, let $H_{m,n}$ have vertex set $\{h, v_1, \dots, v_{n-1}\}$ with (i) all $n-1$ hub edges $\{h, v_i\}$, and (ii) a spoke graph on $\{v_1, \dots, v_{n-1}\}$ with $\left\lfloor \frac{(m-2)(n-1)}{2} \right\rfloor$ edges in which every v_i has spoke degree at most $m-2$. Such a spoke graph exists by Lemma A.9.



(a) Shared-clique graph: three K_4 's glued at one hub v^* ($m = 4$, $k = 3$). Each non-hub vertex has degree $m-1 = 3$.



(b) Hub-and-spokes $H_{m,n}$: the hub edges $\{h, v_i\}$ (light) plus a sparse spoke graph (dark) of spoke-degree at most $m-2$.

Figure A.2. The two extremal families. (a) The shared-clique graph of Construction A.7 works when $(m-1) \mid (n-1)$: every pair contains a non-hub vertex of degree exactly $m-1$, so $\lambda^{\max} \leq m-1$ by the degree bound, while the edge count hits $\frac{m(n-1)}{2}$. (b) The hub-and-spokes graph $H_{m,n}$ of Construction A.8 reaches the same bound for every $n \geq m$: each spoke vertex v_i has total degree $1 + (m-2) = m-1$, and choosing the spoke graph as disjoint cliques K_{m-1} recovers (a). The hub is drawn in orange in both.

Lemma A.9 (Near-regular graphs exist). For $N \geq 1$ and $0 \leq r \leq N-1$ there is a graph on N vertices with $\left\lfloor \frac{rN}{2} \right\rfloor$ edges and maximum degree at most r .

Proof. If rN is even we build an r -regular circulant graph on vertex set $\{0, \dots, N-1\}$: for even r connect each i to $i \pm j \pmod{N}$ for $j = 1, \dots, r/2$; for odd r (then N is even) use $j = 1, \dots, (r-1)/2$ together with the diameter distance $N/2$, which contributes degree one. Both are r -regular with $rN/2$ edges. If rN is odd, then r and N are both odd: take the $(r-1)$ -regular circulant with distances $1, \dots, (r-1)/2$ and add the near-perfect matching that pairs vertex i with $i + (N+1)/2 \pmod{N}$ for $i = 0, \dots, (N-3)/2$, which covers all vertices but one. The total is $\frac{(r-1)N}{2} + \frac{N-1}{2} = \frac{rN-1}{2} = \left\lfloor \frac{rN}{2} \right\rfloor$ edges, with one vertex of degree $r-1$ and all others of degree r . $\square$

Theorem A.10 (Extremal graphs exist). For all $n \geq m \geq 2$ the graph $H_{m,n}$ has $\left\lfloor \frac{m(n-1)}{2} \right\rfloor$ edges and $\lambda^{\max} \leq m-1$.

Proof. The edge count is $(n-1) + \left\lfloor \frac{(m-2)(n-1)}{2} \right\rfloor = \left\lfloor \frac{m(n-1)}{2} \right\rfloor$, since $(m-2)(n-1)$ and $m(n-1)$ have the same parity. For the connectivity, any two distinct vertices include at least one spoke vertex v_i , whose total degree is $1+(m-2) = m-1$; [Lemma A.3](#) then gives $\lambda(u, v) \leq m-1$. When $(m-1) \mid (n-1)$, choosing the spoke graph as k disjoint cliques K_{m-1} recovers [Construction A.7](#), so the hub-and-spokes family genuinely generalises the shared-clique one. $\square$

Together the upper bound and [Theorem A.10](#) prove [Theorem 1.2](#). The proof also reveals which graphs are extremal: equality forces both inequalities in the chain to be tight.

Theorem A.11 (Characterisation of equality). *Let G be simple with $\lambda_G^{\max} \leq m-1$ and $|E(G)| = \left\lfloor \frac{m(n-1)}{2} \right\rfloor$, and let T be a Gomory–Hu tree of G . If $m(n-1)$ is even, then every tree edge has weight exactly $m-1$, every tree edge is also an edge of G , and every edge of G joins vertices at tree distance at most 2. If $m(n-1)$ is odd, there is exactly one unit of slack, located in exactly one of three places: one tree edge has weight $m-2$, or one tree edge is not an edge of G , or one edge of G joins vertices at tree distance 3; everything else is tight.*

Proof. Write the chain of the upper-bound proof as three nonnegative slacks: $S_1 = (m-1)(n-1) - \sum_t c(t)$, counting tree weights below $m-1$; $S_2 = \sum_{e: \text{dist}_T(e) \geq 2} (\text{dist}_T(e) - 2)$, counting edges beyond tree distance two; and $S_3 = (n-1) - |E_1|$, counting tree edges that support no edge of G . Retracing the proof, $\sum_e \text{dist}_T(e) = 2|E| - |E_1| + S_2$ and $\sum_e \text{dist}_T(e) = \sum_t c(t) = (m-1)(n-1) - S_1$, so altogether $2|E| = m(n-1) - (S_1 + S_2 + S_3)$. If $m(n-1)$ is even, equality $|E| = m(n-1)/2$ forces $S_1 = S_2 = S_3 = 0$. If $m(n-1)$ is odd the floor absorbs exactly one unit, so $S_1 + S_2 + S_3 = 1$ and exactly one of the three slacks equals one. $\square$

For $m = 2$ the characterisation says the extremal graphs are exactly the spanning trees: every tree edge must carry weight one and lie in G , so G contains its own Gomory–Hu tree and has only $n-1$ edges to spend. For $m = 4$ and $n \equiv 1 \pmod{3}$ it is consistent with the trees of K_4 blocks glued at shared cut vertices — the case of [Construction A.7](#) that the search of [Chapter 3](#) rediscovers.

A.3 The directed arc value at $m = 2$

Three short lemmas feed the induction. The first shows a hub forces a linear bound for every m , and is the proof behind the hub branch of [Conjecture 1.13](#).

Lemma A.12 (Two-hop lemma). *Let D be a simple digraph with $\lambda_D^{\max} \leq m-1$ containing a vertex r with $d^+(r) = n-1$. Then every $v \neq r$ has at most $m-2$ in-arcs from $V \setminus \{r, v\}$.*

Proof. If $u_1, \dots, u_{m-1}$ were distinct in-neighbours of v other than r , the paths $r \rightarrow v$ and $r \rightarrow u_i \rightarrow v$ for $i = 1, \dots, m-1$ would be m arc-disjoint routes, since the u_i are distinct and $d^+(r) = n-1$ supplies every starting arc. That contradicts $\lambda_D(r, v) \leq m-1$. $\square$

Theorem A.13 (Hub upper bound). *If a simple digraph D with $\lambda_D^{\max} \leq m - 1$ has a vertex r with $d^+(r) = n - 1$, then $|A(D)| \leq m(n - 1)$.*

Proof. Split the arcs into those leaving r (exactly $n - 1$), those entering r (at most $n - 1$), and the internal arcs avoiding r . By [Lemma A.12](#) each of the $n - 1$ other vertices receives at most $m - 2$ internal arcs, so there are at most $(m - 2)(n - 1)$ internal arcs, and the total is at most $m(n - 1)$. $\square$

Lemma A.14 (No transitive triangle). *A simple digraph with $\lambda^{\max} \leq 1$ contains no three vertices x, y, z with all of $(x, y), (y, z), (x, z)$ present, since $x \rightarrow z$ and $x \rightarrow y \rightarrow z$ would be two arc-disjoint routes.*

Lemma A.15 (Deletion monotonicity). *For any digraph D and vertex v_0 , $\lambda^{\max}(D - v_0) \leq \lambda^{\max}(D)$, since arc-disjoint paths in $D - v_0$ are arc-disjoint paths in D .*

Proof of Theorem 1.9. Write $M(n) = \max(2(n - 1), \lfloor n^2/4 \rfloor)$. A direct computation shows $2(n - 1) \geq \lfloor n^2/4 \rfloor$ exactly for $n \leq 7$, with equality at $n = 7$, so $M(n) = 2(n - 1)$ for $n \leq 7$ and $M(n) = \lfloor n^2/4 \rfloor$ for $n \geq 7$. The lower bound $\ell_2^{\text{dir}}(n) \geq M(n)$ is witnessed by the directed hub construction ([Construction 1.8](#) at $m = 2$, the bidirected star) and the one-directional bipartite construction ([Construction 1.7](#)).

For the upper bound we show by strong induction on n that $\lambda_D^{\max} \leq 1$ forces $|A(D)| \leq M(n)$.

Base cases $2 \leq n \leq 7$. For $n = 2$ there are at most 2 arcs and $M(2) = 2$. For $n = 3$, suppose $|A(D)| \geq 5$: then D misses at most one arc of the complete bidirected triangle, and whichever arc is missing, three of the remaining arcs form a transitive triangle, contradicting [Lemma A.14](#); hence $|A(D)| \leq 4 = M(3)$. For $4 \leq n \leq 7$ the pruned exhaustive search of [Chapter 2](#) walks every simple digraph with $\lambda^{\max} \leq 1$ and reports the maxima 6, 8, 10, 12, matching $M(n)$; the transcript is reproduced below, and the cut-counting certifier independently confirms the values within its reach through the $m = 2$ reduction of [Theorem A.24](#).

Inductive step $n \geq 8$. Let D be a simple digraph on n vertices with $\lambda_D^{\max} \leq 1$ and put $a := |A(D)|$. Choose v_0 minimising the total degree $d(v) = d^+(v) + d^-(v)$. Since $\sum_v d(v) = 2a$, averaging gives $d(v_0) \leq 2a/n$. By [Lemma A.15](#) the digraph $D - v_0$ on $n - 1 \geq 7$ vertices satisfies the induction hypothesis, so $a - d(v_0) \leq M(n - 1) = \lfloor \frac{(n-1)^2}{4} \rfloor$. Combining,

$$a \leq \frac{2a}{n} + \left\lfloor \frac{(n-1)^2}{4} \right\rfloor \implies a \leq \frac{n}{n-2} \left\lfloor \frac{(n-1)^2}{4} \right\rfloor.$$

It remains to check that the right-hand side is at most $\lfloor n^2/4 \rfloor$. For even $n = 2k$ ($k \geq 4$): $\lfloor (2k-1)^2/4 \rfloor = k(k-1)$, and $\frac{2k}{2k-2} \cdot k(k-1) = k^2 = \lfloor n^2/4 \rfloor$ exactly. For odd $n = 2k+1$ ($k \geq 4$): $\lfloor (2k)^2/4 \rfloor = k^2$, and the polynomial division $(2k+1)k^2 = (2k-1)(k^2+k) + k$ gives

$$\frac{n}{n-2} \left\lfloor \frac{(n-1)^2}{4} \right\rfloor = \frac{(2k+1)k^2}{2k-1} = k(k+1) + \frac{k}{2k-1},$$

and since $0 < \frac{k}{2k-1} < 1$ and a is an integer, $a \leq k(k+1) = \lfloor n^2/4 \rfloor$. In both parities $a \leq M(n)$, completing the induction. $\square$

Two remarks on the shape of this proof. First, the minimum-degree deletion closes the step on its own; no separate hub case is needed, because the averaging bound holds whether or not D contains a hub, although [Theorem A.13](#) shows directly that a hub digraph never exceeds the linear branch. Second, for even n the bound is exactly tight against the one-directional bipartite construction, which is $\frac{n}{2}$ -regular in total degree, so every step of the chain is achieved by the conjectured extremiser; for odd n the slack $\frac{k}{2k-1} < 1$ is absorbed by integrality. The proof is long not because it is deep but because the two regimes of $M(n)$ must be carried through the arithmetic separately, and that bookkeeping is precisely what [Chapter 2](#) hands to the machine at the base cases.

Proof of [Theorem 1.10](#). By Whitney's inequality $\kappa^{\max} \leq \lambda^{\max}$, every digraph feasible for the vertex problem is feasible for the arc problem, so $k_2^{\text{dir}}(n) \leq \ell_2^{\text{dir}}(n)$. Conversely the hub and one-directional bipartite constructions have $\kappa^{\max} = 1$, so they are feasible for the vertex problem and give the matching lower bound. Hence $k_2^{\text{dir}}(n) = \ell_2^{\text{dir}}(n) = \max(2(n-1), \lfloor n^2/4 \rfloor)$. $\square$

A.4 Structural propositions on the directed frontier

These are the proved ingredients of the reduction discussed in [Chapter 4](#). The upper bound for $m \geq 3$ is conditional only on the backward-arc lemma stated there.

Proposition A.16 (Mutually unreachable out-neighbourhoods). *Let D be a simple digraph with $\lambda_D^{\max} \leq 1$. For any vertex u and distinct out-neighbours v, w of u , there is no directed path from v to w in $D - u$.*

Proof. If P were a directed v - w path avoiding u , then $u \rightarrow w$ and $u \rightarrow v \xrightarrow{P} w$ would be two arc-disjoint u -to- w routes: the first uses only the arc (u, w) , the second starts with (u, v) and continues along P , none of whose arcs touch u . This contradicts $\lambda_D(u, w) \leq 1$. The restriction to $D - u$ is not cosmetic, and [Figure A.3](#) draws both the argument and the reason for the restriction: in the digraph with arcs (u, v) , (v, u) , (u, w) every local connectivity is 1, yet $v \rightarrow u \rightarrow w$ reaches w from v through u . $\square$

Proposition A.17 (Disjoint second out-neighbourhoods). *Let D be a simple digraph with $\lambda_D^{\max} \leq 1$, let $u \in V$, and let $v_1 \neq v_2$ be out-neighbours of u . Then $(N^+(v_1) \setminus \{u\}) \cap (N^+(v_2) \setminus \{u\}) = \emptyset$.*

Proof. A common out-neighbour $w \neq u$ would give the two arc-disjoint routes $u \rightarrow v_1 \rightarrow w$ and $u \rightarrow v_2 \rightarrow w$, contradicting $\lambda_D(u, w) \leq 1$. $\square$




Figure A.3. Why Proposition A.16 must exclude paths through u . (a) For out-neighbours v, w of u in a digraph with $\lambda^{\max} \leq 1$, no $v \rightsquigarrow w$ path can avoid u : such a path would combine with the lone arc $u \rightarrow w$ into two arc-disjoint routes. (b) The restriction to $D - u$ is essential. In the three-vertex digraph $\{(u, v), (v, u), (u, w)\}$ all connectivities equal 1, yet v reaches w via $v \rightarrow u \rightarrow w$ — a route that reuses the vertex u , which the proposition allows.

Proposition A.18 (Two-hop budget on bipartite partitions). *Let D be a simple digraph with $\lambda_D^{\max} \leq m - 1$. If V admits a partition $A \cup B$ with every arc from A to B present, then every $b \in B$ has in-degree at most $m - 2$ from inside B , and consequently the number of arcs inside B is at most $(m - 2)|B|$.*

Proof. Fix $a \in A$ and $b \in B$, and let $b'_1, \dots, b'_d$ be the in-neighbours of b inside B , where $d = \deg_B^-(b)$. The direct arc $a \rightarrow b$ together with the two-step detours $a \rightarrow b'_i \rightarrow b$ are $1 + d$ arc-disjoint a -to- b routes, all present because $A \rightarrow B$ is complete. Feasibility forces $1 + d \leq m - 1$, so $d \leq m - 2$, and summing over B bounds the internal arcs by $(m - 2)|B|$. $\square$

Combining Proposition A.18 with the best partition gives, for any extremal digraph that has a complete $A \rightarrow B$ partition and no arc from B back to A , the bound $|A(D)| \leq |B|(n - |B| + m - 2)$, since the $|A||B|$ arcs across the partition and the at most $(m - 2)|B|$ arcs inside B are all there are: under the no-back-arc hypothesis a route between two vertices of B never leaves B , so the two-hop budget applies to B 's own arcs. Maximised at $|B| = \lceil (n + m - 2)/2 \rceil$, this is $\lfloor (n + m - 2)^2/4 \rfloor$, the second branch of Conjecture 1.13, uniformly in m , and the remaining gap is exactly the backward-arc hypothesis.

A.5 The directed multigraph problem at large n

The body proves $L_m^{\text{dir}}(n) = 2(n - 1)(m - 1)$ for $n \leq 6$ (Theorem 2.1). This section records what is known beyond, where the one-directional bipartite construction takes over. Throughout, the scaling reduction of Chapter 2 is used in the following form.

Lemma A.19 (Scaling reduction). *Let D be a directed multigraph with $\mu(u, v) \leq m - 1$ on every ordered pair and $\lambda^{\max}(D) \leq m - 1$. Then $w := \mu/(m - 1)$ takes values in $[0, 1]$, satisfies $\text{maxflow}(s, t; w) \leq 1$ for all $s \neq t$, and $|A(D)| = (m - 1) \sum_{u \neq v} w(u, v)$.*

Proof. Dividing all capacities by $m - 1$ divides every flow value by $m - 1$, and by Menger the

maximum flow under μ is $\lambda_D(s, t) \leq m - 1$. □

Lemma A.20 (Arithmetic identity). *For all $n \geq 2$, $\left\lfloor \frac{(n-1)^2}{4} \right\rfloor + \lfloor n/2 \rfloor = \left\lfloor \frac{n^2}{4} \right\rfloor$.*

Proof. For $n = 2k$ the left side is $k(k-1) + k = k^2$; for $n = 2k+1$ it is $k^2 + k = k(k+1)$. Both match the right side. □

Proposition A.21 (Bipartite upper bound for even $n \geq 8$, conditional). *Let $n = 2k \geq 8$ be even. Suppose every fractional weighting w' on $n-1$ vertices with values in $[0, 1]$ and all pairwise maximum flows at most one has total weight at most $\left\lfloor \frac{(n-1)^2}{4} \right\rfloor$. Then every directed multigraph D on n vertices with $\lambda^{\max}(D) \leq m-1$ satisfies $|A(D)| \leq (m-1) \left\lfloor \frac{n^2}{4} \right\rfloor$.*

Proof. By [Lemma A.19](#) it suffices to bound $M := \sum_{u \neq v} w(u, v)$ by $\lfloor n^2/4 \rfloor$ for the scaled weighting w . Pick v_0 minimising the weighted total degree $d(v_0) = \sum_u (w(u, v_0) + w(v_0, u))$; averaging gives $d(v_0) \leq 2M/n$. Deleting v_0 can only lower maximum flows, so the hypothesis applies to the restricted weighting and $M - d(v_0) \leq \lfloor (n-1)^2/4 \rfloor = k(k-1)$. Combining, $M(1 - \frac{2}{n}) \leq k(k-1)$, hence $M \leq \frac{k}{k-1} \cdot k(k-1) = k^2 = \lfloor n^2/4 \rfloor$. □

The small cases $n \in \{4, 6\}$ genuinely do not satisfy the conclusion, and must not: there $2(n-1) > \lfloor n^2/4 \rfloor$ and the double star beats the bipartite weighting, which is why the certified values of [Theorem 2.1](#) sit on the linear branch. For odd n the same deletion gives $M \leq k(k+1) + \frac{k}{2k-1}$, an overshoot of less than one that integrality no longer absorbs, because fractional weights need not have integer totals. The induction would close if some vertex always had weighted degree at most $(m-1)k$ before scaling, which is the missing minimum-degree statement.

Conjecture A.22 (Minimum-degree lemma for odd n). *In any directed multigraph D on $n = 2k+1$ vertices ($k \geq 3$) with $\lambda^{\max}(D) \leq m-1$ and $m \geq 3$, some vertex has total degree $d(v) \leq (m-1)k$, counting arcs with multiplicity.*

At $m = 2$ the statement is a proposition rather than a conjecture, through a reduction that also settles the whole multigraph problem at $m = 2$.

Proposition A.23 (Parallel-arc elimination at $m = 2$). *A directed multigraph with $\lambda^{\max} \leq 1$ has no parallel arcs, since two parallel arcs are already two arc-disjoint routes. Hence it is a simple digraph.*

Theorem A.24 (Directed multigraphs at $m = 2$). *For all $n \geq 2$, $L_2^{\text{dir}}(n) = \ell_2^{\text{dir}}(n) = \max(2(n-1), \lfloor \frac{n^2}{4} \rfloor)$.*

Proof. Immediate from [Proposition A.23](#) and [Theorem 1.9](#). □

Proposition A.25 (Minimum degree at $m = 2$, odd n). *In any directed multigraph on $n = 2k+1 \geq 7$ vertices with $\lambda^{\max} \leq 1$, some vertex has $d(v) \leq k$.*

Proof. By [Proposition A.23](#) and [theorem 1.9](#), $|A(D)| \leq \max(2(n-1), \lfloor n^2/4 \rfloor) = \lfloor n^2/4 \rfloor = k(k+1)$, the maximum being the quadratic branch because $n \geq 7$. The average total degree is $\frac{2k(k+1)}{2k+1} < k+1$, and some integer degree is at most k . The hypothesis $n \geq 7$ is needed: on the linear branch the statement can fail, as the bidirected star on three vertices has minimum total degree 2 > 1. $\square$

For $m \geq 3$ the two extreme regimes of [Conjecture A.22](#) are settled. In the *purely saturated* regime, where every multiplicity is 0 or $m-1$, the scaled weighting is the indicator of a simple digraph with $\lambda^{\max} \leq 1$, its degrees are integers, and the argument of [Proposition A.25](#) applies verbatim after multiplying back by $m-1$. In the *all-unsaturated* simple regime at $n=7$, $m=3$, the program certifies by a cut-counting infeasibility that no simple digraph satisfies all two-hop inequalities with every total degree at least 7. The open case is the *mixed* regime, where some multiplicity lies strictly between 0 and $m-1$: there the scaled degrees are genuinely fractional, the averaging bound $\frac{2k(k+1)}{2k+1}$ lands strictly between k and $k+1$, and no integrality rescues it. This is the single identified gap that separates the certified values of [Theorem 2.1](#) from the full conjecture

$$L_m^{\text{dir}}(n) = (m-1) \max(2(n-1), \lfloor \frac{n^2}{4} \rfloor),$$

whose lower bound is unconditional: the double star realises $2(n-1)(m-1)$ and the one-directional bipartite multigraph, every arc at multiplicity $m-1$, realises $(m-1) \lfloor n^2/4 \rfloor$ with $\lambda^{\max} = m-1$.

Lemma A.26 (Attachment lemma). *Let $m \geq 2$ and let D_0 be the one-directional complete bipartite multigraph with parts A and B , $|A| = p \geq 2$, $|B| = q \geq 2$, and $\mu(a, b) = m-1$ for every $a \in A$, $b \in B$, with no other arcs. Let D be obtained from D_0 by adding one new vertex v together with any multiset of arcs incident to v (multiplicities at most $m-1$). If $\lambda_D^{\max} \leq m-1$, then $d(v) \leq (m-1) \max(p, q)$.*

[Figure A.4](#) sets up the proof: the new vertex v , the four classes its incident arcs fall into, and one of the route patterns that the hypothesis forbids.

Proof. Write $i_A = \sum_a \mu(a, v)$, $i_B = \sum_b \mu(b, v)$, $o_A = \sum_a \mu(v, a)$, $o_B = \sum_b \mu(v, b)$, so that $d(v) = i_A + i_B + o_A + o_B$. Three route patterns are forbidden outright, because each adds an m -th arc-disjoint route on top of the $m-1$ parallel arcs of a bipartite pair:

- ☆ $\mu(a, v) \geq 1$ and $\mu(v, b) \geq 1$: the path $a \rightarrow v \rightarrow b$ together with the $m-1$ arcs $a \rightarrow b$ gives $\lambda(a, b) \geq m$.
- ☆ $\mu(a, v) \geq 1$ and $\mu(v, a') \geq 1$ with $a' \neq a$: the path $a \rightarrow v \rightarrow a' \rightarrow b$ (any $b \in B$) gives $\lambda(a, b) \geq m$.
- ☆ $\mu(b, v) \geq 1$ and $\mu(v, b') \geq 1$ with $b' \neq b$: the path $a \rightarrow b \rightarrow v \rightarrow b'$ (any $a \in A$) gives $\lambda(a, b') \geq m$.




Figure A.4. The attachment lemma (Lemma A.26): the new vertex v glued onto $(m-1) \cdot B_{p,q}$. Each $a \in A$ sends $m-1$ parallel arcs to each $b \in B$ (the bent bundle). The arcs at v split into four classes by direction and side: i_A (into v from A), o_A (out of v to A), i_B , o_B . The red dashed route $a \rightarrow v \rightarrow b$ is forbidden: laid on top of the $m-1$ existing $a \rightarrow b$ arcs it makes an m -th arc-disjoint $a-b$ route, so $\lambda(a, b) \geq m$. Ruling out this and two sibling patterns caps $d(v) \leq (m-1) \max(p, q)$.

Case I: $i_A \geq 1$. By (1), $o_B = 0$. If $o_A \geq 1$, then by (2) all in-arcs from A and all out-arcs to A involve one single vertex a^* , and the flow estimate $\lambda(a^*, v) \geq \mu(a^*, v) + \sum_b \min(\mu(a^*, b), \mu(b, v)) = \mu(a^*, v) + i_B \leq m-1$ gives $d(v) = \mu(a^*, v) + i_B + \mu(v, a^*) \leq 2(m-1)$. If $o_A = 0$, then v is a pure sink and for every $a \in A$ the same estimate reads $\mu(a, v) + i_B \leq m-1$; summing over A , $d(v) = i_A + i_B \leq p(m-1) - (p-1)i_B \leq p(m-1)$. **Case II:** $i_A = 0$, $o_B \geq 1$. Symmetric to Case I under arc reversal, with (3) in place of (2): either $d(v) \leq 2(m-1)$, or v is a pure source and $\lambda(v, b) \geq \mu(v, b) + o_A \leq m-1$ for every b gives $d(v) \leq q(m-1) - (q-1)o_A \leq q(m-1)$. **Case III:** $i_A = o_B = 0$. Then $\lambda(a, v) \geq i_B$ and $\lambda(v, b) \geq o_A$ force $d(v) = i_B + o_A \leq 2(m-1)$. Since $p, q \geq 2$, all cases give $d(v) \leq (m-1) \max(p, q)$. $\square$

Corollary A.27 (Attachment equality). *In the setting of Lemma A.26 with $\max(p, q) \geq 3$, suppose $d(v) = (m-1) \max(p, q)$. If $q > p$, then v is a pure source with $\mu(v, b) = m-1$ for every $b \in B$, so $D = (m-1) \cdot B_{p+1,q}$ with v joining the source side; if $p > q$, symmetrically a pure sink at full multiplicity; if $p = q$, one of the two.*

Proof. Rerun the cases of Lemma A.26. The mixed cases cap $d(v)$ at $2(m-1) < (m-1) \max(p, q)$. The pure-sink case gives $d(v) \leq p(m-1) - (p-1)i_B$, which reaches $(m-1) \max(p, q)$ only when $p \geq q$, $i_B = 0$ and every $\mu(a, v) = m-1$; the pure-source case symmetrically forces $o_A = 0$ and every $\mu(v, b) = m-1$ when $q \geq p$. Feasibility of the resulting digraph is the bipartite construction itself. $\square$

Theorem A.28 (Conditional odd step, $m \in \{3, 4\}$). *Fix $k \geq 4$ and $m \in \{3, 4\}$, and assume (i) $L_m^{\text{dir}}(2k) = (m-1)k^2$, and (ii) every directed multigraph attaining it is the one-directional complete bipartite multigraph $(m-1) \cdot B_{k,k}$. Then $L_m^{\text{dir}}(2k+1) = (m-1)k(k+1)$.*

Proof. The lower bound is $(m-1) \cdot B_{k,k+1}$. For the upper bound let D on $n = 2k+1$ vertices have $\lambda^{\max} \leq m-1$ and $|A(D)| = (m-1)k(k+1) + j$ with $j \geq 1$. Deleting any vertex v leaves a

feasible multigraph on $2k$ vertices, so by (i) $d(v) \geq |A(D)| - (m-1)k^2 = (m-1)k + j$ for every v . The average degree is

$$\frac{2|A(D)|}{2k+1} = (m-1)k + j + \frac{(m-1)k - (2k-1)j}{2k+1},$$

by direct computation. If $(m-1)k - (2k-1)j < 0$ the average lies below the minimum, a contradiction. Otherwise the correction term is nonnegative and, because $m \leq 4$ and $j \geq 1$, strictly below one: $(m-1)k - (2k-1)j \leq (m-3)k + 1 \leq k + 1 < 2k + 1$. The average therefore lies in $[(m-1)k + j, (m-1)k + j + 1)$ while every degree is at least $(m-1)k + j$, so some vertex v has $d(v) = (m-1)k + j$ exactly, and $D - v$ has exactly $(m-1)k^2$ arcs. By (ii), $D - v = (m-1) \cdot B_{k,k}$, and [Lemma A.26](#) caps $d(v) \leq (m-1)k < (m-1)k + j$, the final contradiction. Hence $j \leq 0$. $\square$

Remark A.29 (The $m = 3$ chain, sharpened). [Theorem A.28](#) removes [Conjecture A.22](#) from the critical path: the odd case no longer needs a minimum-degree statement, only the extremal characterisation one level down, and for $m = 3$ that characterisation propagates along the same induction in both parities. *Odd levels* ($n = 2k + 1 \geq 9$): an extremiser has $|A| = 2k(k+1)$ arcs and every degree at least $2k$ (deletion bound); all degrees $\geq 2k + 1$ is impossible since $(2k+1)^2 > 4k(k+1) = \sum_v d(v)$, so some $d(v) = 2k$ exactly, $D - v$ is the even extremiser $2 \cdot B_{k,k}$, and [Corollary A.27](#) (the $p = q$ case) forces v to rejoin as a full-multiplicity pure source or sink: $D = 2 \cdot B_{k+1,k}$ or $2 \cdot B_{k,k+1}$. *Even levels* ($n = 2k \geq 10$): equality in the averaging step forces $2k$ -regularity, hence $D - v$ extremal at $2k - 1$ for every v ; odd uniqueness one level down leaves $2 \cdot B_{k,k-1}$ or $2 \cdot B_{k-1,k}$, and [Corollary A.27](#) with $d(v) = 2k = 2 \max(k-1, k)$, $q > p$, rebuilds exactly $2 \cdot B_{k,k}$. *The seam at $n = 8$* : the same regularity argument runs from $n = 7$, where the branches tie at 24 arcs and several extremisers coexist. On the linear branch the extremisers are richer than the double star alone: every *bidirected spanning tree at full multiplicity $m-1$* is feasible with $2(n-1)(m-1)$ arcs (the only arcs crossing the bipartition under a tree edge are that edge's own $m-1$ copies, so every local connectivity is exactly $m-1$), and exhaustive enumeration (`enumerate_extremal_directed_multigraphs`) confirms that at $n = 4, 5$, $m = 3$ these doubled trees are the *only* extremisers (2 and 3 of them, matching the tree shapes). A deleted vertex of the 8-regular D leaves a 24-arc extremiser with maximum degree at most 8: among doubled trees only the doubled path qualifies (tree degree ≤ 2), the double star (hub degree 24) being excluded. If $D - v$ were the doubled path P_7 , the degree count would force v to attach with weight 4 to each path end, i.e. bidirected at full multiplicity, making D the doubled bidirected cycle C_8 — infeasible, since the two doubled directions around the cycle give $\lambda = 4 > 2$ between any pair. [Figure A.5](#) draws this collapse of the doubled path into the infeasible doubled cycle. So every deletion is bipartite-type, the attachment equality applies, and $D = 2 \cdot B_{4,4}$ again. The whole $m = 3$ problem therefore rests on two finite integral statements about multigraphs with multiplicities in $\{0, 1, 2\}$: (a) $L_3^{\text{dir}}(7) = 24$, equivalently $M^*(7) = 12$; and (b) every 24-arc feasible multigraph on 7 vertices with maximum degree at most 8 is $2 \cdot B_{3,4}$, $2 \cdot B_{4,3}$, or the doubled bidirected path P_7 . Both targets sit at sizes the certifier and the enumerator can realistically reach with the strengthened relaxation

now wired into `certify_directed_multigraph` (`use_deletion_cuts`: deletion bounds from the certified $M^*(5)$, $M^*(6)$ and the degree-pair inequality $d^+(s) + d^-(t) - w(s, t) \leq n - 1$, all proved valid). For $m \geq 4$ one hole remains: at an odd extremal level the counting no longer forces a vertex of degree $(m - 1)k$ (all degrees $\geq (m - 1)k + 1$ is arithmetically consistent once $k(m - 3) \geq 1$), so the uniqueness statement does not yet propagate through odd levels, although the *value* at odd levels follows from even uniqueness whenever that is available. For $m \geq 5$ the same escape climbs one level: a hypothetical extremiser carrying $j \geq 1$ surplus arcs could have every degree strictly above the deletion bound (all degrees $\geq (m - 1)k + j + 1$ is consistent exactly when $(2k - 1)j \leq (m - 1)k - 2k - 1$, which admits $j = 1$ once $m \geq 5$), so no vertex deletion is forced to land on an extremal multigraph and even the *value* step stalls. This is why [Theorem A.28](#) is stated for $m \in \{3, 4\}$.




Figure A.5. The seam argument of [Remark A.29](#). A degree count forces a deleted vertex of the 8-regular extremiser D to leave the doubled bidirected path P_7 (left, each path edge a pair of opposing arcs). To reach degree 8 the new vertex v must attach with full multiplicity to the two ends 1 and 7 (red), bidirected. That closes the path into the doubled bidirected cycle C_8 (right), which is *infeasible*: the two directions around the cycle give two arc-disjoint routes each way, so $\lambda = 4 > 2$ between any pair. The contradiction rules out the path branch, forcing every deletion to be bipartite-type and hence $D = 2 \cdot B_{4,4}$.

A.6 The hypergraph bounds

The body's hypergraph claims rest on two theorems: a Menger theorem for Berge routes, which also certifies the helper-point checker of [Chapter 2](#), and a hypergraph Gomory–Hu theorem from the recent literature.

Theorem A.30 (Menger for hypergraphs). *For a hypergraph $\mathcal{H}$ and vertices u, v , the maximum number of hyperedge-disjoint Berge u – v paths equals the minimum size of a set C of hyperedges whose removal separates u from v .*

Proof. Build the flow network of [Figure 2.4](#): for each hyperedge e a helper arc $e^- \rightarrow e^+$ of capacity one, and arcs $x \rightarrow e^-$ and $e^+ \rightarrow x$ of unbounded capacity for each $x \in e$. A Berge path $u, e_1, v_1, \dots, e_k, v$ becomes the directed walk $u \rightarrow e_1^- \rightarrow e_1^+ \rightarrow v_1 \rightarrow \dots \rightarrow v$, and since the e_i are distinct, hyperedge-disjoint Berge paths use distinct helper arcs, so they translate to arc-disjoint flow paths. Conversely an integral maximum flow decomposes into paths, each of

which, suppressing the helper nodes, is a Berge path, and no two share a hyperedge because each helper arc has capacity one. Finally, any finite cut of the network consists of helper arcs only, and a set of helper arcs separates u from v exactly when the corresponding hyperedges do. Max-flow min-cut now gives the claim. $\square$

Theorem A.31 (Hypergraph Gomory–Hu; Bérczi et al. [2]). *Every r -uniform hypergraph $\mathcal{H}$ on vertex set V has a weighted tree T^* on V such that $\lambda_{\mathcal{H}}(u, v)$ equals the minimum weight on the u – v tree path, and each tree edge t induces a vertex partition $(S_t, V \setminus S_t)$ with $|\delta_{\mathcal{H}}(S_t)| = c^*(t)$, where $\delta_{\mathcal{H}}(S)$ is the set of hyperedges meeting both sides.*

Proof of Proposition 1.14. Upper bound. Let $\mathcal{H}$ be r -uniform with $\lambda_{\mathcal{H}}^{\max} \leq m - 1$ and let T^* be its Gomory–Hu tree. Every tree weight is a local connectivity, so $\sum_t c^*(t) \leq (m - 1)(n - 1)$. For a hyperedge e let $\text{ST}(e)$ be the smallest subtree of T^* containing the r vertices of e ; it has at least $r - 1$ edges, and for each of its edges t the hyperedge e meets both sides of the induced partition, so $e \in \delta_{\mathcal{H}}(S_t)$. Summing over hyperedges and exchanging the order of summation,

$$(r - 1) |E(\mathcal{H})| \leq \sum_e |\{t : e \in \delta_{\mathcal{H}}(S_t)\}| = \sum_t |\delta_{\mathcal{H}}(S_t)| = \sum_t c^*(t) \leq (m - 1)(n - 1),$$

and dividing by $r - 1$ with integrality gives $|E(\mathcal{H})| \leq \left\lfloor \frac{(m-1)(n-1)}{r-1} \right\rfloor$. *Lower bound.* When $(r - 1) \mid (n - 1)$, take a hub h , partition the other $n - 1$ vertices into groups of size $r - 1$, form the star hypertree of hyperedges $\{h\} \cup G_i$, and give each hyperedge multiplicity $m - 1$. Every non-hub vertex lies in exactly $m - 1$ hyperedges, which form a cut isolating it, so by Theorem A.30 every pair has Berge connectivity at most $m - 1$, while the count is $\frac{(m-1)(n-1)}{r-1}$. For general n under the hypothesis $m - 1 \leq \binom{n-2}{r-2}$, Theorem A.32 below attains the floor exactly, without even needing repeated hyperedges. $\square$

The multiplicities in the lower bound can in fact be avoided: for $r \geq 3$, simple r -uniform hypergraphs already attain the same bound, in sharp contrast to the graph case $r = 2$, where simplicity costs the factor that separates Mader’s $\lfloor m(n - 1)/2 \rfloor$ from the multigraph value $(m - 1)(n - 1)$.

Theorem A.32 (Simplicity is free for $r \geq 3$). *Let $r \geq 3$, $n \geq r$, and $2 \leq m$ with $m - 1 \leq \binom{n-2}{r-2}$. Then the maximum number of hyperedges in a simple r -uniform hypergraph on n vertices with Berge edge-connectivity at most $m - 1$ is exactly $\left\lfloor \frac{(m-1)(n-1)}{r-1} \right\rfloor$.*

Proof. The upper bound is Proposition 1.14, since a simple hypergraph is a multihypergraph. For the lower bound, fix a hub h and let $V' = V \setminus \{h\}$. By Lemma A.33 applied with rank $r - 1$ and degree ceiling $d = m - 1$ on the $n - 1$ vertices of V' , there is an $(r - 1)$ -uniform simple hypergraph $\mathcal{G}^*$ on V' with maximum degree at most $m - 1$ and exactly $\left\lfloor \frac{(m-1)(n-1)}{r-1} \right\rfloor$ hyperedges. Adding h to each hyperedge produces distinct r -sets, so the result $\mathcal{H}$ is simple, and every non-hub vertex w lies in $\deg_{\mathcal{G}^*}(w) \leq m - 1$ hyperedges, which form a cut isolating w . By Theorem A.30, $\lambda_{\mathcal{H}}^{\max} \leq m - 1$. $\square$

Lemma A.33 (Sparse uniform hypergraphs exist). *Let $r \geq 2$, $n \geq r$, and $0 \leq d \leq \binom{n-1}{r-1}$. There exists an r -uniform simple hypergraph on n vertices with maximum degree at most d and exactly $\lfloor \frac{dn}{r} \rfloor$ hyperedges.*

Proof. Put $e := \lfloor \frac{dn}{r} \rfloor \leq \frac{n}{r} \binom{n-1}{r-1} = \binom{n}{r}$. By Baranyai's theorem on equitable partitions of the complete r -uniform hypergraph [1], the $\binom{n}{r}$ possible r -sets split into two classes of sizes e and $\binom{n}{r} - e$ such that within each class every vertex is covered either $\lfloor re/n \rfloor$ or $\lceil re/n \rceil$ times. The class of size e has maximum degree at most $\lceil re/n \rceil \leq \lceil d \rceil = d$. $\square$

A.7 The hypergraph vertex problem

The whole vertex problem translates into a graph problem through one observation, the one the rest of this section rests on and which Figure A.6 draws in full. Write $I(\mathcal{H})$ for the bipartite *incidence graph* of a hypergraph $\mathcal{H}$: one node per vertex, one node per hyperedge, an edge when the vertex lies in the hyperedge. A Berge path with distinct hyperedges is exactly a path of $I(\mathcal{H})$ between two vertex-class nodes, hyperedge-disjoint Berge paths use disjoint hyperedge-nodes, and internally vertex-disjoint ones use disjoint interior vertex-nodes. Hence

$$\kappa_{\mathcal{H}}(u, v) = \kappa_{I(\mathcal{H})}(u, v),$$


the maximum number of internally disjoint u - v paths in $I(\mathcal{H})$, and the hypergraph vertex problem asks for the maximum number of degree- r nodes on one side of a bipartite graph whose other side holds n nodes, subject to no two of those n nodes being joined by m internally disjoint paths. Figure A.6a traces a single Berge path through both pictures, and Figure A.6b shows the star hypertree of Theorem A.34, whose incidence graph is a tree.

Theorem A.34 (Hypergraph vertex value at $m = 2$). *For $r \geq 2$ and $n \geq 1$, the maximum number of hyperedges of an r -uniform multihypergraph on n vertices with $\kappa^{\max} \leq 1$ is*


$$k_2^{(r)}(n) = \left\lfloor \frac{n-1}{r-1} \right\rfloor,$$

attained by the star hypertree. In particular $k_2^{(r)}(n) = \ell_2^{(r)}(n)$: the vertex and edge problems agree at $m = 2$ for every uniformity, and repeated hyperedges never help.

Proof. We claim $\kappa_{\mathcal{H}}^{\max} \leq 1$ holds if and only if $I(\mathcal{H})$ is a forest. If $I(\mathcal{H})$ contains a cycle $v_1, e_1, v_2, e_2, \dots, v_\ell, e_\ell, v_1$ (it alternates classes, so $\ell \geq 2$, all v_i and e_i distinct), then v_1, e_1, v_2 and $v_1, e_\ell, v_\ell, e_{\ell-1}, \dots, e_2, v_2$ are two Berge v_1 - v_2 paths with disjoint hyperedge sets and disjoint interior vertex sets, so $\kappa(v_1, v_2) \geq 2$. (At $\ell = 2$ this is the repeated pair: two hyperedges sharing two vertices.) Conversely, two such paths between u and v concatenate to a cycle of $I(\mathcal{H})$. This proves the claim. A forest on $n + q$ nodes has at most $n + q - 1$ edges, while $I(\mathcal{H})$ has exactly rq edges, so $rq \leq n + q - 1$, i.e. $q \leq (n-1)/(r-1)$, and q is an integer. The star hypertree has $\lfloor (n-1)/(r-1) \rfloor$ hyperedges and a tree as incidence graph (each block vertex



(a) A 3-uniform hypergraph $\mathcal{H}$ (left) and its bipartite incidence graph $I(\mathcal{H})$ (right): circles for vertices, hollow squares for hyperedges, an edge when the vertex lies in the hyperedge. The Berge path $1 e_1 3 e_2 5$ is highlighted green in both pictures — in $\mathcal{H}$ it threads two hyperedges, in $I(\mathcal{H})$ it is an ordinary path $1 - e_1 - 3 - e_2 - 5$ alternating sides.



(b) A star hypertree: every hyperedge $\{h\} \cup G_i$ shares only the hub h (orange), and the other $r - 1 = 2$ vertices of each block sit in a single hyperedge. Its incidence graph is a tree, so $\kappa^{\max} \leq 1$.

Figure A.6. The translation the vertex problem rests on. Internally vertex-disjoint Berge paths in $\mathcal{H}$ correspond exactly to internally disjoint paths in $I(\mathcal{H})$, so $\kappa_{\mathcal{H}}(u, v) = \kappa_{I(\mathcal{H})}(u, v)$. **(top)** A hypergraph and its incidence graph with one Berge path drawn in both. **(bottom)** The star hypertree of [Theorem A.34](#), whose incidence graph is a tree.

lies in one hyperedge, every cycle would need two hyperedges sharing two vertices), so the floor is attained. $\square$

Proposition A.35 (General lower bound). *For $r \geq 3$, $m \geq 2$ and $m - 1 \leq \binom{n-2}{r-2}$,*

$$k_m^{(r)}(n) \geq \left\lfloor \frac{(m-1)(n-1)}{r-1} \right\rfloor,$$

attained by a simple hypergraph.

Proof. The simple construction of [Theorem A.32](#) has Berge edge-connectivity at most $m - 1$, and Whitney's inequality $\kappa \leq \lambda$ holds pairwise (an internally vertex-disjoint family is in particular hyperedge-disjoint under our definition), so the same hypergraph is feasible for the vertex problem. $\square$

The $m = 3$ case also closes completely. The engine is a rank bound for graphs in which one side of a partition must stay below local 3-connectivity; the hypergraph theorem then falls out of the incidence translation. Throughout, $\text{rank}(G) = |E(G)| - |V(G)| + 1$ denotes the cycle rank of a connected multigraph.

Lemma A.36 (Incidence rank lemma). *Let G be a connected multigraph with vertex set $X \cup Z$ such that (i) Z is independent, (ii) no edge incident to Z is parallel to another edge, (iii) every $z \in Z$ has degree at least 2, and (iv) $\kappa_G(x, x') \leq 2$ for all distinct $x, x' \in X$. Then $\text{rank}(G) \leq |X| - 1$.*

Proof. Induction on $|V| + |E|$. *Base* $|V| \leq 2$. A single vertex lies in X by (iii), and $0 \leq |X| - 1$. On two vertices, (i)–(iii) force both into X , parallel edges are internally disjoint paths, so (iv) caps the multiplicity at 2 and $\text{rank} \leq 1 = |X| - 1$.

Step 1: a cut vertex. Let $B_1, \dots, B_c$ ($c \geq 2$) be the blocks. A bridge block has rank $0 \leq |X \cap B| - 1$, since by (i) it has an endpoint in X . Every other block is 2-connected, hence has minimum degree 2, so it satisfies (i)–(iv) and the induction hypothesis. Summing, and writing $b(v)$ for the number of blocks containing a cut vertex v , the block-cut tree identity $\sum_{v \text{ cut}} (b(v) - 1) = c - 1$ gives

$$\text{rank}(G) = \sum_i \text{rank}(B_i) \leq \sum_i (|X \cap B_i| - 1) = |X| - 1 - \sum_{z \in Z \text{ cut}} (b(z) - 1) \leq |X| - 1.$$

Step 2: a Z -vertex of degree two. If G is 2-connected and some $z_0 \in Z$ has degree exactly 2, its two edges reach distinct vertices of X (by (i) and (ii)); suppress z_0 , replacing it by one new X – X edge. Rank, $|X|$, all local connectivities between remaining vertices, and the hypotheses are preserved, and the graph shrinks.

Step 3: an X -vertex of degree two. If G is 2-connected, every Z -degree is at least 3 (Step 2 exhausted), and some $x_0 \in X$ has degree 2, delete x_0 . The result is connected, its rank is exactly

$\text{rank}(G) - 1$, the Z -degrees stay at least 2, and $|X|$ drops by one (note $|X| \geq 2$: with $|X| \leq 1$, (i) and (ii) would force every z to have degree at most 1). Induction gives $\text{rank}(G) - 1 \leq |X| - 2$.

Step 4: the remaining case is impossible. Now G is 2-connected with $|V| \geq 3$ and every degree at least 3. First, G is simple: a parallel class lies inside X by (ii), and if $\mu(x, x') \geq 2$ then a third internally disjoint route exists — walk from x to any third vertex w inside $G - x'$, then from w to x' inside $G - x$; the concatenation visits x only first and x' only last, so it contains an $x-x'$ path with at least one interior vertex — giving $\kappa(x, x') \geq 3$, against (iv). If G is 3-connected, Menger puts every pair at $\kappa \geq 3$, so $|X| \leq 1$ and every z has degree at most 1 by (i), absurd. Otherwise decompose G into its triconnected components (Tutte [19]; Hopcroft–Tarjan [10]), whose shape and the leaf at which the argument bites are sketched in Figure A.7: a tree of pieces, each a cycle of length at least three, a bond, or a 3-connected simple graph on at least four vertices, glued along virtual edges, where a vertex avoiding all virtual edges of its piece lies in that piece alone and keeps its degree, and where the far side of any virtual edge $\{a, b\}$ supplies a real $a-b$ path internally avoiding the piece. Since G is neither a cycle (degrees) nor a bond (simple) nor 3-connected, the tree has at least two leaves; let L be one, with unique virtual edge $\{a, b\}$. If L is a cycle, a vertex of L off $\{a, b\}$ has degree 2 in G , absurd. If L is a bond, it holds at least two real parallel edges, absurd in a simple graph. If L is 3-connected, then any two of its vertices are joined by three internally disjoint paths inside L , at most one through the virtual edge, which the far-side path replaces: so $\kappa_G(u, v) \geq 3$ for all $u, v \in V(L)$, forcing $|X \cap V(L)| \leq 1$ by (iv). But $|V(L)| \geq 4$ leaves a Z -vertex $z^* \notin \{a, b\}$; its neighbours lie in $X \cap V(L)$ by (i), at most one vertex, reached by at most one edge since G is simple — degree at most 1, absurd. $\square$




Figure A.7. The triconnected (Tutte/SPQR) decomposition in Step 4 of Lemma A.36, schematic: G splits into a tree of pieces (blobs) glued along virtual edges, and the analysis works at a *leaf* L with its single virtual edge $\{a, b\}$ (dashed red). The far side of the tree supplies a *real* $a-b$ path (green) that internally avoids L . If L is 3-connected, any two of its vertices are joined by three internally disjoint paths inside L ; at most one uses the virtual edge, and the far-side path replaces it, so $\kappa_G \geq 3$ for every pair in L . With $|V(L)| \geq 4$ that forces a hyperedge-node $z^* \notin \{a, b\}$ of degree at most one — the contradiction that closes the lemma.

Theorem A.37 (Hypergraph vertex value at $m = 3$). *For $r \geq 2$, every r -uniform multihypergraph on n vertices with $\kappa^{\max} \leq 2$ has at most $\left\lfloor \frac{2(n-1)}{r-1} \right\rfloor$ hyperedges. For $r \geq 3$ with $2 \leq \binom{n-2}{r-2}$ the*

bound is attained by a simple hypergraph, so

$$k_3^{(r)}(n) = \left\lfloor \frac{2(n-1)}{r-1} \right\rfloor = \ell_3^{(r)}(n).$$

Proof. Apply [Lemma A.36](#) to each connected component of the incidence graph $I(\mathcal{H})$, with X the vertex class and Z the hyperedge class: Z is independent, incidences are simple even when hyperedges repeat, hyperedge-nodes have degree $r \geq 2$, components without an X -node are impossible, and $\kappa_I(u, v) = \kappa_{\mathcal{H}}(u, v) \leq 2$ for vertex pairs. Summing $\text{rank} \leq |X_c| - 1$ over the C components, $rq - (n + q) + C \leq n - C$, hence $q(r-1) \leq 2(n-1) + 2(1-C) \leq 2(n-1)$. The lower bound is [Proposition A.35](#) at $m = 3$. $\square$

Remark A.38 (Edges as gates at $r = 2$, and the boundary of the method). At $r = 2$ the theorem speaks of multigraphs under the hyperedge-as-gate convention, in which parallel edges are distinct one-step routes; it gives the cactus value $2(n-1)$, complementing the body's convention for graphs, where parallel adjacencies count once. The proof of [Lemma A.36](#) leans on the triconnected decomposition exactly where $\kappa \leq 2$ lives; for $m = 4$ the analogous question asks for a rank bound relative to $\kappa \leq 3$, where no comparably clean decomposition exists, and the graph case ($m = 5$, Sørensen–Thomassen) warns that the pattern $k_m^{(r)} = \ell_m^{(r)}$ must fail eventually. Exhaustive computation confirms $k_4^{(3)}(5) = 6 = \lfloor 3 \cdot 4/2 \rfloor$, so the agreement survives at least one step beyond the theorem; where it first breaks for $r \geq 3$ is open.

Proposition A.39 (Directed hypergraphs: first bounds). *Model a directed r -uniform hyperedge as a tail vertex together with $r-1$ head vertices, a Berge route entering at the tail and leaving at a head (the program's convention). If every pair of vertices has at most $m-1$ arc-disjoint directed Berge routes, then*

$$|E(\mathcal{H})| \leq \frac{(m-1)n(n-1)}{r-1},$$

and for $m-1 \leq \binom{n-\lceil n/2 \rceil - 1}{r-2}$ there are feasible constructions with $\max_{1 \leq \alpha \leq n-1} \alpha \left\lfloor \frac{(m-1)(n-\alpha)}{r-1} \right\rfloor \approx (m-1)n^2/(4(r-1))$ hyperedges, so the value is quadratic in n .

Proof. Upper bound: a hyperedge with tail u serves the $r-1$ ordered pairs (u, h) , h a head. If some ordered pair (u, v) were served by m distinct hyperedges, those would be m arc-disjoint one-step routes, so each ordered pair is served at most $m-1$ times and $(r-1)|E| \leq (m-1)n(n-1)$. Lower bound: split V into tails A , $|A| = \alpha$, and heads B . By [Lemma A.33](#) there is a simple $(r-1)$ -uniform hypergraph $\mathcal{G}^*$ on B with maximum degree $m-1$ and $\lfloor (m-1)|B|/(r-1) \rfloor$ edges; give every tail $a \in A$ the hyperedges (a, H) , $H \in \mathcal{G}^*$. No vertex of B is a tail, so every route is a single step and $\lambda(a, b) = \deg_{\mathcal{G}^*}(b) \leq m-1$, while pairs inside A or starting in B have no routes. $\square$

A.8 The random threshold

The probabilistic ingredients are two Chernoff bounds, quoted in the form we use them.

Theorem A.40 (Chernoff bounds; see [16, Cor. 4.4, Thm. 4.4]). *Let X be a sum of independent $\{0, 1\}$ -valued random variables with mean μ . For $\delta > 0$, $\Pr[X \geq (1 + \delta)\mu] \leq \exp(-\frac{\min(\delta, \delta^2)\mu}{4})$, and for $\delta \in (0, 1)$, $\Pr[X \leq (1 - \delta)\mu] \leq \exp(-\frac{\delta^2\mu}{2})$.*

Corollary A.41. *Let $X \sim \text{Binomial}(n - 1, p)$ with $p = cm/n$ for a constant $c \in (0, 1)$. Then for all sufficiently large m , $\Pr[X \geq m] \leq e^{-\alpha m}$ with $\alpha = \alpha(c) > 0$.*

Proof. The mean is $\mu = (n - 1)p \leq cm$, so $m \geq (1 + \delta)\mu$ with $\delta \geq \frac{1-c}{c} > 0$, and Theorem A.40 applies with $\mu \geq cm/2$ for $n \geq 2$, giving $\alpha = \min(\frac{1-c}{c}, (\frac{1-c}{c})^2) \frac{c}{8}$. $\square$

Proof of Theorem 1.15. Recall the hypotheses: $m = m(n) \geq 2$ with $m/\ln n \rightarrow \infty$ and $m = o(n)$, and $p = cm/n$ for a constant $c > 0$.

Case $c > 1$. The edge count $|E|$ of $G(n, p)$ is binomial with mean $\mu = \binom{n}{2}p = \frac{cm(n-1)}{2}$. With $\delta = 1 - \frac{1}{c}$ the lower-tail bound of Theorem A.40 gives

$$\Pr\left[|E| \leq \frac{m(n-1)}{2}\right] \leq \exp\left(-\frac{(c-1)^2 m(n-1)}{4c}\right) \rightarrow 0,$$

so with high probability $|E| > \left\lfloor \frac{m(n-1)}{2} \right\rfloor$, and the contrapositive of Mader's theorem (Theorem 1.2) forces a pair with $\lambda \geq m$. Hence $\Pr[\lambda^{\max} \geq m] \rightarrow 1$.

Case $c < 1$. Each degree is $\text{Binomial}(n - 1, p)$, so by Corollary A.41 and a union bound over the n vertices,

$$\Pr[\Delta(G) \geq m] \leq n e^{-\alpha m} = e^{\ln n - \alpha m} \rightarrow 0,$$

since $m/\ln n \rightarrow \infty$. By the degree bound (Lemma A.3) $\lambda^{\max} \leq \Delta(G)$, so $\Pr[\lambda^{\max} \geq m] \rightarrow 0$. $\square$

Remark A.42 (The vertex and directed analogues). Below the threshold, $\kappa^{\max} \leq \lambda^{\max}$ gives the vertex statement for free, and in $D(n, p)$ the out-degree version of the same union bound applies. Above the threshold, the minimum degree exceeds m with high probability by the lower-tail bound, and the theorems of Bollobás [3] for $G(n, p)$ and its directed analogue [6] state that in this regime the global connectivities equal the minimum degree (respectively minimum semi-degree) with high probability, so $\kappa^{\max}$ and the directed $\lambda^{\max}$ also cross m at $p^* = m/n$. The growth hypothesis $m/\ln n \rightarrow \infty$ is what lets the union bound beat the factor n ; for fixed m the bound is vacuous and finer tools (Poisson approximation of low-degree vertex counts) would be needed.

A.9 Computational transcripts

The cut-counting optimisation formulation that certifies Theorem 2.1 is given in Chapter 2. Here are the solver transcripts that back it, together with the exhaustive-search output that settles the directed base cases. The complete commented source accompanies the document as a single

self-contained file, `erdos915_unified.py` in the `program/` directory, which runs with only `numpy`, `networkx`, `scipy`, and `matplotlib`.

```

1 Cut-MILP certification of the directed multigraph problem
2 M*(n) is the scaled optimum; L_m^dir(n) = (m-1) M*(n).
3
4 n=3: status=OPTIMAL, M*=4.000, target=2(n-1)=4, proof=True, time=0.0s, L_3^
    ↳ dir(n)=8
5 n=4: status=OPTIMAL, M*=6.000, target=2(n-1)=6, proof=True, time=0.3s, L_3^
    ↳ dir(n)=12
6 n=5: status=OPTIMAL, M*=8.000, target=2(n-1)=8, proof=True, time=5.9s, L_3^
    ↳ dir(n)=16
7 n=6: status=OPTIMAL, M*=10.000, target=2(n-1)=10, proof=True, time=1259.9s,
    ↳ L_3^dir(n)=20

```

Listing A.1. Solver output: $M^*(n) = 2(n-1)$, optimal with zero gap, for $n = 3, 4, 5, 6$.

The cut-counting optimisation closes the gap up to $n = 5$ inside a short budget and reaches $n = 6$ in about twenty minutes, after which the branch count defeats it. The base cases of the directed induction ([Theorem 1.9](#)) run to $n = 7$, and those are settled instead by the pruned exhaustive search over all simple digraphs with $\lambda^{\max} \leq 1$. That search is exact and complete at these sizes, and it agrees with $\ell_2^{\text{dir}}(n) = M(n)$ throughout.

```

1 Exhaustive search over all simple digraphs with lambda^max <= 1 (directed,
    ↳ m=2).
2 Branch and bound; reported max arc count = ell_2^dir(n) = max(2(n-1), floor
    ↳ (n^2/4)).
3
4 n=2: max= 2, formula= 2, proved (complete), nodes=5, 0.0s
5 n=3: max= 4, formula= 4, proved (complete), nodes=22, 0.0s
6 n=4: max= 6, formula= 6, proved (complete), nodes=537, 0.0s
7 n=5: max= 8, formula= 8, proved (complete), nodes=17823, 0.1s
8 n=6: max=10, formula=10, proved (complete), nodes=788101, 9.1s
9 n=7: max=12, formula=12, proved (complete), nodes=45122129, 718.1s
10
11 The cut-MILP certifier of the multigraph problem closes n <= 6 (n = 6 in
    ↳ about
12 21 minutes); the directed m = 2 base cases through n = 7 that the induction
13 requires are settled by this exhaustive search instead.

```

Listing A.2. Exhaustive-search output for the base cases $2 \leq n \leq 7$: the maximum arc count equals $M(n) = 2, 4, 6, 8, 10, 12$, proved complete at each n .

A.10 How the program checks itself

The invariants are checked by the self-test at the foot of `erdos915_unified.py`, run simply with `python erdos915_unified.py`. It confirms the checker against the proven $m = 2$ values 2, 4, 6, 8, the named constructions against their predicted sizes and connectivities, the Gomory–Hu shortcut against brute-force max-flow, the Berge connectivities of the hypergraph checker, the

Monte Carlo appearance threshold and Whitney's inequality on every sample, the cut-counting optima for small n , the search reaching the certified optimum, and the driver returning the proven value or a feasible witness in each mode. Every check passes. A single companion script, `make_figures.py`, regenerates every figure in the thesis from the same file, with the mapping listed in `program/README.md`.

Appendix B

Supplementary Figures

This appendix collects the supplementary figures that the program produces but that the chapters reference rather than reproduce in full. Each one is named where it belongs in the body, and the file name printed in its caption matches the output of `make_figures.py` and the figure table in `program/README.md`. Figures already shown in the running text are not repeated here.

B.1 Search traces across the variants

Figure 3.2 in Chapter 3 follows one cooling run in detail. The runs below repeat that experiment for the other variants. Each plots every visited graph by its binding connectivity (horizontal) against its size (vertical), colouring each point by the step at which it was visited, from the hot early exploration in warm tones to the converged tail in blue. In every case the walk strays past the feasibility boundary while hot and is driven back onto the densest feasible graph as the temperature falls, exactly as in the body figure. The seeds are fixed, so each picture reproduces verbatim.

Search trace: simple undirected, $n = 7, m = 3$




Figure B.1. Search trace for the simple undirected edge problem at $n = 7, m = 3$. File `trace_simple_undirected_n7_m3.png`.

Search trace: simple directed, $n = 5, m = 3$




Figure B.2. Search trace for the simple directed arc problem at $n = 5, m = 3$. File `trace_simple_directed_n5_m3.png`.

Search trace: multi undirected, $n = 5, m = 3$




Figure B.3. Search trace for the multigraph undirected edge problem at $n = 5, m = 3$. File `trace_multi_undirected_n5_m3.png`.

Search trace: multi directed, $n = 5, m = 3$




Figure B.4. Search trace for the multigraph directed arc problem at $n = 5, m = 3$. File `trace_multi_directed_n5_m3.png`.

Search trace: multi directed vertex-sep, $n = 4, m = 3$




Figure B.5. Search trace for the multigraph directed *vertex-disjoint* problem at $n = 4, m = 3$, where the binding quantity is $\kappa^{\max}$. File `trace_multi_directed_n4_m3_vertex.png`.

B.2 The variant landscape at $m = 6$ and in three dimensions

These are the $m = 6$ companions to the $m = 3$ enumeration views of [Chapter 4](#), together with a three-dimensional view of the appearance threshold. They confirm that raising the forbidden threshold relocates the feasibility boundary without changing the qualitative picture.




Figure B.6. How the full small- n enumeration distributes by binding connectivity $\lambda^{\max}$ across all twelve variants at $m = 6$, the companion to [Figure 4.5](#). Blue bars are feasible graphs ($\lambda^{\max} \leq 5$), red bars infeasible ones, and the dashed line marks the feasibility boundary. File `conn_dist_m6.png`.




Figure B.7. The same enumeration at $m = 6$ viewed by edge count, the companion to Figure 4.6. Feasible graphs are blue, infeasible ones red, and the dotted vertical line marks the extremal value at that size. File edges_dist_m6.png.




Figure B.8. The appearance threshold of Figure 4.2 as a three-dimensional bar chart, for three representative variants (simple undirected edge, simple directed arc, three-uniform hypergraph edge) at $n = 12$, $m = 3$. Each panel plots the fraction of $G(n, p)$ samples (height) reaching each binding connectivity $\lambda^{\max}$ (depth) as the density p varies (width), so the mass migrates from low to high connectivity as p crosses the proved threshold $p^* = m/n$. File threshold_3d.png.

Bibliography

- [1] Z. Baranyai. On the factorization of the complete uniform hypergraph. In *Infinite and Finite Sets (Colloq., Keszthely, 1973)*, Vol. I, volume 10 of *Colloq. Math. Soc. Janos Bolyai*, pages 91–108. North-Holland, 1975.
- [2] K. Bérczi, K. Chandrasekaran, T. Király, and S. Kulkarni. Splitting-off in hypergraphs. In *Proceedings of the 51st International Colloquium on Automata, Languages, and Programming (ICALP 2024)*, Leibniz International Proceedings in Informatics (LIPIcs), 2024. arXiv:2307.08555.
- [3] B. Bollobás. The evolution of random graphs—the giant component. *Transactions of the American Mathematical Society*, 286(1):257–274, 1984. Shows $\kappa(G(n, p)) = \delta(G(n, p))$ with high probability.
- [4] B. Bollobás and P. Erdős. Extremal problems in graph theory. *Matematikai Lapok*, 13:143–152, 1962. Original statement of Problem 915.
- [5] P. Erdős. Erdős problems. <https://www.erdosproblems.com/915>. Problem 915.
- [6] A. Frieze and M. Karoński. *Introduction to Random Graphs*. Cambridge University Press, 2016. Chapter 7 covers connectivity in $G(n, p)$ and $D(n, p)$; the directed analogue of Bollobás’s $\kappa = \delta$ result appears in Section 7.3.
- [7] R. E. Gomory and T. C. Hu. Multi-terminal network flows. *Journal of the Society for Industrial and Applied Mathematics*, 9(4):551–570, 1961. Introduces the Gomory-Hu tree structure.
- [8] M. J. H. Heule. Schur number five. In *Proceedings of the Thirty-Second AAAI Conference on Artificial Intelligence (AAAI-18)*, pages 6598–6606. AAAI Press, 2018.
- [9] M. J. H. Heule, O. Kullmann, and V. W. Marek. Solving and verifying the Boolean Pythagorean triples problem via cube-and-conquer. In *Theory and Applications of Satisfiability Testing (SAT 2016)*, volume 9710 of *Lecture Notes in Computer Science*, pages 228–245. Springer, 2016. The certified SAT proof is roughly 200 terabytes in DRAT format.
- [10] J. E. Hopcroft and R. E. Tarjan. Dividing a graph into triconnected components. *SIAM Journal on Computing*, 2(3):135–158, 1973.
- [11] J. L. Leonard. On a conjecture of Bollobás and Erdős. *Periodica Mathematica Hungarica*, 2(1–4):191–194, 1972. Proves $\ell_m(n) = k_m(n)$ for $m \leq 4$.
- [12] W. Mader. Ein Extremalproblem des Zusammenhangs von Graphen. *Mathematische Zeitschrift*, 131:223–231, 1973. Solves Erdős Problem 915 for edge-disjoint paths.

- [13] B. D. McKay and A. Piperno. Practical graph isomorphism, II. *Journal of Symbolic Computation*, 60:94–112, 2014. Describes the nauty/Traces canonical-labelling tools underlying geng.
- [14] B. D. McKay and S. P. Radziszowski. $R(4, 5) = 25$. *Journal of Graph Theory*, 19(3):309–322, 1995. Exhaustive-generation determination of a Ramsey number.
- [15] K. Menger. Zur allgemeinen Kurventheorie. *Fundamenta Mathematicae*, 10:96–115, 1927. The original statement of Menger’s theorem.
- [16] W. Mulzer. Five proofs of Chernoff’s bound with applications. *Bulletin of the EATCS*, 124, 2018. Accessible survey of Chernoff bound variants.
- [17] A. A. Razborov. Flag algebras. *The Journal of Symbolic Logic*, 72(4):1239–1282, 2007.
- [18] B. A. Sørensen and C. Thomassen. On k -rails in graphs. *Journal of Combinatorial Theory, Series B*, 17(2):143–159, 1974. Determines $k_5(n) = \lfloor 8n/3 \rfloor - 3$.
- [19] W. T. Tutte. *Connectivity in Graphs*. University of Toronto Press, Toronto, 1966.
- [20] H. Whitney. Congruent graphs and the connectivity of graphs. *American Journal of Mathematics*, 54(1):150–168, 1932.

