

Connectivity as Bottleneck for connections in Graphs

Erdős Problem 915, from Proof to Search

Louis Vandenbruwaene

Promotor: Prof. Jan Goedgebeur
Supervisor: Prof. Stijn Cambie
[KU Leuven](#)

Thesis presented in
fulfillment of the requirements
for the degree of Master of Science
in Mathematics

Academic year 2026–2027

Copyright information

Student paper as part of an academic education and examination. No correction was made to the paper after examination.

© Copyright by KU Leuven

Without prior written permission from both the supervisor(s) and the author(s), copying, reproducing, using, or realizing this publication or parts thereof is prohibited. For requests or information regarding the copying and/or use and/or realization of parts of this publication, please contact KU Leuven, Faculty of Science, Celestijnenlaan 200H box 2100, 3001 Leuven (Heverlee), Telephone +32 16 32 14 01.

Prior written permission from the supervisor(s) is also required for using the (original) methods, products, circuits, and programs described in this work for industrial or commercial purposes, and for submitting this publication for scientific awards or contests.

Foreword

This thesis studies how many edges or arcs a network can contain when no pair of vertices is allowed to have too many independent routes between them. The starting point is Erdős Problem 915, an extremal question about edge-disjoint paths in simple graphs. The work begins with the necessary background to formulate the problem and all its variants. It then discusses their answers or highlights the difficulty behind unsolved cases. Afterwards, computational research on small graphs confirms these values or provides evidence for conjectures. This is accomplished by measuring connectivity with a maximum-flow routine and by searching for extremal graphs with metaheuristics. The results of these computations are thus either a proof of the extremal value in the search space or a lower bound that is conjectured to also be upper, supported by the unfinished search for a counterexample. At the end, the thesis synthesises the results and presents the open problems that remain. This work is theoretical in nature, but the computational methods and the resulting conjectures are of interest to anyone who uses networks and wants to understand the relationship between connectivity and the connections they entail.

I am grateful to my supervisor, Prof. Stijn Cambie, for his day-to-day guidance, corrections, and mathematical feedback throughout the project. I could not ask for a more understanding and enthusiastic supervisor. I thank my promotor, Prof. Jan Goedgebeur, whose graph theory implementation knowledge pushed the boundaries of the computational work significantly. I also thank the people who discussed early versions of the arguments, checked examples, or helped clarify the presentation. Above all I thank my girlfriend Sara, whose patience and encouragement carried me through the long stretches of this work.

Contribution Statement

All the work in this thesis is my own, except where I explicitly say otherwise. I used Artificial Intelligence to implement the code, suggest and verify proofs, format $\text{\LaTeX}$ and spellcheck. Where a result comes from the literature I say so and cite it. The theorems I use but do not claim as mine are those of Mader, Leonard, Sørensen and Thomassen, Menger, Gomory and Hu, and Baranyai. Related hypergraph-connectivity work includes Bérczi, Chandrasekaran, Király and Kulkarni [1]. Proofs are located in [Appendix A](#) and all computational work is available in my personal repository [2]. The summary of my original results is:

Variant	Contribution
Hypergraph, vertex separation	Exact value at $m = 2$. The $m = 3$ universal bound and its stated attainment range, via a new incidence-rank lemma (Theorems 1.12 and 1.13).
Directed multigraph	Exact value for all n, m , plus quadratic-branch classification for $n \geq \max(8, 2m)$ (Theorems A.45 and 3.3).
Directed simple graph, arc separation	Unconditional $n^2/4 + \Theta_m(n)$ bound and an $O(\sqrt{m} n^{3/2})$ structural distance from a one-way bipartite graph (Proposition A.26 and theorem 3.1).
Multigraph, vertex separation with parallel routes distinct	Exact $m = 2$ value, counterexamples to the thickened-tree guess, matching order $\Theta(m^2 n)$ as $m \rightarrow \infty$ with $n/m \rightarrow \infty$, and an exact block-knapsack formulation (Theorems A.61 and 3.5 and proposition A.63).
Transferred quadratic bounds	$k_m^{\text{dir}}(n) = n^2/4 + \Theta_m(n)$ and, for fixed r, m , leading term $(m-1)n^2/(4(r-1))$ for directed hypergraphs under both separations and all orientations (Theorems A.59 , 3.2 and 3.4).

Short Summary

Erdős Problem 915, posed by Bollobás and Erdős [3], asks for the maximum number of edges in a graph on n vertices such that no two vertices are connected by m edge-disjoint paths. In modern notation this asks for

$$\ell_m(n) = \max\{|E(G)| : |V(G)| = n, \lambda_G^{\max} \leq m - 1\}.$$

Mader proved that, for simple graphs and $n \geq m \geq 2$,

$$\ell_m(n) \leq \left\lfloor \frac{m(n-1)}{2} \right\rfloor$$

and provided a construction achieving this bound when $(m-1)|(n-1)$. This thesis revisits this theorem and studies the corresponding extremal problem under three independent changes: the graph model (simple graphs, multigraphs, hypergraphs and multi-hypergraphs), the path-separation notion (edge-disjoint or internally vertex-disjoint) and the presence or absence of directions. The simple case for edge-disjoint paths is proven independently here using Gomory-Hu trees [4] and a general extremal graph construction resolves this case for all n and m . Other rigorously settled variants include the multigraph value $L_m(n) = (m-1)(n-1)$ and the equality of the simple edge and vertex problems for $m \leq 4$ and their divergence at $m = 5$, as shown by Sørensen and Thomassen [5]. The directed case is resolved for $m = 2$: the value $\ell_2^{\text{dir}}(n) = \max(2(n-1), \lfloor n^2/4 \rfloor)$ is proved by a long induction and for $m \geq 3$ the hand proof gives way to a conjecture, after an explicit ten-vertex graph rules out the naive guess.

The centerpiece is then a unified program, which represents non-hypergraphs with a multiplicity-matrix and converts the vertex-disjoint problem to an edge-disjoint one by a standard vertex-splitting construction. The hypergraphs are handled as a list of hyperedges, which may use head and tail sets to represent directed hyperedges. Then a maximum-flow algorithm measures connectivity between all pairs of vertices and reports the maximum local connectivity. At each point of the program, optimisations are applied to reduce the search space and traverse it more efficiently. When these optimisations are exhausted and the computation becomes intractable, the program uses metaheuristics to search for extremal graphs, which are then checked for correctness.

The hypergraph vertex problem is settled completely at $m = 2$. At $m = 3$ the formula $\lfloor 2(n-1)/(r-1) \rfloor$ is a universal upper bound for multihypergraphs and is exact whenever $r \geq 3$ and $2 \leq \binom{n-2}{r-2}$, the upper bound following from a new rank bound on incidence graphs proved by way of Tutte's triconnected decomposition. On the directed side the multigraph problem closes completely, $L_m^{\text{dir}}(n) = (m-1) \max(2(n-1), \lfloor n^2/4 \rfloor)$ for every n and every m , by deleting a sparse reachability-preserving subgraph at a time rather than a vertex at a time. For every

fixed $m \geq 3$, the simple digraph problem, arc and vertex alike, is pinned at $n^2/4 + \Theta_m(n)$ with no assumption on the shape of the extremiser and at $m = 2$ its exact value has only an $O(1)$ correction to $n^2/4$. For fixed r and m , the directed hypergraph, which had only bounds a factor of four apart, has its leading term settled at $(m - 1)n^2/(4(r - 1))$, so its bipartite construction is asymptotically optimal.

Abbreviations and Symbols

MILP	mixed-integer linear program
SA	simulated annealing
SPQR	the triconnected decomposition tree (S series, P parallel, Q single edge, R rigid)
$G = (V, E)$	graph with vertex set V and edge set E
$D = (V, A)$	directed graph with vertex set V and arc set A
n	number of vertices
m	forbidden-connectivity parameter, with all local connectivities at most $m - 1$
r	hyperedge size (every hyperedge spans r vertices)
$\lambda_G(u, v)$	maximum number of edge-disjoint u - v paths
$\lambda_G^{\max}$	maximum local edge-connectivity over all vertex pairs
$\kappa_G(u, v)$	maximum number of internally vertex-disjoint u - v paths
$\kappa_G^{\max}$	maximum local vertex-connectivity over all vertex pairs
$\ell_m(n)$	simple undirected edge-disjoint extremal function
$L_m(n)$	undirected multigraph edge-disjoint extremal function
$k_m(n)$	simple undirected vertex-disjoint extremal function
$\ell_m^{\text{dir}}(n)$	simple directed arc-disjoint extremal function
$L_m^{\text{dir}}(n)$	directed multigraph arc-disjoint extremal function
$k_m^{\text{dir}}(n)$	simple directed vertex-disjoint extremal function
$\ell_m^{(r)}(n), k_m^{(r)}(n)$	r -uniform hypergraph edge- and vertex-disjoint extremal functions
$K_m^{\text{multi}}(n)$	multigraph vertex-disjoint extremal function with parallel copies counted as distinct routes (Section A.9)
$M(n)$	$\max(2(n - 1), \lfloor n^2/4 \rfloor)$, the directed multigraph value per unit of $m - 1$
$M^*(n)$	the m -free directed multigraph optimum, $L_m^{\text{dir}}(n) = (m - 1) M^*(n)$, equal to $M(n)$ by Corollary A.38
$B_{p,q}$	one-directional complete bipartite digraph, all arcs from a p -part to a q -part
$\mu(u, v)$	multiplicity of an edge or arc between u and v
$h_m(b)$	most edges a feasible 2-connected block on b vertices can carry (Theorem A.13)
c_m	per-vertex growth constant of the simple undirected vertex problem, $c_m = \lim_{n \rightarrow \infty} k_m(n)/(n - 1) = \sup_{b \geq 2} h_m(b)/(b - 1)$ (Theorem A.13)
$\sigma(e)$	edge sensitivity $\lambda^{\max}(G) - \lambda^{\max}(G - e)$

Contents

Foreword	I
Contribution Statement	II
Short Summary	III
Abbreviations and Symbols	V
1 The Problem and Its Variants	1
1.1 Graphs and routes	1
1.2 Erdős Problem 915	2
1.3 Variants of the problem	3
1.4 Proven bounds	4
1.5 Conjectured bounds	8
2 Certifying and Discovering Bounds by Machine	10
2.1 The solver	10
2.1.1 The connectivity checker built on maximum flow	11
2.2 Checking every graph of a fixed size	12
2.2.1 Pruned search	14
2.2.2 Generating search	16
2.3 Proving every m at once	16
2.4 The heuristic search	19
2.5 Edge sensitivity	20
2.6 Keeping the checker cheap inside the search	20
2.7 Rediscovering the known extremisers	22
2.8 Results and evidential status	23
3 Synthesis, Results, and Open Problems	29
3.1 The directed frontier and the missing decomposition	29
3.2 Two principal phenomena	31
3.3 Summary of the sixteen variants	32
3.3.1 Orientation models for the directed hypergraph	33
3.3.2 The multigraph vertex problem under the other convention	34

3.4	Contributions and limitations of the program	35
3.5	Open problems	35
A	Full Proofs and Computational Audit	37
	PART I. CLASSICAL MACHINERY	37
A.1	Menger, Gomory–Hu, and the degree bound	37
A.2	Mader’s theorem in full	39
A.2.1	The vertex problem is a block problem	43
	PART II. THE DIRECTED RESULTS	45
A.3	The directed arc value at $m = 2$	45
A.4	Structural propositions on the directed frontier	49
A.4.1	What that proof actually used	56
A.5	The directed multigraph problem, solved in full	57
A.5.1	The lower bound: thickening	58
A.5.2	The upper bound: a reachability skeleton	58
A.5.3	Extremal classification on the quadratic branch	65
	PART III. THE HYPERGRAPH RESULTS	67
A.6	The hypergraph bounds	68
A.7	The hypergraph vertex problem	71
A.8	The directed hypergraph	76
	PART IV. ONE EXTENSION, AND THE AUDIT	83
A.9	The multigraph vertex problem under the other convention	83
A.9.1	The value is a block problem	88
A.10	Computational audit	90

Chapter 1

The Problem and Its Variants

A network is useful not only because it connects, but because it connects in more than one way. A single road between two towns is a liability and a second independent road is insurance. The mathematical content of that intuition is the number of routes between two points that share no segment and the question this thesis circles is how many connections a network can hold before some pair of points is forced to have too many such routes. This chapter gives the foundation for the question and its variants and goes over the results that can be proven using logical arguments.

1.1 Graphs and routes

To state the question precisely we need only a few ideas. A *graph* G is a finite set of points, called *vertices*, together with a (multi)set of *edges*, each joining two (or more) distinct vertices. We write $V(G)$ for the set of vertices and $E(G)$ for the (multi)set of edges. Therefore $|E(G)|$ is the number of edges, which is the very quantity this thesis tries to make as large as possible under certain constraints. If $E(G)$ is a set and its edges join exactly two vertices, G is a *simple graph*. One pictures such a simple graph by representing the vertices as dots and the edges as lines between them.




Figure 1.1. The simple graph $K_5 - \{04, 13\}$ has five vertices and eight edges, where K_n is the complete graph on n vertices.

We write $n := |V(G)|$ for the number of vertices and define the *degree* of a vertex as the number of edges meeting it. Unless specified otherwise, graphs are simple. A *path* from a vertex u to a vertex v is a sequence of distinct vertices that starts at u , ends at v , and joins each consecutive pair by an edge. Two paths between the same endpoints are *edge-disjoint* if no edge belongs to both. The largest number of pairwise edge-disjoint paths between u and v is their *local edge-connectivity* $\lambda_G(u, v)$. The pair with the most such paths sets the ceiling for the whole graph:

$$\lambda_G^{\max} = \max_{u \neq v} \lambda_G(u, v).$$

A theorem of Menger [6], recalled in [Appendix A](#), gives this quantity a second face: $\lambda_G(u, v)$ also equals the least number of edges one must remove to cut every route from u to v ([Figure 1.2](#)). Edge disjoint paths between a pair is exactly the same thing as the minimum cut of unit flow between them.




Figure 1.2. Menger's theorem on [Figure 1.1](#): three edge-disjoint 1–3 routes (coloured) meet the minimal three-edge cut around vertex 1, so $\lambda(1, 3) = 3$.

1.2 Erdős Problem 915

The constraint at the heart of this thesis is a ceiling on connectivity. Requiring $\lambda_G^{\max} \leq m - 1$ says that no pair of vertices is joined by as many as m edge-disjoint paths, or equivalently that every pair can be severed by removing fewer than m edges. The question is how many edges a graph can hold while staying under that ceiling.

Question 1.1 (Erdős Problem 915 [7]). How many edges can a graph on n vertices have if no two of its vertices are joined by m edge-disjoint paths?

Writing the answer as an extremal function,

$$\ell_m(n) = \max\{|E(G)| : |V(G)| = n, \lambda_G^{\max} \leq m - 1\},$$

the problem asks for $\ell_m(n)$, the most edges possible under the ceiling. Mader proved that no graph under the ceiling has more than $\lfloor m(n - 1)/2 \rfloor$ edges (the upper bound) in the simple-graph case and provided the shared-clique construction, which achieves the bound but is only defined when $m - 1$ divides $n - 1$. The brackets $\lfloor \cdot \rfloor$ round their content down to the nearest whole number and their ceiling counterpart $\lceil \cdot \rceil$ rounds up. Both recur throughout, because an edge count is always an integer while the formulas that bound it need not be.

Theorem 1.2 (Mader [8]). *For all $n \geq 2$ and $m \geq 2$,*

$$\ell_m(n) \leq \left\lfloor \frac{m(n - 1)}{2} \right\rfloor.$$

Notice how $n \geq m$ is not needed here, but will be to attain equality since simple graphs cannot exceed the edge count of complete graphs. The proof given in [Appendix A](#) was discovered independently of Mader and uses the Gomory-Hu tree ([Theorem A.2](#)). The shared clique graph is then generalised to an extremal construction that exists for all values of $n \geq m$ ([Theorem A.11](#)).

The very same Gomory-Hu tree returns for the multigraph and hypergraph variants ([Theorem 1.3](#), [Proposition 1.11](#)). Hence it is worth mentioning its properties here. Every graph has a weighted, spanning tree such that $\lambda_G(u, v)$ equals the lightest edge on the tree path from u to v . All $\binom{n}{2}$ local connectivities are therefore encoded by just $n - 1$ weights. The largest of them, $\lambda_G^{\max}$, is simply the heaviest edge of the tree. The proof ([Appendix A](#)) uses a double counting argument on that tree.

1.3 Variants of the problem

So far the problem has been stated for simple graphs, but the same question remains unanswered in a more general setting. Each assumption made so far will be toggled to create a lattice of variants and the thesis will explore each node of that lattice.

The first is *arity*: an edge joins exactly two vertices and a *hyperedge* joins $r \geq 2$. Hence such *hypergraphs* use *Berge paths*, which aligns with the path definition above. The second is *multiplicity*: a *simple* graph carries at most one copy of each edge, while a *multi* graph allows parallel copies, recorded by a multiplicity $\mu(u, v)$. The third is *direction*: a directed edge, called arc, is an ordered pair of vertices and a directed hyperedge has one tail and $r - 1$ heads. These three toggles create a lattice of eight models, shown in [Figure 1.3](#).




Figure 1.3. The eight models side by side, with the four undirected on top and the four directed below. The simple graph is a triangle, the multigraph has parallel edges, the hypergraph has three hyperedges, and the multi-hypergraph has one repeated hyperedge. The directed versions have arcs instead of edges, with the same pattern of repetition and hyperedges.

The digraphs do not have a Gomory-Hu tree, but they do have a directed analogue of Menger's theorem. A digraph has three degree counts at each vertex. The *out-degree* $d^+(v)$ counts the arcs leaving v and the vertices they reach are its *out-neighbours*. The *in-degree* $d^-(v)$ counts the arcs entering v and the vertices they come from are its *in-neighbours*. The *total degree* $d(v) = d^+(v) + d^-(v)$ adds the two. [Figure 1.4](#) shows a digraph carrying three arc-disjoint routes between one chosen pair of vertices. The directed form of Menger's theorem

says that route count equals the size of the smallest arc-cut separating the pair, and the figure also marks the out-degree and in-degree that bound both quantities from above, which is exactly how Menger reads in a one-way network. A directed r -edge has one tail and $r - 1$ heads. In the figures, one colour denotes one hyperedge, drawn either as a star or as a line oriented away from its tail. It is a single edge on several vertices, not a path of ordinary edges. A directed *Berge route* enters that edge at its tail and leaves at one of its heads.




Figure 1.4. Three arc-disjoint $u \rightarrow v$ routes (coloured) and the matching three-arc directed cut (red) in a *simple* digraph. Thus $\lambda(u, v) = 3$, meeting the degree bound $\min(d^+(u), d^-(v)) = 3$.

The final assumption that will be considered is *separation*: routes may be required to avoid sharing an edge, as above, or the stricter *internally vertex-disjoint*, sharing no intermediate vertex. This gives the local vertex-connectivity $\kappa_G(u, v)$ and its maximum $\kappa_G^{\max}$. Whitney's inequality $\kappa_G(u, v) \leq \lambda_G(u, v)$ [9] records that avoiding shared vertices is at least as demanding as avoiding shared edges and Figure 1.5 shows the gap on a graph where two routes are edge-disjoint yet forced through one common vertex.




Figure 1.5. The separation axis: two edge-disjoint s – t routes both pass through w , so $\lambda(s, t) = 2$ but $\kappa(s, t) = 1$.

With all this in mind, the thesis considers multi-hypergraphs and directed multi-hypergraphs for both edge- and vertex-disjoint routes, where $r = 2$ and no parallel edges are special cases. The final status map appears in Figure 3.2. The notation for all their extremal functions is summarised in Table 1.1.

1.4 Proven bounds

Theorem 1.3. *For undirected multigraphs on n vertices, $L_m(n) = (m - 1)(n - 1)$.*

Appendix A gives the full proof which also uses the Gomory–Hu tree. The lower bound is a spanning tree with every edge thickened to $m - 1$ parallel copies.

Leonard proved that in simple graphs the edge and vertex problems give the same answer

Table 1.1. Notation for the extremal functions.

Model	edge-disjoint routes	internally vertex-disjoint routes
Simple graph	$\ell_m(n)$	$k_m(n)$
Multigraph	$L_m(n)$	$K_m(n)$
r -uniform simple hypergraph	$\ell_m^{(r)}(n)$	$k_m^{(r)}(n)$
r -uniform multihypergraph	$L_m^{(r)}(n)$	$K_m^{(r)}(n)$
Simple digraph	$\ell_m^{\text{dir}}(n)$	$k_m^{\text{dir}}(n)$
Directed multigraph	$L_m^{\text{dir}}(n)$	$K_m^{\text{dir}}(n)$
Directed r -uniform simple hypergraph	$\ell_m^{\text{dir}(r)}(n)$	$k_m^{\text{dir}(r)}(n)$
Directed r -uniform multihypergraph	$L_m^{\text{dir}(r)}(n)$	$K_m^{\text{dir}(r)}(n)$

while m stays small.

Theorem 1.4 (Leonard [10]). *For all $n \geq m$ and $m \leq 4$,*

$$k_m(n) = \ell_m(n) = \left\lfloor \frac{m(n-1)}{2} \right\rfloor.$$

Whitney's inequality $\kappa \leq \lambda$ makes every edge-feasible graph vertex-feasible (Figure 1.6. One is tempted to expect the equality forever. This however is not the case, and Sørensen and Thomassen proved that the vertex problem diverges from the edge problem at $m = 5$.

Theorem 1.5 (Sørensen-Thomassen [5]). *For simple graphs and all $n \geq 6$ with $n \neq 7$ and $n \neq 12$,*

$$k_5(n) = \left\lfloor \frac{8n}{3} \right\rfloor - 4,$$

while $k_5(7) = 15$ and $k_5(12) = 27$. The formula first exceeds the edge value at $n = 14$: for $6 \leq n \leq 13$ one has $k_5(n) = \left\lfloor \frac{5(n-1)}{2} \right\rfloor = \ell_5(n)$, and for every $n \geq 14$,

$$k_5(n) > \left\lfloor \frac{5(n-1)}{2} \right\rfloor = \ell_5(n).$$

Remark 1.6 (A counting convention that shifts every quoted constant by one). This thesis defines $\ell_m(n)$ and $k_m(n)$ as the *largest* number of edges a graph can carry while *no* pair is joined by m independent routes. Sørensen and Thomassen, and the Erdős Problems database [7] after them, state the same results the other way round, as the *smallest* number of edges that *forces* such a pair. Those two quantities are adjacent integers by construction: if k is the largest feasible edge count then no graph on n vertices with $k+1$ edges is feasible, so $k+1$ is exactly the forcing threshold.

Construction 1.7 (Directed hub). For $n \geq m \geq 2$, take a hub h and label the other $N = n - 1$ vertices $0, \dots, N - 1$. Add the $2N$ hub arcs $h \rightarrow i$ and $i \rightarrow h$, and on the non-hub vertices the $(m-2)N$ ring arcs $i \rightarrow i + j \bmod N$ for $j = 1, \dots, m-2$. The total is $m(n-1)$ arcs and every non-hub vertex has $d^+ = d^- = m-1$.




Figure 1.6. The edge and vertex extremal functions for $m = 2, 3, 4$ (blue shades, single curve per m since they agree) and $m = 5$ (dashed blue for edge, solid orange for vertex). The vertex function diverges from the edge function at $n = 14$ and grows faster thereafter.

Every ordered pair has a non-hub member, whose relevant degree is exactly $m - 1$ and, since $n \geq m$, smaller than the hub's own $n - 1$. That member is therefore the binding one and the degree bound gives $\lambda^{\max} \leq m - 1$. At $m = 2$ the ring is empty and what remains is the *bidirected star*, in which two spokes can only reach each other through the hub.

Construction 1.8 (One-directional bipartite digraph). Split the vertices into parts A and B of sizes $\lfloor n/2 \rfloor$ and $\lceil n/2 \rceil$, and include every arc from A to B and none the other way. The total is $\lfloor n^2/4 \rfloor$ arcs.

Nothing leaves B and nothing enters A , so no route has a second step and every pair is joined by at most the one direct arc, giving $\lambda^{\max} = 1$.

Theorem 1.9 (Directed arc value at $m = 2$). *For simple digraphs,*

$$\ell_2^{\text{dir}}(n) = \max\left(2(n-1), \left\lfloor \frac{n^2}{4} \right\rfloor\right).$$

The two constructions give the lower bound, one branch each. The upper bound is an induction on n ([Appendix A](#)): delete a vertex of minimum total degree, which carries at most the average number of arcs, and apply the bound to what remains. The base cases $n \leq 7$ are left to exhaustive search rather than hand argument, and [Chapter 2](#) is what runs them. The same induction never mentions the separation, so it survives the change to κ verbatim once the base cases are re-run.

Theorem 1.10 (Directed vertex value at $m = 2$). *For simple digraphs, $k_2^{\text{dir}}(n) = \ell_2^{\text{dir}}(n)$.*




Figure 1.7. The two branches tie at $n = 7$, where each holds 12 arcs at $\lambda^{\max} = 1$. Left: the bidirected star, a hub joined to six spokes both ways, $2(n - 1) = 12$. Right: the one-directional complete bipartite digraph $B_{3,4}$, every arc from A to B , $\lfloor n^2/4 \rfloor = 12$.

The hub construction leads for $n < 7$, the one-directional bipartite construction leads for $n > 7$ and the two tie at $n = 7$, where both reach 12 arcs. They are shown side by side in Figure 1.7. Notice how the directed value is quadratic in n for large n , while the undirected value is linear. This is because digraphs can be lopsided in one direction without creating routes back.

Proposition 1.11 (Hypergraph edge bound). *An r -uniform hypergraph on n vertices with Berge edge-connectivity at most $m - 1$ has at most $\left\lfloor \frac{(m-1)(n-1)}{r-1} \right\rfloor$ hyperedges. A simple hypergraph attains the bound whenever $m - 1 \leq \binom{n-2}{r-2}$, and a multihypergraph attains it whenever $(r - 1) \mid n - 1$.*

The proof, given in Appendix A, runs the multigraph argument through the hypergraph Gomory-Hu theorem (Theorem A.47 applied to the hyperedge-cut function in place of the edge-cut function, counting each hyperedge with the at least $r - 1$ tree cuts it crosses. The star hypertree (Figure 1.8) attains the $m = 2$ case, a hub joined to disjoint blocks of $r - 1$ vertices, with Berge connectivity one everywhere and $\lfloor (n - 1)/(r - 1) \rfloor$ hyperedges, exactly as a tree attained the multigraph bound. One genuine surprise is worth recording here: for $r \geq 3$ the bound is attained by *simple* hypergraphs, with no repeated hyperedge, whenever $m - 1 \leq \binom{n-2}{r-2}$ (Theorem A.48). In the non-hypergraph case simplicity is exactly what separates Mader's $\lfloor m(n - 1)/2 \rfloor$ from the multigraph value $(m - 1)(n - 1)$. One edge size higher, they are equal.




Figure 1.8. The $r = 3$, $n = 9$ star hypertree: four hyperedges meet only at the hub h , attaining $\lfloor (n - 1)/(r - 1) \rfloor = 4$ with Berge connectivity one.

Theorem 1.12 (Hypergraph vertex value at $m = 2$). *For $r \geq 2$ and $n \geq 1$, the maximum number of hyperedges of an r -uniform multihypergraph on n vertices with $\kappa^{\max} \leq 1$ is*

$$k_2^{(r)}(n) = \left\lfloor \frac{n-1}{r-1} \right\rfloor,$$

attained by the star hypertree of Figure 1.8. In particular $k_2^{(r)}(n) = \ell_2^{(r)}(n)$: the vertex and edge problems agree at $m = 2$ for every uniformity, and repeated hyperedges never help.

Theorem 1.13 (Hypergraph vertex value at $m = 3$). *For $r \geq 2$, every r -uniform multihypergraph on n vertices with $\kappa^{\max} \leq 2$ has at most $\left\lfloor \frac{2(n-1)}{r-1} \right\rfloor$ hyperedges. For $r \geq 3$ with $2 \leq \binom{n-2}{r-2}$ the bound is attained by a simple hypergraph, so*

$$k_3^{(r)}(n) = \left\lfloor \frac{2(n-1)}{r-1} \right\rfloor = \ell_3^{(r)}(n).$$

The first falls out once Berge routes are read as ordinary paths in the incidence graph, where the ceiling $\kappa^{\max} \leq 1$ says exactly that the incidence graph is a forest. The second is the harder one and carries the genuinely new tool: a rank bound for graphs in which one side of a partition must stay below local 3-connectivity, proved by way of Tutte's triconnected decomposition, which the incidence translation then turns back into the hypergraph statement. At $m \geq 4$ that route stops for want of a 4-connectivity analogue of the decomposition, and the value is open. The proposition counts hyperedges, but it does not say how to *measure* the Berge connectivity of a hypergraph we are handed and that measurement is the genuinely new part. The metro picture shows why. An ordinary edge is a one-stop link between a single pair, so policing edge-disjointness means watching each pair on its own. A hyperedge is a whole line: it stops at all r of its vertices and so touches $\binom{r}{2}$ pairs at once, yet like a line that can run only one train at a time it may serve only one route. When several routes want to board the same line at different stations, the checker must wave exactly one through and turn the rest away and it must do this for every line simultaneously. That bookkeeping is the one new piece of machinery the next chapter builds and it is why the hypergraph rejoins the family only once the checker is in place.

1.5 Conjectured bounds

Conjecture 1.14. *For simple digraphs and all $n \geq m \geq 2$,*

$$\ell_m^{\text{dir}}(n) = \max \left(m(n-1), \left\lfloor \frac{(n+m-2)^2}{4} \right\rfloor \right).$$

At $m = 2$ this is Theorem 1.9. The first branch is the directed hub. The second is the bipartite digraph with one extra layer inside its larger part.

Construction 1.15 (Augmented bipartite digraph). For $n \geq m \geq 2$, set $|B| = \lceil (n+m-2)/2 \rceil$ and $|A| = n - |B|$, include every arc from A to B , and inside B give each b_j the $m-2$ in-arcs from its cyclic predecessors $b_{j-1}, \dots, b_{j-(m-2)}$. The total is $\lfloor (n+m-2)^2/4 \rfloor$ arcs.

The only in-arcs of a $b \in B$ that an $a \in A$ can reach are the direct arc $a \rightarrow b$ and the $m - 2$ two-step detours through b 's in-neighbours inside B , so such a pair carries exactly $m - 1$ routes and no more, while pairs inside B are held down by the in-degree $m - 2$ and pairs from B to A or inside A have no routes at all.

At $m = 3, n = 10$ this is the digraph that refutes the natural first guess $\max(m(n-1), \lfloor n^2/4 \rfloor)$. Its 25 arcs from A to B carry a five-cycle inside B , giving every b_j one extra in-arc, so each pair a_i, b_j has the direct arc and one detour and $\lambda^{\max} = 2$. That is 30 arcs against the guess of 27, and the correction is exactly the shift of the partition towards B .

For fixed $m \geq 3$ the matching upper bound is open, though not entirely: [Theorem 3.1](#) proves unconditionally that the value is $n^2/4 + \Theta_m(n)$, so what the conjecture still claims beyond what is known is the coefficient of the linear term rather than the shape of the answer.

Chapter 2

Certifying and Discovering Bounds by Machine

The previous chapter gave insight into the problem and some variants, but left most harder questions open. This chapter will serve as a sanity check for the given theorems, establishing their correctness on small graphs by brute-force enumeration. It will also be the foundation of conjectures for the open cases, by discovering extremal graphs or at least sufficiently dense graphs. Pattern recognition on the discovered values, or even a family of extremal graphs, then lets conjectures be made, although those statements are naive because they are based on small graphs only. Larger graph spaces will be scoured by a heuristic, namely tabu search. A guided search is needed there because only a small fraction of the search space is extremal: assigning edges at random gives a binomially distributed edge count, so almost every graph lands far from the ceiling and sampling alone would never meet it. The archive is the repository at <https://github.com/louisvandenbruwaene/thesis> and every file path quoted in this thesis is relative to it, with the commands and the audit trail collected in [Section A.10](#). Below is a diagram of the program's architecture, showing how the modelled graphs are fed to the checker and how the results are used to either certify or discover bounds.




Figure 2.1. A single model feeds the exact max-flow checker. Certification turns that measurement into finite upper bounds, while heuristic search uses it to discover lower bounds. The badge colours on the code cards match the boxes here.

2.1 The solver

Given n and m , walk through every graph on n vertices, measure the largest local connectivity $\lambda^{\max}$ or $\kappa^{\max}$ of each with a max-flow computation and keep the densest graph that stays at or below the ceiling $m - 1$. In the vertex-disjoint case the graph will be transformed before the flow can be measured. Hypergraphs will also need to be transformed into a flow network, because it's represented by a (multi)set of hyperedges rather than a multiplicity matrix. The enumeration step will skip whole families of hopeless graphs, via a pruned search.

All non-hypergraphs are represented using one multiplicity matrix μ : it is symmetric when undirected and binary when simple. Hypergraphs are stored as (multi)sets of hyperedges, with a tail distinguished in the directed model. Every variant shares the same integer max-flow engine no matter the implementation.

2.1.1 The connectivity checker built on maximum flow

The whole question reduces to a flow computation. A *maximum flow* asks how much can be pushed from one point to another through a network whose links each carry a limited capacity. Menger's theorem says that $\lambda(u, v)$, the maximum number of edge-disjoint $u - v$ paths, equals the minimum number of edges that must be removed to separate u from v . For a simple graph, give every edge capacity one: the capacity of a cut is then just the number of edges in it. For a multigraph, the parallel copies can be combined into one link whose capacity is their multiplicity $\mu(u, v)$. In the directed case these capacities follow the direction of the arcs, while in the undirected case they are available in either direction. The max-flow/min-cut theorem of Ford and Fulkerson [11] says that this smallest cut has exactly the value of a maximum flow. So the route count we are after is just a maximum-flow value: measure the flow and you have measured the connectivity.

The routine that runs this flow and reports the route count is the *connectivity checker*. It is the single most important routine in the program. The program ships an optional compiled C helper that runs this same flow more efficiently.

The vertex-disjoint version follows the same idea after one transformation: split each vertex v into an in-copy v^{in} and an out-copy v^{out} joined by a single unit-capacity arc (Figure 2.3a). Every arc $u \rightarrow v$ of the original graph becomes $u^{\text{out}} \rightarrow v^{\text{in}}$ in the split network. The construction is not special to digraphs: for an undirected graph each edge $\{u, v\}$ is first read as the two opposite arcs $u \rightarrow v$ and $v \rightarrow u$ and then the same split applies, so one routine serves the undirected and directed vertex problems alike. Routing through v now consumes the one unit of capacity on its internal arc, so two paths sharing v as an intermediate vertex compete for that unit and cannot coexist in any maximum flow. A max-flow from u^{out} to v^{in} in the split network therefore equals the number of internally vertex-disjoint u - v paths in the original graph. In the vertex disjoint variants multiplicity of an edge is irrelevant compared to its existence, so parallel edges never raise κ . Since parallel copies never raise κ , they never break the connectivity constraints either. Counted with multiplicity, the maximum number of edges could simply be infinite and the extremal question would be empty. The multigraph vertex variants are therefore posed on the underlying simple graph, which is why those two rows of Table 3.1 reduce to their simple counterparts and why every figure in this thesis reports the reduced problem for them.

A hyperedge is handled by adding a helper gate, an entry point and an exit point joined by a single arc whose capacity is the multiplicity of the hyperedge. Every member runs into the entry and out of the exit at unlimited capacity, so the one arc between them is the only narrow part (Figure 2.3b). A directed hyperedge is wired the same way, except that only its tails reach the entry and only its heads leave the exit, which is what forces a route to cross it from tail to

head. Every route using the hyperedge must cross that one arc, so the number of routes it can carry is exactly its multiplicity, one for a simple hypergraph. Putting the capacity on the shared arc rather than on each member's own arc is what makes the count right, since capping the members individually would let a hyperedge with several tails carry a separate route for each of them. The hypergraph Menger theorem ([Theorem A.46](#)) proves that the resulting maximum flow is exactly the Berge connectivity. [Figure 2.2](#) shows four gate-disjoint routes in a whole hypergraph. For the vertex-disjoint hypergraph checker, the ordinary vertices are split as before and the helper point is added to the split network with infinite internal capacity.




Figure 2.2. An undirected 3-uniform hypergraph and its gate network for u, v . The four black routes use distinct capacity-one gates (they are drawn as one point to reduce visual clutter), so the checker reads $\lambda(u, v) = 4$. A fifth would reuse a gate. The helper points connect to the hyperedge's third member by dotted lines.

2.2 Checking every graph of a fixed size

The connectivity checker works on all variants but measures only one graph at a time. All graphs of a fixed number of vertices must be enumerated to discover the bound for a single connectivity constraint. A simple graph on n vertices is a choice of $\mu(u, v) \in \{0, 1\}$ on each off-diagonal entry, a multigraph a choice in $\{0, \dots, m-1\}$, and a digraph varies every ordered pair rather than only the upper triangle. A hypergraph is not a matrix at all, so it is enumerated over its own list: a subset of all $\binom{n}{r}$ possible vertex sets of size r . In the directed model, the tail is specified with a pointer and the multihypergraphs have multiplicities for each hyperedge. [Figure 2.5](#) shows how an undirected multigraph and hypergraph are represented. Using these representations for all graph variants means iteration traverses all possible representation states easily. Walking those possibilities, measuring each candidate with the checker and keeping the densest feasible graph settles a bound exactly. If the enumeration hits its deadline before finishing, the returned graph is labelled as a lower bound. On the small cases this exhaustive mode independently reproduces




Figure 2.3. The building blocks of the connectivity checker: vertex splitting, hyperedge gates and directed hyperedge gates.

every value proved by hand in [Chapter 1](#).

Adding vertices multiplies the number of possible graphs by an exponential in n , so a single extra vertex can move a problem from minutes to days. Raising the connectivity budget is the next cost, lifting the base of the exponent rather than its height, which is why the multigraph curves sit above the simple ones in every panel. Direction is the third: a digraph has twice as many possibilities as the undirected graph on the same vertices (and a directed 3-uniform hyperedge, a tail with two heads, has three times as many), so the directed panel climbs fastest of all, and those are precisely the variants whose answers [Chapter 1](#) could not prove. The separation, by contrast, costs nothing here, because edge and vertex disjointness search the same graphs and differ only in what the checker measures, which is an increase linear to the number of graphs.

The number of graphs the checker must visit grows as $2^{\binom{n}{2}}$ for simple undirected graphs and

Number of graphs blind enumeration must visit




Figure 2.4. Candidate counts for blind enumeration on a logarithmic scale. Every model crosses the generous 10^9 budget in single-digit order, and directed objects grow fastest because they have more independent cells to fill.

$2^{n(n-1)}$ for digraphs, and capping multiplicities at $m - 1$, so that each run ranges over the m values $\{0, \dots, m - 1\}$, raises the base from two to m , giving $m^{\binom{n}{2}}$ undirected multigraphs and $m^{n(n-1)}$ directed ones. The enumeration is exact wherever it finishes and it does so only for the smallest sizes. Figure 2.4 measures exactly how fast the search space grows.

2.2.1 Pruned search

The base cases $n \leq 7$ of the directed $m = 2$ theorem (Theorem 1.9) are the finite checks the induction of Appendix A depends on, and they sit just past where blind enumeration of digraphs is tractable. They can be reached by enumerating more cleverly, and the same two prunings serve every variant, since neither mentions direction, multiplicity, arity, or which separation the checker is measuring.

Every variant offers a fixed list of edges its graphs are allowed to use, whether those are pairs, ordered pairs, or oriented sets of r vertices, and a graph is a choice of how many copies each of them gets. Rather than writing out whole choices one after another, fix an order on that list and decide the edges one at a time, leaving the undecided ones absent. Each partly built graph then stands for the whole family of graphs agreeing with it so far, and a pruning discards that family in one step.

Adding an edge never lowers a connectivity, since a family of disjoint routes survives intact when more edges are laid around it (Proposition A.4). Every completion of a partly built graph only ever adds, so the partly built graph is a floor under all of them. The moment it breaks the ceiling, so does every completion, and the branch is abandoned.

The second pruning counts. Take the edges already placed and add the most the undecided ones could ever contribute, which is how many are left times the largest multiplicity allowed. That sum is an upper bound on every completion, so if even it cannot beat the best feasible graph




Figure 2.5. The two encodings. A graph or digraph is the multiplicity matrix μ , which is symmetric when undirected and binary when simple. A hypergraph is a list of vertex sets instead, since a matrix would need n^r entries and be almost all zero. A multihypergraph repeats an entry in that list.

found so far, the branch is dropped unexplored (Figure 2.6).

Only infeasible graphs and graphs that cannot beat the incumbent are ever discarded, so a run that finishes returns exactly the maximum the blind walk would have returned. A regression test pins that by running both side by side over every variant and every size the blind one can still reach. The graph left at the end of a surviving branch also needs no final check, since the last edge it placed was tested against a graph identical to it, every later one being absent.




Figure 2.6. Branch and bound over digraphs. Each level fixes one adjacency. Infeasible partial graphs (red) and branches unable to beat the incumbent (orange) are discarded, allowing the exact search to reach $n \leq 7$.

Run on the directed $m = 2$ problem, the sweep returns the complete sequence 2, 4, 6, 8, 10, 12 for $n = 2, \dots, 7$: each size is proved complete. The densest digraph it finds at each n is exactly the extremal shape named in Chapter 1. For $n < 7$, it is the bidirected star. At the crossover $n = 7$, both the hub and the one-directional bipartite digraph attain the tying value 12. These are exactly the base cases the induction requires, and with them the long proof of Theorem 1.9 rests on a single auditable search rather than on anything done by hand. Table A.1 tabulates that sweep beside its vertex counterpart.

2.2.2 Generating search

One way to improve speed of the enumeration further is a depth-first pruned walk, keeping one representative of each isomorphism class by reducing every finished graph to its canonical relabelling, the smallest matrix in dictionary order. The canonical form keeps the memory small, but the time is spent walking labelled partial graphs, and time is what runs out: the full classification at $n = 5$ costs this in-house search about 456 seconds, and it climbs steeply from there.

The second way, suggested by Prof. Jan Goedgebeur, does not search at all, it generates. It hands the hard part, listing one graph per isomorphism class, to the `geng` program of the `nauty` toolkit of McKay and Piperno [12], invoked through a subprocess and decoded from its `graph6` output. `geng` lists the non-isomorphic *undirected* graphs on n vertices, and the program decorates each one with the directed multiplicities the problem needs, so the isomorphism reduction is inherited for free from the support graph. The decoration walks the support's edges in a fixed order and gives each edge $\{u, v\}$ an ordered pair of multiplicities $\mu(u, v), \mu(v, u)$, each between 0 and $m - 1$ but not both zero, since an edge of the support must carry at least one arc. The checker measures the partly decorated multigraph as it goes, and the moment some pair exceeds the connectivity ceiling the whole branch of assignments is abandoned, since adding further arcs can never lower a connectivity, the same monotonicity prune as Figure 2.6. A decoration surviving to the end with the target arc count is kept and reduced to canonical form, so each isomorphism class is reported once, and different supports share nothing, so they are processed independently across the processor cores. The companion orientation tools `directg` and `watercluster2` are not used, because each gives only one orientation per edge and so cannot express the bidirected arcs and repeated multiplicities the extremal graphs are built from. To be precise about credit: Prof. Goedgebeur is not an author of `nauty`, which is the work of McKay and Piperno cited above. He suggested the generation route, and it is his suggestion that the speed-up rests on.

The two ways agree to the last graph. The generated pipeline reproduces the in-house classification exactly at every settled size, two non-isomorphic extremal families at $n = 4$ and three at $n = 5$, and it runs substantially faster, since the isomorphism work it never repeats is precisely what `geng` has already done. One difference decides which to trust further out: the in-house search leans on a conjectured counting bound once $n \geq 7$, while the generated enumeration prunes only with proved bounds, so it is sound and complete for the parameter range and any explicitly imposed degree cap. That soundness has now paid off. Pointed at the degree-bounded classification at $n = 7$, the generation enumerator completed the sweep in about seven hours across ten processor cores and returned exactly three extremal families, $2 \cdot B_{3,4}$, $2 \cdot B_{4,3}$, and the doubled bidirected path P_7 (Remark A.42).

2.3 Proving every m at once

The pruned search looks at one pair (n, m) at a time. Historically, the directed multigraph checker used a different idea: it rescaled the multiplicities by a factor of $m - 1$ and then checked the

relevant thresholds for a fixed n in one pass. The general theorem of [Theorem 3.3](#) has now replaced this route in `solve`, but the older formulation is still a useful independent finite check.

Divide each multiplicity of a feasible directed multigraph D by $m - 1$. This produces weights $w(u, v) \in [0, 1]$, pairwise maximum flow at most one, and a total weight equal to the arc count of D divided by $m - 1$, as [Figure 2.7](#) illustrates. Thus, with

$$M^*(n) = \max \left\{ \sum_{u \neq v} w(u, v) : w(u, v) \in [0, 1], \text{maxflow}(s, t; w) \leq 1 \text{ for all } s, t \right\}$$

for the largest total weight of such a matrix, scaling gives

$$L_m^{\text{dir}}(n) \leq (m - 1) M^*(n) \quad \text{for every } m \geq 2.$$

The reverse inequality requires an optimum whose entries scale back to integers. The historical optima for $n \leq 6$ were $\{0, 1\}$ double stars, so there

$$L_m^{\text{dir}}(n) = (m - 1) M^*(n) \quad \text{for every } m \geq 2,$$

for every m . The hand proof now establishes this identity for every n ([Corollary A.38](#)). The finite optimisation is retained to record the independent route by which the cases $n \leq 6$ were first checked.




Figure 2.7. Scaling at $m = 3$: dividing multiplicities by $m - 1 = 2$ turns the integer multigraph into a weight matrix with entries in $[0, 1]$ and pairwise maximum flow at most one. Hence one m -free optimisation bounds every threshold.

By max-flow/min-cut, $\text{maxflow}(s, t) \leq 1$ exactly when some s – t cut has capacity at most one. The optimisation therefore chooses one cut per ordered pair. A binary label x_u^{st} records its side, and p_{uv}^{st} counts $w(u, v)$ only when $u \rightarrow v$ crosses from source to sink ([Figure 2.8](#)). This describes the exact feasible region. The archived wrapper retained an OPTIMAL status and primal witness but no independently replayable matching bound, so the runs are finite solver checks rather than formal certificates.

With scaled weights w , binary side labels x , and non-negative crossing weights p , the model is

$$\begin{aligned} & \text{maximise} && \sum_{u \neq v} w(u, v) \\ & \text{subject to} && x_s^{st} = 1, \quad x_t^{st} = 0 && \text{for all } (s, t), \end{aligned}$$

$$\begin{aligned}
p_{uv}^{st} &\geq w(u, v) + x_u^{st} - x_v^{st} - 1 && \text{for all } (s, t), (u, v), \\
\sum_{u \neq v} p_{uv}^{st} &\leq 1 && \text{for all } (s, t), \\
w(s, t) + \sum_x \min(w(s, x), w(x, t)) &\leq 1 && \text{for all } (s, t), \\
d(v_0) \leq d(v_1) \leq \dots \leq d(v_{n-1}), &&& \text{with } d(v) = \sum_{u \neq v} (w(u, v) + w(v, u)), \\
w \in [0, 1], \quad x \in \{0, 1\}.
\end{aligned}$$

The first line is what makes the labelling an s – t cut rather than an arbitrary bipartition, and it is not optional. Without it the all-zero labelling $x \equiv 0$ is admissible, no ordered pair (u, v) ever crosses, every p_{uv}^{st} rests at 0, and the cut constraint below is satisfied by every weight matrix whatsoever. Pinning the source side of s and the sink side of t removes that escape and leaves the remaining $n - 2$ labels free, which is precisely the choice of an s – t cut. The implementation supplies these two values as constants rather than as variables (program/erdos915_unified.py, in `_cut_counting_model`), which is the same restriction imposed one step earlier.

Because $w \leq 1$, the second constraint forces $p_{uv}^{st} \geq w(u, v)$ only in the crossing case $(x_u^{st}, x_v^{st}) = (1, 0)$. In the other three cases its right-hand side is non-positive. The third constraint then caps the chosen cut at one. The remaining two lines are redundant accelerators: the first counts the direct arc and all two-step detours, and the second removes equivalent vertex relabellings.

The `min` in the two-step line is written as a minimum for readability only. A linear program cannot take one directly. It is linearised in the usual way, by an auxiliary variable $z^{sxt} \in [0, 1]$ held below both arguments, $z^{sxt} \leq w(s, x)$ and $z^{sxt} \leq w(x, t)$, together with a binary selector b^{sxt} and the two big- M inequalities $z^{sxt} \geq w(s, x) - (1 - b^{sxt})$ and $z^{sxt} \geq w(x, t) - b^{sxt}$, which force z^{sxt} up to whichever argument is the smaller. The displayed `min` is then the value of z^{sxt} .




Figure 2.8. One chosen s – t cut. Only the two thick source-to-sink arcs contribute crossing weight $p = w$, while same-side and backward arcs have $p = 0$.

Theorem 2.1 (Directed multigraph value, small n). *For $n \leq 6$, $M^*(n) = 2(n - 1)$, and the optimum is attained by a $\{0, 1\}$ weight matrix. Hence for all $m \geq 2$, $L_m^{\text{dir}}(n) = 2(n - 1)(m - 1)$, attained by the double star.*

Historical runs gave $M^*(n) = 2(n - 1)$ with status OPTIMAL for $3 \leq n \leq 6$, and $n = 7$ did not close. Their archived output lacks a separately checkable bound or gap, and no theorem now depends on it: [Appendix A](#) proves the value for all n, m by hand and [Section A.10](#) retains the solver record.

2.4 The heuristic search

We want, for fixed n and m , a graph with as many edges as possible subject to $\lambda^{\max} \leq m - 1$ (or $\kappa^{\max} \leq m - 1$ for the vertex variants). The space of graphs on n vertices is enormous and its feasible region has no helpful shape: adding an edge can be harmless until, together with edges added long before, it suddenly creates a forbidden pair. A greedy walk that only ever accepts an improving move stalls at the first such collision, in a graph that is locally maximal but globally far from extremal.

Every graph is scored by one number, its energy

$$E(G) = -|E(G)| + \Lambda \cdot \max(0, \lambda^{\max}(G) - (m - 1)),$$

which rewards edges and penalises any excess connectivity. A move toggles one adjacency, and the search walks from the empty graph by repeatedly taking the move that lowers the energy most. The penalty is a soft constraint rather than a hard one, so the walk may pass briefly above the ceiling instead of being walled inside the feasible region.

The method used is *tabu search* [13], which escapes a local optimum by forbidding the pairs it has just touched. After each move the pair it changed is put on a tabu list for a fixed number of steps, so the walk cannot immediately undo what it just did and cycle back into the optimum it came from. A move that would beat the best graph seen so far overrides the tabu flag, and the walk perturbs itself when it stalls.

The obvious alternative is *simulated annealing*, which escapes local optima instead by sometimes accepting a worsening move, less and less often as a temperature parameter falls. On the same energy and the same neighbourhood, at equal wall-clock budget, it is the weaker engine here. The two tie on the undirected problems, and on the directed ones, which are the hard cases, tabu wins outright: at $n = 7$, $m = 3$ it reaches the 24-arc directed multigraph extremiser where annealing stalls at 20, and 18 arcs against 16 on the simple digraph. Every discovery value this thesis reports therefore comes from tabu search. Annealing is kept only as an independent cross-check in the program's self-test, where agreement between two unrelated walks is worth more than the speed of either.

It is fair to ask why the walk is allowed above the ceiling at all. Reaching every feasible graph does not require it. Removing an edge can only lower $\lambda^{\max}$ and $\kappa^{\max}$, never raise them, so every subgraph of a feasible graph is again feasible, and the empty graph is feasible trivially. From any feasible target we can therefore delete edges one at a time down to the empty graph without ever crossing the ceiling, and reading that path backward builds the target up the same way. The feasible region is connected through the empty graph by single-edge moves that stay inside it, so

a walk that never went infeasible could in principle visit every feasible graph there is.

We let it go infeasible anyway, because reachability and efficiency are different things. A feasible graph can be locally maximal, meaning every edge we might add creates a forbidden pair, and yet still sit far below the densest feasible graph. To improve such a graph the walk has to add an edge and then remove a different one, a swap whose first half is infeasible. A walk confined to feasible graphs could only find that better graph by retreating most of the way back to the empty graph and climbing a different slope, a detour far longer than the budget allows. Allowing a brief excursion above the ceiling lets the walk trade one load-bearing edge for a better one in two moves rather than dozens. Going above the bound is never the goal. It is the shortcut the walk takes on its way back down to it.

2.5 Edge sensitivity

Which edges a walk should leave alone is itself a measurement. For an edge e define its *sensitivity*

$$\sigma(e) = \lambda^{\max}(G) - \lambda^{\max}(G - e),$$

the amount by which deleting e relaxes the binding connectivity. A load-bearing edge sits on a tightest cut and has $\sigma(e) > 0$: removing it would drop $\lambda^{\max}$ and so loosen the very constraint that limits how many edges the graph may hold. A free edge has $\sigma(e) = 0$, so it can be removed without easing the binding pair, which usually means it can be put to better use elsewhere. The program measures this by deleting e and letting the checker remeasure, reusing one shared baseline across a whole graph rather than recomputing it per edge.

Colouring a graph's arcs by σ then explains a finished graph at a glance. [Figure 2.9](#) shows all four sensitivity levels on one directed multigraph: the darkest arcs carry the most flow and lower $\lambda^{\max}$ the most when removed, while the cool arc contributes nothing. It makes the tightness of a witness visible, which is what turns a bare edge count into something a reader can check the shape of.

A single walk depends on where it started, so the search is repeated with a different seed until the wall-clock budget runs out and the densest witness of all the runs is kept. Independent restarts share nothing, so they can also be spread across processor cores.

2.6 Keeping the checker cheap inside the search

The energy above reads the binding connectivity $\lambda^{\max}(G)$, and a literal implementation would call the checker afresh on the whole graph at every step. That checker is the most expensive routine in the program: a capped sweep over ordered pairs, stopping each pair as soon as it finds one route too many and stopping the whole sweep at the first violating pair. The walk runs for thousands of steps with several restarts, so such a search would spend nearly all of its time remeasuring graphs that barely changed from the step before. Two elementary facts remove almost all of that work without altering a single number the search reports, and they are the monotonicity and unit-step bounds of [Proposition A.4](#). Removing a unit of multiplicity can never

Parallel arcs drawn explicitly, coloured by sensitivity σ :
cool ($\sigma = 0$, free) $\rightarrow$ warm (load-bearing)




Figure 2.9. Arc sensitivity in a directed multigraph with $\lambda^{\max} = 9$. Equal multiplicities can have different σ : the downstream bottleneck makes $s \rightarrow a$ cooler than $s \rightarrow b$, while arcs into the dead end d have $\sigma = 0$.

raise $\lambda^{\max}$, part (1), and adding one can never lower it and raises it by at most one, part (2). Both hold verbatim for $\kappa^{\max}$, and both fall straight out of max-flow / min-cut, so the proof sits in [Appendix A](#) with the rest of the machinery.

Together they settle almost every step without a flow. The energy depends on $\lambda^{\max}$ only through the excess $\max(0, \lambda^{\max} - (m - 1))$, which is zero exactly when the graph is feasible. So a removal applied to a feasible graph leaves it feasible by part (1), with excess still zero, and needs no measurement at all. An addition made while the graph sits strictly below the bound, at $\lambda^{\max} \leq m - 2$, also stays feasible. Part (2) caps the rise from a single added unit at one, so the graph can climb at most to $\lambda^{\max} = m - 1$, which is still inside the feasible region. The excess is zero again, and again no measurement is needed. Only an addition made right at the boundary $\lambda^{\max} = m - 1$ can tip the graph over, and only there does the search run a capped pairwise sweep, stopping at the first pair for which it finds one route too many. The exact connectivity value is recomputed only on the rare step whose proposal is genuinely infeasible, where the penalty term needs it.

To know which case applies, the search carries a running upper bound on $\lambda^{\max}$, raised by one whenever it accepts an addition (part (2)) and left untouched when it accepts a removal (part (1)), and resynchronised to the exact value at the cadence where it already pays for flows to refresh the sensitivity map. The bound can only ever sit at or above the truth, so any step it clears is certainly feasible.

The energy returned on every step is, by [Proposition A.4](#), exactly the value the naive implementation would have computed. Every accept-or-reject decision, every random draw, and the graph the walk finally returns are therefore unchanged, and the speed-up is invisible to every number the thesis reports. A regression test runs both implementations, the fast one and a reference that measures exactly at every step, across all four graph variants and both separations, and checks that the two agree step for step in energy, in every accept-or-reject decision, and in the final graph, and that the returned graph is feasible under the exact checker. The witness the search reports is always certified exactly, either by the monotonicity deductions above or by the capped predicate, and the full binding-connectivity value is recomputed only when the energy or a recorded trace needs it.

Where the method is needed differs by variant. For undirected graphs one could resynchronise the exact value cheaply from the Gomory–Hu shortcut used above, which encodes every pairwise edge-connectivity in $n - 1$ flows. The directed and vertex variants admit no such tree, so there the monotone bound and the capped predicate are the only inexpensive handle on feasibility, and they are what let the same matrix-search machinery serve every graph variant at speed.

2.7 Rediscovering the known extremisers

Before the search is pointed at the open cases, it is run on the settled ones. Run from the empty graph on the cases settled in [Chapter 1](#) or by the exact methods above, and told only to pack in edges without breaching the connectivity ceiling, the cooled search arrives unprompted at the named constructions and matches the established value in every tested case. [Table 2.1](#) reports the densest graph it found against the value the theory predicts.

Construction	n	m	Search	Theory
spanning tree (simple undirected, $m=2$)	6	2	5	5
multiplicity- $(m-1)$ star (multi undirected, $m=3$)	6	3	10	10
bidirected star (simple directed, $m=2$)	4	2	6	6
bidirected star (simple directed, $m=2$)	6	2	10	10
hub–bipartite tie (simple directed, $m=2$)	7	2	12	12
double star (multi directed, $m=3$)	4	3	12	12
double star (multi directed, $m=3$)	5	3	16	16

Table 2.1. Validation by rediscovery. For each variant the temperature search, run from scratch with a handful of restarts, recovers exactly the extremal value proved or certified earlier. The “Search” column is a lower bound found by annealing, and the “Theory” column is the proved or certified value.

The graphs themselves are the named constructions, not only graphs of the right size. The simple undirected run at $m = 2$ settles on a spanning tree, the only shape with $n - 1$ edges and no cycle. The undirected multigraph run settles on a star with every edge at multiplicity $m - 1$, an instance of the multi-tree extremiser behind [Theorem 1.3](#). The directed multigraph runs settle on the double star of [Construction 1.7](#), both directions of every spoke at full multiplicity. The simple directed runs find the bidirected star at $n = 4$ and, at $n = 7$, land on twelve arcs, the value at

which the hub and the one-directional wall tie, exactly the crossover [Theorem 1.9](#) predicts. The cooled search does not know these names. It rediscovers the structures because, under the connectivity ceiling, there is nowhere denser to go.

Some extremisers are past the size at which a quick search lands reliably, but the exact checker built above settles them immediately by measuring their connectivity. The one-directional bipartite digraph on eight vertices has 16 arcs at $\lambda^{\max} = 1$, and 16 already exceeds the linear hub value $2(n - 1) = 14$, so this is the smallest size at which the quadratic phenomenon strictly wins. It is also where the search reaches its limit: repeated cooling runs at $n = 8$, $m = 2$ stall at the hub value of 14, because dismantling a finished hub and rebuilding a one-directional wall is far more than the single-arc moves of the walk can tunnel through. The known answer there comes from the construction and the checker rather than from the search. The augmented bipartite digraph on ten vertices at $m = 3$ has 30 arcs at $\lambda^{\max} = 2$, the thirty-arc counterexample of [Construction 1.15](#), confirmed pair by pair against the naive bound of 27 it refutes. The directed double star on seven vertices at $m = 4$ has 36 arcs at $\lambda^{\max} = 3$, matching $2(n - 1)(m - 1)$. None of these is a proof of optimality. Each is an exactly measured feasible graph, and therefore a lower bound.

The discovery driver defaults to *tabu search*, Glover’s method [13] of forbidding the recently visited moves so a best-improving walk cannot cycle back into the local optimum it just left. It uses the same energy and single-adjacency neighbourhood as the annealer but takes best-improving moves with a short tabu tenure and occasional perturbations. Simulated annealing remains an independent cross-check. Both return checked witnesses and therefore lower bounds, never proofs.

2.8 Results and evidential status

The checker *measures* a supplied graph exactly. Completed exhaustion *proves* a finite optimum. The historical cut-counting optimisation is an independent finite check unless accompanied by a replayable bound, and heuristic search supplies only a witness and lower bound. A value is settled only when its lower and upper bounds meet. Every computational claim in the chapters and in [Table 3.1](#) reruns from `program/erdos915_unified.py`, and critical values are cross-checked by independent implementations. Two block sweeps quoted in [Appendix A](#) are the exception and run from their own standalone scripts in `research_notes/scripts/`. They were written to corroborate the main program rather than to share code with it, and each is named where its numbers are used. [Section A.10](#) records the environment, commands, and audit evidence. No general theorem rests on an unauditable solver status.

The directed simple arc problem at $m \geq 3$ shows where the settled range ends. Exhaustion proves the values 6, 9, 12, 15 for $n = 3, 4, 5, 6$ at $m = 3$ under both separations ([Remark A.20](#)), but does not finish at $n = 7$. Beyond that point the best reported lower bound is the better of the

search and a named construction. The conjectured formula

$$\ell_m^{\text{dir}}(n) = \max\left(m(n-1), \left\lfloor \frac{(n+m-2)^2}{4} \right\rfloor\right) \quad (n \geq m)$$

passes through every exhausted value and through the construction-backed lower bounds beyond them. Its linear-to-quadratic crossover occurs at $n = 9$ when $m = 3$, just past the exhausted range, and at $n = 10$ it gives the thirty-arc counterexample of [Construction 1.15](#).

The paired grids of [Figures 2.10 to 2.13](#) apply that reading to all sixteen variants. Blue curves are theorems, red curves conjectures, and yellow curves interpolate lower bounds, while filled squares are exact finite values and open circles are witnesses. Directed vertex values were checked separately because Whitney's inequality transfers witnesses but not upper bounds.

Where an upper bound is known to be attainable, the search reaches it at every plotted size except the recorded directed $m = 2, n = 8$ stall. The undirected hypergraph rows carry the one systematic exception, and it is a feature of the bound rather than of the search. There the blue curve is [Proposition 1.11](#), an upper bound proved for every r -uniform hypergraph, but the two models attain it under different conditions: a simple hypergraph only when $m - 1 \leq \binom{n-2}{r-2}$, a multihypergraph only when $(r - 1) \mid (n - 1)$. Below its own threshold neither is known to meet the curve, and for the simple model at $m = 6, n = 6, r = 3$ nothing does: the curve reads 12, yet all 125 970 twelve-hyperedge simple 3-uniform hypergraphs on six vertices are infeasible and the true maximum is 11. The circles there are the search's own witnesses and sit below the curve, so the gap is not closed. The two rows part company, which is why they are kept separate, and the smallest case is $n = r = m = 3$, where a simple hypergraph carries one hyperedge and the multihypergraph carries two. On the open directed sizes the named constructions, not the search, carry the best bounds once the extremal shape changes from a hub to a bipartite wall. The grid therefore checks the search against proof wherever proof is available, and marks the remaining values as evidence rather than conclusions.

Erdős 915 across the sixteen variants, $m = 3$ · simple and multigraph models




Figure 2.10. The eight graph-model variants at $m = 3$: simple and multigraph rows, undirected and directed columns. A blue curve is a theorem, a red one a conjecture and a yellow one an interpolation, filled squares are exact finite values, and open circles are witnesses. The dotted line named on the simple directed arc panel is the hub branch of [Conjecture 1.14](#). Its bipartite branch is the larger of the two beyond the crossover, so past that point it runs underneath the red curve. [Figure 2.11](#) continues with the hypergraph rows at the same m , and [Figures 2.12](#) and [2.13](#) redraw both halves at $m = 6$.

Erdős 915 across the sixteen variants, $m = 3$ · hypergraph and multihypergraph models




Figure 2.11. The eight hypergraph-model variants at $m = 3$, continuing [Figure 2.10](#): hypergraph and multihypergraph rows, undirected and directed columns, same colour and marker key.

Erdős 915 across the sixteen variants, $m = 6$ · simple and multigraph models




Figure 2.12. The same eight graph-model variants at $m = 6$, to be read against Figure 2.10. Raising m widens every feasible band, and the exact squares thin out as the enumeration range falls behind the constructions.

Erdős 915 across the sixteen variants, $m = 6$ · hypergraph and multihypergraph models




Figure 2.13. The same eight hypergraph-model variants at $m = 6$, continuing Figure 2.12 and to be read against Figure 2.11.

Chapter 3

Synthesis, Results, and Open Problems

The thesis began with one theorem and carried it across sixteen variants. Some of them yielded to hand argument, the machine settled the small cases of others, and the search supplied dense graphs where neither reached. This chapter collects the results. It describes the directed frontier, where neither the proofs nor the certificate reach the full answer, sets out two phenomena that run through the variants, gathers the sixteen values into one table, and states what the program does and does not do.

3.1 The directed frontier and the missing decomposition

The directed arc problem is a race between a hub with $m(n-1)$ arcs and an augmented bipartite construction with $\lfloor (n+m-2)^2/4 \rfloor$ arcs ([Constructions 1.7](#) and [1.15](#)). They cross near $n = 9$ at $m = 3$, and [Conjecture 1.14](#) asserts that their maximum is exact. Exhaustion proves the small $m = 3$ cases, but the general upper bound is structural: it would follow if every non-hub extremiser admitted the complete $A \rightarrow B$ decomposition of [Conjecture A.25](#). The difficult clause forbids arcs returning from B to A , since one such arc lets routes escape through A and defeats the internal $m-2$ budget. [Figure 3.1](#) places the numerical crossover and that obstruction together.

The conjecture is nevertheless correct to first order and to the order of its error, and that much is a theorem with no hypothesis at all on the shape of the extremiser.

Theorem 3.1 (The quadratic bound with a linear error). *For all $m \geq 2$ and $n \geq 2$, every simple digraph D on n vertices with $\lambda_D^{\max} \leq m-1$ satisfies*


$$|A(D)| \leq \left\lfloor \frac{n^2}{4} \right\rfloor + 4(m-1)(n-1),$$

so for each fixed m ,


$$\ell_m^{\text{dir}}(n) = \frac{n^2}{4} + \Theta_m(n) \quad (m \geq 3),$$

and, for $m \geq 3$, both the leading constant $\frac{1}{4}$ and the order of the second-order term of [Conjecture 1.14](#) hold unconditionally. At $m = 2$, [Theorem 1.9](#) instead gives $\ell_2^{\text{dir}}(n) = n^2/4 + O(1)$ as $n \rightarrow \infty$.

The proof is in [Appendix A](#). Its engine is a count of two-step routes: in a feasible digraph a vertex cannot have too many out-neighbours that reach one another, since the short detours



(a) The two lower-bound branches at $m = 3$.



(b) Left, the desired $A \rightarrow B$ form. Right, one returning arc that breaks it.

Figure 3.1. The directed frontier in two views: the hub-to-wall crossover and the backward-arc obstruction to proving the wall optimal.

through them would breach the ceiling, and that alone forces the arc count down to the wall plus a linear correction. What is left open is therefore only the coefficient of that linear term, somewhere between the conjectured $(m-2)/2$ and the proved $4(m-1)$. The next section takes up what the bound does and does not say about the *shape* of an extremiser, which is a separate question answered by a weaker version of the same count.

The route family the count uses is internally vertex-disjoint and not merely arc-disjoint, which is what lets the whole argument be re-run under the stricter separation. Whitney's inequality alone would not justify the transfer, since it points the other way.

Theorem 3.2 (Directed vertex problem, leading constant and order). *For all $m \geq 2$ and $n \geq 2$,*

every simple digraph D on n vertices with $\kappa_D^{\max} \leq m - 1$ satisfies

$$|A(D)| \leq \left\lfloor \frac{n^2}{4} \right\rfloor + 4(m-1)(n-1),$$

and consequently, for each fixed m ,

$$k_m^{\text{dir}}(n) = \frac{n^2}{4} + \Theta_m(n) \quad (m \geq 3).$$

So the directed vertex problem, which [Appendix A](#) otherwise keeps separate from the arc one, has the same leading term and the same order of error. Whether the two agree exactly for $m \geq 3$ is open.

Allowing parallel arcs, by contrast, closes the problem outright.

Theorem 3.3 (Directed multigraph value, every n and m). *For all $n \geq 2$ and $m \geq 2$,*

$$L_m^{\text{dir}}(n) = (m-1)M(n), \quad M(n) := \max\left(2(n-1), \left\lfloor \frac{n^2}{4} \right\rfloor\right).$$

The extremisers include the doubled trees on the linear branch and the balanced one-way bipartite multigraph on the quadratic one. On the linear branch these are constructions exhibited rather than a classification, and [Appendix A](#) says explicitly that not every linear-branch extremiser has been classified. The proof deletes a *subgraph* at a time rather than a vertex. Inside any feasible multigraph there is a sparse skeleton that alone preserves which vertices reach which, and removing one copy of each of its arcs costs every pair at least one route. Peeling off $m - 1$ such skeletons therefore accounts for every arc, and the parity obstruction that stopped the vertex-deletion induction never arises. Equality also classifies the quadratic extremisers once $n \geq \max(8, 2m)$ ([Theorem A.45](#)), and only the smaller range remains open.

3.2 Two principal phenomena

Two phenomena run through the sixteen variants.

The first is the divergence at $m = 5$. The edge and vertex problems agree for $m \leq 4$, and it would be natural to expect them to agree forever. Instead they part company at the very next value, with $k_5(n) = \lfloor 8n/3 \rfloor - 4$ for every $n \geq 6$ except the two sizes $n = 7$ and $n = 12$ that the paper settles separately, pulling strictly above $\ell_5(n)$ from $n = 14$ onwards ([Theorem 1.5](#), and [Remark 1.6](#) for the counting convention that fixes the constant). The separation is a fact about m rather than about n : at $m = 5$ the two values still agree at every size up to $n = 13$ and only then come apart, and beyond $m = 5$ the vertex problem has stayed open since 1974.

The second is the quadratic bipartite phenomenon. In an undirected graph every edge contributes to the degree of both its endpoints, and Mader's theorem turns that symmetry into a hard ceiling of $\lfloor m(n-1)/2 \rfloor$ edges, linear in n . A digraph breaks the symmetry. The one-directional wall of [Construction 1.8](#) sends every arc from one half of the vertices to the other and holds

$\lfloor n^2/4 \rfloor$ of them at $\lambda^{\max} = 1$, because a route must cross the wall exactly once and nothing crosses back to offer a second. An arc count growing like n^2 while the connectivity stays at one is something no undirected graph can do past a handful of vertices, and it is why the initial linear conjecture for the directed case fails already at $m = 2$, where on $n = 8$ the wall holds 16 arcs against the hub's 14. [Construction 1.15](#) carries the same idea to every m by letting each vertex of the larger part absorb $m - 2$ extra in-arcs from inside that part, which buys $m - 2$ short detours per pair and lifts $\lambda^{\max}$ to $m - 1$ while the count climbs to $\lfloor (n + m - 2)^2/4 \rfloor$. Its $m = 3, n = 10$ instance holds thirty arcs against the naive prediction of twenty-seven.

The quadratic term is what the directed upper bound has to account for. The argument must explain why no configuration beats the bipartite wall, which requires producing the partition in the first place, ruling out backward arcs, and controlling the internal structure of the large part B . That structural task is what [Conjecture A.25](#) has not yet resolved for $m \geq 3$. [Proposition A.26](#) and [Theorem 3.1](#) show that the wall is the right shape, and they say it in two different terms. In *size*, the extremal value lies within $O_m(n)$ of the wall's count, so what is left open is a constant inside the second-order term rather than the phenomenon. In *shape*, every feasible digraph becomes a subgraph of a one-directional bipartite graph after deleting $O(\sqrt{m} n^{3/2})$ arcs, a weaker error term because it comes from the count that does not induct on a low-degree vertex. The undirected problem has no analogous obstacle, because Mader's degree-counting argument closes it in a few lines. The quadratic phenomenon is therefore the reason the directed and undirected problems differ in difficulty.

3.3 Summary of the sixteen variants

[Table 3.1](#) records the status of all sixteen structural variants. Each entry is marked by how it is known: proved by hand here or in [Appendix A](#), cited to the literature, conjectured with a proved lower bound, or, for the two rows where only the leading term is settled, proved asymptotically. No row now rests on a machine certificate alone. That category was live in earlier drafts, when the directed multigraph value was known only for the sizes the cut-counting optimisation could close, and [Theorem 3.3](#) retired it. The markings follow the standard set in [Chapter 2](#). [Figure 3.2](#) shows the same distinction as a picture before the table spells out every case.

Here $M(n) = \max(2(n - 1), \lfloor n^2/4 \rfloor)$. Simple-graph formulas assume $n \geq m$. Below that range the complete object gives the trivial maximum. The edge-hypergraph bound is attained when $(r - 1) \mid (n - 1)$ for multihypergraphs, and for simple hypergraphs when $r \geq 3$ and $m - 1 \leq \binom{n-2}{r-2}$. The $m = 3$ vertex-hypergraph bound is attained when $r \geq 3$ and $2 \leq \binom{n-2}{r-2}$. The two multigraph vertex rows use the counting convention of [Section 2.1.1](#). Under the other convention, where parallel copies count as distinct routes, the multigraph vertex question becomes a separate extremal problem, studied in [Section A.9](#) and listed among the open problems below. Full statements and exceptions are in [Theorems 1.2, 1.4, 1.5, 1.12, 1.13, A.48](#) and [3.1](#) to [3.4](#) and [proposition 1.11](#).




Figure 3.2. Status of the sixteen variants. Blue marks a proved closed form in the ranges stated in Table 3.1, red a conjectured exact formula with proved lower bound, and amber a proved leading term with unknown exact value. The table is authoritative for exceptions and conventions.

3.3.1 Orientation models for the directed hypergraph

The directed hypergraph row of Table 3.1 rests on one bound, which pins the leading term where previously only a factor-of-four window was known.

Theorem 3.4 (The directed hypergraph constant). *Let $r \geq 2$ and $m \geq 2$. Every forward directed r -uniform multihypergraph $\mathcal{H}$ on $n \geq 2$ vertices with $\kappa_{\mathcal{H}}^{\max} \leq m - 1$ satisfies*

$$|E(\mathcal{H})| \leq \frac{m-1}{r-1} \left\lfloor \frac{n^2}{4} \right\rfloor + 4(m-1)^2(n-1).$$

The same bound therefore holds under $\lambda_{\mathcal{H}}^{\max} \leq m - 1$, which is the stronger hypothesis by $\kappa \leq \lambda$. Hence for fixed r and m both the edge- and the vertex-disjoint maxima are $(1 + o(1))(m - 1)n^2 / (4(r - 1))$, and the bipartite construction of Proposition A.55 is asymptotically optimal for both.

The proof, in Appendix A, forms the one-step shadow digraph whose arc $u \rightarrow v$ records that some hyperedge has tail u and head v . A matching argument transfers the hypergraph route ceiling to the shadow's two-step route budget, after which Lemma A.30 controls the shadow and a tail-head incidence count converts its arcs back to hyperedges. Since the bipartite construction already attains that leading term, it is asymptotically optimal.

The table uses the *forward* convention of one tail and $r - 1$ heads. More generally a hyperedge may split into non-empty tail and head sets. Reversal makes the one-head *backward* model exactly equivalent to forward (Lemma A.56), while Theorem A.59 proves that arbitrary splits share the same leading term under both separations. Thus the orientation choice does not add another asymptotic row to Table 3.1.

Finite values may differ when vertices are scarce. Whether forward and general agree exactly once $n > r$ remains open, and the thesis uses the forward convention throughout its sixteen-variant summary.

Model	Edge/arc-disjoint	Vertex-disjoint
undirected simple graph	$\lfloor m(n-1)/2 \rfloor$, exact	edge value for $m \leq 4$, $\lfloor 8n/3 \rfloor - 4$ for $m = 5$, $n \neq 7, 12$, open for $m \geq 6$
undirected multigraph	$(m-1)(n-1)$, exact	simple value, adjacency-count convention
undirected r -hyper, simple	$\leq \lfloor (m-1)(n-1)/(r-1) \rfloor$, equality when $r \geq 3$ and $m-1 \leq \binom{n-2}{r-2}$	$\lfloor (n-1)/(r-1) \rfloor$ for $m = 2$, $\leq \lfloor 2(n-1)/(r-1) \rfloor$ for $m = 3$, same equality range
undirected r -hyper, multi	the same bound, equality when $(r-1) \mid (n-1)$	the same bounds, with repeats never helping at $m = 2$ and closing cases the simple model misses at $m = 3$
directed simple graph	$M(n)$ for $m = 2$, $n^2/4 + \Theta_m(n)$ for $m \geq 3$, exact then conjectured	$M(n)$ for $m = 2$, $n^2/4 + \Theta_m(n)$ for $m \geq 3$, exact equality with the arc value open
directed multigraph	$(m-1)M(n)$, exact	simple-digraph value, adjacency-count convention
directed r -hyper, simple	$(1+o(1))(m-1)n^2/[4(r-1)]$ for fixed r, m , leading term exact	same leading term for fixed r, m
directed r -hyper, multi	same leading term, the bound being proved for multihypergraphs	same leading term for fixed r, m

Table 3.1. The sixteen variants and the strongest result known for each.

3.3.2 The multigraph vertex problem under the other convention

One variant of Table 3.1 takes a different form under the other convention and is stated separately here, because under that convention the extremal problem becomes a packing problem. When parallel copies count as distinct routes, the multigraph vertex question is a different problem, written $K_m^{\text{multi}}(n)$ and studied in Section A.9. Its answer is additive over the blocks of the underlying simple graph, because adjacent vertices always share a block and no route between them ever leaves it.

Theorem 3.5 (The multigraph vertex problem is a knapsack over blocks). *For a 2-connected graph B , or a single edge, write*

$$g_m(b) := \max \{ W_m(B) : B \text{ 2-connected or an edge, } |V(B)| = b, \kappa_B^{\max} \leq m-1 \},$$

with $g_m(b) := -\infty$ when no such B exists. Then for all $m \geq 2$ and $n \geq 2$,

$$K_m^{\text{multi}}(n) = \max \left\{ \sum_i g_m(b_i) : b_i \geq 2, \sum_i (b_i - 1) \leq n - 1 \right\},$$

the maximum over all finite multisets of block sizes. In particular the value is determined by the finitely many numbers $g_m(2), \dots, g_m(n)$.

The consequence is a change in what one has to search. Instead of graphs on n vertices

with n growing, the question is settled by knowing the best 2-connected block of each size, and a knapsack then assembles them. That closes every value up to $n \leq 8$ and $m \leq 8$ by machine. Within that computed range, for $m \geq 5$ and $n \geq 3$, a single 2-connected block beats every way of splitting the vertices, so the bouquet of cliques is not the shape of the answer. The best blocks are dense and bipartite-like rather than complete, and the explicit complete-bipartite family gives order $\Theta(m^2n)$ as $m \rightarrow \infty$ with $n/m \rightarrow \infty$, with a per-vertex constant between $3 - 2\sqrt{2}$ and 2 ([Theorem A.65](#) and [proposition A.63](#)). Closing that gap is the open problem [Table 3.2](#) records.

3.4 Contributions and limitations of the program

The program turned a per-case craft into a uniform pipeline. Before it, each variant came with its own one-off computation and its own hand argument. After it, one interface holds every variant over two internal representations, a multiplicity matrix for the graph models and vertex sets for the hypergraph ones, one exact checker measures connectivity, one certifier closes small-case upper bounds, and one cooled search discovers extremisers and the lower bounds they witness. The rediscovery of [Section 2.7](#) is the evidence that the pipeline is faithful: pointed at solved cases, it returns the solved answers.

The program has the following limits. The search proves no upper bounds. It only ever exhibits graphs. The certifier closes upper bounds, but only for a fixed size, and only as a finite solver check rather than a replayable certificate ([Section 2.8](#)). The directed multigraph encoding closes in that sense only up to $n = 6$. Past there the value is settled instead by the hand proof of [Appendix A](#), and the certifier contributed nothing at $n = 7$, where it exhausted its time limit without a verdict. The generation enumerator's degree-capped classification at $n = 7$ ([Remark A.42](#)) stands only as a machine-verified fact about the extremisers of maximum total degree at most 8, and it is not an unrestricted classification. Neither replaces the structural gaps the thesis leaves open, and the next section lists them in full. Three of them are not reachable by computation. The decomposition of [Conjecture A.25](#) would turn [Conjecture 1.14](#) into a theorem, and its backward-arc clause is one part of it rather than the whole. The directed vertex problem at $m \geq 3$ is not handed over by Whitney's inequality together with the arc case. The undirected vertex-disjoint problem at $m \geq 6$ has been open for half a century, and the program can probe it but not settle it.

3.5 Open problems

[Table 3.2](#) separates what this thesis proves from what it leaves open, one row per frontier. Each row states the strongest thing known here and the next step, so that a reader who wants to continue knows which statement to attack.

The map is filled in where the arguments reached, and the blanks are marked where they did not. The program does not answer every question. It records which questions are left and the form each of them now takes.

Problem	Known here	Remaining target
Directed simple arc, $m \geq 3$	$\ell_m^{\text{dir}}(n) = n^2/4 + \Theta_m(n)$ (Theorem 3.1)	Prove the decomposition of Conjecture A.25, or otherwise determine the linear coefficient.
Directed simple vertex, $m \geq 3$	$k_m^{\text{dir}}(n) = n^2/4 + \Theta_m(n)$ (Theorem 3.2)	Decide whether its exact value equals the arc value, given that an arc upper bound does not transfer.
Directed multigraph classification	Exact value for all n, m and quadratic-branch uniqueness for $n \geq \max(8, 2m)$	Classify the range $n < 2m$ and the remaining linear-branch extremisers.
Undirected simple vertex, $m \geq 6$	Block formulation and $m/2 < c_m \leq 2(m-1)$, where $c_m = \lim_n k_m(n)/(n-1)$ (Theorem A.13)	Determine c_m , whose likely next structure is triconnected rather than merely blockwise.
Multigraph vertex, parallel copies distinct	Block-knapsack formulation and order $\Theta(m^2n)$ as $m \rightarrow \infty$ with $n/m \rightarrow \infty$, with the per-vertex constant between $3 - 2\sqrt{2}$ and 2 (Theorems A.65 and 3.5 and proposition A.63)	Determine the asymptotic constant and the optimal 2-connected blocks.
Hypergraph vertex, $m \geq 4$	Exact at $m = 2$ and, in the stated range, $m = 3$ (Theorems 1.12 and 1.13)	Locate the first failure for $r \geq 3$ and replace the rank-two decomposition used at $m = 3$.
Directed hypergraphs	For fixed r, m , leading term $(m-1)n^2/(4(r-1))$ for all orientations and both separations	Determine finite exact values and whether forward and general agree once $n > r$.

Table 3.2. The remaining problems, separated from the results already proved in this thesis.

Appendix A

Full Proofs and Computational Audit

This appendix carries every proof in the thesis. The body states the results and says in a few sentences what makes each one work, and the argument itself is here, together with the supporting lemmas and propositions that no chapter states in its own right, and with figures beside the arguments that need them.

How to read this appendix. It runs in four parts, and they are not of equal weight. PART I is classical machinery, [Theorems A.1](#) and [A.2](#) and Mader’s theorem, and may be skipped by a reader who trusts the cited results. PARTS II and III carry the original arguments, and a reader with time for three of them should read [Theorem 3.3](#) (the directed multigraph value, closed for every n and m by deleting a reachability skeleton rather than a vertex), [Lemma A.53](#) with [Theorem 1.13](#) (the incidence rank bound proved through Tutte’s triconnected decomposition, which settles the hypergraph vertex problem at $m = 3$), and [Lemma A.30](#) with the three theorems it drives, [Theorems 3.1](#), [3.2](#) and [3.4](#), which pin six of the sixteen variants asymptotically with no hypothesis on the shape of the extremiser. PART IV holds one self-contained extension and the computational audit.

Part I. Classical machinery

The two theorems every later argument runs on, and Mader’s theorem proved in full through the Gomory–Hu tree. Cited results, reproved here so the thesis stands on its own.

A.1 Menger, Gomory–Hu, and the degree bound

We use two classical results in their capacitated form. Unit capacities represent simple graphs, integer capacities represent multiplicities, and cut capacity is the total capacity crossing the cut.

Theorem A.1 (Menger, as max-flow min-cut [6]). *In any capacitated digraph, and for distinct vertices u, v , the maximum flow from u to v equals the minimum capacity of a cut separating u from v . At unit capacities this is the combinatorial statement we cite throughout: the maximum number of pairwise edge-disjoint u – v paths equals the minimum number of edges whose removal separates them,*

$$\lambda_G(u, v) = \min\{|C| : C \subseteq E(G), G - C \text{ has no } u\text{--}v \text{ path}\}.$$

Undirected graphs become symmetric networks, multigraph multiplicities become capacities, hypergraphs use the gate network of [Figure 2.3b](#), and vertex-disjoint routes use the vertex splitting of [Figure 2.3a](#). Thus one maximum-flow theorem supports every checker variant.

Theorem A.2 (Gomory–Hu [\[4\]](#)). *Every undirected capacitated graph G has a weighted tree T on the same vertex set, its Gomory–Hu tree, such that for all u, v the value $\lambda_G(u, v)$ equals the minimum edge weight on the u – v path in T , and each tree edge e corresponds to a minimum cut of G whose capacity is the weight of e . All $\binom{n}{2}$ pairwise values are thereby stored in the $n - 1$ weights of a single tree.*

The theorem applies to undirected simple graphs, multigraphs, and hyperedge cuts ([Theorem A.47](#)), but not to asymmetric directed cuts or vertex connectivity. Its standard submodular-cut construction is proved in [\[4\]](#). Here we use only the displayed path-minimum identity.

We also use the immediate endpoint bound.

Lemma A.3 (Degree bound). *For every pair of vertices of a graph G , $\lambda_G(u, v) \leq \min(\deg u, \deg v)$. In a digraph D , $\lambda_D(u, v) \leq \min(d^+(u), d^-(v))$.*

Proof. Every edge-disjoint u – v path uses a distinct edge at u , so there are at most $\deg u$ of them, and likewise at most $\deg v$. In the directed case each arc-disjoint path leaves u by a distinct out-arc and enters v by a distinct in-arc. $\square$

[Chapter 2](#) leans on two monotonicity facts to keep the search’s connectivity checks cheap. Both follow directly from max-flow / min-cut.

Proposition A.4 (Monotonicity of the binding connectivity). *Let G be a graph or directed multigraph, and let G' be obtained from G by changing one adjacency by a single unit of multiplicity. Then*

- (1) (monotonicity) *removing the unit cannot raise $\lambda^{\max}$, and adding it cannot lower $\lambda^{\max}$, and*
- (2) (unit step) *adding the unit raises $\lambda^{\max}$ by at most one.*

Both statements hold verbatim for the vertex measure $\kappa^{\max}$.

Proof. Fix an ordered pair (s, t) . By the max-flow / min-cut theorem $\lambda_G(s, t)$ is the smallest capacity of any cut separating s from t , where the capacity of a cut is the total multiplicity of the arcs that cross it from the source side to the sink side.

For part (1), reducing one adjacency by a unit lowers the capacity of every cut that the changed arc crosses and leaves all other cut capacities untouched, so no cut capacity rises. The smallest cut capacity therefore cannot rise either, which gives $\lambda_{G'}(s, t) \leq \lambda_G(s, t)$. Taking the maximum over all pairs yields $\lambda^{\max}(G') \leq \lambda^{\max}(G)$. Adding a unit is the same statement read backwards.

For part (2), write G'' for G with one unit added to a single adjacency. That extra unit of capacity crosses any fixed cut at most once, so every cut has capacity at most one greater in G''

than in G . The smallest cut capacity thus grows by at most one, giving $\lambda_{G''}(s, t) \leq \lambda_G(s, t) + 1$, and together with part (1) we have $\lambda_G(s, t) \leq \lambda_{G''}(s, t) \leq \lambda_G(s, t) + 1$. The maximum over pairs gives $\lambda^{\max}(G'') \leq \lambda^{\max}(G) + 1$.

For $\kappa^{\max}$ the argument is identical on the vertex-split network of Figure 2.3a, where changing an adjacency changes the capacity of its single arc $u^{\text{out}} \rightarrow v^{\text{in}}$ in the same way. Parallel copies leave that capacity at one and so change $\kappa^{\max}$ not at all, which only strengthens both parts. $\square$

A.2 Mader's theorem in full

Let T be a Gomory–Hu tree of G and let $\text{dist}_T(e)$ count the tree edges on the path between the endpoints of $e \in E(G)$, as Figure A.1 illustrates. The upper bound is the following double count. Simplicity enters only because at most $n - 1$ graph edges can join adjacent vertices of T .




Figure A.1. A graph G and a Gomory–Hu tree T on the same vertices. Each graph edge crosses $\text{dist}_T(e)$ tree cuts, and at most $n - 1$ edges of a simple graph have tree distance one.

Lemma A.5 (Distance-one edges). *At most $n - 1$ edges of a simple graph G satisfy $\text{dist}_T(e) = 1$.*

Proof. An edge $e \in E(G)$ has $\text{dist}_T(e) = 1$ if and only if its endpoints are adjacent in T . The tree T has exactly $n - 1$ edges, and since G is simple each pair of adjacent tree vertices supports at most one edge of G . $\square$

Lemma A.6 (Sum of tree distances). $\sum_{e \in E(G)} \text{dist}_T(e) \geq 2|E(G)| - (n - 1)$.

Proof. Partition $E(G)$ into $E_1 = \{e : \text{dist}_T(e) = 1\}$ and $E_{\geq 2} = \{e : \text{dist}_T(e) \geq 2\}$. Then

$$\sum_{e \in E(G)} \text{dist}_T(e) \geq |E_1| + 2|E_{\geq 2}| = 2|E(G)| - |E_1| \geq 2|E(G)| - (n - 1),$$

using Lemma A.5 in the last step. $\square$

Lemma A.7 (Double-counting identity). *Let G carry unit capacities, so that each weight $c(t)$ of a Gomory–Hu tree T is the plain number of edges of G crossing the cut that t induces. Then*

$$\sum_{t \in E(T)} c(t) = \sum_{e \in E(G)} \text{dist}_T(e).$$

This is the form used for Mader's theorem, where G is a simple graph and so unit-capacitated by default.

Proof. By [Theorem A.2](#) each tree edge t induces a cut of G of capacity $c(t)$, so $c(t) = \sum_{e \in E(G)} \mathbf{1}_{e \text{ crosses } t}$. Exchanging the order of summation,

$$\sum_{t \in E(T)} c(t) = \sum_{e \in E(G)} \sum_{t \in E(T)} \mathbf{1}_{e \text{ crosses } t} = \sum_{e \in E(G)} \text{dist}_T(e),$$

where the last equality holds because an edge $\{u, v\}$ crosses exactly those tree cuts induced by the tree edges on the unique u – v path in T . $\square$

Proof of [Theorem 1.2](#), upper bound. Let G be a simple graph on n vertices with $\lambda_G^{\max} \leq m - 1$ and let T be a Gomory–Hu tree of G . Each tree edge $t = \{x, y\}$ has weight $c(t) = \lambda_G(x, y) \leq m - 1$, so summing over the $n - 1$ tree edges, $\sum_t c(t) \leq (m - 1)(n - 1)$. Combining with [Lemmas A.6](#) and [A.7](#),

$$2|E(G)| - (n - 1) \leq \sum_{e \in E(G)} \text{dist}_T(e) = \sum_{t \in E(T)} c(t) \leq (m - 1)(n - 1).$$

Rearranging gives $2|E(G)| \leq m(n - 1)$, and since $|E(G)|$ is an integer, $|E(G)| \leq \left\lfloor \frac{m(n-1)}{2} \right\rfloor$. $\square$

The factor two comes from charging all but at most $n - 1$ edges to at least two tree cuts. Parallel edges destroy that refinement, leaving the multigraph bound $(m - 1)(n - 1)$ of [Theorem 1.3](#).

Proof of [Theorem 1.3](#). For the lower bound, take a spanning tree of K_n , that is, a connected cycle-free subgraph that touches all n vertices and so has exactly $n - 1$ edges, and give every tree edge multiplicity $m - 1$, meaning $m - 1$ parallel copies. The result has $(m - 1)(n - 1)$ edges. Any two vertices are joined by a unique path in the tree, of some length $k \geq 1$. To separate them one must cut some edge of that path, and the cheapest choice is the lightest tree edge on it, which still carries $m - 1$ copies, so $\lambda(u, v) = m - 1$ for every pair and $\lambda^{\max} = m - 1$. The graph sits right at the ceiling.

For the upper bound, suppose $\lambda_G^{\max} \leq m - 1$ and build a Gomory–Hu tree T of G ([Theorem A.2](#)), the same $(n - 1)$ -edge summary of all connectivities used for Mader’s theorem. Each tree edge e stands for a minimum cut of G , of capacity $c(e) = \lambda_G(u, v) \leq m - 1$ for the pair (u, v) it represents. Now charge each edge of G to a tree edge: an edge $\{x, y\}$ crosses the cut of the lightest tree edge on the tree path from x to y , so it is counted inside that tree edge’s capacity. Summing the $n - 1$ capacities therefore counts every edge of G at least once, which already gives $|E(G)| \leq \sum_{e \in T} c(e) \leq (m - 1)(n - 1)$. Unlike Mader’s simple-graph count there is no second charge to halve, because parallel edges are now allowed and nothing forces an edge to be counted twice. The tree-of-multiplicity- $(m - 1)$ construction above meets this bound exactly, which is why the answer is the clean product $(m - 1)(n - 1)$ and not a floor. $\square$

Construction A.8 (Hub-and-spokes graph). For $n \geq m \geq 2$, let $H_{m,n}$ have vertex set $\{h, v_1, \dots, v_{n-1}\}$ with (i) all $n - 1$ *hub edges* $\{h, v_i\}$, and (ii) a *spoke graph* on $\{v_1, \dots, v_{n-1}\}$ with $\left\lfloor \frac{(m-2)(n-1)}{2} \right\rfloor$ edges in which every v_i has spoke degree at most $m - 2$. Such a spoke graph

exists by [Lemma A.9](#). This is the general extremal construction. If $(m-1) \mid (n-1)$, take the spoke graph to be disjoint copies of K_{m-1} , and adjoining h turns them into copies of K_m with one shared vertex, the shared-clique construction.

Lemma A.9 (Near-regular graphs exist). *For $N \geq 1$ and $0 \leq r \leq N-1$ there is a graph on N vertices with $\lfloor \frac{rN}{2} \rfloor$ edges and maximum degree at most r .*

Proof. If rN is even we build an r -regular circulant graph on vertex set $\{0, \dots, N-1\}$: for even r connect each i to $i \pm j \pmod{N}$ for $j = 1, \dots, r/2$, and for odd r (then N is even) use $j = 1, \dots, (r-1)/2$ together with the diameter distance $N/2$, which contributes degree one. Both are r -regular with $rN/2$ edges. If rN is odd, then r and N are both odd: take the $(r-1)$ -regular circulant with distances $1, \dots, (r-1)/2$ and add the near-perfect matching that pairs vertex i with $i + (N+1)/2 \pmod{N}$ for $i = 0, \dots, (N-3)/2$, which covers all vertices but one. The total is $\frac{(r-1)N}{2} + \frac{N-1}{2} = \frac{rN-1}{2} = \lfloor \frac{rN}{2} \rfloor$ edges, with one vertex of degree $r-1$ and all others of degree r . $\square$

Theorem A.10 (Extremal graphs exist). *For all $n \geq m \geq 2$ the graph $H_{m,n}$ has $\lfloor \frac{m(n-1)}{2} \rfloor$ edges and $\lambda^{\max} \leq m-1$.*

Proof. The edge count is $(n-1) + \lfloor \frac{(m-2)(n-1)}{2} \rfloor = \lfloor \frac{m(n-1)}{2} \rfloor$, since $(m-2)(n-1)$ and $m(n-1)$ have the same parity. For the connectivity, any two distinct vertices include at least one spoke vertex v_i , whose total degree is at most $1 + (m-2) = m-1$. [Lemma A.3](#) then gives $\lambda(u, v) \leq m-1$. $\square$

The upper bound above proves [Theorem 1.2](#), and it uses no relation between n and m : the Gomory–Hu tree has $n-1$ edges whatever m is. [Theorem A.10](#) adds that the bound is attained once $n \geq m$, which is where that hypothesis is needed and where it belongs. Below it the complete graph is itself feasible and the bound is valid but slack, with equality returning exactly at $n = m$. The proof also reveals which graphs are extremal: equality forces both inequalities in the chain to be tight.

Theorem A.11 (Characterisation of equality). *Let G be simple with $\lambda_G^{\max} \leq m-1$ and $|E(G)| = \lfloor \frac{m(n-1)}{2} \rfloor$, and let T be a Gomory–Hu tree of G . If $m(n-1)$ is even, then every tree edge has weight exactly $m-1$, every tree edge is also an edge of G , and every edge of G joins vertices at tree distance at most 2. If $m(n-1)$ is odd, there is exactly one unit of slack, located in exactly one of three places: one tree edge has weight $m-2$, or one tree edge is not an edge of G , or one edge of G joins vertices at tree distance 3. Everything else is tight.*

Proof. Write the chain of the upper-bound proof as three non-negative integer slacks, one for each inequality it uses:

$$S_1 = (m-1)(n-1) - \sum_t c(t), \quad S_2 = \sum_{e: \text{dist}_T(e) \geq 2} (\text{dist}_T(e) - 2), \quad S_3 = (n-1) - |E_1|.$$

In words, S_1 counts tree weights below $m - 1$, S_2 counts edges beyond tree distance two, and S_3 counts tree edges that support no edge of G . Retracing the proof gives two expressions for the same quantity,

$$\sum_e \text{dist}_T(e) = 2|E| - |E_1| + S_2 \quad \text{and} \quad \sum_e \text{dist}_T(e) = \sum_t c(t) = (m-1)(n-1) - S_1.$$

Substituting $|E_1| = (n-1) - S_3$ into the first and equating it with the second,

$$2|E| - (n-1) + S_2 + S_3 = (m-1)(n-1) - S_1, \\ \text{so} \quad 2|E| = m(n-1) - (S_1 + S_2 + S_3).$$

The whole characterisation reads off that last identity. If $m(n-1)$ is even, equality $|E| = m(n-1)/2$ forces $S_1 = S_2 = S_3 = 0$. If $m(n-1)$ is odd the floor absorbs exactly one unit, so $S_1 + S_2 + S_3 = 1$ and exactly one of the three slacks equals one. $\square$

For $m = 2$ the characterisation says the extremal graphs are exactly the spanning trees: every tree edge must carry weight one and lie in G , so G contains its own Gomory–Hu tree and has only $n - 1$ edges to spend. For $m = 4$ and $n \equiv 1 \pmod{3}$ it is consistent with the trees of K_4 blocks glued at shared cut vertices, the case of [Construction A.8](#) that the search of [Chapter 2](#) rediscovers.

Remark A.12 (Audit of the counting convention). This remark completes [Remark 1.6](#), which fixes the convention. What follows is the bookkeeping that convention forces on every constant quoted from the literature.

The shift is visible in every entry the database and this thesis both state. It gives $k_2(n) = n$ where a tree has $n - 1$ edges, $k_3(2n) = 3n - 1$ against $\lfloor 3(2n-1)/2 \rfloor = 3n - 2$ here, $k_4(n) = 2n - 1$ against $2n - 2$, $\ell_5(2n) = 5n - 2$ against $5n - 3$, $\ell_6(n) = 3n - 2$ against $3n - 3$, and it writes the Bollobás–Erdős conjecture as $1 + \binom{m}{2}n$ where the convention here gives $\binom{m}{2}n$. Every one is exactly one apart, and the values in this thesis are the ones an exhaustive search returns: $k_3(4) = 4$, $k_3(5) = 6$ and $k_4(5) = 8$ were each confirmed by walking every graph of that size.

One trap is worth naming, because an earlier draft of this thesis fell into it. The comparison $k_5 > \ell_5$ is invariant under the shift, since *both* functions move by one, so no internal check can detect a half-converted statement. What does detect it is evaluating the two sides in different conventions, which manufactures a spurious divergence one size early. Reading $\lfloor 8 \cdot 13/3 \rfloor - 3 = 31$ against this thesis's $\ell_5(13) = 30$ appears to separate the two problems at $n = 13$, but the 31 is a forcing count and the 30 is an avoiding count. Done consistently, $k_5(13) = 30 = \ell_5(13)$ and the separation begins at $n = 14$, which is also what [\[5\]](#) proves directly by showing $f_5(n) = \lfloor 5(n-1)/2 \rfloor + 1$ throughout $6 \leq n \leq 13$.

The range likewise comes from the paper rather than from the database, which records only the clean tail $n \geq 13$. Theorem 4 there covers every $n \geq 6$ apart from $n = 7$ and $n = 12$, and supplies those two values separately, which is where the $k_5(7) = 15$ and $k_5(12) = 27$ of [Theorem 1.5](#) come from.

A.2.1 The vertex problem is a block problem

The *vertex* problem $k_m(n)$ is the one open since 1974, and this short section records its block structure. It changes the object being searched: from graphs on n vertices, with n growing, to 2-connected graphs alone.

Theorem A.13 (The vertex problem is a knapsack over blocks). *For $b \geq 2$ let*

$$h_m(b) := \max\{|E(B)| : B \text{ 2-connected or a single edge, } |V(B)| = b, \kappa_B^{\max} \leq m-1\}.$$

Then for all $m \geq 2$ and $n \geq 2$,

$$k_m(n) = \max\left\{\sum_i h_m(b_i) : b_i \geq 2, \sum_i (b_i - 1) \leq n - 1\right\}.$$

Consequently $k_m(n) \geq k_m(n_1) + k_m(n_2)$ whenever $n = n_1 + n_2 - 1$, and

$$c_m := \lim_{n \rightarrow \infty} \frac{k_m(n)}{n-1} = \sup_{b \geq 2} \frac{h_m(b)}{b-1}$$

exists, the limit being approached from below.

Proof. Let G be feasible. Adjacent vertices u, v lie in a common block B , and no u - v path leaves B : leaving means crossing a cut vertex, and returning means crossing that same cut vertex a second time, which no path does. So $\kappa_G(u, v) = \kappa_B(u, v)$, every block inherits feasibility, and since every edge lies in exactly one block, $|E(G)| = \sum_B |E(B)| \leq \sum_B h_m(|V(B)|)$. Within one component the block sizes satisfy $\sum_B (|V(B)| - 1) = (\text{order of the component}) - 1$, and summing over components loses one further vertex apiece, so $\sum_B (|V(B)| - 1) \leq n - 1$. That is $\leq$.

For $\geq$, take blocks $B_1, \dots, B_t$ with $\sum (b_i - 1) \leq n - 1$, pick one vertex, and attach all the B_i to it, pairwise disjoint otherwise. The result has $1 + \sum (b_i - 1) \leq n$ vertices; add isolated vertices if necessary to bring its order to exactly n . Its nontrivial blocks are exactly the B_i , so pairs inside a block keep their κ and pairs split between two blocks are separated by the shared vertex and have $\kappa \leq 1 \leq m - 1$. It is feasible with $\sum_i |E(B_i)|$ edges.

Superadditivity is the special case of gluing two extremal graphs at a single vertex. Writing $a_n := k_m(n)$, the sequence (a_{n+1}) is superadditive in $n - 1$, so Fekete's lemma gives $a_n/(n - 1) \rightarrow \sup_n a_n/(n - 1)$, and $a_n/(n - 1) \geq h_m(b)/(b - 1)$ whenever $(b - 1) \mid (n - 1)$, while $a_n \leq \max_b h_m(b)/(b - 1) \cdot (n - 1)$ from the display. So the limit is $\sup_b h_m(b)/(b - 1)$. $\square$

Three things follow at once.

At $m = 3$ it *proves* Theorem 1.4's value. A block with $\kappa^{\max} \leq 2$ is an edge or a cycle, since a block that is neither contains a theta subgraph and a theta gives three internally disjoint paths. So $h_3(2) = 1$ and $h_3(b) = b$ for $b \geq 3$, the rate $b/(b - 1)$ is largest at $b = 3$, and the knapsack packs triangles: $k_3(n) = 3 \lfloor (n - 1)/2 \rfloor + ((n - 1) \bmod 2) = \left\lfloor \frac{3(n-1)}{2} \right\rfloor$.

At $m = 4$ the picture is $h_4(b) = 2(b - 1)$ for every $b \geq 4$, so the rate is exactly 2 at every block size and $k_4(n) = 2(n - 1)$. This one is a repackaging rather than a second proof. The upper bound $h_4(b) \leq k_4(b) = 2(b - 1)$ is [Theorem 1.4](#) itself read on a block, so it cannot turn round and reprove it, and nothing as elementary as the theta argument above is on offer for the blocks with $\kappa^{\max} \leq 3$. What the reformulation adds is the extremal block, one the appendix has already built: the hub-and-spokes graph $H_{4,b}$ of [Construction A.8](#), a hub joined to every other vertex with a cycle on the rest. Unlike the shared-clique graph it is a single block, and it meets $2(b - 1)$ at every size. An exhaustive sweep of the 2-connected graphs confirms that it is optimal, and the unique optimum, at every b from 4 to 8.

At every m it identifies the Bollobás–Erdős construction, as many copies of K_m sharing one vertex as fit, as the case $b = m$: $h_m(m) = \binom{m}{2}$ and the rate is exactly $m/2$. Their conjecture is precisely the assertion $c_m = m/2$, that no block beats K_m . Exhaustive computation over all 2-connected graphs on at most nine vertices, of which there are 194066 at the top size alone, confirms that none does, for every $m \leq 8$: the rate $m/2$ is attained and never exceeded. That sweep runs from its own standalone script, `research_notes/scripts/simple_vertex_blocks.py`, rather than from the main program, so it corroborates the checker instead of sharing code with it. Since the rate of a bouquet of blocks is a weighted average of the rates of its blocks, a counterexample must contain a single 2-connected block that beats $m/2$ on its own, and by the computation such a block needs at least ten vertices.

The reason the computation does not reach further is structural rather than a matter of processor time. The conjecture is false: Leonard disproved it at $m = 5$ with an explicit graph on 57 vertices [\[14\]](#), and Mader proved that for every $m \geq 6$ the difference $k_m(n) - mn/2$ is unbounded [\[8\]](#), so $c_m > m/2$ there. What replaces it is a lower bound of Sørensen and Thomassen [\[5\]](#): for each fixed m ,

$$c_m \geq \frac{m(m-1)-2}{2m-3} = \frac{m}{2} + \frac{1}{4} + O(1/m),$$

so the conjectured rate is wrong, but only by a constant.

Their witness is built by a recursion, described here in the language of this section, since it says why the blocks above cannot find it. Start from $G^0 = K_m$ minus an edge, and form G^j from two fresh copies of G^0 together with G^{j-1} , identifying one vertex of each with one vertex of the next, the three identifications arranged in a *cycle*. Each step adds $2m - 3$ vertices and $m(m - 1) - 2$ edges, which is the rate above. The arrangement is cyclic, and three graphs glued in a cycle at three vertices have no cut vertex at all, so G^j is 2-connected and is a *single block*. [Theorem A.13](#) therefore sees it as one indivisible object of size $m + (2m - 3)j$ and gets no purchase on it. What the graph does have is 2-cuts, one pair of identification vertices separating each copy from the rest, so the decomposition that would see it is the triconnected one, along 2-cuts rather than cut vertices, which is the same Tutte/SPQR machinery [Lemma A.53](#) already uses on incidence graphs. That is the concrete next step this reformulation points at.

It also explains the sizes. The recursion beats the rate $m/2$ only once $j > 2/(m - 4)$, so the smallest member of the family that outruns a bouquet of K_m 's has 26 vertices at $m = 5$, 24 at

$m = 6$ and 18 at $m = 7$, against the nine that exhaustive enumeration reaches. The construction was rebuilt and checked here at $m = 5, 6, 7$ and $j \leq 3$: the vertex and edge counts match the formulas exactly, every member is 2-connected, and every member has $\kappa^{\max} = m - 1$, so each is feasible and none contains the forbidden m internally disjoint paths.

Remark A.14 (A degeneracy question, and what it would give). A general density bound is $k_m(n) < 2(m - 1)n$: Mader's density theorem [15] would otherwise produce an m -connected subgraph. Feasibility says much more than that. It gives, for instance, that any two adjacent vertices have at most $m - 2$ common neighbours, since each common neighbour is a route of length two beside the edge.

The natural strengthening is that a feasible graph always has a vertex of degree at most $m - 1$. Since feasibility passes to induced subgraphs, that would make every feasible graph $(m - 1)$ -degenerate and give

$$k_m(n) \leq (m - 1)n - \binom{m}{2},$$

halving the constant in the only general upper bound known. It holds for $m = 3$, where it is classical: a graph of minimum degree at least three contains a subdivision of K_4 , and two branch vertices of that subdivision are joined by three internally disjoint paths. Exhaustive search over all graphs with minimum degree at least m finds no feasible one for $m \leq 6$ and $n \leq 10$. It is stated here as a question rather than a conjecture, since the sizes reached are well below those at which the Bollobás–Erdős conjecture itself first fails.

Part II. The directed results

The exact value at $m = 2$, the structural propositions that bound the frontier at $m \geq 3$, and the directed multigraph problem closed for every n and every m .

A.3 The directed arc value at $m = 2$

Three short lemmas feed the induction. The first shows a hub forces a linear bound for every m , and is the proof behind the hub branch of [Conjecture 1.14](#). The three lemmas below and the proof of [Theorem 1.9](#) that they support are the author's own, and its finite base cases $2 \leq n \leq 7$ were verified by the author's program, through the exhaustive search of [Chapter 2](#) whose completed range is recorded in [Table A.1](#).

Lemma A.15 (Two-hop lemma). *Let D be a simple digraph with $\lambda_D^{\max} \leq m - 1$ containing a vertex r with $d^+(r) = n - 1$. Then every $v \neq r$ has at most $m - 2$ in-arcs from $V \setminus \{r, v\}$.*

Proof. If $u_1, \dots, u_{m-1}$ were distinct in-neighbours of v other than r , that is, distinct vertices each sending an arc to v , the paths $r \rightarrow v$ and $r \rightarrow u_i \rightarrow v$ for $i = 1, \dots, m - 1$ would be m arc-disjoint

routes, since the u_i are distinct and $d^+(r) = n - 1$ supplies every starting arc. That contradicts $\lambda_D(r, v) \leq m - 1$. $\square$

Theorem A.16 (Hub upper bound). *If a simple digraph D with $\lambda_D^{\max} \leq m - 1$ has a vertex r with $d^+(r) = n - 1$, then $|A(D)| \leq m(n - 1)$.*

Proof. Split the arcs into those leaving r (exactly $n - 1$), those entering r (at most $n - 1$), and the internal arcs avoiding r . By [Lemma A.15](#) each of the $n - 1$ other vertices receives at most $m - 2$ internal arcs, so there are at most $(m - 2)(n - 1)$ internal arcs, and the total is at most $m(n - 1)$. $\square$

Lemma A.17 (No transitive triangle). *A simple digraph with $\lambda^{\max} \leq 1$ contains no three vertices x, y, z with all of $(x, y), (y, z), (x, z)$ present, since $x \rightarrow z$ and $x \rightarrow y \rightarrow z$ would be two arc-disjoint routes.*

Lemma A.18 (Deletion monotonicity). *Write $D - v_0$ for the digraph obtained from D by deleting the vertex v_0 together with all arcs incident to it. For any digraph D and vertex v_0 ,*

$$\lambda^{\max}(D - v_0) \leq \lambda^{\max}(D) \quad \text{and} \quad \kappa^{\max}(D - v_0) \leq \kappa^{\max}(D),$$

since arc-disjoint paths in $D - v_0$ are still arc-disjoint paths in D , and internally vertex-disjoint paths in $D - v_0$ are still internally vertex-disjoint paths in D . The two statements are proved by the same one-line observation because deleting a vertex only removes routes, never creates one, and it cannot make two surviving routes share anything they did not share before.

In words: deleting a vertex can never manufacture independent routes. Every route surviving in $D - v_0$ was already a route in D , so no pair comes out of the deletion better connected than it went in. This single guarantee is what every vertex-deletion induction in this section rests on, and it holds for both separations alike.

Proof of [Theorem 1.9](#). Write $M(n) = \max(2(n - 1), \lfloor n^2/4 \rfloor)$. A direct computation shows $2(n - 1) \geq \lfloor n^2/4 \rfloor$ exactly for $n \leq 7$, with equality at $n = 7$, so $M(n) = 2(n - 1)$ for $n \leq 7$ and $M(n) = \lfloor n^2/4 \rfloor$ for $n \geq 7$. The lower bound $\ell_2^{\text{dir}}(n) \geq M(n)$ is witnessed by the directed hub construction ([Construction 1.7](#) at $m = 2$, the bidirected star) and the one-directional bipartite construction ([Construction 1.8](#)).

For the upper bound we show by strong induction on n that $\lambda_D^{\max} \leq 1$ forces $|A(D)| \leq M(n)$.

Base cases $2 \leq n \leq 7$. All six are settled uniformly by the pruned exhaustive search of [Chapter 2](#), which walks every simple digraph with $\lambda^{\max} \leq 1$ on up to seven vertices and reports the maxima 2, 4, 6, 8, 10, 12, matching $M(n)$ at each size. The transcript is recorded in [Table A.1](#), and the cut-counting certifier independently confirms the values within its reach through the $m = 2$ reduction of [Corollary A.39](#). The two smallest are also immediate by hand, as a check on the machine rather than as a separate argument. At $n = 2$ there are at most 2 arcs. At $n = 3$ a fifth arc would leave at most one arc of the complete bidirected triangle missing, so three of the

remaining arcs form a transitive triangle, which [Lemma A.17](#) forbids. The point of the theorem is precisely that $n = 4, 5, 6, 7$ are *not* immediate by hand, which is what makes handing the case-check to the program the right move.

Inductive step $n \geq 8$. Let D be a simple digraph on n vertices with $\lambda_D^{\max} \leq 1$ and put $a := |A(D)|$. Choose v_0 minimising the total degree $d(v) = d^+(v) + d^-(v)$. Since $\sum_v d(v) = 2a$, averaging gives $d(v_0) \leq 2a/n$. By [Lemma A.18](#) the digraph $D - v_0$ on $n - 1 \geq 7$ vertices satisfies the induction hypothesis, so $a - d(v_0) \leq M(n - 1) = \left\lfloor \frac{(n-1)^2}{4} \right\rfloor$. Combining,

$$a \leq \frac{2a}{n} + \left\lfloor \frac{(n-1)^2}{4} \right\rfloor \implies a \leq \frac{n}{n-2} \left\lfloor \frac{(n-1)^2}{4} \right\rfloor.$$

It remains to check that this right-hand side never exceeds $\lfloor n^2/4 \rfloor$, and the two parities behave differently enough to treat them in turn.

For even $n = 2k$ with $k \geq 4$, first simplify $\lfloor (n-1)^2/4 \rfloor$. Expanding $(2k-1)^2 = 4k^2 - 4k + 1$ and dividing by 4 gives $k^2 - k + \frac{1}{4}$, whose floor is $k(k-1)$. The bound then reads

$$a \leq \frac{2k}{2k-2} k(k-1) = \frac{k}{k-1} k(k-1) = k^2,$$

and since $\lfloor n^2/4 \rfloor = \lfloor (2k)^2/4 \rfloor = k^2$ as well, the bound lands exactly on the target with no slack left over.

For odd $n = 2k+1$ with $k \geq 4$, the same simplification gives $\lfloor (n-1)^2/4 \rfloor = \lfloor (2k)^2/4 \rfloor = k^2$, so the bound is $a \leq \frac{2k+1}{2k-1} k^2$. To compare this fraction with the target, split off its integer part by dividing $(2k+1)k^2$ by $2k-1$. The remainder is exactly k , because $(2k-1)(k^2+k) = 2k^3 + k^2 - k$ and $(2k+1)k^2 = 2k^3 + k^2$ differ by k , so

$$\frac{n}{n-2} \left\lfloor \frac{(n-1)^2}{4} \right\rfloor = \frac{(2k+1)k^2}{2k-1} = k(k+1) + \frac{k}{2k-1}.$$

The leftover fraction $\frac{k}{2k-1}$ lies strictly between 0 and 1 for every $k \geq 2$ (the induction here is already past the base cases, so $k \geq 4$), so the bound puts a at most an integer $k(k+1)$ plus a positive amount below one. Since a is itself an integer it cannot reach the next integer up, so $a \leq k(k+1)$. The target $\lfloor n^2/4 \rfloor = \lfloor (2k+1)^2/4 \rfloor = k^2 + k = k(k+1)$ agrees, so for odd n it is integrality, not the raw arithmetic, that closes the last fraction of a unit. In both parities $a \leq \lfloor n^2/4 \rfloor \leq M(n)$, completing the induction. $\square$

Two remarks on the shape of this proof. First, the minimum-degree deletion closes the step on its own, and no separate hub case is needed, because the averaging bound holds whether or not D contains a hub, although [Theorem A.16](#) shows directly that a hub digraph never exceeds the linear branch. Second, for even n the bound is exactly tight against the one-directional bipartite construction, which is $\frac{n}{2}$ -regular in total degree, so every step of the chain is achieved by the conjectured extremiser, and for odd n the slack $\frac{k}{2k-1} < 1$ is absorbed by integrality. The proof is long not because it is deep but because the two regimes of $M(n)$ must be carried through the

arithmetic separately, and that bookkeeping is precisely what [Chapter 2](#) hands to the machine at the base cases.

Remark A.19 (Which way Whitney runs). It is worth pausing on the direction of Whitney's inequality before using it, because the tempting reading is the wrong one. From $\kappa_D(u, v) \leq \lambda_D(u, v)$ for every pair it follows that $\kappa_D^{\max} \leq \lambda_D^{\max}$, hence that

$$\{D : \lambda_D^{\max} \leq m - 1\} \subseteq \{D : \kappa_D^{\max} \leq m - 1\}.$$

Every digraph feasible for the *arc* problem is feasible for the *vertex* problem, and not the other way round. The vertex-feasible family is the larger of the two, so the inequality between the extremal numbers it hands over for free is

$$k_m^{\text{dir}}(n) \geq \ell_m^{\text{dir}}(n),$$

and an upper bound for the arc problem says nothing at all about the vertex problem. The gap is not vacuous: two routes that share an interior vertex but no arc are counted by λ and not by κ , so there really are digraphs in the difference of the two families. Whitney therefore gives one thing, that an arc construction is a vertex *lower* bound, and the vertex upper bound has to be proved separately. The proof below does that, and [Chapter 3](#) treats the two separations the same way for $m \geq 3$, where the vertex problem stays open even though the arc conjecture is stated.

Proof of Theorem 1.10. Write $M(n) = \max(2(n-1), \lfloor n^2/4 \rfloor)$ as before.

Lower bound. The directed hub and the one-directional bipartite construction both have $\lambda^{\max} = 1$, so by [Remark A.19](#) both have $\kappa^{\max} \leq 1$ and are feasible for the vertex problem. Hence $k_2^{\text{dir}}(n) \geq M(n)$.

Upper bound. We show by strong induction on n that $\kappa_D^{\max} \leq 1$ forces $|A(D)| \leq M(n)$. The induction is the one that proved [Theorem 1.9](#), and every step of it survives the change of separation, because the only two properties of the ceiling that argument uses are that it is inherited by vertex deletion and that it holds at the base sizes.

The base cases $2 \leq n \leq 7$ are settled by the same pruned exhaustive search of [Chapter 2](#), run with the vertex feasibility test in place of the arc one. It walks every simple digraph with $\kappa^{\max} \leq 1$ on up to seven vertices and reports the maxima 2, 4, 6, 8, 10, 12, matching $M(n)$ at each size. [Table A.1](#) records this run beside the arc one.

For the inductive step $n \geq 8$, let D be a simple digraph on n vertices with $\kappa_D^{\max} \leq 1$ and put $a := |A(D)|$. Choose v_0 of minimum total degree, so $d(v_0) \leq 2a/n$ by averaging. By [Lemma A.18](#), which holds for κ exactly as it does for λ , the digraph $D - v_0$ on $n-1 \geq 7$ vertices again satisfies $\kappa^{\max} \leq 1$, so the induction hypothesis gives $a - d(v_0) \leq M(n-1) = \lfloor (n-1)^2/4 \rfloor$. From here the two parity computations are verbatim those of [Theorem 1.9](#): they use only the two displayed inequalities and the integrality of a , no property of the separation. They yield $a \leq \lfloor n^2/4 \rfloor \leq M(n)$, completing the induction.

Hence $k_2^{\text{dir}}(n) = M(n) = \ell_2^{\text{dir}}(n)$. The two problems have the same answer at $m = 2$, but this is a computed coincidence of the two searches meeting at every base size, not a formal

consequence of Whitney's inequality. $\square$

Remark A.20 (The directed vertex problem at $m \geq 3$). The proof above transfers the $m = 2$ induction but nothing more, and for $m \geq 3$ the vertex problem is separate: the arc conjecture, and the decomposition of [Conjecture A.25](#) that would prove it, both concern the smaller family and leave $k_m^{\text{dir}}(n)$ untouched. What can be said is computational. Running the exhaustive search of [Chapter 2](#) under both feasibility tests at $m = 3$ gives

n	2	3	4	5	6
$\ell_3^{\text{dir}}(n)$	2	6	9	12	15
$k_3^{\text{dir}}(n)$	2	6	9	12	15

with every entry proved complete, so the two separations agree at every size the exhaustion reaches, and from $n = 3$ onwards both match the conjectured $\max(m(n-1), \lfloor (n+m-2)^2/4 \rfloor)$. The table stops at $n = 6$ because that is where the exhaustion stops: at $n = 7$ neither sweep closes inside an hour and a half, and each returns the construction's 18 as a bare lower bound. At $n = 2$ the conjecture's formula gives 3 where only 2 arcs exist at all, the usual degeneracy of the formula below $n = m$. Agreement over five sizes is evidence and not a proof, and the natural next question is structural rather than numerical: to separate the two problems one needs a digraph where some pair carries strictly more arc-disjoint routes than internally disjoint ones *and* where that slack pays for extra arcs. The two demands pull against each other, which is perhaps why no small example does both.

Lemma A.21 (The augmented bipartite digraph is feasible). *For $m \geq 2$ and $n \geq m$, the digraph of [Construction 1.15](#) satisfies $\lambda^{\max} = m - 1$ and has $\lfloor (n+m-2)^2/4 \rfloor$ arcs. Hence $\ell_m^{\text{dir}}(n) \geq \lfloor (n+m-2)^2/4 \rfloor$.*

Proof. The arc count is $|B|(|A| + m - 2)$, one arc from each of the $|A|$ vertices of A plus $m - 2$ internal in-arcs, for each of the $|B|$ vertices of B . Now $|A| + m - 2 = \lfloor (n+m-2)/2 \rfloor$ and $|B| = \lceil (n+m-2)/2 \rceil$, and the product of the two halves of an integer N is $\lfloor N^2/4 \rfloor$, which gives the stated total at $N = n + m - 2$.

For the connectivity, fix $a \in A$ and $b \in B$. Two kinds of route run from a to b : the direct arc, and the two-step detours $a \rightarrow b' \rightarrow b$ that pass through an in-neighbour b' of b inside B . There are $m - 2$ detours, so $m - 1$ routes in all, and no two of them share an arc. They are also all there are, because the only in-arcs of b that a can reach are exactly those $m - 1$ arcs. Deleting them therefore cuts every route from a to b . Pairs inside B are limited by the in-degree $m - 2$, pairs inside A and pairs from B to A have no routes at all, so $\lambda^{\max} = m - 1$. $\square$

A.4 Structural propositions on the directed frontier

These are the proved ingredients of the reduction discussed in [Chapter 3](#). The section ends with what the reduction still assumes, stated as [Conjecture A.25](#), and with the one quadratic bound that assumes nothing, [Proposition A.26](#).

Proposition A.22 (Mutually unreachable out-neighbourhoods). *Let D be a simple digraph with $\lambda_D^{\max} \leq 1$. For any vertex u and distinct out-neighbours v, w of u , there is no directed path from v to w in $D - u$.*

Proof. If P were a directed $v \rightsquigarrow w$ path avoiding u , then $u \rightarrow w$ and $u \rightarrow v \xrightarrow{P} w$ would be two arc-disjoint u -to- w routes: the first uses only the arc (u, w) , the second starts with (u, v) and continues along P , none of whose arcs touch u . This contradicts $\lambda_D(u, w) \leq 1$. The restriction to $D - u$ is not cosmetic, and Figure A.2 draws both the argument and the reason for the restriction: in the digraph with arcs $(u, v), (v, u), (u, w)$ every local connectivity is 1, yet $v \rightarrow u \rightarrow w$ reaches w from v through u . $\square$




Figure A.2. Why Proposition A.22 must exclude paths through u . (a) For out-neighbours v, w of u in a digraph with $\lambda^{\max} \leq 1$, no $v \rightsquigarrow w$ path can avoid u : such a path would combine with the lone arc $u \rightarrow w$ into two arc-disjoint routes. (b) The restriction to $D - u$ is essential. In the three-vertex digraph $\{(u, v), (v, u), (u, w)\}$ we have $\lambda^{\max} = 1$, yet v reaches w via $v \rightarrow u \rightarrow w$, a route that reuses the vertex u , which the proposition allows.

Proposition A.23 (Disjoint second out-neighbourhoods). *Let D be a simple digraph with $\lambda_D^{\max} \leq 1$, let $u \in V$, and let $v_1 \neq v_2$ be out-neighbours of u . Then $(N^+(v_1) \setminus \{u\}) \cap (N^+(v_2) \setminus \{u\}) = \emptyset$.*

Proof. A common out-neighbour $w \neq u$ would give the two arc-disjoint routes $u \rightarrow v_1 \rightarrow w$ and $u \rightarrow v_2 \rightarrow w$, contradicting $\lambda_D(u, w) \leq 1$. $\square$

Proposition A.24 (Two-hop budget on bipartite partitions). *Let D be a simple digraph with $\lambda_D^{\max} \leq m - 1$. If V admits a partition $A \cup B$ with $A \neq \emptyset$ and every arc from A to B present, then every $b \in B$ has in-degree at most $m - 2$ from inside B , and consequently the number of arcs inside B is at most $(m - 2)|B|$.*

In words: once every arc from A to B is present, a vertex of B can afford very few in-arcs from inside B . Each such arc creates a fresh two-step detour on top of the direct arc that every vertex of A already has, and the ceiling allows only $m - 2$ of them.

Proof. Fix $a \in A$ and $b \in B$, and let $b'_1, \dots, b'_d$ be the in-neighbours of b inside B , where $d = \deg_B^-(b)$. The direct arc $a \rightarrow b$ together with the two-step detours $a \rightarrow b'_i \rightarrow b$ are $1 + d$ arc-disjoint a -to- b routes, all present because $A \rightarrow B$ is complete. Feasibility forces $1 + d \leq m - 1$, so $d \leq m - 2$, and summing over B bounds the internal arcs by $(m - 2)|B|$. $\square$

We now combine [Proposition A.24](#) with the best choice of partition. Suppose every arc of D either runs from A to B or stays inside B : the partition $A \rightarrow B$ is complete, no arc lies inside A , and crucially *no arc runs from B back to A* . Write $b = |B|$, so that $|A| = n - b$. The arcs then come in exactly two kinds. The arcs crossing the partition number $|A||B| = (n - b)b$, one for each ordered A – B pair. The arcs inside B number at most $(m - 2)b$: since no route between two vertices of B ever needs to leave B (there is no arc back to A to leave by), [Proposition A.24](#) applies to B on its own and caps the in-degree of each $b \in B$ from inside B at $m - 2$. There are no other arcs, so, adding the two kinds,

$$|A(D)| \leq (n - b)b + (m - 2)b = b(n + m - 2 - b).$$

The right-hand side is a downward parabola in b . Writing $s = n + m - 2$, it is $b(s - b) = -b^2 + sb$, maximised at $b = s/2$, so over integers the best partition takes $b = \lceil s/2 \rceil$ (equivalently $\lfloor s/2 \rfloor$), and the maximum value is

$$\max_b b(s - b) = \left\lfloor \frac{s^2}{4} \right\rfloor = \left\lfloor \frac{(n + m - 2)^2}{4} \right\rfloor.$$

This is the quadratic branch of [Conjecture 1.14](#), reached uniformly in m . The count is often summarised as resting on the absence of backward arcs alone, and that summary is too narrow. Four things were granted at once: that a partition $V = A \cup B$ exists at all, that every arc from A to B is present, that no arc lies inside A , and that no arc runs from B back to A . Only the last of the four is the backward-arc condition. The first three describe the shape of the extremiser, and nothing proved so far forces an arbitrary extremal digraph to have that shape. The gap is therefore a structural one:

Conjecture A.25 (The missing decomposition). *Let n be large enough that the quadratic branch of [Conjecture 1.14](#) strictly exceeds the linear one, that is, $\lfloor (n + m - 2)^2/4 \rfloor > m(n - 1)$. Then every extremal digraph D on n vertices with $\lambda_D^{\max} \leq m - 1$ admits a partition $V(D) = A \cup B$ into two non-empty parts such that every arc from A to B is present, no arc lies inside A , and no arc runs from B to A .*

The hypothesis on n has to be there, because the obvious phrasing is too narrow. On the linear branch the extremisers are *not* just the hub. At $m = 2$, every bidirected spanning tree is feasible and has exactly $2(n - 1)$ arcs, since between two vertices the only route follows the unique tree path, so the whole extremal set on that branch is the set of bidirected trees. An exhaustive classification confirms it: at $n = 5$ and $n = 6$ there are exactly 3 and 6 extremal digraphs up to isomorphism, which are precisely the numbers of trees on 5 and 6 vertices, and none of them admits the decomposition. Excluding “the hub” would leave all the other trees behind, so the hypothesis is on the size rather than on the shape.

Given [Conjecture A.25](#) the count above closes [Conjecture 1.14](#) for $m \geq 3$. The backward-arc clause is the part where the natural delete-and-compensate exchange fails to be monotone, which is why it is the clause usually named, but the partition itself is not free either, and [Chapter 3](#)

states the position that way. Testing the conjecture by exhaustion is harder than it looks: the crossover sits at $n = 8$ for $m = 2$ and at $n = 9$ for $m = 3$, so every size an exhaustive sweep can reach at $m = 3$ still lies on the linear branch, where the statement says nothing. That is the practical reason this conjecture has resisted a computational probe rather than a proof.

What can be proved without any of the four is the leading constant, and the argument is short enough to give in full. It also says something the conjecture does not: after deleting a lower-order number of arcs, every feasible digraph becomes a subgraph of a one-directional bipartite digraph.

Proposition A.26 (Unconditional quadratic bound and stability). *Let D be a simple digraph on $n \geq 2$ vertices with $\lambda_D^{\max} \leq m - 1$. Then*

$$|A(D)| \leq \left\lfloor \frac{n^2}{4} \right\rfloor + \sqrt{m} n^{3/2}.$$

Moreover D can be made one-directional bipartite, meaning every remaining arc runs from one part to the other and none returns, by deleting at most $\sqrt{m} n^{3/2}$ arcs. Consequently, for each fixed m ,

$$\ell_m^{\text{dir}}(n) = \frac{n^2}{4} + O_m(n^{3/2}),$$

so the leading constant $\frac{1}{4}$ of [Conjecture 1.14](#) holds unconditionally, with no hypothesis on the structure of the extremiser.

Proof. Step 1, a budget on two-step routes. Fix an ordered pair (u, v) with $u \neq v$, and let $p_2(u, v)$ count the vertices x with both $u \rightarrow x$ and $x \rightarrow v$ present. Such an x is automatically distinct from u and from v , since D has no loops, so each one gives a genuine two-step route $u \rightarrow x \rightarrow v$. Two of these routes with different midpoints $x \neq x'$ share no arc, since the first uses (u, x) and (x, v) and the second uses (u, x') and (x', v) , and neither uses the arc (u, v) , which has no midpoint at all. So the direct arc, when present, together with all the two-step routes, is a family of pairwise arc-disjoint u -to- v routes, and feasibility caps its size:

$$\mathbf{1}_{(u,v) \in A(D)} + p_2(u, v) \leq \lambda_D(u, v) \leq m - 1.$$

Summing over all $n(n - 1)$ ordered pairs gives $|A(D)| + \sum_{u \neq v} p_2(u, v) \leq (m - 1) n(n - 1)$.

Step 2, from pairs to degrees. Write Δ for the number of digons of D , meaning the pairs of vertices carrying an arc in each direction. Three small facts turn Step 1 into a statement about degrees alone.

(2a) *Walks of length two, counted by their midpoint.*

$$\sum_{u, v \in V} p_2(u, v) = \sum_{x \in V} d^-(x) d^+(x),$$

where the left-hand sum runs over *all* ordered pairs, the diagonal $u = v$ included. Fixing a vertex x , a walk $u \rightarrow x \rightarrow v$ is nothing more than an independent choice of an in-neighbour u of x and an out-neighbour v of x , so x is the midpoint of exactly $d^-(x) d^+(x)$ of them.

(2b) *What the diagonal counts.* A diagonal term $p_2(u, u)$ counts the closed walks $u \rightarrow x \rightarrow u$, and each digon contributes two such walks, one based at each of its endpoints. Hence

$$\sum_{u \in V} p_2(u, u) = 2\Delta.$$

(2c) *Digons are few.* Distinct digons use disjoint pairs of arcs, so $2\Delta \leq |A(D)|$.

Splitting the all-pairs sum of (2a) along its diagonal and feeding in Step 1 now gives the inequality the rest of the argument runs on:

$$\begin{aligned} \sum_{x \in V} d^+(x) d^-(x) &= \sum_{u \neq v} p_2(u, v) + 2\Delta && \text{by (2a) and (2b),} \\ &\leq [(m-1)n(n-1) - |A(D)|] + 2\Delta && \text{by Step 1,} \\ &\leq [(m-1)n(n-1) - |A(D)|] + |A(D)| && \text{by (2c),} \\ &= (m-1)n(n-1). \end{aligned}$$

The point of the last two lines is that the digon term is absorbed exactly by the slack Step 1 left behind, so nothing is given away in passing from pairs to degrees. It is worth reading the conclusion in words: a vertex with a large in-degree *and* a large out-degree is expensive, because it manufactures two-step routes for many pairs at once, and the budget for those routes is only quadratic in n .

Step 3, the smaller half-degree is small on average. Since $\min(a, b) \leq \sqrt{ab}$ for non-negative a, b , and by the Cauchy-Schwarz inequality in the form $\sum_x \sqrt{a_x} \leq \sqrt{n \sum_x a_x}$,

$$\begin{aligned} \sum_{x \in V} \min(d^+(x), d^-(x)) &\leq \sum_{x \in V} \sqrt{d^+(x) d^-(x)} \leq \sqrt{n \sum_{x \in V} d^+(x) d^-(x)} \\ &\leq \sqrt{n \cdot (m-1)n(n-1)} = n\sqrt{(m-1)(n-1)} \leq \sqrt{m} n^{3/2}. \end{aligned}$$

Step 4, deleting the smaller half of every vertex. Call x *source-like* if $d^+(x) \geq d^-(x)$ and *sink-like* otherwise, and write S and T for the two classes, so $V = S \cup T$. Delete every in-arc of a source-like vertex and every out-arc of a sink-like vertex. A source-like x loses $d^-(x) = \min(d^+(x), d^-(x))$ arcs and a sink-like x loses $d^+(x) = \min(d^+(x), d^-(x))$ arcs, so the number of deleted arcs is at most $\sum_x \min(d^+(x), d^-(x))$, which Step 3 bounds by $\sqrt{m} n^{3/2}$. An arc (a, b) survives only if a is not sink-like and b is not source-like, that is, only if $a \in S$ and $b \in T$. So what remains runs entirely from S to T , which is the one-directional bipartite shape, and it has at most $|S||T| \leq \lfloor n^2/4 \rfloor$ arcs. Adding the two counts gives the bound.

The asymptotic statement follows because the augmented bipartite construction ([Construction 1.15](#)) already gives $\ell_m^{\text{dir}}(n) \geq \lfloor (n+m-2)^2/4 \rfloor = n^2/4 + O_m(n)$, and $n^{3/2}$ dominates n . $\square$

The error term $n^{3/2}$ is larger than the $(m-2)n/2$ that [Conjecture 1.14](#) predicts, so the proposition as it stands does not separate the conjectured formula from its near neighbours. It can be

sharpened to the right order, and the extra idea is small enough to be worth giving, because it explains where the $\sqrt{n}$ was being wasted.

Step 3 is lossy exactly when every vertex carries a middling degree. The Cauchy-Schwarz step is tight only if $d^+(x)d^-(x)$ is about $(m-1)n$ at every vertex, which forces every degree down to about $\sqrt{mn}$. But a digraph with $\Theta(n^2)$ arcs has average degree $\Theta(n)$, and at a vertex of large degree the constraint $d^+(x)d^-(x) \leq (m-1)n$ bites much harder: with $d^+ + d^-$ of order n , the smaller of the two must be of order m rather than $\sqrt{mn}$. The dense case and the tight case of Step 3 are therefore incompatible, and the way to exploit that is a dichotomy on the minimum degree, with vertex deletion to handle the sparse side.

Lemma A.27 (Small side of a high-degree vertex). *For any vertex x of a digraph with total degree $d(x) = d^+(x) + d^-(x) > 0$,*

$$\min(d^+(x), d^-(x)) \leq \frac{2 d^+(x) d^-(x)}{d(x)}.$$

Proof. Write $\mu = \min(d^+(x), d^-(x))$ and $M = \max(d^+(x), d^-(x))$, so $\mu + M = d(x)$ and $\mu M = d^+(x)d^-(x)$. Since $\mu \leq M$ we have $M \geq d(x)/2$, hence $d^+(x)d^-(x) = \mu M \geq \mu d(x)/2$, which rearranges to the claim. $\square$

In words: a feasible one-way simple graph is never more than a linear amount denser than the balanced bipartite wall, which holds $\lfloor n^2/4 \rfloor$ arcs. The wall's quadratic term is therefore the leading term, and what remains in question is how large the linear correction is. Nothing here assumes anything about the shape of D .

Proof of Theorem 3.1. Induction on n . For $n = 2$ a simple digraph has at most 2 arcs and the right-hand side is $1 + 4(m-1) \geq 5$, so the base holds. Let $n \geq 3$ and split on the minimum total degree.

Case 1: some vertex v has $d(v) < n/2$. Since $d(v)$ is an integer, $d(v) \leq \lfloor n/2 \rfloor$. By Lemma A.18 the digraph $D - v$ on $n - 1$ vertices is still feasible, so the induction hypothesis applies to it and

$$|A(D)| \leq \left\lfloor \frac{(n-1)^2}{4} \right\rfloor + 4(m-1)(n-2) + \left\lfloor \frac{n}{2} \right\rfloor = \left\lfloor \frac{n^2}{4} \right\rfloor + 4(m-1)(n-2),$$

using Lemma A.31 for the last step, and this is below the claimed bound.

Case 2: every vertex has $d(x) \geq n/2$. By Lemma A.27 and Step 2 of Proposition A.26,

$$\sum_{x \in V} \min(d^+(x), d^-(x)) \leq \sum_{x \in V} \frac{2 d^+(x) d^-(x)}{d(x)} \leq \frac{4}{n} \sum_{x \in V} d^+(x) d^-(x) \leq \frac{4}{n} (m-1) n(n-1),$$

which is $4(m-1)(n-1)$. Step 4 of Proposition A.26 deletes at most that many arcs and leaves at most $\lfloor n^2/4 \rfloor$, so the bound holds in this case too.

For the two-sided asymptotic statement, [Construction 1.15](#) gives

$$\ell_m^{\text{dir}}(n) \geq \left\lfloor \frac{(n+m-2)^2}{4} \right\rfloor = \frac{n^2}{4} + \frac{(m-2)n}{2} + O_m(1),$$

so the second-order term is linear in n from both sides. $\square$

What is left open is therefore a constant factor in one coefficient, not a shape. [Conjecture 1.14](#) says the second-order coefficient is exactly $(m-2)/2$, and the theorem pins it between that and $4(m-1)$, a factor of $8(m-1)/(m-2)$: sixteen at $m=3$, and tending to eight as m grows. Closing that gap is where the structure of [Conjecture A.25](#) lives, and [Remark A.28](#) rules out the most obvious way to try.

Remark A.28 (Case 2's coefficient needs a new inequality). One might hope to cut Case 2's constant using the observation that in the conjectured extremiser, one whole side has $\min(d^+, d^-) = 0$, so summing [Lemma A.27](#)'s worst case at every vertex looks wasteful. That observation alone does not cut the coefficient. The value four is the best universal coefficient obtainable from the two facts used in Case 2, the budget $\sum_x d^+(x) d^-(x) \leq (m-1)n(n-1)$ of Step 2 in [Proposition A.26](#) and the floor $d(x) \geq n/2$.

Think of the budget as money and of $t_x = \min(d^+(x), d^-(x))$ as what each vertex buys with it. At a vertex with $d(x) = s \geq n/2$, one also has $s \geq 2t$. Hence the cheapest possible cost for a given t is the two-branch function

$$c(t) = \min_{s \geq \max(n/2, 2t)} t(s-t) = \begin{cases} t(n/2 - t), & 0 \leq t \leq n/4, \\ t^2, & t \geq n/4. \end{cases}$$

On the first branch the value per unit cost is $t/c(t) = 1/(n/2 - t)$, which increases to $4/n$, whereas on the second it is $1/t$, which decreases from $4/n$. Thus $t \leq (4/n)c(t)$ for every t , with equality in the continuous relaxation at $t = n/4$. Summing and using the shared budget $Q = (m-1)n(n-1)$ gives

$$\sum_{x \in V} t_x \leq \frac{4}{n} \sum_{x \in V} d^+(x) d^-(x) \leq \frac{4Q}{n} = 4(m-1)(n-1).$$

Conversely, fix m and take n large. Assign $t = n/4$ to as many abstract vertices as the budget permits, assign one further $t \in [0, n/4]$ so that its continuous cost absorbs the remainder, and assign zero to the rest. This uses at most $16(m-1) + 1 \leq n$ vertices. Since the remainder is within $O_m(n)$ of the full cost $n^2/16$, the final t is $n/4 - O_m(\sqrt{n})$. The resulting total is $4(m-1)n - O_m(\sqrt{n})$, matching the coefficient four asymptotically. Therefore no smaller universal coefficient follows from these two inequalities alone.

The crude per-vertex bound therefore loses nothing within this relaxation, and the reason is that its two sources of slack close at that very point. [Lemma A.27](#) is an equality only when $d^+(x) = d^-(x)$, and the floor substitution $d(x) \rightarrow n/2$ is an equality only when $d(x) = n/2$. Both hold at once exactly when $d^+(x) = d^-(x) = n/4$, which is the profile the relaxed optimum

concentrates on. Cutting the constant therefore needs a second and independent inequality, one that constrains interference among the handful of expensive near-balanced vertices themselves rather than a sharper reading of the one already in hand.

A.4.1 What that proof actually used

Neither the proposition nor the theorem above ever used feasibility twice. Each used it once, in Step 1, to cap a single explicit family of routes, and everything after that is arithmetic on in-degrees and out-degrees. Promoting that one cap from a deduction to a hypothesis costs nothing here and buys two further results below, one of them for a problem [Chapter 3](#) lists as untouched by the arc case.

Definition A.29 (Two-step budget). Let D be a simple digraph and let $C \geq 1$. Call D C -budgeted if every ordered pair (u, v) of distinct vertices satisfies

$$\mathbf{1}_{(u,v) \in A(D)} + p_2(u, v) \leq C,$$

where $p_2(u, v)$ counts, as in [Proposition A.26](#), the vertices $x \notin \{u, v\}$ with both (u, x) and (x, v) present.

Lemma A.30 (The counting core). *Every C -budgeted simple digraph D on $n \geq 2$ vertices satisfies*

$$|A(D)| \leq \left\lfloor \frac{n^2}{4} \right\rfloor + 4C(n-1).$$

Proof. Read the proofs of [Proposition A.26](#) and [Theorem 3.1](#) again with C written wherever $m-1$ stood. Step 1 of the proposition is now the hypothesis rather than a deduction, and it was the only place feasibility ever entered, since Steps 2, 3 and 4 manipulate in-degrees and out-degrees and nothing else. The theorem's induction needs one further point, that the hypothesis is inherited by induced subdigraphs, which [Lemma A.18](#) supplied there by way of feasibility. Here it is immediate: deleting a vertex only deletes arcs and removes candidate midpoints, so neither term of the budget can rise. With those two observations the induction runs verbatim and delivers the stated bound. $\square$

The first application is the directed *vertex* problem. [Theorem 1.10](#) settles it at $m = 2$, and for $m \geq 3$ this thesis has been careful to record that it does not follow from the arc case: Whitney's $\kappa \leq \lambda$ makes the vertex-feasible family the larger of the two, so an arc upper bound restricts nothing there. That remains true of the *bound*. What does carry across is the route-counting, because the family Step 1 exhibits happens to be disjoint in the stronger sense as well, and once that is noticed the argument never needs the arc value at all.

Proof of Theorem 3.2. Fix an ordered pair (u, v) of distinct vertices. The direct arc (u, v) , when it is present, is a route with no interior vertex at all, and a two-step route $u \rightarrow x \rightarrow v$ has interior exactly $\{x\}$, one vertex, distinct from both u and v . Distinct midpoints give disjoint interiors, and the direct arc's interior is disjoint from all of them for want of any element, so these $\mathbf{1}_{(u,v) \in A(D)} +$

$p_2(u, v)$ routes are pairwise internally vertex-disjoint and not merely arc-disjoint. Feasibility caps the family at $\kappa_D(u, v) \leq m - 1$, so D is $(m - 1)$ -budgeted and [Lemma A.30](#) applies.

For the lower bound, the augmented bipartite digraph of [Construction 1.15](#) has $\lambda^{\max} \leq m - 1$, hence $\kappa^{\max} \leq m - 1$ by Whitney, so for $n \geq m$ it witnesses $k_m^{\text{dir}}(n) \geq \lfloor (n + m - 2)^2/4 \rfloor$. For fixed $m \geq 3$ this is $n^2/4 + \Theta_m(n)$, and at $m = 2$ [Theorem 1.10](#) gives the sharper $n^2/4 + O(1)$ asymptotic. $\square$

The vertex problem at $m \geq 3$ is therefore open in the same narrow sense as the arc problem rather than in the wide one. Its leading constant is $\frac{1}{4}$ unconditionally, the term after it is linear in n , and since $\ell_m^{\text{dir}}(n) \leq k_m^{\text{dir}}(n)$ with both now inside $n^2/4 + \Theta_m(n)$, the two values agree to first order. What is left is the coefficient of the linear term, the same quantity [Conjecture A.25](#) is needed for on the arc side, together with the conjecture that the two values coincide exactly. It is worth being clear about what did and did not transfer. The upper bound of [Theorem 3.1](#) did not: it is proved over the smaller family and says nothing here. The route family behind it did, and that is a statement about a construction rather than about a bound, which is why Whitney's inequality does not block it.

A.5 The directed multigraph problem, solved in full

The body proves $L_m^{\text{dir}}(n) = 2(n - 1)(m - 1)$ for $n \leq 6$ by an optimisation that scales m out of the problem entirely ([Theorem 2.1](#)), and the section above, on the simple digraph at $m = 2$, closes every n by repeatedly deleting one vertex of least degree and averaging. The same argument almost settles the multigraph case at every m . Dividing every multiplicity by $m - 1$, as in the scaling step of [Chapter 2](#), turns a feasible multigraph into a weighting with values in $[0, 1]$ whose pairwise maximum flows are all at most one. Deleting a vertex of least weighted degree from that weighting then drives the same averaging argument through cleanly on the linear branch, and on the quadratic branch as well whenever n is even. What breaks it is a single fractional remainder at odd n : a leftover of size $\frac{k}{2k-1}$ that an integer count could round away but a genuinely fractional weight cannot. Finding a specific vertex that is provably light enough to absorb that remainder turns out to resist every direct attempt.

This section abandons that attempt and proves the full value a different way. Instead of deleting one vertex at a time, it deletes one whole *sparse subgraph* at a time, chosen so that the deletion is guaranteed to lower every pair's connectivity by at least one in a single stroke. That change removes the fractional step altogether, along with the restriction to a particular m or a particular parity of n .

Lemma A.31 (Arithmetic identity). *For all $n \geq 2$, $\left\lfloor \frac{(n-1)^2}{4} \right\rfloor + \lfloor n/2 \rfloor = \left\lfloor \frac{n^2}{4} \right\rfloor$.*

Proof. For $n = 2k$ the left side is $k(k - 1) + k = k^2$, and for $n = 2k + 1$ it is $k^2 + k = k(k + 1)$. Both match the right side. $\square$

A.5.1 The lower bound: thickening

The lower bound rests on one observation: multiplying every arc's multiplicity by a fixed integer multiplies every cut's capacity, and hence every pairwise connectivity, by that same integer.

Lemma A.32 (Thickening). *Let D_0 be a directed multigraph and $k \geq 1$ an integer. Write $k \cdot D_0$ for the multigraph obtained by multiplying every multiplicity $\mu(u, v)$ by k . Then $\lambda_{k \cdot D_0}(u, v) = k \cdot \lambda_{D_0}(u, v)$ for every ordered pair $u \neq v$, and $|A(k \cdot D_0)| = k |A(D_0)|$.*

Proof. By [Theorem A.1](#), $\lambda_{D_0}(u, v)$ equals the minimum capacity, over all u - v cuts, of the total multiplicity crossing that cut. Multiplying every multiplicity by k multiplies the capacity of every cut by k at once, so it multiplies the minimum over all cuts by k as well, which is precisely $\lambda_{k \cdot D_0}(u, v) = k \cdot \lambda_{D_0}(u, v)$. The arc count scales by k because every one of its summands, each multiplicity, does. $\square$

Construction A.33 (Thickened tree and thickened bipartite digraph). For $m \geq 2$, take any spanning tree T of K_n and give every edge multiplicity $m - 1$ in each direction. Between two vertices, T offers only its one tree path, so this is $m - 1$ copies of the $m = 2$ bidirected tree of [Theorem 1.9](#), feasible at $\lambda^{\max} = 1$ there, and [Lemma A.32](#) with $k = m - 1$ gives $\lambda^{\max} = m - 1$ here and an arc count of $2(n - 1)(m - 1)$. Alternatively, take the one-directional complete bipartite digraph of [Construction 1.8](#) (balanced parts A, B with $|A| = \lfloor n/2 \rfloor$, $|B| = \lceil n/2 \rceil$, feasible at $\lambda^{\max} = 1$) and give every arc multiplicity $m - 1$: the same lemma gives $\lambda^{\max} = m - 1$ and $(m - 1) \lfloor n^2/4 \rfloor$ arcs.

Taking whichever of the two constructions is larger gives $L_m^{\text{dir}}(n) \geq (m - 1)M(n)$ for every n and m , which is the lower bound half of [Theorem 3.3](#). The rest of this section proves the matching upper bound.

A.5.2 The upper bound: a reachability skeleton

Consider first what a single vertex deletion gives at $m = 2$: removing a vertex can only remove routes passing through it, so no pair's connectivity ever rises, and the $m = 2$ induction rests on that guarantee together with an averaging bound on how much one well-chosen vertex carries. The multigraph problem at a general m needs those same two ingredients: a deletion that is guaranteed to lower every connectivity, paired with a bound on how much it removes. There, though, the natural analogue of “one vertex” breaks down. Once multiplicities are free to sit anywhere between 0 and $m - 1$, no single vertex is guaranteed to carry a controllable share of the arcs, which is exactly the fractional remainder above. The fix is to stop insisting on a vertex and delete a whole *subgraph* instead: one general enough to always exist, sparse enough to bound, and specific enough that removing it is guaranteed to lower every pairwise connectivity by at least one.

Definition A.34 (Reachability skeleton). Let D be a directed multigraph. A spanning subdigraph $R \subseteq D$, using at most one copy of each arc it uses (no parallel copies, even if D has them), is a *reachability skeleton* of D if, for every pair of vertices $u \neq v$, u reaches v by some directed path in R exactly when u reaches v by some directed path in D .

A reachability skeleton keeps every yes-or-no fact about who can reach whom and discards everything else, in particular every parallel copy of every arc. It is a caricature of D : thin, but faithful on the one question connectivity actually depends on. The next two lemmas turn that faithfulness into the whole induction. The first says a reachability skeleton always exists and is never too big. The second says deleting it always lowers every connectivity by at least one.

Lemma A.35 (A sparse reachability skeleton always exists). *Every loopless directed multigraph D on $n \geq 2$ vertices has a reachability skeleton R with $|A(R)| \leq M(n)$.*

Building R takes three steps: handle the traffic *inside* each strongly connected piece of D , handle the traffic *between* pieces, then count. A *strongly connected component* (an SCC) is a maximal set of vertices any two of which reach each other, and every directed multigraph partitions uniquely into its SCCs, since mutual reachability is an equivalence relation. [Figure A.3](#) carries one small running example through all three steps, and every object named below appears in it concretely.

Step 1: inside each component. Fix one SCC of order $n_i \geq 1$ and pick any vertex r inside it as a root. Since r reaches every vertex of the component, a standard fact about digraphs gives a spanning *out-arborescence*: a tree of $n_i - 1$ arcs of D , every one pointing away from r , along which r alone reaches every other vertex of the component (grow it one hop at a time, always available since the component is strongly connected). Symmetrically, because every vertex of the component reaches r , there is a spanning *in-arborescence* of $n_i - 1$ arcs, every one pointing toward r , along which every vertex reaches r . Taking both trees together, any two vertices u, v of the component are joined by the route $u \rightarrow r \rightarrow v$, first the in-arborescence, then the out-arborescence, so the $2(n_i - 1)$ arcs of the two trees alone reconstruct *every* reachability relation inside this one component.

Step 2: between components. Contract each SCC to a single point and, between two contracted points, keep one arc whenever D has at least one arc from the first component to the second (dropping any further parallel arcs between the same two components at this stage). Call the result Q_0 , a digraph on q points, one per component. Two different SCCs can never reach each other in both directions, since that would make them one strongly connected component after all, so Q_0 has no directed cycle: it is a DAG (a directed acyclic graph). Reachability between two different components of D is, by this very construction, exactly reachability between the two corresponding points of Q_0 . Thin Q_0 down to an *inclusion-minimal* spanning subdigraph $Q \subseteq Q_0$ that still has the same reachability relation, deleting one arc at a time for as long as some arc is redundant, which must stop since Q_0 has only finitely many arcs.

The reason to thin Q_0 down to Q is that Q 's *underlying undirected graph*, meaning Q with the arrows erased, has no triangle. Suppose three points x, y, z of Q were pairwise joined by some arc of Q . Since $Q \subseteq Q_0$ and Q_0 has no cycle, Q has none either, so between any two of x, y, z the arc runs in only one direction, never both. Three points pairwise joined with a consistent direction on each pair admit only one pattern, up to relabelling: a linear order, say $x \rightarrow y, y \rightarrow z$, and $x \rightarrow z$. But then $x \rightarrow z$ is redundant, since x already reaches z via y , so deleting it would

leave the reachability relation of Q unchanged, contradicting that Q is inclusion-minimal. So no three points of Q can be pairwise joined at all: that is triangle-freeness of the underlying graph.

A triangle-free graph on q vertices has at most $\lfloor q^2/4 \rfloor$ edges, Mantel's theorem [16], the $r = 2$ case of the more general theorem of Turán [17]. Applied to Q ,

$$|A(Q)| \leq \left\lfloor \frac{q^2}{4} \right\rfloor.$$

Step 3: assembling R and counting. Build R from the two arborescences of Step 1 inside every SCC, together with one witness arc of D for every arc of Q (some arc of D realises it, by the very definition of Q_0). By Step 1, every pair inside one component reaches each other in R exactly as in D . By Step 2, and because within-component reachability is already secured, every pair in two different components reaches each other in R exactly as in D too: to travel from u to v across the components Q 's path visits, reach each witness arc's tail inside its own component (Step 1), cross the witness arc, and repeat. Since $R \subseteq D$ throughout, R can never certify a reachability D does not already have, so R is a genuine reachability skeleton. Writing $n_1, \dots, n_q$ for the SCC orders, so $\sum_i n_i = n$, its arc count is at most the $2(n_i - 1)$ arborescence arcs of each component plus one witness per arc of Q . The first count is an upper bound rather than an equality, since the in- and out-arborescence of one component may share an arc, which only helps. So

$$|A(R)| \leq \sum_{i=1}^q 2(n_i - 1) + |A(Q)| \leq 2(n - q) + \left\lfloor \frac{q^2}{4} \right\rfloor =: f(q).$$

It remains to bound $f(q)$ over the range $1 \leq q \leq n$ that the number of components can actually take. Its increments are

$$f(q+1) - f(q) = -2 + \left\lfloor \frac{q+1}{2} \right\rfloor,$$

using the identity $\lfloor (q+1)^2/4 \rfloor - \lfloor q^2/4 \rfloor = \lfloor (q+1)/2 \rfloor$, checked on the two parities of q (at $q = 2j$ both sides are j , at $q = 2j+1$ both are $j+1$). Since $\lfloor (q+1)/2 \rfloor$ never decreases as q grows, neither do the increments of f , which is to say f is convex. A convex function on an interval attains its largest value at one of the two ends, because a value at an interior point is a weighted average of the two ends or less. Hence

$$f(q) \leq \max(f(1), f(n)) \quad \text{for } 1 \leq q \leq n.$$

At $q = 1$ the whole digraph is one component, so $f(1) = 2(n-1) + 0$. At $q = n$ every component is a single vertex, so $f(n) = 0 + \lfloor n^2/4 \rfloor$. Hence $f(q) \leq \max(2(n-1), \lfloor n^2/4 \rfloor) = M(n)$ for every q in range, which is the lemma.

Lemma A.36 (Deleting a skeleton lowers every connectivity by one). *Let D be a directed multigraph with $\lambda^{\max}(D) \leq k$ for some $k \geq 1$, and let R be a reachability skeleton of D . Write $D - R$*




Figure A.3. A small instance of [Lemma A.35](#). In D (left) three bidirected pairs $A = \{a_1, a_2\}$, $B = \{b_1, b_2\}$, $C = \{c_1, c_2\}$ are the strongly connected components, joined by $a_2 \rightarrow b_1$, $b_2 \rightarrow c_1$, and $a_1 \rightarrow c_2$. Contracting each pair to a point gives a condensation Q_0 on three points, carrying all three arcs $A \rightarrow B$, $B \rightarrow C$ and $A \rightarrow C$. That is exactly the transitively-closed triangle the proof rules out of the thinned Q . The arc $A \rightarrow C$ is redundant, since A already reaches C through B , so the transitive reduction Q keeps only $A \rightarrow B$ and $B \rightarrow C$. The underlying graph of Q is then a path on three points, triangle-free as claimed. The reachability skeleton R (right) reflects exactly this: every SCC keeps both of its internal arcs (an in- and an out-arborescence on two vertices is just the pair itself), while of the two component-crossing arcs of D only the two witnesses of Q 's two arcs survive. The direct arc $a_1 \rightarrow c_2$ is dropped, yet a_1 still reaches c_2 in R : first $a_1 \rightarrow a_2$, then $a_2 \rightarrow b_1$, then $b_1 \rightarrow b_2$, then $b_2 \rightarrow c_1$, then $c_1 \rightarrow c_2$.

for the multigraph obtained by deleting, from each multiplicity $\mu(u, v)$, the one copy (if any) that R uses on that pair. Then $\lambda^{\max}(D - R) \leq k - 1$.

In words: subtracting one copy of each skeleton arc costs every pair at least one route (a single pair can lose more, if some of its own routes happened to run through the deleted arcs too). The skeleton preserves reachability, so however many arc-disjoint routes survive the deletion, one further route can always be run entirely inside the arcs that were deleted, and that extra route is what the original ceiling has to pay for.

Proof. Fix an ordered pair $u \neq v$ and suppose $D - R$ carries $t \geq 1$ pairwise arc-disjoint u - v paths. These use arcs of $D - R$ and hence of D , since $D - R \subseteq D$, so u reaches v in D . Because R is a reachability skeleton, u then also reaches v in R : some directed path P from u to v uses only arcs of R . Every arc of R has been subtracted out of $D - R$, so P shares no arc with any of the t paths already found inside $D - R$. Together, P and those t paths are $t + 1$ pairwise arc-disjoint u - v paths in D , so $t + 1 \leq \lambda_D(u, v) \leq k$, that is, $t \leq k - 1$. Pairs with $t = 0$ trivially satisfy $t \leq k - 1$ once $k \geq 1$, so this holds for every pair, giving $\lambda^{\max}(D - R) \leq k - 1$. $\square$

Proof of Theorem 3.3. The lower bound is [Construction A.33](#). For the upper bound, fix n and prove by induction on $k = m - 1 \geq 0$ that every directed multigraph D on n vertices with $\lambda^{\max}(D) \leq k$ has $|A(D)| \leq k M(n)$. At $k = 0$, every pair has $\lambda_D(u, v) \leq 0$, so no arc (u, v) can even be present, since a single arc is already one route, and $|A(D)| = 0 = 0 \cdot M(n)$.

For $k \geq 1$, take D with $\lambda^{\max}(D) \leq k$ and let R be a reachability skeleton, which exists with $|A(R)| \leq M(n)$ by [Lemma A.35](#). By [Lemma A.36](#), $D - R$ has $\lambda^{\max}(D - R) \leq k - 1$, and it is again a directed multigraph on the same n vertices, so the induction hypothesis gives

$|A(D - R)| \leq (k - 1)M(n)$. Since R uses at most one copy of each arc it touches, the arcs of D split exactly into those of R and those left in $D - R$, so

$$|A(D)| = |A(R)| + |A(D - R)| \leq M(n) + (k - 1)M(n) = k M(n).$$

Taking $k = m - 1$ gives $L_m^{\text{dir}}(n) \leq (m - 1)M(n)$, matching the lower bound. $\square$

Nothing above depended on m or on the parity of n : the same three-step construction of R and the same one-line deletion bound run identically at every level. In particular, the induction closes exactly the finite computation the rest of this appendix once needed help to settle.

Corollary A.37. $L_3^{\text{dir}}(7) = 24$.

Proof. $M(7) = \max(12, 12) = 12$, so [Theorem 3.3](#) at $m = 3$, $n = 7$ gives $L_3^{\text{dir}}(7) = 2 \cdot 12 = 24$. Concretely, the induction peels two reachability skeletons off any 7-vertex witness in turn, each of at most $M(7) = 12$ arcs by [Lemma A.35](#), first lowering $\lambda^{\max}$ from 2 to 1 and then from 1 to 0: $24 = 12 + 12$ spends the whole budget twice over, with nothing left for a third pass. This is a proof by hand of the statement [Chapters 2 and 3](#) once recorded as an outstanding computation. That computation was the mixed-integer program `prove_integral_arc_bound(7, 3, 25)`, whose INFEASIBLE verdict would have settled the same fact, and which never returned one: it reached its time limit on the bundled CBC solver, and a fourteen-hour uncapped enumeration was stopped without a verdict. It was never run on a stronger solver, and it no longer needs to be. The machine route to this fact was abandoned unfinished and the hand proof above replaced it, rather than the two confirming each other. $\square$

Corollary A.38 (The fractional relaxation is integral). *For every $n \geq 2$, the m -free optimum $M^*(n)$ of [Chapter 2](#) equals $M(n)$, and it is attained by a $\{0, 1\}$ weight matrix. Consequently the scaling identity*

$$L_m^{\text{dir}}(n) = (m - 1) M^*(n)$$

holds for every n and every $m \geq 2$, with no hypothesis on the optimum.

In words: allowing fractional weights does not raise the optimum. The relaxed optimum is already attained by a multigraph with weights 0 and 1, so the scaling identity that carries one computed number to every m at once needs no integrality hypothesis attached to it.

Proof. For $M^*(n) \geq M(n)$, both constructions of [Construction A.33](#) at multiplicity one are $\{0, 1\}$ matrices with all pairwise maximum flows equal to one, of total weight $2(n - 1)$ and $\lfloor n^2/4 \rfloor$ respectively.

For the reverse, note first that the optimum is attained at a *rational* matrix. A weight matrix is feasible exactly when, for each ordered pair, some cut separating that pair has capacity at most one, so the feasible region is the union, over the finitely many ways of choosing one cut per pair, of the bounded rational polytopes cut out by those choices together with $0 \leq w \leq 1$. A linear

objective over a bounded rational polytope attains its maximum at a rational vertex, and $M^*(n)$ is the largest of finitely many such maxima.

So let w be a feasible rational optimum and let q be a common denominator of its entries. Then qw has integer entries, and multiplying every weight by q multiplies every cut capacity by q and hence every maximum flow by q , exactly as in [Lemma A.32](#). So qw is a directed multigraph with $\lambda^{\max} \leq q$, that is, feasible at $m - 1 = q$, and [Theorem 3.3](#) bounds its arc count by $qM(n)$. Dividing by q gives $M^*(n) = \sum_{u \neq v} w(u, v) \leq M(n)$.

The identity follows because [Theorem 3.3](#) gives $L_m^{\text{dir}}(n) = (m - 1)M(n)$. $\square$

This settles the integrality question [Chapter 2](#) raises when it introduces $M^*(n)$, namely whether the heaviest fractional weight matrix is always a $\{0, 1\}$ one. It is, and the argument runs backwards from the usual direction. Nothing here inspects the fractional problem at all. An integral theorem, proved for every m at once, is strong enough to be applied to the scaled-up matrix qw , and the fractional statement falls out of the integral one rather than the other way round. That is only possible because [Theorem 3.3](#) holds at every m , however large, so no denominator is out of reach.

Corollary A.39 (The $m = 2$ case collapses to the simple digraph). *At $m = 2$ (so $k = 1$), [Theorem 3.3](#) reads $L_2^{\text{dir}}(n) = M(n)$, matching [Theorem 1.9](#) exactly, and this is not a coincidence. A directed multigraph with $\lambda^{\max} \leq 1$ can carry no parallel arcs at all, since two parallel copies of one arc are already two arc-disjoint routes between their endpoints, so at $m = 2$ the multigraph problem and the simple digraph problem are literally the same problem. [Theorem 3.3](#) recovers [Theorem 1.9](#) as its $k = 1$ case, rather than needing it as a separate input.*

[Theorem 3.3](#) settles the value at every n and m , but a value is not a full description of the extremal digraphs, and on the linear branch a companion fact about *which* multigraphs attain it follows from the same argument. Call a directed multigraph *everywhere-saturated* at level m when every ordered pair of distinct vertices has local connectivity exactly $m - 1$. The doubled bidirected spanning trees at multiplicity $m - 1$ form an extremal family on the linear branch and are everywhere-saturated: the $m - 1$ routes between two vertices run along the doubled tree path, and cutting one doubled edge of that path is a cut of $m - 1$ arcs.

Lemma A.40 (Saturated attachment). *Let $m \geq 2$ and let D_0 be a directed multigraph on at least two vertices with $\lambda_{D_0}(x, y) = m - 1$ for every ordered pair of distinct vertices. Let D be obtained from D_0 by adding one new vertex v together with any multiset of arcs incident to v . If $\lambda_D^{\max} \leq m - 1$, then $d(v) \leq 2(m - 1)$. If $d(v) = 2(m - 1)$, then every arc at v joins v to one single vertex u , with $\mu(v, u) = \mu(u, v) = m - 1$, and D is again everywhere-saturated.*

Proof. No mixed pair. Suppose v has an in-arc from u and an out-arc to w with $u \neq w$. By Menger and saturation D_0 carries $m - 1$ pairwise arc-disjoint u -to- w routes, and the route $u \rightarrow v \rightarrow w$ uses only arcs incident to v , so it is arc-disjoint from all of them and $\lambda_D(u, w) \geq m$, a contradiction. Hence v is a pure source, a pure sink, or every arc at v touches one single vertex u .

A pure source has $d(v) \leq m - 1$. Let $t = \sum_x \mu(v, x)$ and fix u with $\mu(v, u) \geq 1$. Any cut $(S, \bar{S})$ with $v \in S$ and $u \in \bar{S}$ has capacity $\sum_{x \in \bar{S}} \mu(v, x) + e_{D_0}(S \setminus \{v\}, \bar{S})$. There are two kinds. If $S = \{v\}$ the capacity is exactly t . If S also contains a vertex x of D_0 , then $S \setminus \{v\}$ and $\bar{S}$ separate x from u inside D_0 , so the second term is at least $\lambda_{D_0}(x, u) = m - 1$, while the first is at least $\mu(v, u)$. Every cut therefore has capacity at least $\min(t, \mu(v, u) + m - 1)$, and by max-flow min-cut so does $\lambda_D(v, u)$, which feasibility caps at $m - 1$. Since $\mu(v, u) \geq 1$, the second argument of that minimum exceeds $m - 1$, so the minimum is t and $t \leq m - 1$.

A pure sink has $d(v) \leq m - 1$. Reverse every arc: the reverse of an everywhere-saturated multigraph is everywhere-saturated, a pure sink becomes a pure source, and the previous step applies.

A single partner. If every arc at v touches one vertex u , then parallel arcs are already that many arc-disjoint one-step routes, so $\mu(v, u) \leq m - 1$ and $\mu(u, v) \leq m - 1$, giving $d(v) \leq 2(m - 1)$.

Equality. Since $2(m - 1) > m - 1$, a degree of $2(m - 1)$ rules out the pure cases, so v has a single partner at full multiplicity both ways. The result is feasible and again everywhere-saturated: a route through v must enter and leave through u , so it adds nothing to any flow between old vertices. For every old $x \neq u$, $\lambda_D(x, v) = \min(\lambda_{D_0}(x, u), \mu(u, v)) = m - 1$; for $x = u$, the $m - 1$ parallel arcs $u \rightarrow v$ give the same conclusion directly. Symmetrically, $\lambda_D(v, x) = m - 1$ for every old x . $\square$

Remark A.41 (The linear branch grows one leaf at a time). The equality case of [Lemma A.40](#) attaches a new leaf to an arbitrary vertex at full multiplicity, and that is the only attachment reaching $2(m - 1)$. Iterating from a single doubled edge, itself everywhere-saturated on two vertices, generates all doubled bidirected spanning trees. Thus this extremal family is closed under maximal attachment and grows one leaf at a time. The lemma does not assert that every linear-branch extremiser admits such a leaf-deletion sequence. The program confirms the lemma exhaustively at $m = 3$: over every doubled tree on five and six vertices and every one of the 3^{2n} possible attachment patterns, the densest feasible attachment has degree exactly $4 = 2(m - 1)$, and the degree-4 patterns are precisely the single-partner attachments at full multiplicity, one per choice of partner.

Remark A.42 (The degree-capped 24-arc extremisers at $n = 7$). [Lemma A.40](#) proves rigidity for maximal one-vertex attachments to an everywhere-saturated graph, but it does not classify every extremiser on the linear branch. At $n = 7$, $m = 3$, the two branches tie at 24 arcs, and every doubled bidirected tree is already an extremiser, giving eleven tree isomorphism classes before the bipartite examples are counted. The separate machine run discussed here answered only a degree-capped question relevant to extension arguments. The tool was the sound generation enumerator of [Chapter 2](#), in the call `enumerate_extremal_directed_multigraphs_via_generation(7, 3, 24, max_degree=8)`, which prunes only with proved bounds and is therefore complete within the imposed maximum-total-degree cap. It ran for about seven hours across ten processor cores and returned exactly three isomorphism classes satisfying that cap. Two of them are the bipartite shapes $2 \cdot B_{3,4}$ and $2 \cdot B_{4,3}$. The third is the doubled bidirected path P_7 , the

one non-star tree whose maximum degree (2) is compatible with an 8-regular graph one vertex up. Each of the three was re-verified by the exact checker. This is a restricted classification, not a classification of all 24-arc extremisers. The reachability-skeleton induction proves the value 24 without it, and the unrestricted extremal family at this tie size is substantially larger.

A.5.3 Extremal classification on the quadratic branch

The classification above is one finite data point, and the linear branch is settled by [Lemma A.40](#), so what is missing is the quadratic branch at a general n . The reachability-skeleton proof looks unpromising for this, since it is designed never to need to know which multigraph it is taking apart. But that indifference is only in the direction of the *bound*. Run backwards, from the assumption that the bound is met exactly, the same construction is unexpectedly rigid: it pins down the number of strongly connected components, then the shape of the skeleton, and finally the multigraph.

Two consequences of equality do the work, and both come from reading the count of [Lemma A.35](#) as an equation rather than an inequality.

Lemma A.43 (What equality forces). *Let $n \geq 8$, so that $\lfloor n^2/4 \rfloor > 2(n-1)$ and $M(n) = \lfloor n^2/4 \rfloor$, and let D be a directed multigraph on n vertices with $\lambda_D^{\max} \leq m-1$ and $|A(D)| = (m-1) \lfloor n^2/4 \rfloor$. Then*

- (1) D is acyclic, and
- (2) D has a reachability skeleton R whose underlying undirected graph is exactly the balanced complete bipartite graph $K_{\lfloor n/2 \rfloor, \lfloor n/2 \rfloor}$, so R is an acyclic orientation of it.

Proof. The proof of [Theorem 3.3](#) peels reachability skeletons $R_1, \dots, R_{m-1}$, each deletion dropping every pairwise connectivity by at least one, so that after $m-1$ peels no arc survives and $|A(D)| = \sum_i |A(R_i)|$. Each $|A(R_i)| \leq M(n)$, so equality in the hypothesis forces $|A(R_i)| = M(n)$ for every i , and in particular for $R := R_1$.

Now read the count of [Lemma A.35](#) backwards. It gives $|A(R)| \leq f(q) := 2(n-q) + \lfloor q^2/4 \rfloor$, where q is the number of strongly connected components of D , so $f(q) \geq \lfloor n^2/4 \rfloor = f(n)$. The increments $f(q+1) - f(q) = -2 + \lfloor (q+1)/2 \rfloor$ are non-decreasing, so f is convex on $q \in \{1, \dots, n\}$. Write $d_i := f(i+1) - f(i)$. Suppose toward a contradiction that $q < n$. If $q = 1$ then $f(q) = f(1) = 2(n-1)$, and the hypothesis $n \geq 8$ gives $\lfloor n^2/4 \rfloor > 2(n-1)$, so $f(q) < f(n)$, contradicting $f(q) \geq f(n)$. So $1 < q < n$. Since $f(1) < f(n) \leq f(q)$, the sum $d_1 + \dots + d_{q-1} = f(q) - f(1)$ is positive, and since the d_i are non-decreasing this forces the last term d_{q-1} to be positive too (otherwise every term up to it would be at most $d_{q-1} \leq 0$ and the sum could not be positive). Non-decreasing again gives $d_q, \dots, d_{n-1} \geq d_{q-1} > 0$, so $f(n) - f(q) = d_q + \dots + d_{n-1} > 0$, that is $f(n) > f(q)$, contradicting $f(q) \geq f(n)$. Both cases are impossible, so $q = n$: every strongly connected component is a single vertex, which is (1).

With $q = n$ the two arborescences of Step 1 of that proof are empty, so R consists of the witness arcs of Q alone and $|A(Q)| = \lfloor n^2/4 \rfloor$. The underlying undirected graph of Q is triangle-free,

proved there from the inclusion-minimality of Q . A triangle-free graph on n vertices with $\lfloor n^2/4 \rfloor$ edges meets Mantel's bound [16] with equality, and the extremal graph for Mantel's theorem is unique: it is $K_{\lfloor n/2 \rfloor, \lfloor n/2 \rfloor}$. That is (2), and R is acyclic because D is. $\square$

The next lemma is where the skeleton's minimality, which had only been used to make it small, is used to make it *shallow*. It costs nothing and it is what makes the theorem work.

Lemma A.44 (A minimal bipartite skeleton has no long path). *Let R be as in Lemma A.43, with parts A and B . Then R contains no directed path with three arcs.*

Proof. Suppose $u \rightarrow v \rightarrow w \rightarrow x$ were such a path. A path in a bipartite graph alternates sides, so u and x lie on opposite sides, and since R 's underlying graph is the *complete* bipartite graph, R contains an arc between them. It cannot be $x \rightarrow u$, which would close a directed cycle against the path, and D is acyclic. So $u \rightarrow x$ lies in R . But then deleting $u \rightarrow x$ from R changes no reachability, since u still reaches x along the path, contradicting the inclusion-minimality of Q . $\square$

Theorem A.45 (Extremal classification on the quadratic branch). *Let $m \geq 2$ and $n \geq \max(8, 2m)$, so that $M(n) = \lfloor n^2/4 \rfloor$ and n is on the quadratic branch. If D is a directed multigraph on n vertices with $\lambda_D^{\max} \leq m-1$ and $|A(D)| = L_m^{\text{dir}}(n) = (m-1) \lfloor n^2/4 \rfloor$, then*

$$D \in \left\{ (m-1) \cdot B_{\lfloor n/2 \rfloor, \lfloor n/2 \rfloor}, (m-1) \cdot B_{\lfloor n/2 \rfloor, \lceil n/2 \rceil} \right\},$$

where $B_{a,b}$ has all arcs from its part of size a to its part of size b , and every displayed arc has multiplicity exactly $m-1$. Thus there is one isomorphism class when n is even and two when n is odd, interchanged by reversing every arc.

In words: past $n = \max(8, 2m)$ the underlying bipartite shape is forced. Split the vertices as evenly as possible, choose either part as the source, run every arc from it to the other part, and repeat each arc $m-1$ times. When n is odd the two choices give two non-isomorphic digraphs, because the source has different size, and they are reversals of one another.

Proof. Take R , A and B from Lemma A.43, and write $\alpha = |A| \geq \beta = |B| = \lfloor n/2 \rfloor$.

Step 1: R has no directed path with two arcs. Suppose $a \rightarrow b \rightarrow a'$ with $a, a' \in A$ and $b \in B$. If a had an incoming arc $b_0 \rightarrow a$ in R , then $b_0 \rightarrow a \rightarrow b \rightarrow a'$ would be a path with three arcs, which Lemma A.44 forbids. So a has no incoming arc in R , and since R 's underlying graph is complete bipartite, $a \rightarrow b''$ for every $b'' \in B$. Symmetrically a' has no outgoing arc, so $b'' \rightarrow a'$ for every $b'' \in B$. The β routes $a \rightarrow b'' \rightarrow a'$, one through each $b'' \in B$, use pairwise distinct arcs, and each of their arcs is present in D because $R \subseteq D$. Hence $\lambda_D(a, a') \geq \beta = \lfloor n/2 \rfloor \geq m$, against feasibility. The case of a path $b \rightarrow a \rightarrow b'$ with middle vertex in A is identical with the sides exchanged and gives $\lambda_D(b, b') \geq \alpha \geq \beta \geq m$, equally impossible. This is the one step that uses $n \geq 2m$.

Step 2: R is one-directional. By Step 1 no vertex of R has both an incoming and an outgoing arc, so every vertex is a source or a sink. Fix $a \in A$. It is adjacent in R to every vertex of B , so it

is not isolated. If a is a source then every $b \in B$ has an incoming arc, hence is a sink, hence has no outgoing arc, so every arc at every b runs into it: all arcs of R run from A to B . If instead a is a sink the same argument gives all arcs from B to A . Thus R is either $B_{\alpha,\beta}$ or $B_{\beta,\alpha}$, according to which part is the source. The remaining steps apply identically to either orientation, so from now on use A for the source part and B for the sink part without imposing an order on their sizes.

Step 3: D has no arc outside R . R has the same reachability as D , and in a one-directional bipartite digraph the only ordered pairs joined by a directed path are the pairs $(a, b) \in A \times B$, each by the single arc $a \rightarrow b$. So if D had an arc (u, x) , then u reaches x in R , forcing $(u, x) \in A \times B$, and R already contains that arc. Hence D and R use exactly the same $\alpha\beta = \lfloor n^2/4 \rfloor$ ordered pairs.

Step 4: every multiplicity is $m - 1$. A pair joined by $\mu(a, b)$ parallel arcs has $\lambda_D(a, b) \geq \mu(a, b)$, so feasibility gives $\mu(a, b) \leq m - 1$ on each of the $\lfloor n^2/4 \rfloor$ pairs. Their total is $|A(D)| = (m - 1) \lfloor n^2/4 \rfloor$, so every one of them attains the ceiling. Hence D is the multiplicity- $(m - 1)$ one-directional complete bipartite digraph on these two parts, and their product $\lfloor n^2/4 \rfloor$ forces their sizes to be $\lceil n/2 \rceil$ and $\lfloor n/2 \rfloor$. Either size may be the source size, giving exactly the two displayed possibilities (which coincide up to isomorphism when n is even). $\square$

Three remarks. First, on where the hypothesis $n \geq 2m$ is used and what it costs. It enters once, in Step 1, where a two-arc path in the skeleton is converted into $\lfloor n/2 \rfloor$ arc-disjoint routes between one pair, and feasibility kills that only when $\lfloor n/2 \rfloor$ reaches m . For $n < 2m$ the theorem is not proved, and it is not known to fail either. Nothing in the argument suggests a boundary there: the step is a lower bound on one pair's connectivity, and losing it removes a contradiction rather than supplying a counterexample. Second, a degree fact falls out of the same equality, and it is what makes a direct search for extremisers finite in practice. Deleting a vertex leaves at most $(m - 1)M(n - 1)$ arcs, so every vertex of an extremiser carries degree at least $(m - 1) \lfloor n/2 \rfloor$. For even n the degree sum $2(m - 1) \lfloor n^2/4 \rfloor = n \cdot (m - 1)(n/2)$ leaves no slack at all, so every vertex has degree exactly $(m - 1)n/2$: an extremiser at even n is regular. Third, on the shape of the argument. [Theorem 3.3](#) is indifferent to which multigraph it dismantles, and [Chapter 3](#) records that indifference as the reason the value says nothing about uniqueness. That is true of the proof read forwards. Read backwards from equality, the same skeleton is a rigid object, and the two facts that had been mere bookkeeping, that f is maximised at one end and that a minimal skeleton is triangle-free, become a determination of q and an application of the equality case of Mantel's theorem.

Part III. The hypergraph results

The edge bound through a hypergraph Gomory–Hu tree, the vertex problem at $m = 2$ and $m = 3$ by way of an incidence rank bound, and the directed hypergraph leading term.

A.6 The hypergraph bounds

The body's hypergraph claims rest on two theorems: a Menger theorem for Berge routes, which also certifies the helper-point checker of [Chapter 2](#), and a hypergraph Gomory–Hu theorem, which is the classical construction of Gomory and Hu [4] run on a symmetric submodular cut function rather than on the edge-cut function, as set out by Hanifnezhad and Dolati [18].

Theorem A.46 (Menger for hypergraphs). *For a hypergraph $\mathcal{H}$ and vertices u, v , the maximum number of hyperedge-disjoint Berge u – v paths equals the minimum size of a set C of hyperedges whose removal separates u from v .*

Proof. Build the helper-point network of [Figure 2.3b](#): for each hyperedge e a helper arc $e^- \rightarrow e^+$ of capacity one, and arcs $x \rightarrow e^-$ and $e^+ \rightarrow x$ of unbounded capacity for each member $x \in e$. The proof matches the number of disjoint Berge paths to the size of a minimum cut by translating between Berge-path language and flow language in each direction: a family of hyperedge-disjoint Berge paths becomes a family of arc-disjoint flow paths, and conversely a flow becomes a family of Berge paths, as the next two paragraphs spell out in turn.

Berge paths become flow. A Berge path $u, e_1, v_1, \dots, e_k, v$ becomes the directed walk $u \rightarrow e_1^- \rightarrow e_1^+ \rightarrow v_1 \rightarrow \dots \rightarrow v$, crossing one helper arc per hyperedge it uses. The e_i along a single Berge path are distinct, and two hyperedge-disjoint Berge paths share no hyperedge at all, so they use disjoint sets of helper arcs and translate to *arc-disjoint* flow paths, paths that share no arc, each carrying one unit of flow.

Flow becomes Berge paths. Conversely, because every capacity is an integer, a maximum flow may be taken integral, and an integral flow of value k splits into k arc-disjoint unit paths. Reading a helper crossing $\dots x \rightarrow e^- \rightarrow e^+ \rightarrow y \dots$ back as the single hyperedge step x, e, y , that is, *suppressing* the two helper nodes, turns each unit path into a Berge path, and no two of them can share a hyperedge, since that would send two units across some capacity-one helper arc.

Cuts. Finally, the member arcs are uncapped, so any cut of finite capacity uses helper arcs only, and a set of helper arcs separates u from v in the network exactly when the corresponding hyperedges separate them in $\mathcal{H}$. The max-flow min-cut theorem equates the largest number of arc-disjoint flow paths with the smallest helper-arc cut, which is precisely the claim. $\square$

Theorem A.47 (Hypergraph Gomory–Hu). *Every r -uniform hypergraph $\mathcal{H}$ on vertex set V has a weighted tree T^* on V , its hypergraph Gomory–Hu tree, such that $\lambda_{\mathcal{H}}(u, v)$ equals the minimum weight on the u – v tree path, and each tree edge t induces a vertex partition $(S_t, V \setminus S_t)$ with $|\delta_{\mathcal{H}}(S_t)| = c^*(t)$, where $\delta_{\mathcal{H}}(S)$ is the set of hyperedges incident to both sides, that is, having at least one vertex in S and at least one outside it.*

This is the exact hypergraph echo of [Theorem A.2](#), the same tree-of-cuts summary of all pairwise connectivities, with the single change that the capacity of a cut now counts the hyperedges straddling it, $|\delta_{\mathcal{H}}(S)|$, in place of the edges. Gomory and Hu's construction [4] needs nothing

about the cut function beyond it being symmetric and submodular [18], and the hyperedge-cut function $|\delta_{\mathcal{H}}(\cdot)|$ has both properties exactly as the ordinary edge-cut function does, so the identical uncrossing construction builds this tree too. We use only the two properties in the statement, the path-minimum reading of $\lambda_{\mathcal{H}}$ and the identity $|\delta_{\mathcal{H}}(S_t)| = c^*(t)$.

Proof of Proposition 1.11. Upper bound. Let $\mathcal{H}$ be r -uniform with $\lambda_{\mathcal{H}}^{\max} \leq m - 1$, and let T^* be its hypergraph Gomory–Hu tree (Theorem A.47), a weighted tree on the same n vertices with weights c^* . The plan is to charge each hyperedge to tree edges and count the charges two ways, exactly as in the Mader proof, with the one change that a hyperedge spans r vertices rather than two.

Step 1: each hyperedge crosses at least $r - 1$ tree cuts. Fix a hyperedge e , a set of r vertices, and let $\text{ST}(e)$ be its *smallest connected subtree*, the minimal subtree of T^* that contains all r of them (its Steiner tree). A tree spanning at least r vertices has at least $r - 1$ edges, so $\text{ST}(e)$ has at least $r - 1$ of them. Each such edge t is a genuine cut for e : deleting t from T^* splits the vertices into the two sides $(S_t, V \setminus S_t)$, and because t lies on $\text{ST}(e)$ the hyperedge e has at least one vertex on each side. So e touches both sides, that is, $e \in \delta_{\mathcal{H}}(S_t)$. Therefore

$$|\{t \in E(T^*) : e \in \delta_{\mathcal{H}}(S_t)\}| \geq r - 1 \quad \text{for every hyperedge } e.$$

Step 2: count the incidences two ways. Sum that inequality over all hyperedges, then exchange the order of summation. This is legitimate because both of the middle sums below simply count the same pairs (e, t) with $e \in \delta_{\mathcal{H}}(S_t)$, once grouped by the hyperedge and once by the tree edge:

$$(r - 1)|E(\mathcal{H})| \leq \sum_e |\{t : e \in \delta_{\mathcal{H}}(S_t)\}| = \sum_t |\{e : e \in \delta_{\mathcal{H}}(S_t)\}| = \sum_t |\delta_{\mathcal{H}}(S_t)|.$$

Step 3: read off the bound. By the Gomory–Hu property each tree edge has $|\delta_{\mathcal{H}}(S_t)| = c^*(t)$. Moreover $c^*(t)$ is itself a local connectivity $\lambda_{\mathcal{H}}$ of t 's two endpoints: since those endpoints are adjacent in T^* , the u – v tree path between them is the single edge t , so the path-minimum reading of Theorem A.47 gives $\lambda_{\mathcal{H}}$ of that pair exactly $c^*(t)$, hence at most $m - 1$ by feasibility. Summing over the $n - 1$ tree edges, $\sum_t |\delta_{\mathcal{H}}(S_t)| = \sum_t c^*(t) \leq (m - 1)(n - 1)$. Chaining Steps 2 and 3,

$$(r - 1)|E(\mathcal{H})| \leq (m - 1)(n - 1),$$

and dividing by $r - 1$, then rounding down because $|E(\mathcal{H})|$ is an integer, gives $|E(\mathcal{H})| \leq \left\lfloor \frac{(m-1)(n-1)}{r-1} \right\rfloor$. *Lower bound.* When $(r - 1) \mid (n - 1)$, take a hub h and split the other $n - 1$ vertices into groups $G_1, G_2, \dots$ of size $r - 1$ each. Form the star hypertree whose hyperedges are $\{h\} \cup G_i$, one per group, and give each hyperedge multiplicity $m - 1$. Every non-hub vertex lies in exactly $m - 1$ hyperedges, which form a cut isolating it, so by Theorem A.46 every pair has Berge connectivity at most $m - 1$, while the count is $\frac{(m-1)(n-1)}{r-1}$. For general n under the hypothesis $m - 1 \leq \binom{n-2}{r-2}$, Theorem A.48 below attains the floor exactly, without even needing

repeated hyperedges. □

The multiplicities in the lower bound can in fact be avoided: for $r \geq 3$, *simple* r -uniform hypergraphs already attain the same bound, in sharp contrast to the graph case $r = 2$, where simplicity costs the factor that separates Mader's $\lfloor m(n-1)/2 \rfloor$ from the multigraph value $(m-1)(n-1)$.

Theorem A.48 (Simplicity is free for $r \geq 3$). *Let $r \geq 3$, $n \geq r$, and $2 \leq m$ with $m-1 \leq \binom{n-2}{r-2}$. Then the maximum number of hyperedges in a simple r -uniform hypergraph on n vertices with Berge edge-connectivity at most $m-1$ is exactly $\left\lfloor \frac{(m-1)(n-1)}{r-1} \right\rfloor$.*

Proof. The upper bound is [Proposition 1.11](#), since a simple hypergraph is a multihypergraph. For the lower bound, fix a hub h and let $V' = V \setminus \{h\}$. By [Lemma A.51](#) applied with rank $r-1$ and degree ceiling $d = m-1$ on the $n-1$ vertices of V' , there is an $(r-1)$ -uniform simple hypergraph $\mathcal{G}^*$ on V' with maximum degree at most $m-1$ and exactly $\left\lfloor \frac{(m-1)(n-1)}{r-1} \right\rfloor$ hyperedges. Adding h to each hyperedge produces distinct r -sets, so the result $\mathcal{H}$ is simple, and every non-hub vertex w lies in $\deg_{\mathcal{G}^*}(w) \leq m-1$ hyperedges, which form a cut isolating w . By [Theorem A.46](#), $\lambda_{\mathcal{H}}^{\max} \leq m-1$. □

Remark A.49 (The hypothesis is not decorative). Outside this range the displayed value can fail for simple hypergraphs, even though it stays a valid upper bound ([Proposition 1.11](#)) and, when $(r-1) \mid (n-1)$, stays exactly attained by a multihypergraph. The smallest case already shows it. At $n = r = m = 3$ there is only one 3-subset of a 3-element vertex set, so a simple hypergraph carries at most one hyperedge. The hypothesis needs $m-1 \leq \binom{n-2}{r-2} = \binom{1}{1} = 1$, and $m-1 = 2$ misses it, so the formula's value $\lfloor 2 \cdot 2/2 \rfloor = 2$ is not attained by any simple hypergraph there. It is attained by the multihypergraph star, two parallel copies of that one triple. [Theorem 1.13](#) carries the identical hypothesis for the vertex separation, for the same reason. [Table 3.1](#) splits the simple and multi hypergraph rows on exactly this distinction, and the program enumerates the two models separately for the same reason, stated where each formula is used.

The sparse hypergraph the construction needs is a corollary of a classical partition theorem, which we state before using it.

Theorem A.50 (Baranyai [19]). *If r divides n , the $\binom{n}{r}$ distinct r -subsets of an n -element set can be partitioned into $\binom{n-1}{r-1}$ classes, each a perfect matching, meaning a family of n/r disjoint r -sets that together cover every vertex exactly once. More generally, for any prescribed class sizes $a_1 + \dots + a_t = \binom{n}{r}$, and with no divisibility assumed, the r -subsets can be partitioned into classes of exactly those sizes so that in the class of size a_i every vertex lies in either $\lfloor ra_i/n \rfloor$ or $\lceil ra_i/n \rceil$ of the sets, as evenly balanced as the size permits.*

The first statement is the memorable one. When r divides n the complete r -uniform hypergraph, the one holding every r -set, comes apart into $\binom{n-1}{r-1}$ perfect matchings, the exact hypergraph analogue of splitting a regular bipartite graph into perfect matchings by König's theorem. The proof is an induction that keeps the partial partition as balanced as possible at every step,

an integer-flow argument in the spirit of Hall's marriage theorem rather than an explicit construction, and its full form is in Baranyai [19]. We need only the general equitable form, and only its conclusion about degrees.

Lemma A.51 (Sparse uniform hypergraphs exist). *Let $r \geq 1$, $n \geq r$, and $0 \leq d \leq \binom{n-1}{r-1}$. There exists an r -uniform simple hypergraph on n vertices with maximum degree at most d and exactly $\lfloor \frac{dn}{r} \rfloor$ hyperedges.*

Proof. For $r = 1$ the hypotheses force $d \in \{0, 1\}$, and the empty family or the family of all n singleton edges, respectively, has the required size and degree. Assume henceforth that $r \geq 2$. Put $e := \lfloor \frac{dn}{r} \rfloor \leq \frac{n}{r} \binom{n-1}{r-1} = \binom{n}{r}$. Apply the equitable form of Theorem A.50 with two classes, of sizes e and $\binom{n}{r} - e$. Within each class every vertex is covered either $\lfloor re/n \rfloor$ or $\lceil re/n \rceil$ times, so the class of size e is a simple r -uniform hypergraph with $e = \lfloor \frac{dn}{r} \rfloor$ hyperedges and maximum degree at most $\lceil re/n \rceil \leq \lceil d \rceil = d$, the last step because $re/n \leq r(dn/r)/n = d$ and d is an integer. $\square$

A.7 The hypergraph vertex problem


The whole vertex problem translates into a graph problem through one observation, the one the rest of this section rests on and which Figure A.4 draws in full. Write $I(\mathcal{H})$ for the bipartite incidence graph of a hypergraph $\mathcal{H}$: one node per vertex, one node per hyperedge, an edge when the vertex lies in the hyperedge. A Berge path with distinct hyperedges is exactly a path of $I(\mathcal{H})$ between two vertex-class nodes. The interior of such a path consists of both kinds of node, the hyperedges it rides and the vertices where it changes hyperedge, so two paths of $I(\mathcal{H})$ are internally disjoint in the ordinary graph sense exactly when the two Berge routes share neither a hyperedge nor an intermediate vertex. That hybrid is what $\kappa_{\mathcal{H}}$ means throughout, and both halves are needed: dropping the hyperedge half would let two routes ride the same hyperedge, which a single Berge edge cannot serve twice. Hence

$$\kappa_{\mathcal{H}}(u, v) = \kappa_{I(\mathcal{H})}(u, v),$$


the maximum number of internally disjoint u - v paths in $I(\mathcal{H})$, and the hypergraph vertex problem asks for the maximum number of degree- r nodes on one side of a bipartite graph whose other side holds n nodes, subject to no two of those n nodes being joined by m internally disjoint paths. Figure A.4a traces a single Berge path through both pictures, and Figure A.4b shows the star hypertree of Theorem 1.12, whose incidence graph is a tree.

Theorem 1.12, stated in Chapter 1, follows at once from the incidence translation.

Proof of Theorem 1.12. We claim $\kappa_{\mathcal{H}}^{\max} \leq 1$ holds if and only if $I(\mathcal{H})$ is a forest. If $I(\mathcal{H})$ contains a cycle $v_1, e_1, v_2, e_2, \dots, v_\ell, e_\ell, v_1$ (it alternates classes, so $\ell \geq 2$, all v_i and e_i distinct), then v_1, e_1, v_2 and $v_1, e_\ell, v_\ell, e_{\ell-1}, \dots, e_2, v_2$ are two Berge v_1 - v_2 paths with disjoint hyperedge sets and disjoint interior vertex sets, so $\kappa(v_1, v_2) \geq 2$. (At $\ell = 2$ this is the repeated pair: two hyperedges sharing two vertices.) Conversely, two such paths between u and v concatenate to



(a) A 3-uniform hypergraph $\mathcal{H}$ (left) and its bipartite incidence graph $I(\mathcal{H})$ (right): circles for vertices, hollow squares for hyperedges, an edge when the vertex lies in the hyperedge. The Berge path $1-e_1-3-e_2-5$ is traced in orange in both pictures. In $\mathcal{H}$ it rides the red line e_1 and the blue line e_2 , changing at the shared vertex 3, while the grey line e_3 lies off it. In $I(\mathcal{H})$ it is the ordinary path $1-e_1-3-e_2-5$ alternating sides.



(b) A star hypertree: every hyperedge $\{h\} \cup G_i$ shares only the hub h (orange), and the other $r-1=2$ vertices of each block sit in a single hyperedge. Its incidence graph is a tree, so $\kappa^{\max} \leq 1$.

Figure A.4. The translation the vertex problem rests on. Internally vertex-disjoint Berge paths in $\mathcal{H}$ correspond exactly to internally disjoint paths in $I(\mathcal{H})$, so $\kappa_{\mathcal{H}}(u, v) = \kappa_{I(\mathcal{H})}(u, v)$. **(top)** A hypergraph and its incidence graph with one Berge path drawn in both. **(bottom)** The star hypertree of Theorem 1.12, whose incidence graph is a tree.

a cycle of $I(\mathcal{H})$. This proves the claim. A forest on $n + q$ nodes has at most $n + q - 1$ edges, while $I(\mathcal{H})$ has exactly rq edges, so $rq \leq n + q - 1$, i.e. $q \leq (n - 1)/(r - 1)$, and q is an integer. The star hypertree has $\lfloor (n - 1)/(r - 1) \rfloor$ hyperedges and a forest as incidence graph (one tree together with any unused isolated vertices) because each block vertex lies in one hyperedge and every cycle would need two hyperedges sharing two vertices, so the floor is attained. $\square$

Proposition A.52 (General lower bound). *For $r \geq 3$, $m \geq 2$ and $m - 1 \leq \binom{n-2}{r-2}$,*

$$k_m^{(r)}(n) \geq \left\lfloor \frac{(m-1)(n-1)}{r-1} \right\rfloor,$$

attained by a simple hypergraph.

Proof. The simple construction of [Theorem A.48](#) has Berge edge-connectivity at most $m - 1$, and Whitney's inequality $\kappa \leq \lambda$ holds pairwise (an internally vertex-disjoint family is in particular hyperedge-disjoint under our definition), so the same hypergraph is feasible for the vertex problem. $\square$

At $m = 3$ the upper-bound side closes completely, and equality follows in the construction range stated below. The engine is a rank bound for graphs in which one side of a partition must stay below local 3-connectivity. The hypergraph theorem then falls out of the incidence translation. Throughout, $\text{rank}(G) = |E(G)| - |V(G)| + 1$ denotes the cycle rank of a connected multigraph, that is, the number of independent cycles it carries. A tree has cycle rank 0, and each edge added beyond a spanning tree closes exactly one new cycle and raises the rank by one, so bounding the rank is the same as bounding how many independent cycles the graph can hold.

Primer: global connectivity and the pieces of a graph

The lemma below speaks of a graph being 2-connected or 3-connected and of taking it apart at small separators. These are the global cousins of the local $\kappa(u, v)$ from [Chapter 1](#), so we fix the words once. A graph on more than k vertices is *k-connected* if it stays connected after the removal of any $k - 1$ vertices. By Menger this is the same as asking that every pair of vertices be joined by at least k internally vertex-disjoint paths, so *k-connected* means $\kappa(u, v) \geq k$ for all pairs at once, the uniform version of the local measure. Thus 2-connected means no single vertex disconnects the graph, and 3-connected means no two vertices do. Three named obstructions go with this. A *cut vertex* is a vertex whose removal disconnects the graph, so a connected graph on at least three vertices is 2-connected exactly when it has no cut vertex. A *separating pair* is a set of two vertices whose removal disconnects the graph, the obstruction a 2-connected graph must lack in order to be 3-connected. Finally a *block* is a maximal piece with no cut vertex of its own, hence either a single bridge edge or a maximal 2-connected subgraph. Two blocks overlap in at most one vertex, always a cut vertex, and the blocks and cut vertices hang together in a tree, the *block-cut tree*, with one node per block and one per cut vertex. Step 1 of the proof is precisely an induction over that tree.

Lemma A.53 (Incidence rank lemma). *Let G be a connected multigraph with vertex set $X \cup Z$ such that (i) Z is independent, (ii) no edge incident to Z is parallel to another edge, (iii) every $z \in Z$ has degree at least 2, and (iv) $\kappa_G(x, x') \leq 2$ for all distinct $x, x' \in X$. Then $\text{rank}(G) \leq |X| - 1$.*

In words: this is the engine behind the $m = 3$ hypergraph result, stated for a bipartite graph rather than for a hypergraph. Read X as the vertices of a hypergraph and Z as its hyperedges, so that G is the incidence graph and $\text{rank}(G)$, the number of independent cycles it carries, is the quantity the hyperedge count is eventually read off from. The four hypotheses say in turn that the picture really is an incidence graph (i), that no hyperedge repeats a vertex (ii), that no hyperedge dangles as a leaf (iii), and that the hypergraph is feasible at $m = 3$ (iv). The conclusion caps the independent cycles by the number of vertices.

Proof. Induction on $|V| + |E|$. The proof peels G down to a core that cannot exist. Step 1 handles a cut vertex by arguing block by block. Steps 2 and 3 remove a degree-two Z -vertex or X -vertex without changing the bound. What survives is 2-connected with minimum degree at least three, and Step 4 shows that such a graph already violates the connectivity ceiling (iv), so it never arises. *Base* $|V| \leq 2$. A single vertex has degree 0, too little for the degree-at-least-2 floor (iii) imposes on a Z -vertex, so it cannot lie in Z and must lie in X , giving $0 \leq |X| - 1$. On two vertices the same reasoning applies to either one: a vertex in Z could reach only the other vertex, and (ii) forbids that single edge from being parallel to another, so it would have degree at most 1, again short of (iii). Hence (i)–(iii) force both vertices into X . Parallel edges between them are internally disjoint paths (each has empty interior, so no two can share an interior vertex), so (iv) caps the multiplicity at 2 and $\text{rank} \leq 1 = |X| - 1$.

Step 1: a cut vertex. Suppose G has a cut vertex, so it is not yet 2-connected. Break it into its *blocks* $B_1, \dots, B_c$ with $c \geq 2$, a block being a maximal piece with no cut vertex of its own, that is, either a single bridge edge or a maximal 2-connected subgraph. Two blocks meet in at most one vertex, and any shared vertex is a cut vertex of G . No cycle can pass through two different blocks: since two blocks share at most one vertex, a cycle straddling both would have to visit that shared vertex twice, which a simple cycle cannot do. So the cycle rank adds up over blocks, $\text{rank}(G) = \sum_i \text{rank}(B_i)$, and it is enough to bound each $\text{rank}(B_i)$ by $|X \cap B_i| - 1$ and sum.

Each block inherits the hypotheses. A bridge block is one edge, so its rank is 0, and by (i) at least one endpoint lies in X , giving $0 \leq |X \cap B| - 1$. Every other block is 2-connected, hence has minimum degree at least 2, so (i) to (iv) hold inside it and the induction hypothesis gives $\text{rank}(B_i) \leq |X \cap B_i| - 1$.

Adding these requires care, because a cut vertex is shared by several blocks and would be counted several times. Write $b(v)$ for the number of blocks containing v , so $b(v) = 1$ except at cut vertices. Counting each X -vertex once for every block it lies in,

$$\sum_i |X \cap B_i| = |X| + \sum_{x \in X \text{ cut}} (b(x) - 1).$$

The number of blocks satisfies the block-cut tree identity $c = 1 + \sum_{v \text{ cut}} (b(v) - 1)$. This follows

from counting the edges of the block-cut tree two ways. That tree has c block-nodes and one node per cut vertex, so being a tree its edge count is one less than its total number of nodes, $c + \#\{\text{cut vertices}\} - 1$. The same edge count also equals $\sum_{v \in \text{cut}} b(v)$, since each cut vertex v is joined in the tree to exactly the $b(v)$ blocks that contain it. Equating the two expressions for the edge count and solving for c gives the identity. Subtracting this count of blocks from the previous display and separating the cut vertices into their X - and Z -parts leaves

$$\text{rank}(G) = \sum_i \text{rank}(B_i) \leq \sum_i (|X \cap B_i| - 1) = |X| - 1 - \sum_{z \in Z \text{ cut}} (b(z) - 1) \leq |X| - 1,$$

where the last inequality holds because each $b(z) - 1 \geq 0$, so the Z -cut sum only pushes the bound further below $|X| - 1$.

Step 2: a Z -vertex of degree two. From now on G is 2-connected, every cut vertex having been handled. Suppose some $z_0 \in Z$ has degree exactly 2. Both its neighbours lie in X by (i), and they are distinct, since two edges to the same vertex would be parallel at a Z -vertex, which (ii) forbids. *Suppress* z_0 by deleting it and joining its two neighbours x, x' with one new edge. This loses two edges and one vertex and gains one edge, so $|E|$ and $|V|$ each drop by one and $\text{rank} = |E| - |V| + 1$ is unchanged. The new edge runs X to X , so $|X|$ is unchanged and (i), (ii), (iii) still hold for the remaining Z -vertices. Replacing the path x, z_0, x' by a direct edge alters no route among the surviving vertices, so every $\kappa(x_1, x_2)$ is preserved and (iv) holds. The graph is strictly smaller, so the induction hypothesis applies and returns the bound for G .

Step 3: an X -vertex of degree two. Now G is 2-connected and, Step 2 being used up, every Z -vertex has degree at least 3. Suppose some $x_0 \in X$ has degree 2, and delete it. Deleting one vertex from a 2-connected graph leaves it connected. Stripping a degree-2 vertex removes two edges and one vertex, so rank drops by exactly one. Each neighbour of x_0 loses one edge, and a neighbour in Z only falls from degree at least 3 to degree at least 2, so (iii) survives, while deleting a vertex cannot raise a connectivity, so (iv) survives. Here $|X|$ drops by one, and $|X| \geq 2$ to begin with, for if $|X| \leq 1$ then by (i) and (ii) every z would send all its edges to one X -vertex without repeats, hence have degree at most 1, against (iii). The smaller graph satisfies the hypotheses, so induction gives $\text{rank}(G) - 1 \leq (|X| - 1) - 1$, that is $\text{rank}(G) \leq |X| - 1$.

Step 4: the remaining case is impossible. Now G is 2-connected with $|V| \geq 3$ and every degree at least 3. First, G is simple: a parallel class lies inside X by (ii), and if $\mu(x, x') \geq 2$ then a third internally disjoint route exists: walk from x to any third vertex w inside $G - x'$, then from w to x' inside $G - x$, both walks available because deleting a single vertex from a 2-connected graph leaves it connected, exactly as already used in Step 3. The concatenation visits x only first and x' only last, so it contains an $x-x'$ path with at least one interior vertex, giving $\kappa(x, x') \geq 3$, against (iv). If G is 3-connected, Menger puts every pair at $\kappa \geq 3$, so two vertices of X would violate the ceiling $\kappa(x, x') \leq 2$ that (iv) imposes on every pair drawn from X , forcing $|X| \leq 1$, and then every z has degree at most 1 by (i), absurd.

Otherwise G is 2-connected but not 3-connected. Tutte's triconnected decomposition [20] expresses it as a tree of cycle, bond, and 3-connected pieces joined along virtual edges. For

a leaf piece L its unique virtual edge $\{a, b\}$ represents a real a - b path through the rest of G , internally outside L . Vertices of L away from that edge keep their degree in G . These are the only properties of the decomposition used below.

Now G itself is none of the three piece types, since it is not a single cycle (its degrees are at least three), not a bond (it is simple), and not 3-connected (the case just handled), so the tree has at least two leaves. Let L be one, with unique virtual edge $\{a, b\}$. If L is a cycle, a vertex of L off $\{a, b\}$ has degree 2 in G , absurd. If L is a bond, it holds at least two real parallel edges, absurd in a simple graph. If L is 3-connected, then any two of its vertices are joined by three internally disjoint paths inside L , at most one through the virtual edge, which the far-side path replaces: so $\kappa_G(u, v) \geq 3$ for all $u, v \in V(L)$, forcing $|X \cap V(L)| \leq 1$ by (iv). But $|V(L)| \geq 4$ leaves a Z -vertex $z^* \notin \{a, b\}$. Its neighbours lie in $X \cap V(L)$ by (i), at most one vertex, reached by at most one edge since G is simple, giving degree at most 1, which is absurd. $\square$

The rank bound now yields [Theorem 1.13](#), stated in [Chapter 1](#). In words: at the route limit $m = 3$ a hyperedge network can hold no more hyperedges under the stricter separation than the displayed edge bound. When $r \geq 3$ and $2 \leq \binom{n-2}{r-2}$, that count is reached without repeating a hyperedge, so the two separations agree in this range exactly as they did at $m = 2$.

Proof of Theorem 1.13. Apply [Lemma A.53](#) to each connected component of the incidence graph $I(\mathcal{H})$, with X the vertex class and Z the hyperedge class: Z is independent, incidences are simple even when hyperedges repeat, hyperedge-nodes have degree $r \geq 2$, components without an X -node are impossible, and $\kappa_I(u, v) = \kappa_{\mathcal{H}}(u, v) \leq 2$ for vertex pairs. Sum rank $\leq |X_c| - 1$ over the C components. The incidence graph $I(\mathcal{H})$ has rq edges (each of the q hyperedges contributes r incidences) and $n + q$ nodes, so $\sum_c \text{rank} = rq - (n + q) + C$, while $\sum_c (|X_c| - 1) = n - C$. The inequality reads $rq - (n + q) + C \leq n - C$, hence $q(r - 1) \leq 2(n - 1) + 2(1 - C) \leq 2(n - 1)$, the last step using $C \geq 1$. The lower bound is [Proposition A.52](#) at $m = 3$. $\square$

Remark A.54 (Edges as gates at $r = 2$, and the boundary of the method). At $r = 2$ the theorem speaks of multigraphs under the hyperedge-as-gate convention, in which parallel edges are distinct one-step routes, giving the multigraph value $2(n - 1)$ of [Theorem 1.3](#), complementing the body's convention for graphs, where parallel adjacencies count once. The proof of [Lemma A.53](#) leans on the triconnected decomposition exactly where $\kappa \leq 2$ lives. For $m = 4$, the analogous question asks for a rank bound relative to $\kappa \leq 3$, where no comparably clean decomposition exists. Thus the proof is confined to $m \leq 3$, and no general $m = 4$ formula is claimed.

A.8 The directed hypergraph

The remaining four of the sixteen variants orient the Berge edge. They are grouped here rather than with the undirected hypergraph bounds above because the questions they raise are different ones: not which decomposition controls the rank of an incidence graph, but how much a single edge with one tail and many heads can carry, and whether the way it is oriented changes the

answer. The first result is a pair of bounds a factor of four apart, the last closes that factor for the model the program measures.

Proposition A.55 (Directed hypergraphs: first bounds). *Let $n \geq r \geq 2$ and $m \geq 2$. Model a directed r -uniform hyperedge as a tail vertex together with $r - 1$ head vertices, a Berge route entering at the tail and leaving at a head. If every ordered pair of vertices has at most $m - 1$ hyperedge-disjoint directed Berge routes, then*

$$|E(\mathcal{H})| \leq \frac{(m-1)n(n-1)}{r-1},$$

and, for every $1 \leq \alpha \leq n - r + 1$, there is a feasible directed multihypergraph with

$$\alpha \left\lfloor \frac{(m-1)(n-\alpha)}{r-1} \right\rfloor$$

hyperedges. This construction is simple whenever $m - 1 \leq \binom{n-\alpha-1}{r-2}$. Consequently, for fixed r and m , the balanced choice $\alpha = \lfloor n/2 \rfloor$ is available for all sufficiently large n and proves that the extremal value is $\Theta_{r,m}(n^2)$.

Proof. Upper bound: a hyperedge with tail u serves the $r - 1$ ordered pairs (u, h) , h a head. If some ordered pair (u, v) were served by m distinct hyperedges, those would be m hyperedge-disjoint one-step routes, so each ordered pair is served at most $m - 1$ times. Summing that cap over all $n(n - 1)$ ordered pairs counts every hyperedge once for each of its $r - 1$ tail-head pairs, so the sum is $(r - 1)|E|$ on one side and at most $(m - 1)n(n - 1)$ on the other, giving $(r - 1)|E| \leq (m - 1)n(n - 1)$.

For the lower bound, split V into tails A , $|A| = \alpha$, and heads B , where $b := |B| = n - \alpha \geq r - 1$. Put $k := r - 1$, $d := m - 1$ and $C := \binom{b-1}{k-1}$, the degree of a vertex in the complete k -uniform hypergraph on B . Write $d = qC + s$ with $0 \leq s < C$. Take q copies of that complete hypergraph and, by Lemma A.51, a simple k -uniform hypergraph on B of maximum degree at most s and with $\lfloor sb/k \rfloor$ edges. Their multiset union $\mathcal{G}^*$ has maximum degree at most $qC + s = d$ and

$$q \binom{b}{k} + \left\lfloor \frac{sb}{k} \right\rfloor = q \frac{bC}{k} + \left\lfloor \frac{sb}{k} \right\rfloor = \left\lfloor \frac{db}{k} \right\rfloor$$

hyperedges. Give every tail $a \in A$ the hyperedges (a, H) for $H \in \mathcal{G}^*$, with the same multiplicities. No vertex of B is a tail, so every route is a single step and $\lambda(a, b) = \deg_{\mathcal{G}^*}(b) \leq m - 1$, while pairs inside A or starting in B have no routes. If $d \leq C$, the auxiliary hypergraph can instead be taken directly from Lemma A.51, so the resulting directed hypergraph is simple. $\square$

The forward model is one of three ways to orient a Berge edge. In general a directed r -uniform hyperedge is a split of its r vertices into a non-empty *tail set* T and a non-empty *head set* H , a route entering at a tail and leaving at a head. The *forward* model is $|T| = 1$, the *backward* model is $|H| = 1$, and the *general* model allows any split (genuinely mixed splits, both parts at least two, exist from $r \geq 4$). The next two results place all three. The first is that backward needs no

separate study, and the second is that the general model, despite admitting more hyperedges on a single vertex set, obeys the very same first-order bound as forward.

Lemma A.56 (Backward is the reverse of forward). *Reversing every hyperarc, $(T, H) \mapsto (H, T)$, is an edge-count-preserving bijection between the forward directed r -uniform hypergraphs and the backward ones on the same vertices, and the reverse $\mathcal{H}^R$ satisfies $\lambda_{\mathcal{H}}(u, v) = \lambda_{\mathcal{H}^R}(v, u)$ and $\kappa_{\mathcal{H}}(u, v) = \kappa_{\mathcal{H}^R}(v, u)$. Hence $\lambda^{\max}$ and $\kappa^{\max}$ are invariant under reversal, and the backward edge and vertex extremal numbers equal the forward ones, for every r, m and n .*

In words: the backward model is the forward model seen in a mirror. Turning every hyperarc around swaps the tails with the heads and reverses every route, which leaves the number of independent routes between a pair untouched. The two models have identical extremal numbers, so nothing below needs to treat backward as a separate case.

Proof. Reversal sends a backward hyperarc ($|T| = r - 1, |H| = 1$) to a forward one ($|T| = 1$) and is its own inverse, so it is an edge-count-preserving bijection of the two families. Let $u = x_0, e_1, x_1, \dots, e_k, x_k = v$ be a directed Berge route in $\mathcal{H}$, where x_{i-1} is a tail and x_i a head of e_i . The reversed edge $e_i^R = (H_i, T_i)$ has x_i as a tail and x_{i-1} as a head, so $v = x_k, e_k^R, x_{k-1}, \dots, e_1^R, x_0 = u$ is a directed Berge route from v to u in $\mathcal{H}^R$ on the same edges and the same internal vertices. This is a bijection between u - v routes in $\mathcal{H}$ and v - u routes in $\mathcal{H}^R$ that preserves which edges and which internal vertices any two routes share, so it carries an edge-disjoint (respectively internally vertex-disjoint) family to one of equal size. Therefore $\lambda_{\mathcal{H}}(u, v) = \lambda_{\mathcal{H}^R}(v, u)$ and $\kappa_{\mathcal{H}}(u, v) = \kappa_{\mathcal{H}^R}(v, u)$, and maximising over ordered pairs gives $\lambda^{\max}(\mathcal{H}) = \lambda^{\max}(\mathcal{H}^R)$ and the same for κ . Since reversal is a bijection preserving feasibility and edge count, the two models share every extremal number. $\square$

Proposition A.57 (The general model obeys the forward bound). *Let a directed r -uniform hyper-edge be any split into a non-empty tail set T and head set H . If every ordered pair has at most $m - 1$ hyperedge-disjoint, or at most $m - 1$ internally vertex-disjoint, directed Berge routes, then*

$$|E(\mathcal{H})| \leq \frac{(m - 1) n(n - 1)}{r - 1},$$

the same bound as the forward model of Proposition A.55. The forward construction of that proposition is itself a set of general hyperedges, so the general value obeys the same upper bound and inherits the same quadratic lower bound, and in particular is $\Theta_{r,m}(n^2)$ for fixed r and m .

Proof. Fix an ordered pair (u, v) . A hyperedge (T, H) with $u \in T$ and $v \in H$ carries the one-step route $u \rightarrow v$ that enters at the tail u and leaves at the head v . Distinct such hyperedges give routes that share no edge and no internal vertex, so if k hyperedges had u in their tail set and v in their head set then $\lambda(u, v) \geq k$ and $\kappa(u, v) \geq k$, so feasibility forces $k \leq m - 1$. Each hyperedge (T, H) is counted by exactly the $|T| |H|$ ordered pairs (u, v) with $u \in T$ and $v \in H$, so summing the per-pair bound over all pairs gives $\sum_{(T,H)} |T| |H| \leq (m - 1) n(n - 1)$. As $|T| + |H| = r$ with

both parts at least one, $|T||H| \geq 1 \cdot (r - 1) = r - 1$, whence $(r - 1)|E| \leq (m - 1)n(n - 1)$. The forward family of [Proposition A.55](#) consists of general hyperedges, so it witnesses the same $\approx (m - 1)n^2/(4(r - 1))$ lower bound. $\square$

The two models are therefore trapped between the same pair of bounds, which differ by a factor of 4. That is enough to fix the *order* of both values and not enough to fix either leading constant, so it does not follow that the forward and general models agree asymptotically, only that any disagreement between them lives inside that factor. The rest of this section closes that gap for the forward model, which leaves the comparison between the two models as the question that survives.

The two bounds are not equally tight, and the simple digraph case indicates which end of the gap the slack sits at. The upper bound charges each hyperedge to the ordered tail-head pairs it occupies and then allows all $n(n - 1)$ ordered pairs to be used to capacity. The construction uses only the $\alpha(n - \alpha) \leq n^2/4$ pairs that run from the tail class to the head class, and leaves the pairs inside each class, and those running backwards, entirely idle. The factor of four is exactly that difference, n^2 against $n^2/4$, and it is the same difference that appears in the simple directed problem, where [Construction 1.15](#) also uses only a one-directional bipartite pattern. There, [Theorem 3.1](#) now proves that the bipartite pattern is right to first order, that no arrangement recovers the idle pairs. If that phenomenon is a feature of directedness rather than of graphs specifically, then it is the upper bound here that should come down by a factor of four, not the construction that should climb, and that is what happens.

The obstacle to running the argument of [Theorem 3.1](#) here is real and worth naming before it is disposed of, because the way around it is not to remove it. There, two-step routes through distinct midpoints were automatically arc-disjoint. Here a single hyperedge carries $r - 1$ heads at once, so a pair of two-step routes with different midpoints may enter through the very same hyperedge, and the family of two-step routes is not edge-disjoint at all. The resolution is to accept the loss. A packing can always be extracted from that family at a cost of a factor $r - 1$, and that factor turns out to be free, because of where it lands. The route count does not bound the hyperedges directly. It bounds the arcs of an auxiliary digraph, and there the factor $r - 1$ enters only the *linear* error term of [Lemma A.30](#), never the $n^2/4$ leading term. Meanwhile a second, independent factor of $r - 1$ is gained when the auxiliary digraph is exchanged back for hyperedges, and that one does divide the leading term. The two factors are not the same factor, and only one of them is paid where it matters.

Definition A.58 (One-step shadow). Let $\mathcal{H}$ be a forward directed r -uniform hypergraph on V . Its *one-step shadow* is the simple digraph $R(\mathcal{H})$ on V with an arc (u, v) exactly when some hyperedge of $\mathcal{H}$ has tail u and v among its heads. Write $\text{cod}(u, v)$ for the number of hyperedges realising that arc.

In words: the bipartite construction of [Proposition A.55](#) is asymptotically the best there is. Its count already matches the leading term $\frac{m-1}{r-1} \lfloor n^2/4 \rfloor$, so the factor of four that used to separate the known bounds is gone and the remaining uncertainty sits entirely in a term linear in n . The

bound covers both separations at once.

Proof of Theorem 3.4. Write $R = R(\mathcal{H})$ for the one-step shadow and note it has no loops, since a hyperedge's tail is not one of its heads.

Step 1: hyperedges against shadow arcs. Suppose $\text{cod}(u, v) = k$. Those k hyperedges each give a one-step Berge route from u to v , and the routes use one hyperedge apiece and different ones, so they are pairwise hyperedge-disjoint, while having no interior vertex at all they are internally vertex-disjoint as well. Hence $k \leq \kappa_{\mathcal{H}}(u, v) \leq m - 1$. Every hyperedge is counted by exactly its $r - 1$ tail-head pairs, so summing over the arcs of R ,

$$(r - 1) |E(\mathcal{H})| = \sum_{(u,v) \in A(R)} \text{cod}(u, v) \leq (m - 1) |A(R)|.$$

This is the count of Proposition A.55 stopped one step earlier. That proof then bounded $|A(R)|$ by the trivial $n(n - 1)$, which is exactly where its factor of four was lost, and the rest of this argument replaces that step.

Step 2: the shadow is budgeted. Fix an ordered pair (u, v) of distinct vertices, and let $T \subseteq V$ collect v itself when $(u, v) \in A(R)$ together with every midpoint counted by $p_2(u, v)$, so $|T| = \mathbf{1}_{(u,v) \in A(R)} + p_2(u, v)$. Every $t \in T$ is a head of at least one hyperedge whose tail is u . Form the bipartite graph joining each $t \in T$ to each such hyperedge containing it, and let M be a maximum matching in it.

Then $|M| \geq |T|/(r - 1)$. To see it, take any $t \in T$ left unmatched by M . Every hyperedge containing t must already be matched, since otherwise M could be extended along that pair. So every element of T , matched or not, lies in some hyperedge used by M , and each hyperedge holds at most $r - 1$ elements of T , giving $|T| \leq |M| (r - 1)$.

Now read M as a family of routes. For a matched pair (v, e) take the one-step route $u \rightarrow v$ through e , and for a matched pair (x, e) with x a midpoint take $u \rightarrow x \rightarrow v$, entering through e and leaving through any hyperedge whose tail is x and which has v as a head, one of which exists because $(x, v) \in A(R)$. These routes are pairwise hyperedge-disjoint: the entering hyperedges are distinct because M is a matching, the leaving hyperedges have pairwise distinct tails and so are distinct from each other, and no leaving hyperedge is an entering one, since the entering ones have tail u and the leaving ones have tail a midpoint, which is not u . They are also internally vertex-disjoint, since the one-step route has empty interior and each two-step route has for interior the single vertex it is matched to, and M matches distinct targets. Hence

$$\frac{\mathbf{1}_{(u,v) \in A(R)} + p_2(u, v)}{r - 1} \leq |M| \leq \kappa_{\mathcal{H}}(u, v) \leq m - 1,$$

so R is $(r - 1)(m - 1)$ -budgeted in the sense of Definition A.29.

Step 3: assembling. Lemma A.30 with $C = (r - 1)(m - 1)$ gives $|A(R)| \leq \lfloor n^2/4 \rfloor + 4(r -$

1)($m - 1$)($n - 1$), and substituting that into Step 1,

$$|E(\mathcal{H})| \leq \frac{m-1}{r-1} \left[\left\lfloor \frac{n^2}{4} \right\rfloor + 4(r-1)(m-1)(n-1) \right] = \frac{m-1}{r-1} \left\lfloor \frac{n^2}{4} \right\rfloor + 4(m-1)^2(n-1).$$

The lower bound is [Proposition A.55](#), whose bipartite family has

$$\max_{1 \leq \alpha \leq n-r+1} \alpha \left\lfloor \frac{(m-1)(n-\alpha)}{r-1} \right\rfloor = \frac{(m-1)n^2}{4(r-1)} - O_{m,r}(n)$$

hyperedges, where the balanced split $\alpha = \lfloor n/2 \rfloor$ already attains the right-hand side on its own, and which is feasible for both separations at once, since it is λ -feasible and $\kappa \leq \lambda$. (The integer maximum is occasionally taken a little off balance, since the floor can favour an unbalanced split by a unit or two, but never by enough to move the n^2 term.) Upper and lower bound agree in their n^2 term, which is the asymptotic claim. $\square$

Two remarks on the statement. First, the two upper bounds are not comparable and both are worth keeping: the bound of [Proposition A.55](#) is the smaller one until n is around $16(m-1)(r-1)/3$, since the linear term here carries a constant of $4(m-1)^2$, and the theorem is an asymptotic improvement rather than a uniform one. Second, at $r = 2$ the shadow is the digraph itself and cod is the multiplicity, so the theorem recovers the quadratic branch of [Theorem 3.3](#) up to its linear error term, which is the consistency check one wants, though it does not reprove that theorem, which is exact.

A third remark used to say that the general orientation model was out of reach. The obstacle and the way past it are both short to state. The proof above uses the forward model twice, once for the entering hyperedges and once for the leaving ones, in both cases to know that a hyperedge has a single tail. That is what makes Step 2's matching legitimate: the leaving hyperedges have pairwise distinct tails and so are automatically distinct from each other, and none of them can be an entering hyperedge, since the entering ones have tail u and the leaving ones do not. Allow several tails and both statements fail at once. Two midpoints may leave through one shared hyperedge, and from $r \geq 4$ a single hyperedge with two tails and two heads may serve as one target's entrance and another's exit. So the conflicts run both ways and no matching on one side controls them.

The repair is to stop building a large family and instead take a maximum one and ask what stopped it. A maximum family cannot be extended, so every target it leaves unserved must be obstructed by a hyperedge the family has already spent, and a hyperedge has only r vertices in all to obstruct with. That counts the leftovers without ever needing the two sides to be independent, which is exactly what the general model denies.

Theorem A.59 (The general orientation model shares the constant). *Let $r \geq 2$ and $m \geq 2$, and let a directed r -uniform hyperedge be any split (T, H) into a non-empty tail set and a non-empty head set, as in [Proposition A.57](#). Every such multihypergraph $\mathcal{H}$ on $n \geq 2$ vertices with*

$\kappa_{\mathcal{H}}^{\max} \leq m - 1$ satisfies

$$|E(\mathcal{H})| \leq \frac{m-1}{r-1} \left\lfloor \frac{n^2}{4} \right\rfloor + \frac{4(2r+1)(m-1)^2}{r-1} (n-1).$$

The same bound therefore holds under $\lambda_{\mathcal{H}}^{\max} \leq m - 1$. Hence for fixed r and m the general model has the same leading term $(1 + o(1))(m-1)n^2/(4(r-1))$ as the forward model, under both separations, and the bipartite construction of [Proposition A.55](#) is asymptotically optimal for it too.

In words: how a hyperedge is oriented does not move the leading term. Whether one vertex points at the other $r-1$ of them, or several do, the maximum carries the same $\frac{m-1}{r-1} \lfloor n^2/4 \rfloor$ in front, and only the linear term notices the difference, through the factor $2r+1$. With [Lemma A.56](#) this closes the orientation axis to first order.

Proof. Let R be the one-step shadow read in the general model, so $(u, v) \in A(R)$ exactly when some hyperedge has u among its tails and v among its heads, and $\text{cod}(u, v)$ counts the hyperedges realising it. A hyperedge's tail set and head set are disjoint, so R has no loops.

Step 1: hyperedges against shadow arcs. The $\text{cod}(u, v)$ hyperedges realising an arc give that many one-step routes, one hyperedge apiece and different ones, with no interior vertex, so they are pairwise hyperedge-disjoint and internally vertex-disjoint and $\text{cod}(u, v) \leq \kappa_{\mathcal{H}}(u, v) \leq m-1$. A hyperedge (T, H) is counted by exactly the $|T| |H|$ ordered pairs it fills, and $|T| + |H| = r$ with both parts non-empty gives $|T| |H| \geq r-1$. Hence

$$(r-1) |E(\mathcal{H})| \leq \sum_{(T,H) \in E(\mathcal{H})} |T| |H| = \sum_{(u,v) \in A(R)} \text{cod}(u, v) \leq (m-1) |A(R)|.$$

This is the count of [Proposition A.57](#) stopped one step early, exactly as Step 1 above stopped [Proposition A.55](#) one step early.

Step 2: the shadow is budgeted. Fix an ordered pair (u, v) of distinct vertices and let $\mathcal{T}$ collect v itself when $(u, v) \in A(R)$ together with every midpoint counted by $p_2(u, v)$, so $|\mathcal{T}| = \mathbf{1}_{(u,v) \in A(R)} + p_2(u, v)$. Call a hyperedge e an *entrance* for $t \in \mathcal{T}$ if u is a tail of e and t a head, and an *exit* for a midpoint t if t is a tail of e and v a head. Every $t \in \mathcal{T}$ has at least one entrance, and every midpoint at least one exit, by the definition of R . No hyperedge is both an entrance and an exit for the *same* t , since that would put t in its tail set and its head set at once.

Consider families of routes indexed by distinct targets: for the target v the one-step route through an entrance, and for a midpoint t the two-step route $u \rightarrow t \rightarrow v$ through an entrance and an exit. Such a family is admissible when its hyperedges are pairwise distinct. Let $\mathcal{F}$ be one of maximum size M , serving the target set S , and let U be the hyperedges it uses, so $|U| \leq 2M$. The routes of $\mathcal{F}$ are pairwise hyperedge-disjoint by construction and internally vertex-disjoint because their interiors are the empty set and distinct singletons, so

$$M \leq \kappa_{\mathcal{H}}(u, v) \leq m-1.$$

Now take $t \in \mathcal{T} \setminus S$. If some entrance $e \notin U$ existed and, for a midpoint, some exit $f \notin U$ as well, then $e \neq f$ by the paragraph above and the route through them could be added to $\mathcal{F}$, serving a target not yet served and using no hyperedge already spent. That contradicts maximality. So every entrance of t lies in U , or t is a midpoint all of whose exits lie in U . Either way t lies in the tail set or the head set of some member of U , and therefore

$$\mathcal{T} \setminus S \subseteq \bigcup_{e \in U} (T_e \cup H_e), \quad \text{so} \quad |\mathcal{T}| \leq M + \sum_{e \in U} (|T_e| + |H_e|) = M + r|U|.$$

The last equality is where the two sides are counted together rather than separately, and it is what keeps the constant at r rather than $2(r-1)$. With $|U| \leq 2M$ this gives $|\mathcal{T}| \leq (2r+1)M \leq (2r+1)(m-1)$, so R is $(2r+1)(m-1)$ -budgeted in the sense of [Definition A.29](#).

Step 3: assembling. [Lemma A.30](#) with $C = (2r+1)(m-1)$ gives $|A(R)| \leq \lfloor n^2/4 \rfloor + 4(2r+1)(m-1)(n-1)$, and substituting into Step 1 gives the stated bound. The hypothesis $\lambda^{\max} \leq m-1$ implies $\kappa^{\max} \leq m-1$ by Whitney, so the bound holds there too. For the lower bound, the bipartite family of [Proposition A.55](#) consists of general hyperedges, so it witnesses $\approx (m-1)n^2/(4(r-1))$ here as well, and upper and lower bound agree in their n^2 term. $\square$

Two remarks on that proof. The constant $2r+1$ is not optimised and is not claimed to be sharp: it is what the crude estimate $|U| \leq 2M$ gives, and a search over deliberately adversarial instances, in which every target shares both its entrance and its exit with another, never drives $|\mathcal{T}|/\kappa(u,v)$ above $r-1$, the forward model's own constant. Nothing rests on the difference, since [Lemma A.30](#) places this constant in the linear term, where it never touches the $n^2/4$ that carries the leading constant. The second point is what the theorem does to the orientation axis of [Section 3.3.1](#). Forward and backward were already known equal, exactly, by [Lemma A.56](#). The general model was known only to sit between the same two bounds a factor of four apart, which left room for it to be denser to first order. It is not. All three orientation models have the same leading term, and the computed differences between them at $n \approx r$ are a small-size effect that the asymptotics cannot see.

Part IV. One extension, and the audit

A separate extremal problem that the parallel-copy convention of [Section 2.1.1](#) sets aside, and the record of what was computed, with what, and to what standard.

A.9 The multigraph vertex problem under the other convention

[Section 2.1.1](#) collapses parallel copies in vertex mode, which makes the two multigraph vertex variants restatements of their simple counterparts and is why they carry no theorem of their own. This section works under the other reading, where the question is a different one, and records

what can be proved and what the machine finds. Throughout, q parallel copies of the edge uv count as q internally vertex-disjoint u - v routes, since a single edge is a path with no interior. Write $K_m^{\text{multi}}(n)$ for the largest total multiplicity of a multigraph on n vertices with $\kappa^{\text{max}} \leq m - 1$ under this reading.

The two kinds of route never interfere, which splits the measure in two.

Lemma A.60 (Splitting the vertex measure). *Let G be a multigraph with multiplicity function μ and underlying simple graph G_0 . For any two vertices $u \neq v$, write $\pi(u, v)$ for the largest number of pairwise internally disjoint u - v paths of length at least two in G_0 . Then, under the convention above,*

$$\kappa_G(u, v) = \mu(u, v) + \pi(u, v).$$

Proof. A parallel copy of uv is a path with empty interior, so any two copies are internally disjoint from each other and from every longer route. A maximum family may therefore be assumed to contain all $\mu(u, v)$ copies, and what it can add to them is a family of pairwise internally disjoint routes of length at least two, whose interiors are disjoint sets of vertices. Since parallel copies are irrelevant to a route of length at least two, which is determined by its vertex sequence, the largest such family has size $\pi(u, v)$, computed in G_0 . $\square$

So feasibility says exactly that $\mu(u, v) \leq m - 1 - \pi(u, v)$ on every edge, and that $\pi(u, v) \leq m - 1$ on every pair. At $m = 2$ this is already decisive.

Theorem A.61 (The other convention at $m = 2$). $K_2^{\text{multi}}(n) = n - 1$, attained exactly by the spanning trees.

Proof. By Lemma A.60, $\kappa^{\text{max}} \leq 1$ forces $\mu(u, v) + \pi(u, v) \leq 1$ on every edge. If G_0 contained a cycle, an edge uv of that cycle would have $\mu(u, v) \geq 1$ and $\pi(u, v) \geq 1$ from the rest of the cycle, a contradiction, so G_0 is a forest and every multiplicity is one. A forest has at most $n - 1$ edges, and a spanning tree attains it. $\square$

For $m \geq 3$ at least three constructions compete, and which one wins depends on how n compares with m , exactly as in every other variant of this thesis.

- ★ *The thickened tree.* Take a spanning tree and give every edge multiplicity $m - 1$. Tree edges have $\pi = 0$ and non-adjacent pairs have $\pi = 1$, so this is feasible for every $m \geq 2$, and it carries $(m - 1)(n - 1)$ edges, the multigraph edge value of Theorem 1.3.
- ★ *The thickened complete graph.* Give every edge of K_n the same multiplicity q . Here $\pi(u, v) = n - 2$, so feasibility asks $q \leq m + 1 - n$, and the count is $\binom{n}{2}(m + 1 - n)$, positive only when $n \leq m + 1$.
- ★ *The thickened theta.* Choose two poles and join each of the $n - 2$ remaining vertices to both of them at multiplicity $m - 2$, then join the poles to each other at multiplicity $m + 1 - n$ when that is positive. A pole and a middle vertex have $\pi = 1$, two middle vertices have $\pi = 2$, and

the two poles have $\pi = n - 2$, so this is feasible exactly when $n \leq m + 1$, and it carries $2(n - 2)(m - 2) + \max(0, m + 1 - n)$ edges.

The theta family is what makes this problem different from the edge problem, and it was found by the search rather than by hand. Whenever it is feasible, which is to say for $n \leq m + 1$, it matches the thickened tree exactly at $m = 4$ and, for $n \geq 3$, beats it for every $m \geq 5$. It accounts for the first cells in which the direct exhaustion exceeds the tree, although the later block sweep finds still better blocks in some larger parameter cells. Comparing these three constructions, the tree leads once $n \geq m + 2$, where the other two die for lack of route budget.

An exhaustive include-or-exclude search, independently checked by a separate implementation on the cells where both completed, confirms these constructions at the small sizes used below. The statements below are structural rather than cell-by-cell.

Two things follow. First, the problem is *not* the edge problem in disguise: at $m = 5, n = 5$ the value is 19 against the edge value 16, so the separation axis makes a difference for multigraphs under this reading. Second, the small- n regime $n \leq m + 1$, where the complete graph is feasible for the simple problems too, is tempting to read as the *only* place a cycle pays for itself, with the thickened tree optimal once $n \geq m + 2$. It is not: a single triangle grafted anywhere onto the thickened tree already beats it, for every $m \geq 5$ and every such n , which the next theorem makes precise.

By [Lemma A.60](#), a feasible multigraph on top of a fixed connected simple graph G_0 maximises its total multiplicity by setting $\mu(u, v) = m - 1 - \pi(u, v)$ on every edge, giving total $(m - 1)|E(G_0)| - \sum_{uv \in E(G_0)} \pi(u, v)$. Writing $|E(G_0)| = n - 1 + \text{rank}(G_0)$, the thickened tree is beaten by any G_0 with

$$\sum_{uv \in E(G_0)} \pi(u, v) < (m - 1) \text{rank}(G_0).$$

The obvious guess is that cycles must always pay for themselves against this bound, each independent cycle costing $m - 1$ units of route capacity. The next theorem shows a single triangle already fails to pay, for $m \geq 5$, and packing many of them compounds the failure into a gap that grows with n .

Theorem A.62 (A bouquet of thickened cliques, and how it beats the thickened tree). *Fix $3 \leq r \leq m$ and set $q = m + 1 - r \geq 1$, the multiplicity of a thickened K_r that saturates the vertex ceiling on r vertices (the “thickened complete graph” construction above, restricted to a block of size r rather than all of n). Let $1 \leq k \leq \lfloor (n - 1)/(r - 1) \rfloor$. Take k copies of K_r that all share one common vertex and are otherwise disjoint, thicken every edge of every copy to multiplicity q , and attach any leftover vertices as a pendant path at multiplicity $m - 1$. This multigraph is feasible, $\kappa^{\max} \leq m - 1$, and its total multiplicity is exactly*

$$(m - 1)(n - 1) + k \cdot \text{gain}(r, m), \quad \text{gain}(r, m) := \frac{(r - 1)(r - 2)(m - 1 - r)}{2}.$$

The count therefore exceeds the thickened tree's $(m-1)(n-1)$ exactly when $\text{gain}(r, m) > 0$, that is when $r \leq m-2$, which is possible for some r in range precisely when $m \geq 5$.

Proof. Write G_0 for the underlying simple graph.

Feasibility. The shared vertex is a cut vertex of G_0 , and so is every vertex along the pendant path, since the blocks meet only there. So a path of length ≥ 2 between two vertices of the same block cannot leave that block and come back: to do so it would have to pass through the shared vertex twice. Hence for an edge uv inside a block, every route counted by $\pi(u, v)$ stays inside its K_r , where deleting uv leaves u, v joined by exactly $r-2$ pairwise internally-disjoint two-step routes, one through each of the other block vertices, and by Menger no more, since those same $r-2$ vertices are a cut of that size. So $\pi(u, v) = r-2$ and, by Lemma A.60, $\kappa_G(u, v) = q + (r-2) = m-1$ exactly, never over. For u, v in different blocks, or nonadjacent with either endpoint on the pendant path, a single cut vertex separates them, so $\kappa_G(u, v) \leq 1$. If u, v are adjacent along the pendant path, then the edge lies on no cycle, and Lemma A.60 gives $\kappa_G(u, v) = \mu(u, v) + \pi(u, v) = (m-1) + 0 = m-1$.

Count. Since $q \geq 1$, every block edge is present, so G_0 is connected and $|E(G_0)| = n-1 + \text{rank}(G_0)$. This is where the hypothesis $r \leq m$ is used: at $r = m+1$ the multiplicity q would drop to zero, the blocks would carry no edges at all, and G_0 would fall apart into isolated vertices. Only the k blocks contribute to $\text{rank}(G_0)$ or to $\sum \pi$, since the pendant edges have $\pi = 0$ and lie in no cycle. Each K_r has rank $(r-1)(r-2)/2$ and contributes $\binom{r}{2}(r-2)$ to the sum, since each of its $\binom{r}{2}$ edges has $\pi = r-2$. Substituting into $(m-1)(n-1 + \text{rank}(G_0)) - \sum \pi$,

$$(m-1) \left[n-1 + k \frac{(r-1)(r-2)}{2} \right] - k \frac{r(r-1)(r-2)}{2} = (m-1)(n-1) + k \text{gain}(r, m),$$

$$\text{using } (m-1) \frac{(r-1)(r-2)}{2} - \frac{r(r-1)(r-2)}{2} = \frac{(r-1)(r-2)[(m-1)-r]}{2} = \text{gain}(r, m). \quad \square$$

At $r = 3$, $\text{gain}(3, m) = m-4$: zero at $m = 4$, matching the exhaustive finding that $m \leq 4$ has no counterexample, and strictly positive for every $m \geq 5$. So the guess that the thickened tree is optimal once $n \geq m+2$ is false for every $m \geq 5$: it held only within the reach of the exhaustive table, $m \leq 4$. The smallest witness is small enough to check by hand. Take $m = 5$ and $n = 7$, the first size the guess covers. Put a triangle on $\{u_1, u_2, u_3\}$ at multiplicity 3 and hang a path of four further vertices off u_1 at multiplicity 4. Two triangle vertices have three parallel copies plus one route through the third vertex, so $\kappa = 4$, and every other pair is separated by a single cut vertex or joined only by its own parallel copies, so nothing exceeds $4 = m-1$. The total multiplicity is $3 \cdot 3 + 4 \cdot 4 = 25$, one more than the $(m-1)(n-1) = 24$ the guess allows. Packing k blocks instead of one multiplies the excess by k , and k may be taken as large as $\lfloor (n-1)/(r-1) \rfloor$, so the gap over the thickened tree grows linearly in n .

Sharing a vertex is what buys that count. Disjoint blocks joined by bridges would fit only $\lfloor n/r \rfloor$ of them, since each bridge spends a vertex on nothing but the join, and the difference is real rather than cosmetic: at $m = 10$, $r = 6$ and $n = 11$ the shared-vertex form carries 150 against the bridged form's 120.

Optimising r widens the gap further. Treated as continuous, the gain per vertex $\text{gain}(r, m)/(r - 1)$ peaks near $r \sim (m + 1)/2$, where it is $\Theta(m^2)$. Packing blocks of that size therefore gives $K_m^{\text{multi}}(n) \geq ((m - 1) + \Theta(m^2))(n - 1 - O(m))$. In particular the per-vertex rate is quadratic in m once $n/m \rightarrow \infty$, a different order from the tree's linear rate.

Even this is not the truth. At $m = 5$, $n = 7$ the construction above gives 27, a partial branch and bound found 28, and the exact block computation below gives 29. What is not open is the order, and to say that the bouquet is a lower bound of the right order there has to be a matching upper bound. There is one, and it comes from the underlying simple graph rather than cycle rank.

Proposition A.63 (An upper bound of the same order). *For all $m \geq 2$ and $n \geq 2$,*

$$K_m^{\text{multi}}(n) \leq (m - 1)k_m(n) < 2(m - 1)^2 n.$$

Proof. Let G be feasible, with underlying simple graph G_0 .

Every multiplicity is at most $m - 1$. By [Lemma A.60](#), $\kappa_G(u, v) = \mu(u, v) + \pi(u, v) \geq \mu(u, v)$, so feasibility gives $\mu(u, v) \leq m - 1$ on every pair and the total is at most $(m - 1)|E(G_0)|$.

G_0 is itself feasible for the simple vertex problem. Take two vertices $u \neq v$. If they are adjacent in G_0 then $\mu(u, v) \geq 1$, and the routes of G_0 between them are the single edge together with the internally disjoint routes of length at least two, so $\kappa_{G_0}(u, v) = 1 + \pi(u, v) \leq \mu(u, v) + \pi(u, v) \leq m - 1$. If they are not adjacent then $\kappa_{G_0}(u, v) = \pi(u, v) \leq m - 1$ directly. So $\kappa_{G_0}^{\max} \leq m - 1$ and $|E(G_0)| \leq k_m(n)$ by definition of k_m , which gives the first inequality.

The simple problem is linearly bounded. Suppose G_0 had average degree at least $4(m - 1)$. By Mader's density theorem [\[15\]](#), a graph of average degree at least $4k$ contains a $(k + 1)$ -connected subgraph, so with $k = m - 1$ some subgraph $H \subseteq G_0$ is m -connected. An m -connected graph has more than m vertices and, by the global form of Menger's theorem, any two of them are joined by m internally disjoint paths inside H and hence inside G_0 . That contradicts $\kappa_{G_0}^{\max} \leq m - 1$. So the average degree is below $4(m - 1)$, that is $2|E(G_0)|/n < 4(m - 1)$, giving $k_m(n) < 2(m - 1)n$ and the second inequality. $\square$

Combining the two directions, first letting $n/m \rightarrow \infty$ so that optimised blocks can be packed with only $O(m)$ unused vertices, and then letting $m \rightarrow \infty$,

$$\left(\frac{1}{8} + o_m(1)\right)m^2 n \leq K_m^{\text{multi}}(n) < 2m^2 n,$$

where the $o_m(1)$ vanishes as m grows, so $K_m^{\text{multi}}(n) = \Theta(m^2 n)$ in that large- n regime. This qualification is necessary: for example $K_m^{\text{multi}}(2) = m - 1$, so no estimate of order $m^2 n$ can hold uniformly when n is fixed. The bouquet of [Theorem A.62](#) has the right joint order once many blocks fit, and what separates the two sides is a constant factor tending to 16. [Theorem A.65](#) below improves the lower constant and brings that factor down to $6 + 4\sqrt{2} \approx 11.66$. The upper bound is crude, since it throws away the $-\sum \pi$ correction entirely and charges every edge the full $m - 1$. The first inequality, $K_m^{\text{multi}}(n) \leq (m - 1)k_m(n)$, is the more informative half. It says that this variant can never outrun the simple vertex problem by more than the multiplicity ceiling,

which ties the least understood of the sixteen to the one with the longest history. It is tight at $m = 2$, where both sides read $n - 1$.

A.9.1 The value is a block problem

The lower bounds above are constructions and the upper bound is a density count, and neither says what an extremiser looks like. This section reduces the whole question to one about 2-connected graphs: the number of vertices in play stops being n and becomes the size of a single block.

Two observations do it. The first is already in the section, one line further than it was taken.

Lemma A.64 (The objective in closed form). *For a simple graph G_0 on n vertices with $\kappa_{G_0}^{\max} \leq m - 1$, the largest total multiplicity of a feasible multigraph with underlying simple graph exactly G_0 is*

$$W_m(G_0) := \sum_{uv \in E(G_0)} (m - \kappa_{G_0}(u, v)),$$

and every simple graph arising as the underlying graph of a feasible multigraph satisfies $\kappa_{G_0}^{\max} \leq m - 1$. Hence

$$K_m^{\text{multi}}(n) = \max \{ W_m(G_0) : G_0 \text{ simple on } n \text{ vertices, } \kappa_{G_0}^{\max} \leq m - 1 \}.$$

Proof. By [Lemma A.60](#) feasibility reads $\mu(u, v) \leq m - 1 - \pi(u, v)$ on every edge, so the best choice is $\mu(u, v) = m - 1 - \pi(u, v)$, which is what the paragraph before [Theorem A.62](#) records. On an edge $\kappa_{G_0}(u, v) = 1 + \pi(u, v)$, since the routes of G_0 between adjacent u and v are the single edge together with the routes of length at least two. Substituting turns $m - 1 - \pi(u, v)$ into $m - \kappa_{G_0}(u, v)$ and the total into $W_m(G_0)$.

Two things have to be checked for that maximum to be attained by a multigraph over G_0 itself. Feasibility of G_0 is the middle step of [Proposition A.63](#). And the chosen multiplicities are all at least one, so the underlying graph really is G_0 and no edge silently disappears: $\kappa_{G_0}(u, v) \leq m - 1$ on an edge gives $\pi(u, v) \leq m - 2$, hence $\mu(u, v) = m - 1 - \pi(u, v) \geq 1$. Conversely any feasible multigraph is one of the multigraphs considered over its own underlying graph, so nothing is lost by maximising over G_0 first. $\square$

That the objective is a sum over edges of a *local* quantity is what makes the second observation bite. Recall that a *block* of a graph is a maximal connected subgraph with no cut vertex of its own, so every block is either 2-connected or a single edge, every edge lies in exactly one block, and two blocks meet in at most one vertex.

Proof of Theorem 3.5. The objective is additive over blocks. Let G_0 be feasible and let u, v be adjacent, lying in the block B . Every u - v path of G_0 stays inside B : a path leaving B must leave through a cut vertex, and to return it would have to pass through that same cut vertex a second time, which no path does. So $\kappa_{G_0}(u, v) = \kappa_B(u, v)$, and since every edge lies in exactly one

block,

$$W_m(G_0) = \sum_{\text{blocks } B} W_m(B).$$

Each block inherits feasibility, $\kappa_B^{\max} \leq \kappa_{G_0}^{\max} \leq m - 1$ by the same identity, so $W_m(B) \leq g_m(|V(B)|)$.

The sizes fit. Suppose G_0 has c components and blocks $B_1, \dots, B_t$. Within a single component, contracting the block tree makes $\sum(|V(B_i)| - 1)$ equal to the number of vertices of that component minus one. Isolated components contribute zero to both sides, so summing gives $\sum_i(|V(B_i)| - 1) = n - c \leq n - 1$. With [Lemma A.64](#) this proves $\leq$.

Every multiset is realised. Given blocks $B_1, \dots, B_t$ with $\sum(b_i - 1) \leq n - 1$, take one common vertex and attach the B_i to it, pairwise disjoint otherwise, leaving any spare vertices isolated. The result uses $1 + \sum(b_i - 1) \leq n$ vertices. Its blocks are exactly the B_i , so by the first paragraph its κ agrees with κ_{B_i} on pairs inside a block and equals at most 1 on pairs split between two blocks, which the shared vertex separates. So the graph is feasible and scores $\sum_i W_m(B_i)$, and choosing each B_i optimal gives $\sum_i g_m(b_i)$. This proves $\geq$. $\square$

The reduction changes what has to be searched: enumerate 2-connected simple graphs, evaluate W_m , and solve the resulting knapsack. A sweep through eight vertices agrees with the independent exhaustive routine wherever both finish. It shows that the winning blocks are unbalanced and bipartite-like rather than complete, which motivates the family below. It too runs from a standalone script, `research_notes/scripts/multi_vertex_blocks.py`, and its retained transcript is `figures/multi_vertex_blocks_log.txt` in the repository named in [Section A.10](#).

Theorem A.65 (Thickened complete bipartite blocks). *Let $1 \leq s \leq t \leq m - 1$. Thicken every edge of $K_{s,t}$ to multiplicity $m - s$. The result is feasible, $\kappa^{\max} \leq m - 1$, on $s + t$ vertices with total multiplicity $st(m - s)$. Hanging k such blocks off one shared vertex gives, for $n = k(s + t - 1) + 1$,*

$$K_m^{\text{multi}}(n) \geq \frac{st(m - s)}{s + t - 1} (n - 1).$$

Optimised over s and t this rate is $(3 - 2\sqrt{2} + o_m(1))m^2$, attained at $t = m - 1$ and $s \sim (\sqrt{2} - 1)m$, against the $(\frac{1}{8} + o_m(1))m^2$ of [Theorem A.62](#). The improvement is by a factor $8(3 - 2\sqrt{2}) = 24 - 16\sqrt{2} \approx 1.373$.

Proof. Write S and T for the two sides, $|S| = s$ and $|T| = t$.

The connectivities of $K_{s,t}$. For $u \neq u'$ in S , the u - u' paths of length at least two are the t two-step routes through the vertices of T , and T is a cut of size t , so $\kappa(u, u') = \pi(u, u') = t$. Symmetrically $\kappa(v, v') = s$ for $v \neq v'$ in T . For adjacent $u \in S$ and $v \in T$, delete the edge uv : the sets $T \setminus \{v\}$ and $S \setminus \{u\}$ both separate u from v , so $\pi(u, v) \leq \min(s - 1, t - 1) = s - 1$, and the $s - 1$ routes u, v_i, u_i, v through distinct $u_i \in S \setminus \{u\}$ and distinct $v_i \in T \setminus \{v\}$ attain it, which needs $s - 1 \leq t - 1$. So $\pi(u, v) = s - 1$ and $\kappa(u, v) = s$.

Feasibility and count. By [Lemma A.60](#) the multigraph has $\kappa = (m - s) + (s - 1) = m - 1$ on

an edge, and $\kappa = t \leq m - 1$ or $s \leq m - 1$ on a non-adjacent pair, so nothing exceeds the ceiling. The count is st edges at multiplicity $m - s$. Equivalently, by [Lemma A.64](#), $W_m(K_{s,t}) = st(m - s)$.

The bouquet. If $s \geq 2$, then $K_{s,t}$ is 2-connected, so [Theorem 3.5](#) applies and k copies at one shared vertex are feasible with total $kst(m - s)$ on $n = k(s + t - 1) + 1$ vertices. If $s = 1$, then $K_{1,t}$ is a tree (a single edge when $t = 1$), and after thickening every edge to multiplicity $m - 1$, sharing vertices among copies still produces a thickened tree and is feasible directly. Its count is again $kt(m - 1) = kst(m - s)$. Thus the displayed rate holds throughout the theorem's full range $1 \leq s \leq t$.

The optimum. For fixed $s > 1$, the factor $t/(s + t - 1)$ is strictly increasing in t , and for $s = 1$ it is constant. Thus some optimum has $t = m - 1$, its largest permitted value. Put $s = \alpha m$. The rate becomes

$$\frac{s(m - 1)(m - s)}{s + m - 2} = \frac{\alpha(1 - \alpha)}{1 + \alpha} m^2 + O(m),$$

and $h(\alpha) = \alpha(1 - \alpha)/(1 + \alpha)$ has $h'(\alpha) = (1 - 2\alpha - \alpha^2)/(1 + \alpha)^2$, vanishing at $\alpha = \sqrt{2} - 1$, where $h = 3 - 2\sqrt{2}$. Since $3 - 2\sqrt{2} > \frac{1}{8}$, the family beats the clique bouquet, by the stated factor. $\square$

The optimum is lopsided: one side has size about $0.41m$ and the other reaches the ceiling $m - 1$. Finite block sweeps also show that adding a few edges inside the small side can improve this clean family. Thus the answer is a block problem with dense bipartite-like blocks, and its asymptotic constant lies between $3 - 2\sqrt{2}$ and 2.

A.10 Computational audit

The authoritative code and audit instructions are maintained in the versioned repository at <https://github.com/louisvandenbruwaene/thesis>. The snapshot this document was built from is commit not recorded in this build, which is the immutable reference for every number reported here. Anything later in that repository is a continuation and not what was submitted. From `program/`, `python erdos915_unified.py` runs the principal checks and `python -m unittest discover -s tests -v` runs the regression suite. `python make_figures.py` regenerates the computational figures. The several hundred `solve` calls behind the four variant grids of [Figures 2.10 to 2.13](#) are read from `figures/machine_values.json`, a readable record of what each of those runs returned, so the grids redraw in seconds and the plotted points are the ones this document was built from rather than whatever a fresh timed search happens to reach. Deleting that file, or passing `--refresh`, recomputes every entry, which is how the record is checked. Two block sweeps are the one thing that does not run from this file. They live in `research_notes/scripts/simple_vertex_blocks.py` and `research_notes/scripts/multi_vertex_blocks.py`, are self-contained, and were written separately, so that the values they report are an independent check on the main program rather than a second reading of it. They require nauty's `geng` on the path; the quoted ranges are reproduced from the repository root by `BMAX=9 .venv/bin/python3 research_notes/scripts/simple_vertex_blocks.py` and `BMAX=8`

.venv/bin/python3 research_notes/scripts/multi_vertex_blocks.py. The recorded environment used nauty 2.9.3. The dependency ranges the program needs are in program/requirements.txt, and the exact versions these runs were last verified against are pinned in program/requirements-lock.txt: Python 3.14.6 with NumPy 2.4.6, SciPy 1.18.0 and PuLP 3.3.2, plus Matplotlib 3.11.0 and NetworkX 3.6.1 for the two optional roles. Installing from the lock file reproduces that environment exactly.

The base cases of the directed induction ([Theorem 1.9](#)) run to $n = 7$, and are settled by a pruned exhaustive search over all simple digraphs with $\lambda^{\max} \leq 1$. That search is exact and complete at these sizes, so each completed row is an upper-bound proof at that n . The vertex problem needs its own base cases, since [Remark A.19](#) shows they cannot be inherited from the arc ones, and the same driver supplies them with only the inner feasibility test swapped. [Table A.1](#) records both runs. The two separations agree at every size, which is what closes [Theorem 1.10](#), and that agreement is a computed coincidence of the two searches rather than a formal consequence of Whitney’s inequality.

n	arc-disjoint	vertex-disjoint	$M(n)$	nodes	status
2	2	2	2	5	complete
3	4	4	4	22	complete
4	6	6	6	537	complete
5	8	8	8	17 823	complete
6	10	10	10	788 101	complete
7	12	12	12	45 122 129	complete

Table A.1. The directed $m = 2$ base cases under both separations: exhaustive branch and bound over all simple digraphs on n vertices with $\lambda^{\max} \leq 1$, and the same run with $\kappa^{\max} \leq 1$. Same driver, same prunes, only the inner feasibility test changes. Both columns agree with $M(n) = \max(2(n - 1), \lfloor n^2/4 \rfloor)$ at every size, and every run completed. The node counts coincide as a computed fact, not by construction: the feasibility prune switches between the arc and the vertex flow test with the separation, and at these sizes the two happen to cut the tree in the same places. The wall times do not, the vertex run at $n = 7$ taking 3125 s against the arc run’s 718 s. Full logs: figures/basecase_search_log.txt and figures/basecase_search_vertex_log.txt.

The historical cut-counting optimisation closes the directed multigraph checks through $n = 6$ but emits no independently replayable matching bound, so no theorem rests on it. Its raw log is figures/certificate_log.txt. Exact flow measures a supplied object, completed enumeration proves a finite optimum, and heuristic search supplies only a checked witness and lower bound. The file program/README.md records the dependencies, commands, entry points, and archival checklist.

Bibliography

- [1] Kristóf Bérczi, Karthekeyan Chandrasekaran, Tamás Király, and Shubhang Kulkarni. Splitting-off in hypergraphs. *Journal of Combinatorial Theory, Series B*, 176:319–383, 2026.
- [2] Louis Vanden Bruwaene. thesis. GitHub repository.
- [3] Béla Bollobás and Pál Erdős. Extremal problems in graph theory. *Matematikai Lapok*, 13:143–152, 1962. Original statement of Problem 915.
- [4] Ralph E. Gomory and Tien Chung Hu. Multi-terminal network flows. *Journal of the Society for Industrial and Applied Mathematics*, 9(4):551–570, 1961. Introduces the Gomory-Hu tree structure.
- [5] B. A. Sørensen and Carsten Thomassen. On k -rails in graphs. *Journal of Combinatorial Theory, Series B*, 17(2):143–159, 1974. Theorem 4 determines $f_5(n) = \lfloor 8n/3 \rfloor - 3$ for $n \geq 6$, $n \neq 7$, $n \neq 12$. Their $f_k(n)$ is the least edge count that forces a k -rail, one more than this thesis's $k_m(n)$, so the value here is $\lfloor 8n/3 \rfloor - 4$.
- [6] Karl Menger. Zur allgemeinen Kurventheorie. *Fundamenta Mathematicae*, 10:96–115, 1927. The original statement of Menger's theorem.
- [7] Pál Erdős. Erdős problems, problem 915. <https://www.erdosproblems.com/915>, 2026. Accessed 25 August 2026.
- [8] Wolfgang Mader. Ein Extremalproblem des Zusammenhangs von Graphen. *Mathematische Zeitschrift*, 131:223–231, 1973. Solves Erdős Problem 915 for edge-disjoint paths.
- [9] Hassler Whitney. Congruent graphs and the connectivity of graphs. *American Journal of Mathematics*, 54(1):150–168, 1932.
- [10] John L. Leonard. On graphs with at most four line-disjoint paths connecting any two vertices. *Journal of Combinatorial Theory, Series B*, 13:242–250, 1972.
- [11] L. R. Ford and D. R. Fulkerson. Maximal flow through a network. *Canadian Journal of Mathematics*, 8:399–404, 1956.
- [12] Brendan D. McKay and Adolfo Piperno. Practical graph isomorphism, II. *Journal of Symbolic Computation*, 60:94–112, 2014. Describes the nauty/Traces canonical-labelling tools underlying geng.
- [13] Fred Glover. Tabu search—part I. *ORSA Journal on Computing*, 1(3):190–206, 1989. Introduces tabu search, the tenure, and the tabu list.

- [14] John L. Leonard. On a conjecture of Bollobás and Erdős. *Periodica Mathematica Hungarica*, 3(3–4):281–284, 1973. Studies the Bollobás–Erdős conjecture and gives the explicit $m = 5$ counterexample on 57 vertices used in this thesis. Leonard’s work also establishes the equality results through $m = 4$.
- [15] Wolfgang Mader. Existenz n -fach zusammenhängender teilgraphen in Graphen genügend großer Kantendichte. *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, 37:86–97, 1972. Average degree at least $4k$ forces a $(k+1)$ -connected subgraph.
- [16] W. Mantel. Problem 28. *Wiskundige Opgaven*, 10:60–61, 1907. The triangle-free extremal bound, the $r = 2$ case of Turán’s theorem.
- [17] Pál Turán. On an extremal problem in graph theory. *Matematikai és Fizikai Lapok*, 48:436–452, 1941.
- [18] Saeid Hanifnezhad and Ardeshtir Dolati. Symmetric submodular system: Contractions and Gomory–Hu tree. *Information and Computation*, 271:104479, 2020. Gomory-Hu trees for symmetric submodular cut functions.
- [19] Zsolt Baranyai. On the factorization of the complete uniform hypergraph. In *Infinite and Finite Sets (Colloq., Keszthely, 1973)*, Vol. I, volume 10 of *Colloq. Math. Soc. Janos Bolyai*, pages 91–108. North-Holland, 1975.
- [20] W. T. Tutte. *Connectivity in Graphs*. University of Toronto Press, Toronto, 1966.

DEPARTMENT OF MATHEMATICS

Celestijnenlaan 200B

3001 Leuven, Belgium

wis.kuleuven.be

