


Four Bartonella genomes


Whole sequences | adjacent genome pairs | casa_natal



Mauve backbone distributed with genoPlotR; blocks ≥ 10 kb on both genomes.
111 pairwise interval links; lengths in kb. Original reference coordinates; reversed ribbons retain alignment orientation.

Four Bartonella genomes


Whole sequences | all genome pairs | casa_natal



Mauve backbone distributed with genoPlotR; blocks ≥ 10 kb on both genomes.
215 pairwise interval links; lengths in kb. Original reference coordinates; reversed ribbons retain alignment orientation.

Four Bartonella genomes


Nine annotated CDS around rpoB in each genome | adjacent pairs



36 annotated CDS; 24 reciprocal-best protein links within the selected windows.
BLASTp: >=50% identity, >=70% coverage of both proteins, E <=1e-20; tied best hits excluded.
Colors identify gene names or abbreviated products; arrows show strand. HP: hypothetical protein.

Four Bartonella genomes


Nine annotated CDS around rpoB in each genome | all pairs



36 annotated CDS; 46 reciprocal-best protein links within the selected windows.
BLASTp: $\geq 50\%$ identity, $\geq 70\%$ coverage of both proteins, $E \leq 1e-20$; tied best hits excluded.
Colors identify gene names or abbreviated products; arrows show strand. HP: hypothetical protein.
These are local windows drawn around a circle, not complete circular gene maps.

A Conserved blocks connect different chromosome-arm arrangements


380 published blocks | five arms per species | colour follows the *An. gambiae* arm



Jiang et al. (2014), doi:10.1186/s13059-014-0459-2; published signed block orders.
Equal-width blocks; track lengths count blocks, not base pairs. Circular placement does not imply circular chromosomes.
Source stephensi map covers ~62% of the assembly; only 32/86 mapped scaffolds had experimental orientation.
Opposite-orientation blocks are not inversion-event counts or evidence of effects on malaria transmission.

B The same block orders in a circular view


Arc lengths count blocks; chromosomes are biologically linear



Jiang et al. (2014), doi:10.1186/s13059-014-0459-2; published signed block orders.
Equal-width blocks; track lengths count blocks, not base pairs. Circular placement does not imply circular chromosomes.
Source stephensi map covers ~62% of the assembly; only 32/86 mapped scaffolds had experimental orientation.
Opposite-orientation blocks are not inversion-event counts or evidence of effects on malaria transmission.

C X chromosome: changes in block order

66 blocks | 44 opposite in orientation in the source table



Jiang et al. (2014), doi:10.1186/s13059-014-0459-2; published signed block orders.
Equal-width blocks; track lengths count blocks, not base pairs. Circular placement does not imply circular chromosomes.
Source stephensi map covers ~62% of the assembly; only 32/86 mapped scaffolds had experimental orientation.
Opposite-orientation blocks are not inversion-event counts or evidence of effects on malaria transmission.