Subgroup	No. of patients	IMNI	no IMNI		HR(95%CI)	p value
Chemotherapy						
NAC	217	103(47.5%)	114(52.5%)	-	0.451(0.259-0.785)	<0.001
Adjuvant	630	287(45.6%)	343(54.4%)	-	0.539(0.389-0.747)	<0.001
Molecular subtype						
TNBC	142	68(47.9%)	74(52.1%)	-	(0.306–1.254)	0.184
other	705	380(54.0%)	324(46.0%)	-	0.511(0.377-0.693)	<0.001
N stage						
N0	106	52(49.0%)	54(51.0%)	-	0.257(0.052-1.268)	0.095
N1	317	151(47.6%)	166(52.4%)	-	0.543(0.316-0.934)	0.027
N2	255	108(42.4%)	147(57.6%)	-	0.741(0.462-1.189)	0.214
N3	169	104(61.5%)	65(38.5%)	-	0.460(0.291-0.727)	0.001
T stage						
T1	358	150(41.9%)	208(58.1%)	-	0.471(0.296-0.747)	0.001
T2	374	182(48.7%)	192(51.3%)	-	0.568(0.380-0.848)	0.006
T3-4	61	34(55.7%)	27(44.3%)	>	0.909(0.402-2.053)	0.818
pCR						
pCR	41	20(48.8%)	21(51.2%)	>	0.311(0.032-2.974)	0.31
Tumor location						
Latera	466	201(43.1%)	264(56.9%)	-	0.584(0.401-0.850)	0.005
Central/media	381	191(50.1%)	190(49.9%)	-	0.483(0.322-0.726)	<0.001
Overall	847	382(45.1%)	465(54.9%)	•	0.532(0.404-0.701)	<0.001
				0.2 1 1.5 Hzard Ratio(HR)		