

MA1521 CALCULUS FOR COMPUTING

Wang Fei

matwf@nus.edu.sg

Department of Mathematics

Office: S17-06-16

Tel: 6516-2937


Chapter 2: Derivatives with Applications	2
Tangent Line	3
Definition	4
Examples	5
Derivative as Function	8
Differentiable Functions	12
Formulas	14
Chain Rule	19
Formulas of Derivatives	22
Implicit Differentiation	24
Calculus of Inverse Functions	30
Second Derivative	34
Higher Derivatives	36
Parametric Equations	38
Log Differentiation	41
Extreme Values	45
Local Max & Min	47
Extreme Value Thm	48
Finding Extrema	49
Closed Int. Method	54
Increasing Test	57
First Derivative Test	60
Concavity	64
Concavity Test	66
Curve Sketching	67
Inequalities	71
Approximation	74
Optimization	77
Examples	78
l'Hôpital's Rule	89

Chapter 2: Derivatives with Applications

2 / 101

The Tangent Line

- Recall that in Chapter 1 we have seen how to find the tangent line of the curve $y = x^2$ at $P(a, a^2)$:



$$\begin{aligned} m_{PQ} &= \frac{\Delta y}{\Delta x} \\ &= \frac{(a+h)^2 - a^2}{h} \end{aligned}$$

- The **slope** of the tangent line can be written as

$$m := \lim_{h \rightarrow 0} m_{PQ} = \lim_{h \rightarrow 0} \frac{(a+h)^2 - a^2}{h}.$$

3 / 101

Definition of Derivative

- The **derivative of a function f at a number a** is

$$f'(a) := \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}.$$

- f is **differentiable at a** if $f'(a)$ exists.
- $f'(a)$ is the **slope** of $y = f(x)$ at $x = a$.
- Let $x = a + h$. Then $h = x - a$, and $h \rightarrow 0 \Leftrightarrow x \rightarrow a$. We may use an equivalent definition:

$$f'(a) := \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}.$$

- The **tangent line** to $y = f(x)$ at $(a, f(a))$ is the line passing through $(a, f(a))$ with slope $f'(a)$:

$$y - f(a) = f'(a)(x - a).$$

4 / 101

Examples of Derivatives

- Let $f(x) = x^2 - 8x + 9$. Find $f'(3)$.

$$\begin{aligned}
 f'(3) &= \lim_{h \rightarrow 0} \frac{f(3+h) - f(3)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{[(3+h)^2 - 8(3+h) + 9] - (3^2 - 8 \cdot 3 + 9)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{(-6 - 2h + h^2) - (-6)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{-2h + h^2}{h} = \lim_{h \rightarrow 0} (-2 + h) = -2.
 \end{aligned}$$

- The tangent line of $y = f(x)$ passing through $(3, -6)$:

$$y - (-6) = f'(3)(x - 3) = -2(x - 3).$$

That is, $2x + y = 0$.

5 / 101

Examples of Derivatives

- Let $f(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right), & \text{if } x \neq 0, \\ 0, & \text{if } x = 0. \end{cases}$ Find $f'(0)$.


$$\begin{aligned}
 f'(0) &= \lim_{h \rightarrow 0} \frac{f(h+0) - f(0)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{h^2 \sin\left(\frac{1}{h}\right) - 0}{h} \\
 &= \lim_{h \rightarrow 0} h \sin\left(\frac{1}{h}\right) \\
 &= 0.
 \end{aligned}$$

$$\begin{array}{ccccccc}
 \text{Note that} & -|h| & \leq & h \sin\left(\frac{1}{h}\right) & \leq & |h| & \text{for all } h \neq 0. \\
 h \rightarrow 0 & \downarrow & & \downarrow & & \downarrow & \\
 & 0 & \Rightarrow & 0 & \Leftarrow & 0 & \\
 & & & \text{(Squeeze Theorem)} & & &
 \end{array}$$

6 / 101

Velocity

- Let $s = s(t)$ be the position function of a particle.
 - instantaneous velocity** at time $t = a$: $s'(a)$;
 - speed** at time $t = a$: $|s'(a)|$.
- Example.** A ball is dropped from a tower 450m above the ground. Find its velocity after 5 seconds.



$$s(t) = \frac{1}{2}gt^2 = 4.9t^2.$$

$$v(5) = s'(5) = 49.$$

$$\frac{\Delta s}{\Delta t} = \frac{4.9(5+h)^2 - 4.9(5)^2}{(5+h) - 5} = \frac{49h + 4.9h^2}{h}$$

$$\text{Velocity at } t = 5: \lim_{h \rightarrow 0} \frac{49h + 4.9h^2}{h} = 49 \text{ m/s.}$$

7 / 101

Example

- Let $f(x) = \frac{1}{x}$. Find $f'(a)$ at each $a \in \mathbb{R} \setminus \{0\}$.

$$\begin{aligned} f'(a) &= \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} \\ &= \lim_{h \rightarrow 0} \frac{\frac{1}{a+h} - \frac{1}{a}}{h} \\ &= \lim_{h \rightarrow 0} \frac{\frac{-h}{(a+h)a}}{h} \\ &= \lim_{h \rightarrow 0} \frac{-1}{(a+h)a} \\ &= -\frac{1}{a^2}. \end{aligned}$$

- f' is therefore a function defined on $\mathbb{R} \setminus \{0\}$.

8 / 101

Derivative as a Function

- The **derivative** of f at point $x = a$:

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}.$$

- The **derivative** of f as a function:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}.$$

$$\circ \quad f'(x) = y' = \frac{dy}{dx} = \frac{df}{dx} = \frac{d}{dx}f(x) = D_x f(x) = \dots$$

$$\circ \quad \boxed{\frac{dy}{dx} := \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x}} \quad (\text{Leibniz, 1646–1716, German})$$

$$\circ \quad f'(a) = \left. \frac{dy}{dx} \right|_{x=a}.$$

- Example.** If $f(x) = \frac{1}{x}$, then $f'(x) = -\frac{1}{x^2}$.

9 / 101

Examples

- Let $f(x) = \frac{1-x}{2+x}$.

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{\frac{1-(x+h)}{2+(x+h)} - \frac{1-x}{2+x}}{h} \\ &= \lim_{h \rightarrow 0} \frac{-3h}{h(2+(x+h))(2+x)} \\ &= \lim_{h \rightarrow 0} \frac{-3}{(2+(x+h))(2+x)} \\ &= \frac{-3}{(2+x)^2}. \end{aligned}$$

- Domain of $f: \mathbb{R} \setminus \{-2\}$; Domain of $f': \mathbb{R} \setminus \{-2\}$.

10 / 101

Examples

- Let $f(x) = \sqrt{x}$, ($x \geq 0$). Find f' .

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{\sqrt{x+h} - \sqrt{x}}{h} \\ &= \lim_{h \rightarrow 0} \frac{(\sqrt{x+h} - \sqrt{x})(\sqrt{x+h} + \sqrt{x})}{h(\sqrt{x+h} + \sqrt{x})} \\ &= \lim_{h \rightarrow 0} \frac{h}{h(\sqrt{x+h} + \sqrt{x})} \\ &= \lim_{h \rightarrow 0} \frac{1}{\sqrt{x+h} + \sqrt{x}} = \frac{1}{2\sqrt{x}} \quad (x > 0). \end{aligned}$$

The domain of f' may be *smaller* than the domain of f .

11 / 101

Differentiable Functions

- Definition.** (We only consider the differentiability at a point or on **open intervals**)

- f is **differentiable at** a if

$$f'(a) := \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} \text{ exists.}$$

- f is **differentiable on** (a, b) if it is differentiable at **every** $c \in (a, b)$.

- Questions.**

- What's the relation between differentiability and continuity?
- What kinds of functions are differentiable?
- How to construct new differentiable functions?
- How the derivative affects the original function?

12 / 101

Differentiability Implies Continuity

- **Theorem.** If f is **differentiable** at a , then f is **continuous** at a .

- **Remark.** The converse of the theorem is false.

- $f(x) = |x|$ is continuous at 0, not differentiable at 0.

- **Proof.** Suppose f is differentiable at a .

That is, $f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = L$.

- $\lim_{x \rightarrow a} (f(x) - f(a)) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} \cdot (x - a) = L \cdot 0 = 0$.

- $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} [(f(x) - f(a)) + f(a)] = 0 + f(a) = f(a)$.

Therefore, f is continuous at a .

13 / 101

Differentiation Formulas

- Let c be a constant. $(c)' = 0$.

- $\frac{d}{dx}(c) = \lim_{h \rightarrow 0} \frac{c - c}{h} = \lim_{h \rightarrow 0} 0 = 0$.

- Let f be a differentiable function, and c be a constant.

$$\begin{aligned}(cf)'(x) &= \lim_{h \rightarrow 0} \frac{cf(x+h) - cf(x)}{h} \\&= \lim_{h \rightarrow 0} c \left[\frac{f(x+h) - f(x)}{h} \right] \\&= c \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\&= cf'(x).\end{aligned}$$

$$\therefore (cf)' = cf'.$$

14 / 101

Differentiation Formulas

- Let f and g be differentiable functions.

$$\begin{aligned}
 (f + g)'(x) &= \lim_{h \rightarrow 0} \frac{(f + g)(x + h) - (f + g)(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{[f(x + h) + g(x + h)] - [f(x) + g(x)]}{h} \\
 &= \lim_{h \rightarrow 0} \frac{[f(x + h) - f(x)] + [g(x + h) - g(x)]}{h} \\
 &= \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h} + \lim_{h \rightarrow 0} \frac{g(x + h) - g(x)}{h} \\
 &= f'(x) + g'(x).
 \end{aligned}$$

$$\therefore (f + g)' = f' + g'.$$

$$\begin{aligned}
 (f - g)' &= [f + (-g)]' = f' + (-g)' = f' + (-g') \\
 &= f' - g'.
 \end{aligned}$$

15 / 101

Differentiation Formulas

- Let f and g be differentiable functions. What is $(fg)'$?

$$\begin{aligned}
 \frac{d}{dx} [f(x)]^2 &= \lim_{h \rightarrow 0} \frac{[f(x + h)]^2 - [f(x)]^2}{h} \\
 &= \lim_{h \rightarrow 0} \frac{[f(x + h) - f(x)] \cdot [f(x + h) + f(x)]}{h} \\
 &= \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h} \cdot \lim_{h \rightarrow 0} [f(x + h) + f(x)] \\
 &= f'(x) \cdot [f(x) + f(x)] \quad (\because f \text{ is continuous}) \\
 &= 2f'(x)f(x).
 \end{aligned}$$

$$\begin{aligned}
 (fg)' &= \frac{1}{2}[(f + g)^2 - f^2 - g^2]' \\
 &= \frac{1}{2}[2(f + g)'(f + g) - 2f'f - 2g'g] \\
 &= (f' + g')(f + g) - f'f - g'g \\
 &= f'g + fg'.
 \end{aligned}$$

16 / 101

Differentiation Formulas

- Let f and g be differentiable functions. What is $(f/g)'$?
Suppose $g(x) \neq 0$.

$$\begin{aligned}
 \left(\frac{1}{g}\right)'(x) &= \lim_{h \rightarrow 0} \frac{\left(\frac{1}{g}\right)(x+h) - \left(\frac{1}{g}\right)(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\frac{1}{g(x+h)} - \frac{1}{g(x)}}{h} = \lim_{h \rightarrow 0} \frac{\frac{g(x) - g(x+h)}{g(x)g(x+h)}}{h} \\
 &= \lim_{h \rightarrow 0} \left[\frac{g(x+h) - g(x)}{h} \cdot \frac{-1}{g(x)g(x+h)} \right] \\
 &= \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \cdot \lim_{h \rightarrow 0} \frac{-1}{g(x)g(x+h)} \\
 &= g'(x) \cdot \frac{-1}{[g(x)]^2} = \left(-\frac{g'}{g^2}\right)(x).
 \end{aligned}$$

17 / 101

Differentiation Formulas

- Let f and g be differentiable functions. What is $(f/g)'$?
Suppose $g(x) \neq 0$.

$$\begin{aligned}
 \left(\frac{f}{g}\right)' &= \left(f \cdot \frac{1}{g}\right)' = f' \cdot \left(\frac{1}{g}\right) + f \cdot \left(\frac{1}{g}\right)' \\
 &= \frac{f'}{g} + \frac{f \cdot (-g')}{g^2} \\
 &= \frac{f'g - fg'}{g^2}.
 \end{aligned}$$

- $(cf)' = cf'$
- $(f \pm g)' = f' \pm g'$
- $(fg)' = f'g + fg'$
- $\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}, \text{ if } g(x) \neq 0.$

18 / 101

The Chain Rule

- Let $F(x) = \sqrt{x^2 + 1}$. Find F' .

$$\begin{aligned}
 F'(x) &= \lim_{h \rightarrow 0} \frac{\sqrt{(x+h)^2 + 1} - \sqrt{x^2 + 1}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{(\sqrt{(x+h)^2 + 1} - \sqrt{x^2 + 1})(\sqrt{(x+h)^2 + 1} + \sqrt{x^2 + 1})}{h(\sqrt{(x+h)^2 + 1} + \sqrt{x^2 + 1})} \\
 &= \lim_{h \rightarrow 0} \frac{(x+h)^2 + 1 - (x^2 + 1)}{h(\sqrt{(x+h)^2 + 1} + \sqrt{x^2 + 1})} \\
 &= \lim_{h \rightarrow 0} \frac{2xh + h^2}{h(\sqrt{(x+h)^2 + 1} + \sqrt{x^2 + 1})} \\
 &= \lim_{h \rightarrow 0} \frac{2x + h}{\sqrt{(x+h)^2 + 1} + \sqrt{x^2 + 1}} \\
 &= \frac{2x}{2\sqrt{x^2 + 1}} = \frac{x}{\sqrt{x^2 + 1}}.
 \end{aligned}$$

19 / 101

The Chain Rule

- Let $F(x) = \sqrt{x^2 + 1}$. Find F' .

Note that $F = f \circ g$, where

- $f(x) = \sqrt{x}$, and $g(x) = x^2 + 1$.

It is known that

- $f'(x) = \frac{1}{2\sqrt{x}}$, and $g'(x) = 2x$.

Question. Can we write F' by making use of f' and g' ?

- If $y = g(x)$, $z = f(y)$, it seems that

$$F'(x) = \frac{dz}{dx} = \frac{dz}{dy} \frac{dy}{dx} = f'(g(x))g'(x) = \frac{2x}{2\sqrt{x^2 + 1}}.$$

Question. Can we always differentiate the composite of differentiable functions using this method?

20 / 101

The Chain Rule

- **Theorem.** Let f and g be differentiable functions.
Then $F = f \circ g$ is differentiable and

$$F' = (f' \circ g)(g').$$

More precisely,

if g is differentiable at a and f is differentiable at $b = g(a)$, then $F = f \circ g$ is differentiable at a and

$$F'(a) = f'(b)g'(a) = f'(g(a))g'(a).$$

In Leibniz notation, if $y = g(x)$ and $z = f(y)$, then

$$\left. \frac{dz}{dx} \right|_{x=a} = \left. \frac{dz}{dy} \right|_{y=b} \left. \frac{dy}{dx} \right|_{x=a}, \quad \text{or in short} \quad \frac{dz}{dx} = \frac{dz}{dy} \frac{dy}{dx}.$$

21 / 101

Formulas of Derivatives

- $\frac{d}{dx} x^a = ax^{a-1}.$
- $\frac{d}{dx} \sin x = \cos x, \quad \frac{d}{dx} \cos x = -\sin x;$
 $\frac{d}{dx} \tan x = \sec^2 x, \quad \frac{d}{dx} \cot x = -\csc^2 x;$
 $\frac{d}{dx} \sec x = \sec x \tan x, \quad \frac{d}{dx} \csc x = -\csc x \cot x.$
- $\frac{d}{dx} \sin^{-1} x = \frac{1}{\sqrt{1-x^2}}, \quad \frac{d}{dx} \cos^{-1} x = -\frac{1}{\sqrt{1-x^2}}.$
 $\frac{d}{dx} \tan^{-1} x = \frac{1}{1+x^2}, \quad \frac{d}{dx} \cot^{-1} x = -\frac{1}{1+x^2}.$
 $\frac{d}{dx} \sec^{-1} x = \frac{1}{x\sqrt{x^2-1}}, \quad \frac{d}{dx} \csc^{-1} x = -\frac{1}{x\sqrt{x^2-1}}.$
- $\frac{d}{dx} e^x = e^x, \quad \frac{d}{dx} \ln |x| = \frac{1}{x}$

22 / 101


Examples

- $\frac{d}{dx} \ln \frac{\sqrt{x+1}}{e^{2x} \cos^4 5x}.$
 - $\ln \frac{\sqrt{x+1}}{e^{2x} \cos^4 5x} = \frac{1}{2} \ln(x+1) - 2x - 4 \ln |\cos 5x|.$
 - $\frac{d}{dx} \ln \frac{\sqrt{x+1}}{e^{2x} \cos^4 5x} = \frac{1}{2(x+1)} - 2 - \frac{4}{\cos 5x} \cdot (-5 \sin 5x)$
 $= \frac{1}{2(x+1)} - 2 + 20 \tan 5x.$
- $\frac{d}{dx} \left(x \sin^{-1} \frac{1}{x} \right).$
 - $\frac{d}{dx} \left(x \sin^{-1} \frac{1}{x} \right) = 1 \cdot \sin^{-1} \frac{1}{x} + x \cdot \frac{1}{\sqrt{1 - (\frac{1}{x})^2}} \cdot (-x^{-2})$
 $= \sin^{-1} \frac{1}{x} - \frac{1}{x \sqrt{1 - \frac{1}{x^2}}}.$

23 / 101

Implicit Differentiation

- How to find the **slope** of the tangent line to the unit circle $x^2 + y^2 = 1$ at a point $P(x_0, y_0)$ on the circle?



$$\circ \quad \overline{AB} \perp \overline{OP} \Rightarrow (\text{slope of } \overline{AB})(\text{slope of } \overline{OP}) = -1$$

$$y'|_P \cdot \frac{y_0}{x_0} = -1 \Rightarrow \boxed{y'|_P = -\frac{x_0}{y_0}}.$$

24 / 101

Implicit Differentiation

- How to find the **slope** of the tangent line to the unit circle $x^2 + y^2 = 1$ at a point $P(x_0, y_0)$ on the circle?
 - Given that $1 = x^2 + y^2$.
Differentiate both sides with respect to x .

$$\frac{d}{dx}(1) = \frac{d}{dx}(x^2 + y^2).$$

That is,

$$\begin{aligned} 0 &= \frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) = 2x + \frac{dy^2}{dy} \frac{dy}{dx} \\ &= 2x + 2y \frac{dy}{dx}. \end{aligned}$$

$$\frac{dy}{dx} = -\frac{2x}{2y} = -\frac{x}{y} \Rightarrow \left[\frac{dy}{dx} \right]_P = -\frac{x_0}{y_0}.$$

25 / 101

Implicit Differentiation

- In general, we may not have a **function** $y = f(x)$.
Instead, it may be an **equation**

$$f(x, y) = 0.$$

We can still find $y' = \frac{dy}{dx}$ as follow:


- 1). Differentiate $f(x, y)$ with respect to x ;
 - 2). Solve the equation $\frac{d}{dx}f(x, y) = 0$ to express $\frac{dy}{dx}$ in terms of x and y .
- The method introduced is called **implicit differentiation**.

Remark. In order to use the method of implicit differentiation, we shall first assume that $\frac{dy}{dx}$ exists.

26 / 101

Examples

- Find $\frac{dy}{dx}$ if $x^3 + y^3 = 3xy$.



- Differentiate

$$x^3 + y^3 = 3xy$$

with respect to x :

$$(x^3)' = 3x^2$$

$$(y^3)' = 3y^2 y'$$

$$(xy)' = x'y + xy' = y + xy'$$

$$3x^2 + 3y^2 y' = 3(y + xy')$$

- Solve y' : $y' = \frac{x^2 - y}{x - y^2}$, $(x, y) \neq (0, 0), (\sqrt[3]{4}, \sqrt[3]{2})$.

27 / 101


Find the Inverse Function

- Let f be a **one to one** function. Then it admits an inverse function f^{-1} . But how to find f^{-1} ?
- Recall that $y = f(x) \Leftrightarrow f^{-1}(y) = x$. We can thus apply the following procedure:
 - Write $y = f(x)$.
 - Solve the equation for x in terms of y : $x = f^{-1}(y)$.
 - Interchanging x and y to express f^{-1} as a function in variable x : $y = f^{-1}(x)$.
- Interchanging x and y has the same effect as the reflection with respect to $y = x$.
 - So the graphs of f and f^{-1} are **symmetric with respect to the straight line $y = x$** .

28 / 101

Examples

- Find inverse of $f(x) = x^3 + 2$ and $g(x) = \sqrt{-1-x}$.
 1. Let $y = f(x) = x^3 + 2$.
 2. Solve x in terms of y : $x = \sqrt[3]{y-2}$.
 3. Interchange x and y : $f^{-1}(x) = y = \sqrt[3]{x-2}$.
- 1. Let $y = g(x) = \sqrt{-1-x}$, ($x \leq -1$).
- 2. Solve x in terms of y : $x = -y^2 - 1$, ($y \geq 0$).
- 3. Interchange x and y : $g^{-1}(x) = -x^2 - 1$, ($x \geq 0$).



29 / 101

Calculus of Inverse Functions

- Let f be a continuous function defined on an interval.
 - If f is increasing, then f is one to one.
 - $a < b \Rightarrow f(a) < f(b) \Rightarrow f(a) \neq f(b)$.
 - Similarly, if f is decreasing, then f is one to one.
 - If f is one to one, must f be increasing or decreasing?
- **Theorem.** Suppose f is a one-to-one and continuous function defined on an interval. Then
 - f is either increasing or decreasing.
- **Theorem.** Suppose f is a one-to-one and continuous function defined on an interval. Then
 - Its inverse function f^{-1} is continuous.


30 / 101

Calculus of Inverse Functions

- **Theorem.** Let f be a one to one continuous function defined on an interval.

- If f is differentiable at a , and $f'(a) \neq 0$,
- then f^{-1} is differentiable at $b = f(a)$,


$$\text{and } (f^{-1})'(b) = \frac{1}{f'(a)}.$$



31 / 101

Example

- Let $f(x) = x^2$ on $[0, 2]$. Find $(f^{-1})'(1)$.
 - Method 1: $(f^{-1})(x) = \sqrt{x}$. Then $(f^{-1})'(x) = \frac{1}{2\sqrt{x}}$.
 - $(f^{-1})'(1) = \frac{1}{2\sqrt{x}} \Big|_{x=1} = \frac{1}{2}$.
 - Method 2: $f(1) = 1$, and $f'(x) = 2x$. Then
 - $(f^{-1})'(1) = \frac{1}{f'(1)} = \frac{1}{2}$.



- The advantage of Method 2 is that we can find $(f^{-1})'$ without knowing the explicit expression of f^{-1} .

32 / 101

Derivative of Trigonometric Functions

- Recall the **inverse sine function** $\sin^{-1} x : [-1, 1] \rightarrow [-\frac{\pi}{2}, \frac{\pi}{2}]$. What is $(\sin^{-1} x)'$?
 - Let $y = \sin^{-1} x \in [-\frac{\pi}{2}, \frac{\pi}{2}]$. Then $\sin y = x$.
 $\cos y \geq 0 \Rightarrow \cos y = \sqrt{1 - \sin^2 y} = \sqrt{1 - x^2}$.
 $\therefore (\sin^{-1} x)' = \frac{1}{(\sin y)'} = \frac{1}{\cos y} = \frac{1}{\sqrt{1 - x^2}}, -1 < x < 1$.
- The **inverse cosine function** $\cos^{-1} x : [-1, 1] \rightarrow [0, \pi]$.
 - Let $y = \cos^{-1} x \in [0, \pi]$. Then $\cos y = x$.
 $\sin y \geq 0 \Rightarrow \sin y = \sqrt{1 - \cos^2 y} = \sqrt{1 - x^2}$.
 $\therefore (\cos^{-1} x)' = \frac{1}{(\cos y)'} = \frac{-1}{\sin y} = \frac{-1}{\sqrt{1 - x^2}}, -1 < x < 1$.
- Exercise.** Prove that $(\tan x)' = \frac{1}{1 + x^2}$.

33 / 101

Second Derivative

- Let f be a function. We can **differentiate** it to get f' .
- f' is a function, we can **differentiate** it to get $(f')'$.
 - $f'' := (f')'$, is called the **second derivative** of f .
 - By Leibniz notation:

$$f''(x) = \frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d^2 y}{dx^2}.$$

- $f' = D(f) \Rightarrow f'' := D^2(f)$.
- Geometric meaning:**
 - f' measures the change of f ,
 - f'' measures the change of f' .

34 / 101

Physical Meaning of Second Derivative

- Let $s = s(t)$ be the position function of an object along a straight line.
 - $s'(t) = v(t)$: the **instantaneous velocity**, it determines the change of the **position**,
 - $s''(t) = v'(t) = a(t)$: the **acceleration**, it determines the change of the **velocity**.
- Example.** Suppose the position of a particle is given by

$$s = s(t) = t^3 - 6t^2 + 9t.$$

- Velocity:** $v(t) = s'(t) = 3t^2 - 12t + 9$.
- Acceleration:** $a(t) = s''(t) = v'(t) = 6t - 12$.

35 / 101

Higher Derivatives

- Let f be a function.
 - Differentiate f to get f' , the **first derivative**.
 - Differentiate f' to get f'' , the **second derivative**.
 - Differentiate f'' to get f''' , the **third derivative**.
 - Differentiate f''' to get f'''' , the **fourth derivative**.
 -
- In general, we define $f^{(0)} := f$, and for positive integer n ,

$$f^{(n)} := (f^{(n-1)})',$$

called the **n th derivative** of f .

Other notations: if $y = f(x)$, then

$$f^{(n)}(x) = y^{(n)} = \frac{d^n y}{dx^n} = D^n f(x).$$

36 / 101

Examples

- Let $f(x) = x \cos x$. Find f' , f'' and f''' .

$$\begin{aligned}f'(x) &= (x \cos x)' = (x)' \cos x + x(\cos x)' \\&= \cos x - x \sin x.\end{aligned}$$

$$\begin{aligned}f''(x) &= (\cos x - x \sin x)' = (\cos x)' - (x \sin x)' \\&= -\sin x - [(x)' \sin x + x(\sin x)'] \\&= -\sin x - \sin x - x \cos x \\&= -2 \sin x - x \cos x.\end{aligned}$$

$$\begin{aligned}f'''(x) &= (-2 \sin x - x \cos x)' = -2(\sin x)' - (x \cos x)' \\&= -2 \cos x - [(x)' \cos x + x(\cos x)'] \\&= -2 \cos x - (\cos x - x \sin x) \\&= -3 \cos x + x \sin x.\end{aligned}$$

37 / 101

Parametric Equations

- Let x and y be functions of variable t .

- $x = x(t)$ and $y = y(t)$.

This is a **parametric equation** of x and y .

- Examples.**

- $x^2 + y^2 = 1$ can be parameterized as

- $x = \cos t$ and $y = \sin t, t \in \mathbb{R}$.

- $y^2 = x^3 + x^2$ can be parameterized as

- $x = t^2 - 1$ and $y = t^3 - t, t \in \mathbb{R}$.

- Question.** Given a parametric equation, $x = x(t)$ and $y = y(t)$, how to find the derivative of y with respect to x ?

38 / 101

Parametric Equations

- Suppose $x = x(t)$ and $y = y(t)$. Then
 - $\frac{dy}{dx} = \frac{dy/dt}{dx/dt}$, provided that $\frac{dx}{dt} \neq 0$.
- **Example.** Find an expression for $\frac{dy}{dx}$ in terms of t if the curve is defined parametrically by
 - $x = \ln t$ and $y = t^2 - e^t$.

Solution.

- $\frac{dx}{dt} = \frac{1}{t}$ and $\frac{dy}{dt} = 2t - e^t$.
- $\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = t(2t - e^t)$
- $\frac{dx}{dy} = \frac{dx/dt}{dy/dt} = \frac{1}{t(2t - e^t)}.$

39 / 101

Parametric Equations

- We can find the second derivative by chain rule:
 - $\frac{d^2y}{dx^2} = \frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{dt}{dx} \cdot \frac{d}{dt} \left(\frac{dy}{dx} \right) = \frac{1}{dx/dt} \cdot \frac{d}{dt} \left(\frac{dy}{dx} \right).$

Example. $x = \ln t$ and $y = t^2 - e^t$.

- $\frac{dx}{dt} = \frac{1}{t}$ and $\frac{dy}{dt} = 2t - e^t$.
- $\frac{d^2y}{dx^2} = \frac{1}{\frac{dx}{dt}} \cdot \frac{d}{dt} \left(\frac{\frac{dy}{dt}}{\frac{dx}{dt}} \right) = t(4t - e^t - te^t).$
- $\frac{d^2x}{dy^2} = \frac{1}{\frac{dy}{dt}} \cdot \frac{d}{dt} \left(\frac{\frac{dx}{dt}}{\frac{dy}{dt}} \right) = -\frac{4t - e^t - te^t}{t^2(2t - e^t)}.$
- **Note.** $\frac{dy}{dx} \cdot \frac{dx}{dy} = 1$ but $\frac{d^2y}{dx^2} \cdot \frac{d^2x}{dy^2} \neq 1.$

40 / 101

Logarithmic Differentiation

- Find $\frac{dy}{dx}$ if $y = \frac{(x^2 + 1)\sqrt{x + 3}}{x - 1}$, $x > 1$.
 - Of course we can use product and quotient law to evaluate. But do we have a shortcut?
 - This is a product of positive functions.
 - Recall that $\ln ab = \ln a + \ln b$ for $a > 0, b > 0$.
- 1. Take logarithmic function both sides:
 - $\ln y = \ln(x^2 + 1) + \frac{1}{2} \ln(x + 3) - \ln(x - 1)$.
- 2. Differentiate with respect to x :
 - $\frac{1}{y} \cdot \frac{dy}{dx} = \frac{2x}{x^2 + 1} + \frac{1}{2(x + 3)} - \frac{1}{x - 1}$.

$$\frac{dy}{dx} = \left[\frac{2x}{x^2 + 1} + \frac{1}{2(x + 3)} - \frac{1}{x - 1} \right] \frac{(x^2 + 1)\sqrt{x + 3}}{x - 1}.$$

41 / 101

Logarithmic Differentiation

- In general, if $y = f_1(x)f_2(x) \cdots f_n(x)$ is a product of nonzero functions, we can find the derivative as following:
 1. Take logarithmic function both sides:
 - $\ln |y| = \ln |f_1(x)| + \ln |f_2(x)| + \cdots + \ln |f_n(x)|$.
 2. Differentiate with respect to x :
 - $\frac{1}{y} \cdot \frac{dy}{dx} = \frac{f'_1(x)}{f_1(x)} + \frac{f'_2(x)}{f_2(x)} + \cdots + \frac{f'_n(x)}{f_n(x)}$.
 3. $\frac{dy}{dx} = \left[\frac{f'_1(x)}{f_1(x)} + \frac{f'_2(x)}{f_2(x)} + \cdots + \frac{f'_n(x)}{f_n(x)} \right] y$

$$= \left[\frac{f'_1(x)}{f_1(x)} + \frac{f'_2(x)}{f_2(x)} + \cdots + \frac{f'_n(x)}{f_n(x)} \right] f_1(x)f_2(x) \cdots f_n(x).$$
- Such method is called **logarithmic differentiation**.

42 / 101

Logarithmic Differentiation

- **Example.** Find $\frac{dy}{dx}$ if $y = \frac{x \cos x}{\sqrt{\csc x}}$.

1. $\ln |y| = \ln |x| + \ln |\cos x| - \frac{1}{2} \ln |\csc x|.$
2. $\frac{1}{y} \cdot \frac{dy}{dx} = \frac{1}{x} + \frac{-\sin x}{\cos x} - \frac{1}{2} \cdot \frac{-\csc x \cot x}{\csc x}.$
3. $\frac{dy}{dx} = \left[\frac{1}{x} - \tan x + \frac{1}{2} \cot x \right] \frac{x \cos x}{\sqrt{\csc x}}.$

- **Example.** Find the derivative of $y = f(x)g(x).$

1. $\ln |y| = \ln |f(x)| + \ln |g(x)|.$
2. $\frac{1}{y} \cdot \frac{dy}{dx} = \frac{f'(x)}{f(x)} + \frac{g'(x)}{g(x)}.$
3. $\frac{dy}{dx} = \left[\frac{f'(x)}{f(x)} + \frac{g'(x)}{g(x)} \right] y = f'(x)g(x) + f(x)g'(x).$

43 / 101

Logarithmic Differentiation

- Evaluate $\frac{d}{dx} x^x$. Exercise: $\frac{d}{dx} (x^x)^x$ and $\frac{d}{dx} x^{(x^x)}.$

- $\frac{d}{dx} x^a = ax^{a-1}.$ Is $\frac{d}{dx} x^x = x \cdot x^{x-1} = x^x ?$
- $\frac{d}{dx} a^x = a^x \ln a.$ Is $\frac{d}{dx} x^x = x^x \ln x ?$
- Let $y = x^x$. Then $\ln y = x \ln x.$
 - $\frac{1}{y} \frac{dy}{dx} = 1 \cdot \ln x + x \cdot \frac{1}{x} = \ln x + 1.$
 - $\frac{dy}{dx} = y(\ln x + 1) = x^x(\ln x + 1).$

- In general, if $y = f(x)^{g(x)}, (f(x) > 0),$ then

- $\ln y = g(x) \ln f(x) \Rightarrow \frac{1}{y} \frac{dy}{dx} = \frac{d}{dx} [g(x) \ln f(x)].$

44 / 101

Absolute Maximum and Minimum Values

- **Definition.** Let f be a function, and D be its domain.


- f has an **global** (or **absolute**) **maximum** at $c \in D$

$$\iff f(c) \geq f(x) \text{ for all } x \in D.$$

- f has an **global** (or **absolute**) **minimum** at $c \in D$

$$\iff f(c) \leq f(x) \text{ for all } x \in D.$$


- The **absolute maximum** and **absolute minimum** are called the **(absolute) extreme values**.



45 / 101

Example

- Let $f(x) = 3x^4 - 16x^3 + 18x^2$ on $[-1, 3.5]$.




- **global max:** highest point $f(-1) = 37$.
- **global min:** lowest point $f(3) = -27$.
- What can we say about other “**turning points**” and “**end points**”?

46 / 101

Local Maximum and Local Minimum

- **Definition.** Let f be a function with domain D .
 - f has a **local** (or **relative**) **maximum** at $c \in D$
 $\iff f(c) \geq f(x)$ for all x near c
 (i.e., for all x in an **open interval** containing c)
 - f has a **local** (or **relative**) **minimum** at $c \in D$
 $\iff f(c) \leq f(x)$ for all x near c
 (i.e., for all x in an **open interval** containing c)



47 / 101


Extreme Value Theorem

- **Extreme Value Theorem.**
 If f is **continuous** on a **finite closed interval** $[a, b]$,
 - then f attains **extreme values** on $[a, b]$.
 Precisely, f attains an
 - **absolute maximum** value $f(c)$ at some $c \in [a, b]$,
 - **absolute minimum** value $f(d)$ at some $d \in [a, b]$.
 (The proof requires the “**compactness**” of finite closed interval. It is omitted in our course.)
- **Note.** Similarly as the “**Intermediate Value Theorem**”, the “**Extreme Value Theorem**” only shows the **existence** of the extreme values.
 We shall introduce a method to find out the exact value of extreme values.


48 / 101

Where are the Extreme Values?

- The extreme value may be obtained at the **endpoints**.



- If the extreme value is not obtained at the end points,



by definition it must occur as a **local max** or a **local min**.

49 / 101

Finding the Extreme Values

- Let f be a **continuous** function on **closed** interval $[a, b]$.
 1. Compute the values at **endpoints**: $f(a)$, $f(b)$.
 2. Find **local max** and **local min** of f on (a, b) .
 3. Compare the values obtained above to seek out the **extreme values**:
 - The **largest** is the **absolute maximum**,
 - The **smallest** is the **absolute minimum**.
- The 1st and the 3rd steps are easy.

How to find the **local max** and **local min** of f on (a, b) ?

- From the graphs, it seems that the local max and local min always occur at the “**turning points**”.

50 / 101

Fermat's Theorem

- **Fermat's Theorem.**

- Suppose f has a local maximum or local minimum at c .
If $f'(c)$ exists, then $f'(c) = 0$.

- Pierre de Fermat (1601–1665), French Lawyer.

- **Fermat's Last Theorem:** $x^n + y^n = z^n$ has no nontrivial integer solution for $n \geq 3$.
- He wrote: "I have discovered a truly remarkable proof which this margin is too small to contain."


- **Note.** We **CANNOT** find the local maximum and local minimum by simply solving $f'(x) = 0$.

- Even if $f'(c) = 0$, f may not have a local maximum or local minimum at c .
- Even if f has a local maximum or a local minimum at c , $f'(c)$ may not exist, and so $f'(c)$ may not be 0.

51 / 101

Examples


- " $f'(c) = 0 \nRightarrow f$ has local max or local min at c ".
 - Let $f(x) = x^3$. Then $f'(x) = 3x^2$ and $f'(0) = 0$.
But f has no local max or local min at 0.
- " f has local max or local min at $c \nRightarrow f'(c) = 0$ ".
 - Let $g(x) = |x|$. Then f is a local minimum at 0.
But $f'(0)$ does not exist.



52 / 101

Critical Number

- Consider the following diagram:



- Definition.** Let f be a function with domain D . Then $c \in D$ is called a **critical number** of f if
 - $f'(c)$ does not exist, or $f'(c)$ exists and equals 0.

53 / 101

Closed Interval Method

- Fermat's Theorem (Rephrased).**

If f has a local maximum or a local minimum at c ,

- then c is a critical number of f .

- Closed Interval Method:**


Let f be a **continuous** function on interval $[a, b]$.

- Find the values of f at **end points**: $x = a$, $x = b$,
- Find the values of f at **critical numbers** of f in (a, b) :
 - number $c \in (a, b)$ at which $f'(c)$ does not exist, or
 - number $c \in (a, b)$ at which $f'(c) = 0$.
- Compare the values of $f(x)$ evaluated in 1) and 2):
 - The largest is the **absolute maximum** value.
 - The smallest is the **absolute minimum** value.

54 / 101

Examples

- Find the extreme values of $f(x) = x^{\frac{3}{5}}(4 - x)$ on $[-1, 3]$.



1) End points: $-1, 3$;

2) Critical numbers: $0, 1.5$.


$$\begin{aligned} f'(x) &= (x^{\frac{3}{5}})'(4 - x) + x^{\frac{3}{5}}(4 - x)' \\ &= \frac{3}{5}x^{-\frac{2}{5}}(4 - x) - x^{\frac{3}{5}} = \frac{4(3 - 2x)}{5x^{\frac{2}{5}}}. \end{aligned}$$

- $f'(x)$ does not exist $\Rightarrow x = 0$,
 - $f'(x) = 0 \Rightarrow x = 1.5$.
- 3) Comparing $f(-1)$, $f(3)$, $f(0)$, $f(1.5)$,
- Absolute maximum: $f(1.5) \approx 3.1886$.
 - Absolute minimum: $f(-1) = -5$.

55 / 101

Examples

- Let $f(x) = \sqrt[3]{x^2 - x}$ be defined on $[-1, 2]$.
 - $f'(x) = \frac{1}{3}(x^2 - x)^{-2/3}(2x - 1) = \frac{2x - 1}{3(x^2 - x)^{2/3}}$.
 - $f'(x)$ does not exist: $x = 0, x = 1$;
 - $f'(x) = 0$: $x = \frac{1}{2}$.




- Absolute maximum: $f(-1) = f(2) = \sqrt[3]{2}$.
- Absolute minimum: $f(\frac{1}{2}) = -\frac{1}{\sqrt[3]{4}}$.


56 / 101

Increasing/Decreasing Test

- Consider the following functions:



- It seems that $\begin{cases} f \text{ "turns"} & \Leftrightarrow f' = 0, \\ f \text{ is increasing} & \Leftrightarrow f' > 0, \\ f \text{ is decreasing} & \Leftrightarrow f' < 0. \end{cases}$



57 / 101

Increasing/Decreasing Test

- Theorem.** Let f be a function such that
 - It is **continuous** on $[a, b]$, **differentiable** on (a, b) .

Then

- $f'(x) = 0$ on $(a, b) \Leftrightarrow f$ is **constant** on $[a, b]$;
- $f'(x) > 0$ on $(a, b) \Rightarrow f$ is **increasing** on $[a, b]$;
- $f'(x) < 0$ on $(a, b) \Rightarrow f$ is **decreasing** on $[a, b]$.

- The **converse** of Increasing/Decreasing Test **fails**:

- f is **increasing** $\nRightarrow f'(x) > 0$;
- f is **decreasing** $\nRightarrow f'(x) < 0$.

Note that f is not necessarily differentiable.

Even if f is differentiable, f' may be zero at some points.


- Let $f(x) = x^3$. Then f is increasing on \mathbb{R} .
 - $f'(x) = 3x^2 \Rightarrow f'(0) = 0$.

58 / 101

Example

- Let $f(x) = 3x^4 - 4x^3 - 12x^2 + 5$.
 - $f'(x) = 12x^3 - 12x^2 - 24x = 12(x+1)x(x-2)$.

Interval	$x+1$	x	$x-2$	$f'(x)$	$f(x)$
$(-\infty, -1)$	-	-	-	-	\searrow
$(-1, 0)$	+	-	-	+	\nearrow
$(0, 2)$	+	+	-	-	\searrow
$(2, \infty)$	+	+	+	+	\nearrow




59 / 101

First Derivative Test

- Let f be a continuous function. Recall that
 - if f has a **local max** or **local min** at c ,
 - then c is a **critical number** of f .

Now suppose c is a **critical number** of f .

How to check if f has a **local max** or **local min** at c ?



60 / 101

First Derivative Test

• First Derivative Test.

Let f be **continuous** and c a **critical number** of f .
Suppose f is **differentiable** near c (except possibly at c).

- If f' changes from **positive** to **negative** at c ,
then f has a **local maximum** at c .
- If f' changes from **negative** to **positive** at c ,
then f has a **local minimum** at c .
- If f' does not change sign at c ,
then f has **no local max/min** at c .

• Proof. If f' changes from **positive** to **negative** at c , then

- f is **increasing** on the left of c , and
- f is **decreasing** on the right of c .

So f has a **local maximum** at c .

Other two cases can be shown similarly. (Exercise)

61 / 101


Examples

• $f(x) = 3x^4 - 4x^3 - 12x^2 + 5$

○ $f'(x) = 12x^3 - 12x^2 - 24x = 12(x+1)x(x-2)$.

Interval	$(-\infty, -1)$	$(-1, 0)$	$(0, 2)$	$(2, \infty)$
$f'(x)$	−	+	−	+
$f(x)$	↘	↗	↘	↗

- local maximum: $x = 0$;
- local minimum: $x = -1, x = 2$.



62 / 101




Examples


- $f(x) = x^{1/3}(x - 4)$. Find its local max and local min.

- Where are the critical numbers?

$$f'(x) = (x^{4/3} - 4x^{1/3})' = \frac{4}{3}x^{1/3} - \frac{4}{3}x^{-2/3} = \frac{4}{3} \frac{x - 1}{x^{2/3}}.$$

- $f'(x)$ does not exist: $x = 0$;
- $f'(x) = 0$: $x = 1$.


Interval	$(-\infty, 0)$	$(0, 1)$	$(1, \infty)$
$f'(x)$	—	—	+
$f(x)$			



63 / 101

Concavity

- Consider two graphs with the same end points:




- They are both increasing functions, but look different.
- We shall define **concavity** to distinguish the two types of (differentiable) functions.

64 / 101

Concavity

- **Definition.** Let f be differentiable on an open interval I .

- If the graph lies **above** all its tangent lines on I , then it is said to be **concave up**.
- If the graph lies **below** all its tangent lines on I , then it is said to be **concave down**.



65 / 101

Concavity Test

- **Theorem.** Let f be **differentiable** on open interval I .
 - The graph is **concave up** $\Leftrightarrow f'$ is **increasing**.
 - The graph is **concave down** $\Leftrightarrow f'$ is **decreasing**.
- Suppose f is twice differentiable on an open interval I .
 - If $f'' > 0$ on I , by Increasing Test f' is increasing, then the graph of f is concave up.
 - If $f'' < 0$ on I , by Decreasing Test f' is decreasing, then the graph of f is concave down.
- **The Concavity Test.** Let f be a **twice differentiable** function on an open interval I .
 - If $f'' > 0$ on I , then the graph of f is **concave up** on I .
 - If $f'' < 0$ on I , then the graph of f is **concave down** on I .

66 / 101

Graph f using f' and f''

- Sketch the graph of $f(x) = x^4 - 4x^3$.

- $f'(x) = 4x^2(x - 3)$.

Interval	$(-\infty, 0)$	$(0, 3)$	$(3, \infty)$
$f'(x)$	—	—	+
$f(x)$	\searrow	\searrow	\nearrow

So $f(x)$ has local minimum at $x = 3$.

- $f''(x) = 12x(x - 2)$.

Interval	$(-\infty, 0)$	$(0, 2)$	$(2, \infty)$
$f''(x)$	+	—	+
Concavity	Up	Down	Up


67 / 101

Graph f using f' and f''

- Sketch the graph of $f(x) = x^4 - 4x^3$.

Interval	$(-\infty, 0)$	$(0, 3)$	$(3, \infty)$
$f(x)$	\searrow	\searrow	\nearrow

Interval	$(-\infty, 0)$	$(0, 2)$	$(2, \infty)$
Concavity	Up	Down	Up



68 / 101

Graph f using f' and f''

- Sketch the graph of $f(x) = x^{2/3}(6-x)^{1/3}$.

◦ $f'(x) = \frac{4-x}{x^{1/3}(6-x)^{2/3}}$.

Interval	$(-\infty, 0)$	$(0, 4)$	$(4, 6)$	$(6, \infty)$
$f'(x)$	-	+	-	-
$f(x)$	\searrow	\nearrow	\searrow	\searrow

◦ $f''(x) = \frac{-8}{x^{4/3}(6-x)^{5/3}}$.

Interval	$(-\infty, 0)$	$(0, 6)$	$(6, \infty)$
$f''(x)$	-	-	+
Concavity	Down	Down	Up


69 / 101

Graph f using f' and f''

- Sketch the graph of $f(x) = x^{2/3}(6-x)^{1/3}$.

Interval	$(-\infty, 0)$	$(0, 4)$	$(4, 6)$	$(6, \infty)$
$f(x)$	\searrow	\nearrow	\searrow	\searrow

Interval	$(-\infty, 0)$	$(0, 6)$	$(6, \infty)$
Concavity	Down	Down	Up



70 / 101

Some Inequalities

- Show that for all positive $x \neq 1$, $2\sqrt{x} > 3 - \frac{1}{x}$.
 - Let $f(x) = 2\sqrt{x} - \left(3 - \frac{1}{x}\right) = 2\sqrt{x} - 3 + \frac{1}{x}$.
 - $f'(x) = \frac{1}{\sqrt{x}} - \frac{1}{x^2} = \frac{1}{x^2}(\sqrt{x^3} - 1)$.
 - $\begin{cases} f'(x) > 0, & \text{if } x > 1, \\ f'(x) < 0, & \text{if } 0 < x < 1. \end{cases}$
 - $\begin{cases} f \text{ is increasing on } [1, \infty), \\ f \text{ is decreasing on } (0, 1]. \end{cases}$
 - Then for any positive $x \neq 1$, $f(x) > f(1) = 0$.
That is,

$$2\sqrt{x} > 3 - \frac{1}{x}.$$

71 / 101

Some Inequalities

- We have seen that $\sin x < x$ for all $0 < x < \frac{\pi}{2}$.
Show that $\frac{2}{\pi}x < \sin x$ when $0 < x < \frac{\pi}{2}$.
 - Let $g(x) = \frac{\sin x}{x}$ on $(0, \frac{\pi}{2}]$.
 - $g'(x) = \left(\frac{\sin x}{x}\right)' = \frac{\cos x(x - \tan x)}{x^2} < 0$.
 - By Increasing Test, g is decreasing on $(0, \frac{\pi}{2}]$.
 - For any $0 < x < \frac{\pi}{2}$, $g(x) > g(\frac{\pi}{2}) = \frac{\sin(\pi/2)}{\pi/2} = \frac{2}{\pi}$.
That is,

$$\sin x > \frac{2x}{\pi}.$$

72 / 101

Some Inequalities

- Recall that $\sin x < x < \tan x$ for all $x \in (0, \frac{\pi}{2})$.
 $\tan x + \sin x$ and $2x$, which one is bigger on $(0, \frac{\pi}{2})$?
- Let $f(x) = \tan x + \sin x - 2x$.
 - $f'(x) = \sec^2 x + \cos x - 2$.
 $f''(x) = 2 \sec^2 x \tan x - \sin x > 0$
 - f' is increasing on $[0, \frac{\pi}{2})$.
 Then for any $x \in (0, \frac{\pi}{2})$, $f'(x) > f'(0) = 0$.
- f is increasing on $[0, \frac{\pi}{2})$.
 Then for any $x \in (0, \frac{\pi}{2})$, $f(x) > f(0)$.

$$\therefore \tan x + \sin x > 2x.$$

73 / 101

Approximation

- Suppose f is continuous. Then $x \rightarrow a \Rightarrow f(x) \rightarrow f(a)$.
 - In other words, if $x \approx a$, then $f(x) \approx f(a)$.
 - For example, $\sqrt{1.1} \approx \sqrt{1} = 1$.
- Question.** Do we have a better approximation under some stronger assumptions?
- Suppose f' is continuous. Then $x \rightarrow a \Rightarrow f'(x) \approx f'(a)$.
 - f can be approximated by the tangent line at a .
 - $f(x) \approx f(a) + f'(a)(x - a)$.
 - For example, let $f(x) = \sqrt{x}$, then $f'(x) = \frac{1}{2\sqrt{x}}$.
 - $f(1.1) \approx f(1) + f'(1)(1.1 - 1) = 1.05$.

74 / 101

Approximation

- Suppose f'' is continuous. Then
 - $x \rightarrow a \Rightarrow f'(x) \rightarrow f'(a)$ and $f''(x) \rightarrow f''(a)$.
 - Approximate f by a quadratic function P :
 - $P(a) = f(a), P'(a) = f'(a), P''(a) = f''(a)$.

Let $P(x) = p + q(x - a) + r(x - a)^2$. Then

$$P(a) = f(a) \Rightarrow p = f(a);$$

$$P'(a) = f'(a) \Rightarrow q = f'(a);$$

$$P''(a) = f''(a) \Rightarrow 2r = f''(a).$$

$$\therefore f(x) \approx f(a) + f'(a)(x - a) + \frac{f''(a)}{2}(x - a)^2.$$

75 / 101

Approximation

- Let $f(x) = \sqrt{x}$. $f'(x) = \frac{1}{2}x^{-1/2}$, $f''(x) = -\frac{1}{4}x^{-3/2}$.
 - $f(x) \approx f(1) + f'(1)(x - 1) + \frac{f''(1)}{2}(x - 1)^2$.
 - $\sqrt{1.1} \approx 1 + \frac{1}{2} \times 0.1 - \frac{1}{8} \times 0.1^2 = 1.04875$.
- In general, assume that $f^{(n)}$ is continuous, then f can be approximated by a polynomial $P(x)$ of degree n :
 - $f(x) \approx f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \cdots + \frac{f^{(n)}(a)}{n!}(x - a)^n$.
 - $P(a) = f(a), P'(a) = f'(a), \dots, P^{(n)}(a) = f^{(n)}(a)$.

76 / 101


Optimization Problems

- What is **optimization problem**?
 - Finding **extreme values** in practical application.
 - **Maximize** areas, volumes, profits, . . . ,
 - **Minimize** distances, costs, times,
- How to **optimize**?
 - Understand the problem.
 - Draw a diagram.
 - Introduce notations.
 - Find relations among the variables.
 - Express the problem as finding the absolute maximum or minimum of a function $f(x)$ on a specified domain.
 - Find the absolute maximum and minimum.
 - Closed Interval Method (on finite closed interval),
 - Increasing/Decreasing Test (works for all cases).

77 / 101

Example 1

- **Example.** A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He needs no fence along the river. What are the dimensions of the field that has the **largest** area?



- Aim: Maximize $A = xy$, where
 - $2x + y = 2400$, $x, y \geq 0$.

78 / 101


Example 1

- Maximize $A = xy$, where $2x + y = 2400$, $x, y \geq 0$.
 - $2x + y = 2400 \Rightarrow y = 2400 - 2x$.
 - $y \geq 0 \Rightarrow x \leq 1200$.
- It is equivalent to
Finding maximum of $A(x) = x(2400 - 2x)$ on $[0, 1200]$.
 - Critical number:
 - $A'(x) = 2400 - 4x$.
 - $A'(x) = 0 \Rightarrow x = 600$. $A(600) = 720\,000$.
 - Endpoints: $x = 0, 1200$. $A(0) = A(1200) = 0$.
 - $A(x)$ has maximum value 720 000 when $x = 600$.
- Conclusion: the field has the largest area 720 000 ft²,
when it has width 600 ft, and length 1200 ft.

79 / 101

Example 2

- A cylindrical can is to be made to hold 1 liter of oil. Find the dimensions that will **minimize** the cost of the metal to manufacture the can.



- Minimize $S = 2\pi rh + 2\pi r^2$, where
 - $V = \pi r^2 h = 1$, $r, h > 0$.

80 / 101

Example 2

- Minimize $S = 2\pi r^2 + 2\pi r h$, $V = \pi r^2 h = 1$, $r, h > 0$.

- $h = \frac{1}{\pi r^2} \Rightarrow S = 2\pi r^2 + \frac{2\pi r}{\pi r^2} = 2\pi r^2 + \frac{2}{r}$.

Find the minimum of $S(r) = 2\pi r^2 + \frac{2}{r}$ for $r > 0$.

- $S'(r) = 4\pi r - \frac{2}{r^2} = \frac{2}{r^2}(2\pi r^3 - 1)$.

- $S'(r) = 0 \Rightarrow r = \frac{1}{\sqrt[3]{2\pi}} = r_0$.

$0 < r < r_0 \Rightarrow S'(r) < 0$; $r > r_0 \Rightarrow S'(r) > 0$.

$S(r)$ is decreasing on $(0, r_0]$, is increasing on $[r_0, \infty)$.

$\therefore S(r)$ has the absolute minimum at $r = r_0$.


- The cost is minimized when we choose

radius $r = r_0 = \frac{1}{\sqrt[3]{2\pi}}$, and height $h = \sqrt[3]{\frac{4}{\pi}}$.

81 / 101

Example 3

- Find the point on the parabola $y^2 = 2x$ that is **closest** to the point $(1, 4)$.



- Minimize $d = \sqrt{(x - 1)^2 + (y - 4)^2}$,

- where $y^2 = 2x$.

82 / 101

Example 3

- Minimize $d = \sqrt{(x-1)^2 + (y-4)^2}$, with $y^2 = 2x$.

- $d(y) = \sqrt{\left(\frac{y^2}{2} - 1\right)^2 + (y-4)^2}$, $(x = \frac{y^2}{2})$.

It is equivalent to minimizing

- $f(y) = (d(y))^2 = \left(\frac{y^2}{2} - 1\right)^2 + (y-4)^2$ on \mathbb{R} .

- $f'(y) = y^3 - 8$. Then $f'(y) = 0 \Rightarrow y = 2$.
- If $y < 2$, $f'(y) < 0$; f is decreasing on $(-\infty, 2]$.
- If $y > 2$, $f'(y) > 0$; f is increasing on $[2, \infty)$.

- So $f(y)$ attains the absolute minimum at $y = 2$.

$\therefore d(y)$ attains the absolute minimum at $y = 2$. ($x = 2$)


- Therefore, the point on $y^2 = 2x$ which is closest to $(4, 1)$ is $(2, 2)$. Moreover, the distance is

$$d = \sqrt{(2-4)^2 + (2-1)^2} = \sqrt{5}.$$

83 / 101

Example 4

- A man launches his boat from point A on a bank of a straight river, 3km wide, and wants to reach point B , 8km downstream on the opposite bank, **as quick as possible**. If he can row 6km/h and run 8km/h, where should he land?



- Minimize $T(x) = \frac{\sqrt{9+x^2}}{6} + \frac{8-x}{8}$, $0 \leq x \leq 8$.

84 / 101

Example 4


- Minimize $T(x) = \frac{\sqrt{9+x^2}}{6} + \frac{8-x}{8}$ on $[0, 8]$.
 - $T'(x) = \frac{x}{6\sqrt{9+x^2}} - \frac{1}{8}$.
 $T'(x) = 0 \Rightarrow 8x = 6\sqrt{9+x^2} \Rightarrow 16x^2 = 81 + 9x^2$
 $\Rightarrow 7x^2 = 81 \Rightarrow x = \frac{9}{\sqrt{7}} (x > 0)$.
 - Compare the values $T(0)$, $T(8)$ and $T(\frac{9}{\sqrt{7}})$:
 - $T(0) = 1.5$, $T(8) = \frac{73}{6} \approx 1.42$, and
 $T(\frac{9}{\sqrt{7}}) = 1 + \frac{\sqrt{7}}{8} \approx 1.33$.
 - Therefore, he should land at $9/\sqrt{7}$ km away downstream from the starting point.

85 / 101

Example 5: Fermat's Principle and Snell's Law


- Fermat's Principle.** The light travels along a path for which the time is minimized.
- Snell's Law.** Let v_1 and v_2 be the velocity of light in air and in water respectively. Use Fermat's Principle to show that

$$\frac{\sin \theta_1}{\sin \theta_2} = \frac{v_1}{v_2}.$$



86 / 101

Example 5: Fermat's Principle and Snell's Law



- Minimize $T(x) = \frac{\sqrt{a^2 + x^2}}{v_1} + \frac{\sqrt{b^2 + (d-x)^2}}{v_2}$ on $[0, d]$.

$$T'(x) = \frac{x}{v_1 \sqrt{a^2 + x^2}} - \frac{d-x}{v_2 \sqrt{b^2 + (d-x)^2}} = \frac{\sin \theta_1}{v_1} - \frac{\sin \theta_2}{v_2}.$$

- As x moves from 0 to d smoothly,

- $\theta_1 \nearrow$ and $\theta_2 \searrow \Rightarrow T'(x) \nearrow$.
- $T'(0) < 0, T'(d) > 0, T'$ is continuous on $[0, d]$.
 \Rightarrow there is a unique $x_0 \in (0, d)$ with $T'(x_0) = 0$.

$T'(x)$ increases smoothly from negative to positive.


- $T'(x) < 0$ on $(0, x_0) \Rightarrow T(x) \searrow$ on $[0, x_0]$,
- $T'(x) > 0$ on $(x_0, d) \Rightarrow T(x) \nearrow$ on $[x_0, d]$.

$\therefore T(x)$ attains the min if $x = x_0$, at which $\frac{\sin \theta_1}{v_1} = \frac{\sin \theta_2}{v_2}$.

87 / 101

Example 5: Fermat's Principle and Snell's Law

- Recall Example 4:



- By Snell's Law, the time is minimized when $\frac{\sin \theta}{v_1} = \frac{\sin \frac{\pi}{2}}{v_2}$.

$$\circ \frac{x/\sqrt{9+x^2}}{6} = \frac{1}{8} \Rightarrow x = \frac{9}{\sqrt{7}}.$$

88 / 101

Limits of Indeterminate Forms

- How do we compute the following limits?

$$\circ \lim_{x \rightarrow 0} \frac{1 - \cos x}{x + x^2}, \lim_{x \rightarrow 0} \frac{x - \sin x}{x^3}, \lim_{x \rightarrow 0} \frac{\sqrt{1-x} - 1 + \frac{x}{2}}{x^2}.$$

Both the numerator and denominator tend to 0 as $x \rightarrow 0$.

They have the **0/0 Indeterminate Form**.

- How to compute the following?

$$\circ \lim_{x \rightarrow \frac{\pi}{2}} \frac{\sec x}{1 + \tan x}, \lim_{x \rightarrow \infty} \frac{x^2 + 3x}{3x^2 + 1}.$$

Both the numerator and denominator tend to $\pm\infty$ as $x \rightarrow \frac{\pi}{2}$ or $x \rightarrow \infty$.

They have the **∞/∞ Indeterminate Form**.

- Question.** Can we evaluate the limits without using ϵ, δ -definition?
 - We may use **differentiation**.

89 / 101

Example

- $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x + x^2}.$

- Let $f(x) = 1 - \cos x$ and $g(x) = x + x^2$.

- $f(0) = g(0) = 0.$
- $f'(x) = \sin x$, and $g'(x) = 1 + 2x.$

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{f(x)}{g(x)} &= \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{g(x) - g(0)} \\ &= \lim_{x \rightarrow 0} \frac{[f(x) - f(0)]/(x - 0)}{[g(x) - g(0)]/(x - 0)} \\ &= \frac{\lim_{x \rightarrow 0} [f(x) - f(0)]/(x - 0)}{\lim_{x \rightarrow 0} [g(x) - g(0)]/(x - 0)} \\ &= \frac{f'(0)}{g'(0)} = \frac{0}{1 + 2 \cdot 0} = 0. \end{aligned}$$

- However, this method does not work if $g'(0) = 0$.

90 / 101

l'Hôpital's Rule

- **l'Hôpital's Rule.** Let f and g be functions such that
 - $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0$, and
 - f and g are **differentiable** near a (except at a), and
 - $g'(x) \neq 0$ near a (except at a).

Then $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$, provided that the limit on the right hand side exists or equals $\pm\infty$.

- Guillaume François Antoine, Marquis de l'Hôpital (1661–1704) French Mathematician.
 - l'Hôpital's rule is published in his “*Analysis of the infinitely small to understand curves*”, the first book on differential calculus.
 - The rule is discovered by Johann Bernoulli (1667–1748), a Swiss Mathematician.

91 / 101

Examples

- Find $\lim_{x \rightarrow 0} \frac{\sqrt{1-x} - 1 + \frac{x}{2}}{x^2}$.

$$\begin{aligned}
 \lim_{x \rightarrow 0} \frac{\sqrt{1-x} - 1 + \frac{x}{2}}{x^2} &= \lim_{x \rightarrow 0} \frac{(\sqrt{1-x} - 1 + \frac{x}{2})'}{(x^2)'} \\
 &= \lim_{x \rightarrow 0} \frac{\frac{-1}{2\sqrt{1-x}} + \frac{1}{2}}{2x} = \lim_{x \rightarrow 0} \frac{\left(\frac{-1}{2\sqrt{1-x}} + \frac{1}{2}\right)'}{(2x)'} \\
 &= \lim_{x \rightarrow 0} \frac{\frac{-1}{2} \cdot \frac{1}{2\sqrt{(1-x)^3}}}{2} \\
 &= -\frac{1}{8}.
 \end{aligned}$$

92 / 101

Examples

- Find $\lim_{x \rightarrow 0} \frac{x - \sin x}{x^3}$.

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{x - \sin x}{x^3} &= \lim_{x \rightarrow 0} \frac{(x - \sin x)'}{(x^3)'} \\ &= \lim_{x \rightarrow 0} \frac{1 - \cos x}{3x^2} \\ &= \lim_{x \rightarrow 0} \frac{(1 - \cos x)'}{(3x^2)'} \\ &= \lim_{x \rightarrow 0} \frac{\sin x}{6x} \\ &= \lim_{x \rightarrow 0} \frac{(\sin x)'}{(6x)'} \\ &= \lim_{x \rightarrow 0} \frac{\cos x}{6} = \frac{1}{6}. \end{aligned}$$

93 / 101

Remarks on l'Hôpital's Rule

- Remark.**

- The condition $x \rightarrow a$ may be replaced by $x \rightarrow a^+$ or $x \rightarrow a^-$. In other words, l'Hôpital's Rule also holds for **one sided limit**.
- l'Hôpital's Rule holds if $x \rightarrow a$ is replaced by $x \rightarrow \infty$ or $x \rightarrow -\infty$. In other words, it holds for **limit at infinity**.
 - If f and g are differentiable for large x , $\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} g(x) = 0$, $g'(x) \neq 0$ for large x .
Then $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \infty} \frac{f'(x)}{g'(x)}$, if the limit on the right hand side exists or equals $\pm\infty$.
- l'Hôpital's Rule holds if $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0$ is replaced by $\lim_{x \rightarrow a} |f(x)| = \lim_{x \rightarrow a} |g(x)| = \infty$.

94 / 101

l'Hôpital's Rule (∞/∞)

- l'Hôpital's Rule. Suppose that

- $\lim_{x \rightarrow a} |f(x)| = \lim_{x \rightarrow a} |g(x)| = \infty$,
- f and g are differentiable near a (except at a),
- $g'(x) \neq 0$ for all x near a (except at a).

Then $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$, provided that the limit on the right side exists or equals $\pm\infty$.

- Example.** Find $\lim_{x \rightarrow \pi/2} \frac{\sec x}{1 + \tan x}$.

$$\begin{aligned} \lim_{x \rightarrow \pi/2} \frac{\sec x}{1 + \tan x} &= \lim_{x \rightarrow \pi/2} \frac{(\sec x)'}{(1 + \tan x)'} \\ &= \lim_{x \rightarrow \pi/2} \frac{\sec x \tan x}{\sec^2 x} \\ &= \lim_{x \rightarrow \pi/2} \sin x = 1. \end{aligned}$$

95 / 101

Examples

- Find $\lim_{x \rightarrow \infty} \frac{x^2 + 3x}{3x^2 + 1}$.

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{x^2 + 3x}{3x^2 + 1} &= \lim_{x \rightarrow \infty} \frac{(x^2 + 3x)'}{(3x^2 + 1)'} = \lim_{x \rightarrow \infty} \frac{2x + 3}{6x} \\ &= \lim_{x \rightarrow \infty} \frac{(2x + 3)'}{(6x)'} = \frac{2}{6} = \frac{1}{3}. \end{aligned}$$

- Find $\lim_{x \rightarrow 1} (1 - x^2) \tan \frac{\pi x}{2}$.

$$\begin{aligned} \lim_{x \rightarrow 1} (1 - x^2) \tan \frac{\pi x}{2} &= \lim_{x \rightarrow 1} \frac{1 - x^2}{\cot \frac{\pi x}{2}} = \lim_{x \rightarrow 1} \frac{(1 - x^2)'}{(\cot \frac{\pi x}{2})'} \\ &= \lim_{x \rightarrow 1} \frac{-2x}{-\frac{\pi}{2} \csc^2 \frac{\pi x}{2}} = \frac{-2}{-\frac{\pi}{2} \cdot 1} = \frac{4}{\pi}. \end{aligned}$$

- Convert $0 \cdot \infty$ or $\infty - \infty$ indeterminate forms to $\frac{0}{0}$ or $\frac{\infty}{\infty}$ indeterminate forms, then apply l'Hôpital's rule.

96 / 101

Correct or Wrong?

- Evaluate $\lim_{x \rightarrow 1} \frac{x^2 + 1}{2x + 1}$.
 - $\lim_{x \rightarrow 1} \frac{x^2 + 1}{2x + 1} = \lim_{x \rightarrow 1} \frac{(x^2 + 1)'}{(2x + 1)'} = \lim_{x \rightarrow 1} \frac{2x}{2} = \lim_{x \rightarrow 1} x = 1$.
 - We **cannot** apply l'Hôpital's rule unless the limits of numerator and denominator are both 0 or both $\pm\infty$.
- Evaluate $\lim_{x \rightarrow \infty} \frac{x + \sin x}{x}$.
 - $\lim_{x \rightarrow \infty} \frac{x + \sin x}{x} = \lim_{x \rightarrow \infty} \frac{(x + \sin x)'}{x'} = \lim_{x \rightarrow \infty} \frac{1 + \cos x}{1} = \lim_{x \rightarrow \infty} (1 + \cos x)$. So the limit does not exist.
 - l'Hôpital's rule is **inconclusive** if $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} \neq L, \pm\infty$. We shall use squeeze theorem for this question.

97 / 101

Correct or Wrong?

- Evaluate $\lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2 + 1}}$.

$$\begin{aligned}
 \lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2 + 1}} &= \lim_{x \rightarrow \infty} \frac{1}{x/\sqrt{x^2 + 1}} \\
 &= \lim_{x \rightarrow \infty} \frac{\sqrt{x^2 + 1}}{x} \\
 &= \lim_{x \rightarrow \infty} \frac{x/\sqrt{x^2 + 1}}{1} \\
 &= \lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2 + 1}} = \dots
 \end{aligned}$$

The l'Hôpital's rule is useful only when the evaluation of $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$ is simpler than the evaluation of

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)}.$$

98 / 101

Properties of the Exponential Function

- We can compute the number e numerically as a limit.

- **Theorem.** $e = \lim_{x \rightarrow 0} (1+x)^{1/x}$.

$$\begin{aligned}\lim_{x \rightarrow 0} (1+x)^{1/x} &= \lim_{x \rightarrow 0} \exp\left(\frac{1}{x} \ln(1+x)\right) \\ &= \exp\left(\lim_{x \rightarrow 0} \frac{\ln(1+x)}{x}\right) \\ &= \exp\left(\lim_{x \rightarrow 0} \frac{(\ln(1+x))'}{(x)'}\right) \\ &= \exp\left(\lim_{x \rightarrow 0} \frac{1/(1+x)}{1}\right) \\ &= \exp(1) = e.\end{aligned}$$

- **Remark.** Let $y = 1/x$. Then $x \rightarrow 0^+ \Leftrightarrow y \rightarrow \infty$.

- $e = \lim_{y \rightarrow \infty} (1+1/y)^y = \lim_{n \rightarrow \infty} (1+1/n)^n$.

99 / 101

Example

- Evaluate $\lim_{x \rightarrow 0^+} x^x$.

$$\begin{aligned}\lim_{x \rightarrow 0^+} x^x &= \lim_{x \rightarrow 0^+} e^{x \ln x} = \exp\left(\lim_{x \rightarrow 0^+} x \ln x\right) \\ &= \exp\left(\lim_{x \rightarrow 0^+} \frac{\ln x}{1/x}\right) = \exp\left(\lim_{x \rightarrow 0^+} \frac{1/x}{-1/x^2}\right) \\ &= \exp\left(\lim_{x \rightarrow 0^+} (-x)\right) = \exp(0) = 1.\end{aligned}$$

- In general, in order to evaluate $\lim_{x \rightarrow a} (f(x)^{g(x)})$, we use

$$\begin{aligned}\lim_{x \rightarrow a} (f(x)^{g(x)}) &= \lim_{x \rightarrow a} \exp(g(x) \ln(f(x))) \\ &= \exp\left(\lim_{x \rightarrow a} g(x) \ln f(x)\right) = \dots\end{aligned}$$

100 / 101

Continuously Compounded Interest

- Initial deposit: A dollars; Interest rate (year): r .
 - Suppose the interest is credited n times per year.
 - After one year, we have $A \left(1 + \frac{r}{n}\right)^n$ dollars.
- It seems that we will get more if n gets larger.
- **Question.** What will we get after one year if $n \rightarrow \infty$, in other words, if the **interest** is **continuously compounded**?

$$\begin{aligned}
 \circ \quad & \lim_{n \rightarrow \infty} A \left(1 + \frac{r}{n}\right)^n = \lim_{x \rightarrow \infty} A \left(1 + \frac{r}{x}\right)^x \\
 & = A \lim_{x \rightarrow \infty} \exp \left(x \ln \left(1 + \frac{r}{x}\right) \right) = A \exp \left(\lim_{x \rightarrow \infty} \frac{\ln \left(1 + \frac{r}{x}\right)}{\frac{1}{x}} \right) \\
 & = A \exp \left(\lim_{x \rightarrow \infty} \frac{\left(1 + \frac{r}{x}\right)^{-1} \frac{-r}{x^2}}{-\frac{1}{x^2}} \right) \\
 & = A \exp \left(\lim_{x \rightarrow \infty} r \left(1 + \frac{r}{x}\right)^{-1} \right) = A \exp(r) = Ae^r.
 \end{aligned}$$

101 / 101