

# MA1521 CALCULUS FOR COMPUTING

Wang Fei

matwf@nus.edu.sg

Department of Mathematics

Office: S17-06-16

Tel: 6516-2937

<b>Chapter 3: Sequences and Series</b>	<b>2</b>
Definition	3
Limit of Sequence	7
Limit Laws	9
Monotonic Sequences	14
Examples	17
Series	19
Geometric Series	20
Telescoping Series	25
Ratio Test	27
Root Test	30
Power Series	33
Convergence of Power Series	35
Convergence Theorem	38
Power Series Representation	43
Differentiation of Power Series	46
Taylor Series	48
Test for Divergence	53
Comparison Test	55
$p$ -Series	57
Limit Comparison Test	65
Alternating Series	68
Absolute Convergence	71

### What is a Sequence?

- Let's look at some examples of sequences:

- Positive integers:  $1, 2, 3, \dots, n, \dots$
- Constant sequence:  $1, 1, 1, \dots, 1, \dots$
- Power sequence:  $2, 4, 8, 16, \dots, 2^n, \dots$
- Alternating sequence:  $\frac{1}{\sqrt{1}}, \frac{-1}{\sqrt{2}}, \dots, \frac{(-1)^{n+1}}{\sqrt{n}}, \dots$

- Definition.** A **sequence** is a list of numbers written in a definite order:


$$a_1, a_2, a_3, \dots, a_n, \dots$$

- $a_1$ : the 1<sup>st</sup> term;  $a_2$ : the 2<sup>nd</sup> term;  $\dots$ ,  $a_n$ : the  $n^{\text{th}}$  **term**.
- The sequence is denoted by  $\{a_n\}_{n=1}^{\infty}$ , or simply  $\{a_n\}$ .
- $\{n\}_{n=1}^{\infty}$ ,  $\{1\}_{n=1}^{\infty}$ ,  $\{2^n\}_{n=1}^{\infty}$  and  $\left\{\frac{(-1)^{n+1}}{\sqrt{n}}\right\}_{n=1}^{\infty}$ .

3 / 78

### What is a Sequence?

- Consider the sequence  $a_1, a_2, a_3, \dots, a_n, \dots$




- It defines a function  $f : \mathbb{Z}^+ \rightarrow \mathbb{R}$ ,  $f(n) = a_n$ .
- Conversely, given a function  $f : \mathbb{Z}^+ \rightarrow \mathbb{R}$ , it defines a sequence  $\{a_n\}_{n=1}^{\infty}$  such that  $a_n = f(n)$ .
- Therefore, we have an alternative definition for sequence:
  - A **sequence** is a function  $\mathbb{Z}^+ \rightarrow \mathbb{R}$ .


4 / 78

## Examples

- $\left\{ \frac{n}{n+1} \right\}_{n=1}^{\infty}$ .



- $\left\{ \cos \frac{n\pi}{6} \right\}_{n=1}^{\infty}$ .



5 / 78

## Examples

- There are some sequences which cannot be defined by giving a simple formula for the terms,  $n \mapsto a_n$ .

- $\sqrt{2}, \sqrt{\sqrt{2}+2}, \sqrt{\sqrt{\sqrt{2}+2}+2}, \dots$

- $a_1 = \sqrt{2}, a_2 = \sqrt{a_1+2}, a_3 = \sqrt{a_2+2}, \dots$
- $a_1 = \sqrt{2}$  and  $a_n = \sqrt{a_{n-1}+2}$  for  $n \geq 2$ .

- $0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, \dots$

- $F_0 = 0, F_1 = 1$  and  $F_n = F_{n-1} + F_{n-2}$  for  $n \geq 2$ .
- It is the **Fibonacci sequence**.

- Leonardo da Pisa, (1170s or 1180s–1250)  
Italian mathematician.


- $F_n = \frac{(1 + \sqrt{5})^n - (1 - \sqrt{5})^n}{2^n \sqrt{5}}$ .

6 / 78

## Limit of Sequence

- Since a sequence can be viewed as a function, we can similarly talk about the **limit of sequence**.

- **Example.**  $\left\{\frac{n}{n+1}\right\}_{n=1}^{\infty}$ .



- $\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \frac{5}{6}, \frac{6}{7}, \frac{7}{8}, \frac{8}{9}, \frac{9}{10}, \frac{10}{11}, \frac{11}{12}, \frac{12}{13}, \frac{13}{14}, \frac{14}{15}, \frac{15}{16}, \dots$
- As  $n$  gets larger, the term  $a_n = \frac{n}{n+1}$  approaches 1.

We may use the similar notation as for function,

$$\lim_{n \rightarrow \infty} \frac{n}{1+n} = 1.$$

7 / 78

## Limit of Sequence

- **Definition.** Let  $\{a_n\}$  be a sequence.

- The **limit** of  $\{a_n\}$  is  $L$  if " $a_n$  is **arbitrarily close** to  $L$  by taking  $n$  **sufficiently large**". It is denoted by  $\lim_{n \rightarrow \infty} a_n = L$ .

- $\{a_n\}$  is called  $\begin{cases} \text{convergent,} & \text{if } \lim_{n \rightarrow \infty} a_n \text{ exists,} \\ \text{divergent,} & \text{otherwise.} \end{cases}$

- **Definition.** Let  $\{a_n\}$  be a sequence.


- The **limit** of  $\{a_n\}$  is  $\infty$  (resp.  $-\infty$ ) if " $a_n$  is **arbitrarily large** (resp. **arbitrarily negatively large**) by taking  $n$  **sufficiently large**". It is denoted by  $\lim_{n \rightarrow \infty} a_n = \infty$  (resp.  $-\infty$ ).

- **Remark.** If  $\lim_{n \rightarrow \infty} a_n = \pm\infty$ , then  $\{a_n\}$  is **divergent**.


8 / 78

## Examples

- We have known that  $\lim_{x \rightarrow \infty} \frac{x}{x+1} = 1$ .  
Can we use this fact to show that  $\lim_{n \rightarrow \infty} \frac{n}{n+1} = 1$ ?



- $\lim_{x \rightarrow \infty} \cos \frac{\pi x}{6}$  does not exist.  
Can we conclude that  $\lim_{n \rightarrow \infty} \cos \frac{\pi n}{6}$  does not exist as well?



9 / 78

## Limit Laws for Sequences

- Theorem.** Let  $f$  be a function and  $\{a_n\}$  be the sequence such that  $a_n = f(n)$  for all  $n$ .
  - If  $\lim_{x \rightarrow \infty} f(x) = L$ , then  $\lim_{n \rightarrow \infty} a_n = L$ .
- Example.** Evaluate  $\lim_{n \rightarrow \infty} \frac{\ln n}{n}$ .
  - Let  $f(x) = \frac{\ln x}{x}$ , ( $x > 0$ ). Then  $f(n) = a_n$  for all  $n$ .  

$$\lim_{x \rightarrow \infty} \frac{\ln x}{x} = \lim_{x \rightarrow \infty} \frac{1/x}{1} = 0 \Rightarrow \lim_{n \rightarrow \infty} \frac{\ln n}{n} = 0.$$
- Example.** Evaluate  $\lim_{n \rightarrow \infty} \sqrt[n]{n}$ .
  - Let  $f(x) = x^{1/x}$ , ( $x > 0$ ). Then  $f(n) = a_n$  for all  $n$ .  

$$\lim_{x \rightarrow \infty} x^{1/x} = \lim_{x \rightarrow \infty} e^{\frac{\ln x}{x}} = \exp \left[ \lim_{x \rightarrow \infty} \frac{\ln x}{x} \right] = e^0 = 1.$$

$$\Rightarrow \lim_{n \rightarrow \infty} \sqrt[n]{n} = 1.$$

10 / 78

## Examples

- We CANNOT use the theorem for the following cases:


- Evaluate  $\lim_{n \rightarrow \infty} \frac{n!}{n^n}$ .

- Let  $f(x) = \dots$ ?

- $n!$  is only defined for natural numbers. It cannot be extended easily to a function on real numbers.

- Evaluate  $\lim_{n \rightarrow \infty} \sin n\pi$ .

- Let  $f(x) = \sin x\pi$ . Then  $f(n) = a_n$  for all  $n$ .



- $\lim_{x \rightarrow \infty} f(x)$  doesn't exist. So  $\lim_{n \rightarrow \infty} a_n$  doesn't exist?
- However,  $\sin n\pi = 0$  for all  $n$ .  $\lim_{n \rightarrow \infty} \sin n\pi = 0$ .

11 / 78

## Limit Laws for Sequences

- **Theorem.** Let  $\{a_n\}$  and  $\{b_n\}$  be convergent sequences.

- $\lim_{n \rightarrow \infty} (a_n \pm b_n) = \lim_{n \rightarrow \infty} a_n \pm \lim_{n \rightarrow \infty} b_n$ .
- $\lim_{n \rightarrow \infty} (c a_n) = c \lim_{n \rightarrow \infty} a_n$ .
- $\lim_{n \rightarrow \infty} (a_n b_n) = \lim_{n \rightarrow \infty} a_n \cdot \lim_{n \rightarrow \infty} b_n$ .
- $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{\lim_{n \rightarrow \infty} a_n}{\lim_{n \rightarrow \infty} b_n}$ , if  $\lim_{n \rightarrow \infty} b_n \neq 0$ .

- **Theorem.**  $\lim_{n \rightarrow \infty} a_n \Leftrightarrow \lim_{n \rightarrow \infty} a_{2n-1} = \lim_{n \rightarrow \infty} a_{2n} = L$ .

- **Theorem.** Suppose  $a_n \leq b_n$  for all integer  $n$ .

- If  $\lim_{n \rightarrow \infty} a_n = L$  and  $\lim_{n \rightarrow \infty} b_n = M$ , then  $L \leq M$ .

- **Squeeze Theorem.** Suppose  $a_n \leq b_n \leq c_n$  for all  $n$ .

- If  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = L$ , then  $\lim_{n \rightarrow \infty} b_n = L$ .

12 / 78

## Limit Laws for Sequences


- **Example.** If  $\lim_{n \rightarrow \infty} |a_n| = 0$  then  $\lim_{n \rightarrow \infty} a_n = 0$ .
  - Note that  $-|a_n| \leq a_n \leq |a_n|$  for all  $n$ ,  

$$\lim_{n \rightarrow \infty} (-|a_n|) = -\lim_{n \rightarrow \infty} |a_n| = 0 = \lim_{n \rightarrow \infty} |a_n|.$$
  - By Squeeze Theorem  $\lim_{n \rightarrow \infty} a_n = 0$ .
  - E.g.,  $\lim_{n \rightarrow \infty} \left| \frac{(-1)^n}{n} \right| = \lim_{n \rightarrow \infty} \frac{1}{n} = 0 \Rightarrow \lim_{n \rightarrow \infty} \frac{(-1)^n}{n} = 0$ .
- **Example.** Evaluate  $\lim_{n \rightarrow \infty} \frac{n!}{n^n}$ .
  - $$\frac{n!}{n^n} = \frac{1 \cdot 2 \cdot 3 \cdots (n-1) \cdot n}{\underbrace{n \cdot n \cdot n \cdots n \cdot n}_{n \text{ times}}} = \frac{1}{n} \cdot \frac{2}{n} \cdot \frac{3}{n} \cdots \frac{n-1}{n} \cdot \frac{n}{n}.$$
  - $$0 \leq \frac{n!}{n^n} \leq \frac{1}{n}. \quad \left. \begin{array}{l} \lim_{n \rightarrow 0} 0 = 0 \\ \lim_{n \rightarrow 0} \frac{1}{n} = 0 \end{array} \right\} \Rightarrow \lim_{n \rightarrow \infty} \frac{n!}{n^n} = 0.$$

13 / 78

## Monotonic Sequences

- Similarly as increasing/decreasing functions, we can talk about increasing/decreasing sequences.
- **Definition.** Let  $\{a_n\}$  be a sequence.
  - $\{a_n\}$  is called **increasing** if  $a_n < a_{n+1}$  for all  $n$ ,
  - $\{a_n\}$  is called **decreasing** if  $a_n > a_{n+1}$  for all  $n$ .
  - $\{a_n\}$  is called **monotonic** if it is either increasing or decreasing.




$\left\{ \frac{n}{n+1} \right\}$  increases;  $\left\{ \frac{10-n}{10} \right\}$  decreases;  $\left\{ \sin \frac{n\pi}{6} \right\}$  neither.

14 / 78

## Examples

- Show that the sequence  $a_n = \frac{3}{n+5}$  is decreasing.



- $n < n+1 \Rightarrow n+5 < (n+1)+5$   
 $\Rightarrow \frac{3}{n+5} > \frac{3}{(n+1)+5} \Rightarrow a_n > a_{n+1}.$
- $a_n - a_{n+1} = \frac{3}{n+5} - \frac{3}{n+6} = \frac{3}{(n+5)(n+6)} > 0.$
- $\frac{a_{n+1}}{a_n} = \frac{3}{n+6} / \frac{3}{n+5} = \frac{n+5}{n+6} < 1, \quad (a_n > 0).$
- Let  $f(x) = \frac{3}{x+5}$ .  $f'(x) = -\frac{3}{(x+5)^2} < 0$  for  $x > 0$ .  
 $f$  is decreasing on  $\mathbb{R}^+ \Rightarrow \{a_n\}$  is decreasing.
- .....

15 / 78

## Examples

- Determine if  $a_n = \frac{n}{n^2+1}$  is increasing or decreasing.

- Let  $f(x) = \frac{x}{x^2+1}$ .
  - $f'(x) = \frac{1-x^2}{(x^2+1)^2} < 0$  for  $x > 1$ .
- $f$  is decreasing on  $[1, \infty) \Rightarrow \{a_n\}$  is decreasing.

- Determine if  $a_n = \frac{n!}{n^n}$  is increasing or decreasing.


$$\begin{aligned} \frac{a_{n+1}}{a_n} &= \frac{(n+1)!/(n+1)^{n+1}}{n!/n^n} = \frac{(n+1)!n^n}{n!(n+1)^{n+1}} \\ &= \frac{n^n}{(n+1)^n} = \left(\frac{n}{n+1}\right)^n < 1. \end{aligned}$$

- Therefore,  $\{a_n\}$  is decreasing.

16 / 78



## Examples



- Consider a segment of length 1.
  - Cut half in the first day.
  - Cut half of the remaining in the second day.
  - In general, cut half of the remaining everyday.
$$a_1 = \frac{1}{2}, a_2 = \frac{1}{4}, a_3 = \frac{1}{8}, \dots, a_n = \frac{1}{2^n}, \dots$$


How much have we cut by the  $n^{\text{th}}$  day?

- We shall evaluate the sum of the first  $n$  terms:
  - $S_n = a_1 + a_2 + \dots + a_n = \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^n}.$

17 / 78

## Examples

- Consider an  $8 \times 8$  chessboard.



- Put 1 grain of rice in the first square of the chessboard.
- Doubling the number in the next square.
- How much rice do we need to fill in the chessboard?
  - $a_1 = 1, a_2 = 2, a_3 = 4, \dots, a_n = 2^{n-1}, \dots$
  - $S_{64} = a_1 + a_2 + \dots + a_{64} = 1 + 2 + 4 + 8 + \dots + 2^{63}.$

18 / 78

## Series

- Let  $\{a_n\}_{n=1}^{\infty}$  be a sequence. Then the sum of the first  $n$  terms of  $\{a_n\}$  forms a new sequence  $\{S_n\}$ .

- $S_1 = a_1$ ;
- $S_2 = a_1 + a_2$ ;
- $S_3 = a_1 + a_2 + a_3$ ;
- .....;
- $S_n = a_1 + a_2 + \cdots + a_n = \sum_{i=1}^n a_i$ .

$\{S_n\}$  is called the sequence of **partial sums** of  $\{a_n\}$ .

- $\lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \sum_{i=1}^n a_i := \sum_{n=1}^{\infty} a_n$ .

This quantity is called an **infinite series**, or simply **series**.

- $\sum_{n=1}^{\infty} a_n$  is  $\begin{cases} \text{convergent,} & \text{if } \{S_n\} \text{ is convergent, (hội tụ)} \\ \text{divergent,} & \text{if } \{S_n\} \text{ is divergent. (phân kì)} \end{cases}$

19 / 78

## Examples

- Let us consider the examples shown at the beginning.
  - Example 1.**  $a_n = \frac{1}{2^n}$ . Then  $\sum_{n=1}^{\infty} a_n$  is convergent.
    - Then  $S_n = \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \cdots + \frac{1}{2^n} = 1 - \frac{1}{2^n}$ .
    - $\sum_{n=1}^{\infty} a_n = \lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \left(1 - \frac{1}{2^n}\right) = 1$ .
  - Example 2.**  $a_n = 2^{n-1}$ . Then  $\sum_{n=1}^{\infty} a_n$  is divergent.
    - Then  $S_n = 1 + 2 + 2^2 + \cdots + 2^{n-1} = 2^n - 1$ .
    - $\sum_{n=1}^{\infty} a_n = \lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} (2^n - 1) = \infty$ .
- They are special cases of **geometric series**.

20 / 78

## The Geometric Series

- Consider the **geometric sequence** ( $a \neq 0$ ).
  - $a_1 = a, a_2 = ar, a_3 = ar^2, \dots, a_n = ar^{n-1}, \dots$
  - $a$  is the **scalar factor**,  $r$  is the **common ratio**.
- $\sum_{n=1}^{\infty} ar^{n-1}$  is called a **geometric series**.
  - $S_n = a + ar + ar^2 + \dots + ar^{n-2} + ar^{n-1}$ .
  - $rS_n = ar + ar^2 + ar^3 + \dots + ar^{n-1} + ar^n$ .

Then  $(1 - r)S_n = a - ar^n = a(1 - r^n)$ .

- $S_n = \begin{cases} \frac{a(1 - r^n)}{1 - r}, & \text{if } r \neq 1, \\ na, & \text{if } r = 1. \end{cases}$
- $\sum_{n=1}^{\infty} ar^{n-1} = \lim_{n \rightarrow \infty} S_n = \begin{cases} \frac{a}{1 - r}, & \text{if } |r| < 1, \\ \text{divergent}, & \text{if } |r| \geq 1. \end{cases}$

21 / 78

## Examples

- Is the series  $\sum_{n=1}^{\infty} 2^{2n} 3^{1-n}$  convergent?
  - $\frac{a_{n+1}}{a_n} = \frac{2^{2(n+1)} 3^{1-(n+1)}}{2^{2n} 3^{1-n}} = \frac{4}{3} > 1$ .
  - Then  $\sum_{n=1}^{\infty} 2^{2n} 3^{1-n} = \sum_{n=1}^{\infty} 4 \left(\frac{4}{3}\right)^{n-1}$  is divergent.
- Is  $\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + \dots + x^n + \dots$  convergent?
  - Geometric series of scalar factor 1, common ratio  $x$ .
  - $\sum_{n=0}^{\infty} x^n = \begin{cases} \frac{1}{1 - x}, & \text{if } |x| < 1, \\ \text{divergent}, & \text{if } |x| \geq 1. \end{cases}$
  - The **Taylor series** for  $\frac{1}{1 - x}$  about 0.

22 / 78

## Examples

- Evaluate  $\frac{1}{\sqrt{11}} + \frac{1}{\sqrt{33}} + \frac{1}{\sqrt{99}} + \frac{1}{\sqrt{297}} + \dots$ .
  - This is a geometric series with common ratio  $r = \frac{1}{\sqrt{3}}$ .
  - $\frac{1}{\sqrt{11}} + \frac{1}{\sqrt{33}} + \frac{1}{\sqrt{99}} + \frac{1}{\sqrt{297}} + \dots = \frac{\frac{1}{\sqrt{11}}}{1 - \frac{1}{\sqrt{3}}}$
- Evaluate  $\sum_{n=1}^{\infty} \frac{3^{n-1} + 3^{n+1}}{5^n}$ .
  - $\sum_{n=1}^{\infty} \frac{3^{n-1}}{5^n} = \sum_{n=1}^{\infty} \frac{1}{5} \left(\frac{3}{5}\right)^{n-1} = \frac{\frac{1}{5}}{1 - \frac{3}{5}} = \frac{1}{2}$ .
  - $\sum_{n=1}^{\infty} \frac{3^{n+1}}{5^n} = \sum_{n=1}^{\infty} \frac{9}{5} \left(\frac{3}{5}\right)^{n-1} = \frac{\frac{9}{5}}{1 - \frac{3}{5}} = \frac{9}{2}$ .
  - Answer =  $1/2 + 9/2 = 5$ .

23 / 78

## Examples

- Recall that
  - Geometric series =  $\frac{\text{leading term}}{1 - \text{common ratio}}$  for  $|\text{ratio}| < 1$ .
- Find the range of  $x$  for which the series converges.
  - $\sum_{n=1}^{\infty} \left(\frac{2x-1}{3}\right)^{n-2} = \frac{\frac{3}{2x-1}}{1 - \frac{2x-1}{3}} = \frac{9}{(2x-1)(4-2x)}$ .
    - It converges  $\Leftrightarrow \left|\frac{2x-1}{3}\right| < 1 \Leftrightarrow -1 < x < 2$ .
  - $\sum_{n=1}^{\infty} \frac{2^{n-1} + 2^n + 2^{n+1}}{(x+1)^n} = \frac{\frac{7}{x+1}}{1 - \frac{2}{x+1}} = \frac{7}{x-1}$ .
    - It converges  $\Leftrightarrow \left|\frac{2}{x+1}\right| < 1 \Leftrightarrow x < -3 \text{ or } x > -1$ .

24 / 78

### Example

- Is the series  $\sum_{n=1}^{\infty} \frac{1}{n(n+1)}$  convergent?

- $a_n = \frac{1}{n(n+1)} = \frac{1}{n} - \frac{1}{n+1}.$

$$\begin{aligned} S_n &= \frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \cdots + \frac{1}{n(n+1)} \\ &= \left( \frac{1}{1} - \frac{1}{2} \right) + \left( \frac{1}{2} - \frac{1}{3} \right) + \left( \frac{1}{3} - \frac{1}{4} \right) \\ &\quad + \cdots + \left( \frac{1}{n-1} - \frac{1}{n} \right) + \left( \frac{1}{n} - \frac{1}{n+1} \right) \\ &= 1 - \frac{1}{n+1}. \end{aligned}$$

- $\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = \lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \left( 1 - \frac{1}{n+1} \right) = 1.$

25 / 78

### Telescoping Series

- The partial sum of a **telescoping series** has only a fixed number of terms after cancelation. Such evaluation is called the **method of differences**.

- Example.** Evaluate  $\sum_{n=1}^{\infty} \frac{n}{(n+1)!}.$

$$\begin{aligned} \frac{n}{(n+1)!} &= \frac{(n+1) - 1}{(n+1)!} = \frac{n+1}{(n+1)!} - \frac{1}{(n+1)!} = \frac{1}{n!} - \frac{1}{(n+1)!}. \\ S_n &= \frac{1}{2!} + \frac{2}{3!} + \cdots + \frac{n}{(n+1)!} \\ &= \left( \frac{1}{1!} - \frac{1}{2!} \right) + \left( \frac{1}{2!} - \frac{1}{3!} \right) + \cdots + \left( \frac{1}{n!} - \frac{1}{(n+1)!} \right) \\ &= \frac{1}{1!} - \frac{1}{(n+1)!}. \end{aligned}$$

$$\sum_{n=1}^{\infty} \frac{n}{(n+1)!} = \lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \left( 1 - \frac{1}{(n+1)!} \right) = 1.$$

26 / 78

## The Ratio Test

- Consider the series  $\sum_{n=1}^{\infty} a_n$ . Can we know its convergence by checking the ratio of consecutive terms?
  - If  $\left| \frac{a_{n+1}}{a_n} \right| = L$  for all  $n$ , then  $\sum_{n=1}^{\infty} |a_n|$  is a geometric series with common ratio  $L$ .
    - $\sum_{n=1}^{\infty} |a_n|$  is convergent  $\Leftrightarrow$  if  $|L| < 1$ .
  - If  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L$ , then  $\sum_{n=1}^{\infty} |a_n|$  is “more or less the same” as the geometric series of common ratio  $L$ .
    - Do we have a result of convergence for  $\sum_{n=1}^{\infty} a_n$  similar as that for the geometric series?

27 / 78

## The Ratio Test

- Theorem.** Let  $\sum_{n=1}^{\infty} a_n$  be a series.  
 Suppose  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L$ , where  $0 \leq L \leq \infty$ .
  - If  $0 \leq L < 1$ , then  $\sum_{n=1}^{\infty} a_n$  is convergent.
  - If  $1 < L \leq \infty$ , then  $\sum_{n=1}^{\infty} a_n$  is divergent.
  - If  $L = 1$ , the convergence of  $\sum_{n=1}^{\infty} a_n$  is inconclusive.
- Note.**
  - The ratio test does not work if  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| \neq L, \infty$ .
  - The ratio test does not work if  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1$ .

28 / 78

## Examples

- $\sum_{n=1}^{\infty} \frac{(-1)^n n^3}{3^n}$  is convergent.
  - $\left| \frac{a_{n+1}}{a_n} \right| = \left| \frac{(-1)^{n+1} (n+1)^3 / 3^{n+1}}{(-1)^n n^3 / 3^n} \right| = \frac{(n+1)^3}{3n^3}.$
  - $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \frac{(n+1)^3}{3n^3} = \lim_{n \rightarrow \infty} \frac{(1 + \frac{1}{n})^3}{3} = \frac{1}{3}.$
- $\sum_{n=1}^{\infty} \frac{n^n}{n!}$  is divergent.
  - $\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)^{n+1} / (n+1)!}{n^n / n!} = \frac{(n+1)^n}{n^n}.$
  - $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right)^n = e.$
  - $\lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right)^n = e.$

29 / 78

## The Root Test

- Let  $\sum_{n=1}^{\infty} a_n$  be a series.
  - If  $\sqrt[n]{|a_n|} = L$ , then  $|a_n| = L^n$ ,
    - $\sum_{n=1}^{\infty} |a_n|$  is a geometric series of common ratio  $L$ .
  - If  $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = L$ , then  $|a_n|$  is “similar” to  $L^n$ ,
    - $\sum_{n=1}^{\infty} |a_n|$  is thus “more or less the same” as  $\sum_{n=1}^{\infty} L^n$ .
  - We can guess that the **root test** should have the same conclusion as the **ratio test**.
    - They should have the same advantage, as well as the same disadvantage.
    - However, sometimes the **root test** works better.

30 / 78

## The Root Test

- **Theorem.** Let  $\sum_{n=1}^{\infty} a_n$  be a series.  
Suppose  $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = L$ , where  $0 \leq L \leq \infty$ .
  - If  $0 \leq L < 1$ , then  $\sum_{n=1}^{\infty} a_n$  is convergent,
  - If  $1 < L \leq \infty$ , then  $\sum_{n=1}^{\infty} a_n$  is divergent.
  - If  $L = 1$ , the convergence of  $\sum_{n=1}^{\infty} a_n$  is inconclusive.
- **Example.**  $\sum_{n=1}^{\infty} \left( \frac{2n+3}{3n+2} \right)^n$  is convergent.
  - $\sqrt[n]{\left( \frac{2n+3}{3n+2} \right)^n} = \frac{2n+3}{3n+2} = \frac{2 + \frac{3}{n}}{3 + \frac{2}{n}} \rightarrow \frac{2}{3}$  as  $n \rightarrow \infty$ .

31 / 78

## Examples

- $\sum_{n=1}^{\infty} \frac{n^n}{3^{1+3n}}$  is divergent.
  - $\lim_{n \rightarrow \infty} \sqrt[n]{\frac{n^n}{3^{1+3n}}} = \lim_{n \rightarrow \infty} \frac{n}{\sqrt[n]{3} \cdot 3} = \infty$ .
- The root test may work better than the ratio test.
  - $1 + 1 + \frac{1}{2} + \frac{1}{2} + \frac{1}{4} + \frac{1}{4} + \frac{1}{8} + \frac{1}{8} + \dots$
  - $a_{2n-1} = a_{2n} = \frac{1}{2^{n-1}}$ .
    - $\sqrt[2n-1]{a_{2n-1}} = \frac{1}{\sqrt[2n-1]{2^{n-1}}} = \frac{1}{2^{\frac{n-1}{2n-1}}} \rightarrow \frac{1}{\sqrt{2}},$
    - $\sqrt[2n]{a_{2n}} = \frac{1}{\sqrt[2n]{2^{n-1}}} = \frac{1}{2^{\frac{n-1}{2n}}} \rightarrow \frac{1}{\sqrt{2}}.$
  - By root test the series is convergent, but the ratio test does not work.

32 / 78




## Power Series

- Consider the geometric series

$$\sum_{n=0}^{\infty} r^n = 1 + r + r^2 + r^3 + \cdots + r^n + \cdots.$$

We have seen that  $\sum_{n=0}^{\infty} r^n = \begin{cases} \frac{1}{1-r}, & \text{if } |r| < 1, \\ \text{divergent}, & \text{if } |r| \geq 1. \end{cases}$

- Viewed as function:  $\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}, -1 < x < 1.$



33 / 78

## Power Series

- A **power series** about 0 is a series of the form

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + \cdots + c_n x^n + \cdots,$$

$c_i$ 's are constants, called **coefficients**, and  $x$  is a variable.

- In general, a **power series about  $a$**  is a series

$$\sum_{n=0}^{\infty} c_n (x - a)^n = c_0 + c_1 (x - a) + c_2 (x - a)^2 + \cdots.$$

- Remark.** By convention we write  $(x - a)^0 = 1$  for all  $x$ .

- In particular,  $\sum_{n=0}^{\infty} c_n (a - a)^n = c_0.$

- $\sum_{n=0}^{\infty} c_n (x - a)^n$  is convergent at  $x = a$  at least.

- How to find all  $x$  so that the power series is convergent?

34 / 78

## Examples

- Check the convergence of  $\sum_{n=0}^{\infty} a_n$ , where  $a_n = \frac{x^n}{\sqrt{n}}$ .
  - To check whether  $\sum_{n=0}^{\infty} a_n$  is convergent, use **ratio test**.
    - $\left| \frac{a_{n+1}}{a_n} \right| = \left| \frac{x^{n+1}/\sqrt{n+1}}{x^n/\sqrt{n}} \right| = \sqrt{\frac{n}{n+1}} |x|$ .
    - $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \sqrt{\frac{n}{n+1}} |x| = 1 \cdot |x| = |x|$ .
  - $\sum_{n=1}^{\infty} \frac{x^n}{\sqrt{n}}$  is  $\begin{cases} \text{convergent,} & \text{if } |x| < 1, \\ \text{divergent,} & \text{if } |x| > 1. \end{cases}$
  - We will learn how to determine the convergence at  $x = \pm 1$  soon.

35 / 78

## Examples

- $\sum_{n=0}^{\infty} \frac{x^n}{2^n}$  is convergent on  $(-2, 2)$ .
  - $\sum_{n=0}^{\infty} \frac{x^n}{2^n} = \sum_{n=0}^{\infty} \left(\frac{x}{2}\right)^n$  is a geometric series of ratio  $\frac{x}{2}$ .  
It is convergent  $\Leftrightarrow \left|\frac{x}{2}\right| < 1 \Leftrightarrow |x| < 2$ .
- $\sum_{n=0}^{\infty} \frac{x^n}{n!}$  is convergent on  $\mathbb{R}$ .
  - $\lim_{n \rightarrow \infty} \left| \frac{x^{n+1}/(n+1)!}{x^n/n!} \right| = \lim_{n \rightarrow \infty} \frac{|x|}{n+1} = 0$ .
- $\sum_{n=0}^{\infty} n!x^n$  is convergent at  $x = 0$  only.
  - $\lim_{n \rightarrow \infty} \left| \frac{(n+1)!x^{n+1}}{n!x^n} \right| = \lim_{n \rightarrow \infty} (n+1)|x| = \begin{cases} \infty, & x \neq 0, \\ 0, & x = 0. \end{cases}$

36 / 78

## Convergence Theorem for Power Series

- **Theorem.** Let  $\sum_{n=0}^{\infty} c_n x^n$  be a power series.

Then its convergence is described by one of the following three possibilities:

- (i) The series is convergent on  $\mathbb{R}$ ;
- (ii) The series is convergent at  $x = 0$  only;
- (iii) There is a number  $R > 0$  such that
  - the series is convergent if  $|x| < R$ ,
  - the series is divergent if  $|x| > R$ .

- **Remark.**

- The convergence at  $x = \pm R$  is inconclusive.
- For case (i), we may write  $R = \infty$ ;
- For case (ii), we may write  $R = 0$ .

37 / 78

## Convergence Theorem for Power Series


- **Theorem.** Let  $\sum_{n=0}^{\infty} c_n (x - a)^n$  be a power series.

- Then for some  $0 \leq R \leq \infty$ 
  - the series is convergent if  $|x - a| < R$ ;
  - the series is divergent if  $|x - a| > R$ .

- **Remark.** The convergence of the power series at  $x = a + R$  and  $x = a - R$  is inconclusive.

- **Definition.**

- $R$  is called the **radius of convergence**.



38 / 78

## Radius of Convergence

- Let  $\sum_{n=0}^{\infty} c_n(x-a)^n$  be a power series.
    - The radius of convergence  $R$  exists ( $0 \leq R \leq \infty$ ), but how to evaluate  $R$ ?
  - Consider the ratio:  $\left| \frac{c_{n+1}(x-a)^{n+1}}{c_n(x-a)^n} \right| = \left| \frac{c_{n+1}}{c_n} \right| \cdot |x-a|$ .
    - Suppose  $\lim_{n \rightarrow \infty} \left| \frac{c_{n+1}}{c_n} \right| = L$ .  
 Then  $\lim_{n \rightarrow \infty} \left| \frac{c_{n+1}(x-a)^{n+1}}{c_n(x-a)^n} \right| = L \cdot |x-a|$ .
    - The series is  $\begin{cases} \text{convergent,} & \text{if } L \cdot |x-a| < 1, \\ \text{divergent,} & \text{if } L \cdot |x-a| > 1. \end{cases}$
- $\therefore R = L^{-1} = \left( \lim_{n \rightarrow \infty} \left| \frac{c_{n+1}}{c_n} \right| \right)^{-1}$ .

39 / 78

## Radius of Convergence

- Consider the root:  $\sqrt[n]{|c_n(x-a)^n|} = \sqrt[n]{|c_n|} \cdot |x-a|$ .
    - Suppose  $\lim_{n \rightarrow \infty} \sqrt[n]{|c_n|} = L$ .  
 Then  $\lim_{n \rightarrow \infty} \sqrt[n]{|c_n(x-a)^n|} = L \cdot |x-a|$ .
    - The series is  $\begin{cases} \text{convergent,} & \text{if } L \cdot |x-a| < 1, \\ \text{divergent,} & \text{if } L \cdot |x-a| > 1. \end{cases}$
- $\therefore R = L^{-1} = \frac{1}{\lim_{n \rightarrow \infty} \sqrt[n]{|c_n|}}$ .
- Remark.**
    - If  $L = 0$ , then  $R = \infty$ ; if  $L = \infty$ , then  $R = 0$ .
    - The formulas hold only when  $\lim_{n \rightarrow \infty} \left| \frac{c_{n+1}}{c_n} \right|$  or  $\lim_{n \rightarrow \infty} \sqrt[n]{|c_n|}$  exists (or equals  $\infty$ ).

40 / 78

## Examples

- $\sum_{n=0}^{\infty} \frac{(2x-5)^n}{n^2}$ .  $c_n = \frac{2^n}{n^2}$ .  $R = 2^{-1} = 1/2$ .
  - $\lim_{n \rightarrow \infty} \frac{c_{n+1}}{c_n} = \lim_{n \rightarrow \infty} \frac{2^{n+1}/(n+1)^2}{2^n/n^2} = \lim_{n \rightarrow \infty} \frac{2n^2}{(n+1)^2}$   
 $= \lim_{n \rightarrow \infty} \frac{2}{(1+\frac{1}{n})^2} = 2$ .
- $\sum_{n=0}^{\infty} \frac{n^2(x-3)^{n+1}}{5^n}$ .  $c_{n+1} = \frac{n^2}{5^n}$ .  $R = (\frac{1}{5})^{-1} = 5$ 
  - $\lim_{n \rightarrow \infty} \frac{c_{n+1}}{c_n} = \lim_{n \rightarrow \infty} \frac{n^2/5^n}{(n-1)^2/5^{n-1}} = \lim_{n \rightarrow \infty} \frac{n^2}{5(n-1)^2}$   
 $= \lim_{n \rightarrow \infty} \frac{1}{5(1-\frac{1}{n})^2} = \frac{1}{5}$ .

41 / 78

## Examples

- $\sum_{n=0}^{\infty} \frac{3^{2n-1}(2x+1)^n}{n!}$ .  $c_n = \frac{3^{2n-1}2^n}{n!}$ .  $R = \infty$ .
  - $\lim_{n \rightarrow \infty} \frac{c_{n+1}}{c_n} = \lim_{n \rightarrow \infty} \frac{3^{2n+1}2^{n+1}/(n+1)!}{3^{2n-1}2^n/n!}$   
 $= \lim_{n \rightarrow \infty} \frac{18}{n+1} = 0$ .
- $\sum_{n=0}^{\infty} \sqrt[n]{n^n} \left(\frac{1}{2}x - 1\right)^n$ .  $c_n = \frac{\sqrt[n]{n^n}}{2^n}$ .  $R = 0$ .
  - $\lim_{n \rightarrow \infty} \sqrt[n]{c_n} = \lim_{n \rightarrow \infty} \sqrt[n]{\frac{\sqrt[n]{n^n}}{2^n}} = \lim_{n \rightarrow \infty} \frac{\sqrt[n]{n}}{2} = \infty$ .

42 / 78

## Power Series Representation

- Recall the geometric series

- $$\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + \cdots = \frac{1}{1-x} \text{ for } |x| < 1.$$
- $$\frac{1}{1-x}$$
 is represented as power series  $\sum_{n=0}^{\infty} x^n$  if  $|x| < 1$ .
- $$\sum_{n=0}^{\infty} x^n$$
 is a **power series representation** of  $\frac{1}{1-x}$ .

- Find a power series representation of  $\frac{1}{1+x^2}$  about 0.

- Note that  $\frac{a}{1-r} = \sum_{n=0}^{\infty} ar^n$ .
- $$\frac{1}{1+x^2} = \frac{1}{1-(-x^2)} = \sum_{n=0}^{\infty} (-x^2)^n = \sum_{n=0}^{\infty} (-1)^n x^{2n}.$$
- The identity holds  $\Leftrightarrow |x^2| < 1 \Leftrightarrow |x| < 1$ .

43 / 78

## Examples

- Find a power series representation of  $\frac{x^3}{x+2}$  at 0.

- $$\frac{x^3}{x+2} = \frac{\frac{x^3}{2}}{1+\frac{x}{2}} = \sum_{n=0}^{\infty} \frac{x^3}{2} \left(-\frac{x}{2}\right)^n = \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} x^{n+3}$$
- The identity holds  $\Leftrightarrow \left|-\frac{x}{2}\right| < 1 \Leftrightarrow |x| < 2$ .

- Find a power series representation of  $\frac{1}{1-x}$  at  $-1$ .

$$\begin{aligned} \frac{1}{1-x} &= \frac{1}{2-(x+1)} = \frac{\frac{1}{2}}{1-\frac{x+1}{2}} = \sum_{n=0}^{\infty} \frac{1}{2} \left(\frac{x+1}{2}\right)^n \\ &= \sum_{n=0}^{\infty} \frac{(x+1)^n}{2^{n+1}}. \end{aligned}$$

- The identity holds  $\Leftrightarrow \left|\frac{x+1}{2}\right| < 1 \Leftrightarrow |x+1| < 2$ .

44 / 78

## Examples

- Find a power series representation of  $\frac{1}{x^2 + 3x + 2}$  at 0.

$$\circ \frac{1}{x^2 + 3x + 2} = \frac{1}{(x+1)(x+2)} = \frac{1}{x+1} - \frac{1}{x+2}.$$

$$\bullet \frac{1}{x+1} = \frac{1}{1-(-x)} = \sum_{n=0}^{\infty} (-x)^n = \sum_{n=0}^{\infty} (-1)^n x^n.$$

$$\begin{aligned} \frac{1}{x+2} &= \frac{\frac{1}{2}}{1 + \frac{x}{2}} = \sum_{n=0}^{\infty} \frac{1}{2} \left( \frac{-x}{2} \right)^n \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} x^n. \end{aligned}$$

$$\circ \text{ Then } \frac{1}{x^2 + 3x + 2} = \sum_{n=0}^{\infty} \left[ 1 - \frac{1}{2^{n+1}} \right] (-1)^n x^n.$$

$\therefore$  The radius of convergence  $R = \min\{1, 2\} = 1$ .

45 / 78

## Differentiation of Power Series

- $\sum_{n=0}^{\infty} c_n x^n$  is a function.
  - Is it **differentiable**? If yes, what is the **derivative**?

- Power series is a “*generalization*” of polynomial.  
Consider polynomial  $P(x) = a_0 + a_1 x + \cdots + a_n x^n$ .

- It is continuous and differentiable,

$$\bullet P'(x) = a_1 + 2a_2 x + \cdots + na_n x^{n-1}.$$

- Theorem.** (Term by Term Differentiation)

Suppose  $\sum_{n=0}^{\infty} c_n x^n$  has radius of convergence  $R > 0$ .

- Then  $f(x) = \sum_{n=0}^{\infty} c_n x^n$  is differentiable on  $|x| < R$ .

$$\bullet f'(x) = \sum_{n=0}^{\infty} (c_n x^n)' = \sum_{n=1}^{\infty} n c_n x^{n-1}.$$

46 / 78

## Examples

- Recall that  $\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$  for  $|x| < 1$ .
  - Differentiate with respect to  $x$ :
    - $\sum_{n=1}^{\infty} nx^{n-1} = \frac{1}{(1-x)^2}$ .  $\sum_{n=1}^{\infty} nx^n = \frac{x}{(1-x)^2}$ .
  - Differentiate again with respect to  $x$ :
    - $\sum_{n=1}^{\infty} n^2 x^{n-1} = \frac{1+x}{(1-x)^3}$ .  $\sum_{n=1}^{\infty} n^2 x^n = \frac{x(1+x)}{(1-x)^3}$ .
  - They converge for  $|x| < 1$ . Let  $x = 1/2$ . We have
    - $\sum_{n=1}^{\infty} \frac{n}{2^n} = \frac{\frac{1}{2}}{(1-\frac{1}{2})^2} = 2 = \frac{1}{2} + \frac{2}{4} + \frac{3}{8} + \frac{4}{16} + \dots$
    - $\sum_{n=1}^{\infty} \frac{n^2}{2^n} = \frac{\frac{1}{2}(1+\frac{1}{2})}{(1-\frac{1}{2})^3} = 6 = \frac{1}{2} + \frac{4}{4} + \frac{9}{8} + \frac{16}{16} + \dots$

47 / 78

## The Coefficient of Power Series Representation

- Suppose  $\sum_{n=0}^{\infty} c_n x^n$  has radius of convergence  $R > 0$ .
  - Then  $f(x) = \sum_{n=0}^{\infty} c_n x^n$  is differentiable if  $|x| < R$ .

$$f'(x) = \sum_{n=1}^{\infty} c_n n x^{n-1},$$

$$f''(x) = \sum_{n=2}^{\infty} c_n n(n-1) x^{n-2},$$

$$f'''(x) = \sum_{n=3}^{\infty} c_n n(n-1)(n-2) x^{n-3},$$

$$\dots = \dots,$$

$$f^{(k)}(x) = \sum_{n=k}^{\infty} c_n n(n-1) \cdots (n-(k-1)) x^{n-k}.$$

- $f^{(n)}(0) = c_n n(n-1) \cdots (n-(n-1)) = c_n n!$

48 / 78



## Taylor Series and Maclaurin Series

- **Theorem.** Suppose  $f$  has a power series representation  $\sum_{n=0}^{\infty} c_n x^n$  of radius of convergence  $R > 0$ ,

- Then  $c_n = \frac{f^{(n)}(0)}{n!}$ , and  $f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$ .

Such series is called the **Maclaurin series** of  $f$ .

- **Theorem.** Suppose  $f$  has a power series representation  $\sum_{n=0}^{\infty} c_n (x - a)^n$  of radius of convergence  $R > 0$ ,

- Then  $c_n = \frac{f^{(n)}(a)}{n!}$  and  $f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$ .

Such series is called the **Taylor series** of  $f$  at  $a$ .

- Power series representation, if exists, is unique ( $R > 0$ ).

49 / 78

## Examples

- $\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + \cdots + x^n + \cdots$ .

- $c_n = 1$  for all  $n$ , and  $c_n = \frac{f^{(n)}(0)}{n!} \Rightarrow f^{(n)}(0) = n!$ .

- $\frac{x^3}{x+2} = \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} x^{n+3} = \sum_{n=3}^{\infty} \frac{(-1)^{n-3}}{2^{n-2}} x^n$ .

- $c_n = \begin{cases} 0, & n \leq 2, \\ \frac{(-1)^{n-3}}{2^{n-2}}, & n \geq 3. \end{cases} \quad f^{(n)}(0) = \begin{cases} 0, & n \leq 2, \\ \frac{(-1)^{n-3} n!}{2^{n-2}}, & n \geq 3. \end{cases}$

- **Note.**  $f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$  holds only if the power series representation of  $f(x)$  exists.

**Example.** Let  $f(x) = \begin{cases} e^{-1/x^2}, & \text{if } x \neq 0, \\ 0, & \text{if } x = 0. \end{cases}$

- $f^{(n)}(0) = 0$  for all  $n$ , but  $f(x)$  is not the zero function.

50 / 78

## Examples

- Find the Maclaurin series of  $f(x) = e^x$ .
  - $f'(x) = e^x, f''(x) = e^x, \dots, f^{(n)}(x) = e^x, \dots$
  - $\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots$
- Find the Taylor series of  $f(x) = e^{2x-1}$  at  $x = 1$ .
  - $e^{2x-1} = e^{2(x-1)+1} = e \cdot e^{2(x-1)} = \sum_{n=0}^{\infty} \frac{e \cdot 2^n (x-1)^n}{n!}$
  - What is  $f^{(2011)}(1)$ ?
    - $f^{(2011)}(1) = 2011! c_{2011} = 2011! \frac{e \cdot 2^{2011}}{2011!} = e \cdot 2^{2011}$

51 / 78

## Examples

- Find the Maclaurin series of  $f(x) = \sin x$ .

$f(x)$	$f'(x)$	$f''(x)$	$f^{(3)}(x)$
$\sin x$	$\cos x$	$-\sin x$	$-\cos x$
$f(0)$	$f'(0)$	$f''(0)$	$f^{(3)}(0)$
0	1	0	-1
$f^{(4)}(x)$	$f^{(5)}(x)$	$f^{(6)}(x)$	$f^{(7)}(x)$
$\sin x$	$\cos x$	$-\sin x$	$-\cos x$
$f^{(4)}(0)$	$f^{(5)}(0)$	$f^{(6)}(0)$	$f^{(7)}(0)$
0	1	0	-1

- $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!}$
- $\cos x = (\sin x)' = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!}$

52 / 78

## Test for Divergence

- Let  $\sum_{n=1}^{\infty} a_n$  be a convergent series.
  - Suppose that  $\sum_{n=1}^{\infty} a_n$  converges to  $L$ .
  - Let  $S_n = a_1 + a_2 + \cdots + a_{n-1} + a_n$ .
    - $S_{n-1} = a_1 + a_2 + \cdots + a_{n-1}$  for  $n \geq 2$ .

Then we have  $S_n - S_{n-1} = a_n$ .

$\therefore \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} S_n - \lim_{n \rightarrow \infty} S_{n-1} = L - L = 0$ .
- We proved: "If  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$ ."
- Test for Divergence.**
  - If  $\lim_{n \rightarrow \infty} a_n$  does not exist or  $\lim_{n \rightarrow \infty} a_n$  exists but  $\neq 0$ ,
  - then  $\sum_{n=1}^{\infty} a_n$  is divergent.

53 / 78

## Examples

- Is the series  $\sum_{n=1}^{\infty} \frac{n^2}{5n^2 + 4}$  convergent?
  - $\lim_{n \rightarrow \infty} \frac{n^2}{5n^2 + 4} = \lim_{n \rightarrow \infty} \frac{1}{5 + 4/n^2} = \frac{1}{5} \neq 0$ .

$\therefore \sum_{n=1}^{\infty} \frac{n^2}{5n^2 + 4}$  is divergent.
- Consider the geometric series  $\sum_{n=1}^{\infty} ar^{n-1}$ , ( $a \neq 0$ ).
  - $\lim_{n \rightarrow \infty} ar^{n-1} = \begin{cases} 0, & \text{if } |r| < 1, \\ a, & \text{if } r = 1, \\ \text{does not exist,} & \text{otherwise.} \end{cases}$

$\therefore \sum_{n=1}^{\infty} ar^{n-1}$  is divergent if  $|r| \geq 1$ .
- Note.** If  $\lim_{n \rightarrow \infty} a_n = 0$ , test for divergence is **inconclusive**.

54 / 78

## Example

- Is the series  $\sum_{n=1}^{\infty} \frac{1}{\sqrt{4^n + 1}}$  convergent?
  - $\lim_{n \rightarrow \infty} \frac{1}{\sqrt{4^n + 1}} = 0 \Rightarrow$  No Conclusion.
  - We see that  $\frac{1}{\sqrt{4^n + 1}} < \frac{1}{\sqrt{4^n}} = \frac{1}{2^n}$ .  
 $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is **convergent** " $\Leftarrow$ " terms of  $\sum_{n=1}^{\infty} \frac{1}{2^n}$  are "**small**".
    - The terms of  $\sum_{n=1}^{\infty} \frac{1}{\sqrt{4^n + 1}}$  are "**smaller**".
    - It seems that  $\sum_{n=1}^{\infty} \frac{1}{\sqrt{4^n + 1}}$  is **convergent** as well.
- Is the "*comparison*" true? Does it hold in general?

55 / 78

## The Comparison Test

- Theorem.** Let  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  be series such that
  - $0 \leq a_n \leq b_n$  for all  $n$ . (Or for all  $n \geq N$ )
 Then
 
$$\begin{cases} \sum_{n=1}^{\infty} b_n \text{ converges} & \Rightarrow \sum_{n=1}^{\infty} a_n \text{ converges.} \\ \sum_{n=1}^{\infty} a_n \text{ diverges} & \Rightarrow \sum_{n=1}^{\infty} b_n \text{ diverges.} \end{cases}$$
- Example.** Is the series  $\sum_{n=1}^{\infty} \frac{5}{2^n + 4n + 3}$  convergent?
  - $\frac{5}{2^n + 4n + 3} \leq \frac{5}{2^n}$  for all  $n$ .
  - $\sum_{n=1}^{\infty} \frac{1}{2^n}$  converges  $\Rightarrow \sum_{n=1}^{\infty} \frac{5}{2^n} = 5 \sum_{n=1}^{\infty} \frac{1}{2^n}$  converges.  
 $\Rightarrow \sum_{n=1}^{\infty} \frac{5}{2^n + 4n + 3}$  converges.

56 / 78

## **$p$ -Series**

- **Question.** For what values of  $p$ , is the  **$p$ -series**  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

- Use the **test for divergence**:

- $\lim_{n \rightarrow \infty} \frac{1}{n^p} = \begin{cases} 0, & \text{if } p > 0, \\ 1, & \text{if } p = 0, \\ \infty, & \text{if } p < 0. \end{cases}$

$\therefore \sum_{n=1}^{\infty} \frac{1}{n^p}$  is divergent if  $p \leq 0$ .

- However, we cannot use the test for divergence to conclude whether  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  is convergent if  $p > 0$ .

57 / 78

## **Harmonic Series**

- The **Harmonic series** is the  $p$ -series when  $p = 1$ :

- $H = \sum_{n=1}^{\infty} \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots + \frac{1}{n} + \cdots$

- Consider the partial sum of the first  $2^n$  terms:

- $H_1 = 1$ ;

- $H_2 = 1 + \frac{1}{2}$ ;

- $H_4 = 1 + \frac{1}{2} + \left(\frac{1}{3} + \frac{1}{4}\right) \geq 1 + \frac{1}{2} + \left(\frac{1}{4} + \frac{1}{4}\right)$

- $H_8 = 1 + \frac{1}{2} + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right)$   
 $> 1 + \frac{1}{2} + \left(\frac{1}{4} + \frac{1}{4}\right) + \left(\frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8}\right)$   
 $= 1 + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} = 1 + \frac{3}{2}$ .

58 / 78

## Harmonic Series

- The **Harmonic series** is the  $p$ -series when  $p = 1$ :

$$\circ H = \sum_{n=1}^{\infty} \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots + \frac{1}{n} + \cdots$$

- Consider the partial sum of the first  $2^n$  terms:

$$\begin{aligned} \bullet H_{2^n} &= 1 + \frac{1}{2} + \left(\frac{1}{3} + \frac{1}{4}\right) + \cdots + \left(\frac{1}{2^{n-1}+1} + \cdots + \frac{1}{2^n}\right) \\ &> 1 + \frac{1}{2} + \left(\frac{1}{4} + \frac{1}{4}\right) + \cdots + \left(\frac{1}{2^n} + \cdots + \frac{1}{2^n}\right) \\ &= 1 + \underbrace{\frac{1}{2} + \frac{1}{2} + \cdots + \frac{1}{2}}_{n \text{ copies}} = 1 + \frac{n}{2}. \end{aligned}$$

$$\circ \lim_{n \rightarrow \infty} \left(1 + \frac{n}{2}\right) = \infty.$$

$$\therefore \sum_{n=1}^{\infty} \frac{1}{n} = \lim_{n \rightarrow \infty} H_{2^n} = \infty. \text{ So } \sum_{n=1}^{\infty} \text{ is } \mathbf{divergent}.$$

59 / 78

## $p$ -Series

- Theorem.** The  $p$ -series

$$\circ \sum_{n=1}^{\infty} \frac{1}{n^p} \text{ is } \begin{cases} \mathbf{convergent} & \text{if } p > 1, \\ \mathbf{divergent} & \text{if } p \leq 1. \end{cases}$$

- Remark.**

$$\circ \text{ If } p \leq 1, \frac{1}{n^p} \geq \frac{1}{n}. \sum_{n=1}^{\infty} \frac{1}{n} \text{ diverges} \Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p} \text{ diverges.}$$

- The proof of the second statement is omitted.

- Can we use ratio test to check its convergence?

$$\circ \left| \frac{a_{n+1}}{a_n} \right| = \frac{1/(n+1)^p}{1/n^p} = \frac{n^p}{(n+1)^p}.$$

$$\circ \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \frac{n^p}{(n+1)^p} = \lim_{n \rightarrow \infty} \frac{1}{(1 + \frac{1}{n})^p} = 1.$$

- However, the convergence of  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  depends on  $p$ .

60 / 78

## The Root Test

- Can the root test do better for  $p$ -series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$ ?
  - $\lim_{n \rightarrow \infty} \sqrt[n]{\frac{1}{n^p}} = \frac{1}{\left(\lim_{n \rightarrow \infty} \sqrt[n]{n}\right)^p} = \frac{1}{1^p} = 1.$ 

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[n]{n} &= \lim_{x \rightarrow \infty} x^{1/x} = \lim_{x \rightarrow \infty} e^{\frac{\ln x}{x}} \\ &= \exp\left(\lim_{x \rightarrow \infty} \frac{\ln x}{x}\right) = \exp\left(\lim_{x \rightarrow \infty} \frac{1/x}{1}\right) = \exp(0) = 1. \end{aligned}$$
- In fact, if  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L$  then  $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|}$  must exist and equal  $L$ .
  - Hence, if one of the ratio test or root test has the limit 1, **DO NOT** try the other test since it does not work too.

61 / 78

## Examples

- **Note.** In order to use the **Comparison Test** for a (positive) series, we shall first “guess”
  - whether it is **convergent** or **divergent**.
  - If we guess it is **convergent**,
    - find a (positive) **convergent** series whose terms are **bigger** than the terms of the given series.
  - If we guess it is **divergent**,
    - find a (positive) **divergent** series whose terms are **smaller** than the terms of the given series.
- **Example.** Is  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  convergent?
 
$$\frac{\ln n}{n} \geq \frac{1}{n} \text{ if } n \geq 3. \quad \sum_{n=1}^{\infty} \frac{1}{n} \text{ diverges} \Rightarrow \sum_{n=1}^{\infty} \frac{\ln n}{n} \text{ diverges.}$$

62 / 78

## Examples

- **Example.** Is  $\sum_{n=1}^{\infty} \frac{\ln n}{n^2}$  convergent?
  - $\frac{\ln n}{n^2} > \frac{1}{n^2} \cdot \sum_{n=1}^{\infty} \frac{1}{n^2}$  converges  $\Rightarrow$  No conclusion!
  - $\frac{\ln n}{n^2} < \frac{n}{n^2} \cdot \sum_{n=1}^{\infty} \frac{1}{n}$  diverges  $\Rightarrow$  No conclusion!

Let's compare  $\ln n$  and  $\sqrt{n}$ :

- $f(x) = \ln x - \sqrt{x}$ .  $f'(x) = \frac{2 - \sqrt{x}}{2x} < 0$  if  $x > 4$ .
- For  $n \geq 4$ ,  $\ln n - \sqrt{n} \leq \ln 4 - \sqrt{4} \approx -0.6 < 0$ .

$$\frac{\ln n}{n^2} < \frac{\sqrt{n}}{n^2} = \frac{1}{n^{3/2}} \text{ for all } n \geq 4.$$

- $\sum_{n=1}^{\infty} \frac{1}{n^{3/2}}$  converges  $\Rightarrow \sum_{n=1}^{\infty} \frac{\ln n}{n^2}$  converges.

63 / 78

## Examples

- Is the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$  convergent?
  - It is "**similar**" to the convergent series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .  
 $\frac{1}{2^n - 1} > \frac{1}{2^n} \Rightarrow$  Inconclusive by comparison test.
  - $\frac{1}{2^n - 1} \leq \frac{1}{2^n - 2^{n-1}} = \frac{1}{2^{n-1}}$ .  
 $\sum_{n=1}^{\infty} \frac{1}{2^{n-1}}$  is convergent  $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{2^n - 1}$  is convergent.
- Is the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{5 + n^5}}$  convergent?
  - It seems that  $\frac{2n^2 + 3n}{\sqrt{5 + n^5}}$  is "**similar**" to  $\frac{2n^2}{\sqrt{n^5}} = \frac{2}{\sqrt{n}}$ .

64 / 78



## The Limit Comparison Test

- **Theorem.** Let  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  be series of positive terms.

(a) Suppose  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c$  is a positive real number.

- $\sum_{n=1}^{\infty} b_n$  is **convergent**  $\Leftrightarrow \sum_{n=1}^{\infty} a_n$  is **convergent**.

(b) Suppose  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 0$ .

- $\sum_{n=1}^{\infty} b_n$  is **convergent**  $\Rightarrow \sum_{n=1}^{\infty} a_n$  is **convergent**.
- $\sum_{n=1}^{\infty} a_n$  is **divergent**  $\Rightarrow \sum_{n=1}^{\infty} b_n$  is **divergent**.

(c) Suppose  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \infty$ .

- $\sum_{n=1}^{\infty} b_n$  is **divergent**  $\Rightarrow \sum_{n=1}^{\infty} a_n$  is **divergent**.
- $\sum_{n=1}^{\infty} a_n$  is **convergent**  $\Rightarrow \sum_{n=1}^{\infty} b_n$  is **convergent**.

65 / 78

## Examples

- Is the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{5 + n^5}}$  convergent?

- $\lim_{n \rightarrow \infty} \frac{(2n^2 + 3n)/\sqrt{5 + n^5}}{1/\sqrt{n}} = \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{\sqrt{\frac{5}{n^5} + 1}} = 2$ .
- $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$  is divergent  $\Rightarrow \sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{5 + n^5}}$  is divergent.

- Is the series  $\sum_{n=2}^{\infty} \frac{1}{(\ln n)^2}$  convergent?

- $\lim_{n \rightarrow \infty} \frac{1/n}{1/(\ln n)^2} = \lim_{n \rightarrow \infty} \frac{(\ln n)^2}{n} = \lim_{x \rightarrow \infty} \frac{(\ln x)^2}{x}$   
 $= \lim_{x \rightarrow \infty} \frac{2 \ln x \cdot \frac{1}{x}}{1} = \lim_{x \rightarrow \infty} \frac{2 \ln x}{x} = \lim_{x \rightarrow \infty} \frac{2}{x} = 0$ .
- $\sum_{n=1}^{\infty} \frac{1}{n}$  is divergent  $\Rightarrow \sum_{n=2}^{\infty} \frac{1}{(\ln n)^2}$  is divergent.

66 / 78

## Examples

- Is the series  $\sum_{n=2}^{\infty} \left( \frac{1}{\sqrt{n}} \sin \frac{1}{n} \right)$  convergent?
  - $\lim_{n \rightarrow \infty} \frac{\frac{1}{\sqrt{n}} \sin \frac{1}{n}}{\frac{1}{\sqrt{n}n}} = \lim_{n \rightarrow \infty} \frac{\sin \frac{1}{n}}{\frac{1}{n}} = \lim_{x \rightarrow 0^+} \frac{\sin x}{x} = 1.$
  - $\sum_{n=2}^{\infty} \frac{1}{n\sqrt{n}}$  converges  $\Rightarrow \sum_{n=2}^{\infty} \frac{1}{\sqrt{n}} \sin \frac{1}{n}$  converges.
- Is the series  $\sum_{n=1}^{\infty} \frac{\sin^2 n}{n^2}$  convergent?
  - $\lim_{n \rightarrow \infty} \frac{(\sin^2 n)/n^2}{1/n^2} = \lim_{n \rightarrow \infty} \sin^2 n.$  No Conclusion!
  - $\frac{\sin^2 n}{n^2} \leq \frac{1}{n^2}.$  So  $\sum_{n=1}^{\infty} \frac{\sin^2 n}{n^2}$  is convergent.

67 / 78


## Alternating Harmonic Series

- How about the series whose terms are not all positive?
- Is **alternating harmonic series**  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$  convergent?

$$+ \frac{1}{1} - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \frac{1}{7} - \frac{1}{8} + \frac{1}{9} - \frac{1}{10} + \frac{1}{11} - \frac{1}{12} + \dots$$



- Let us check the graph of  $S_n = a_1 + a_2 + \dots + a_n$ :



68 / 78

## The Alternating Series Test

- **Definition.** An **alternating series** is a series whose terms are alternatively *positive* and *negative*.
- **Leibniz Alternating Series Test.**
  - Let  $\sum_{n=1}^{\infty} a_n$  be an **alternating series**. Suppose
    - $\lim_{n \rightarrow \infty} |a_n| = 0$ , and  $\{|a_n|\}$  is **decreasing**.
  - Then the series  $\sum_{n=1}^{\infty} a_n$  is **convergent**.
- **Example.**
  - The **alternating Harmonic series**  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$  is convergent;
  - although the **Harmonic series**  $\sum_{n=1}^{\infty} \frac{1}{n}$  is divergent.

69 / 78

## Examples

- Is the alternating series  $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{n^2}{n^3 + 1}$  convergent?
  - $|a_n| = \frac{n^2}{n^3 + 1} \Rightarrow \lim_{n \rightarrow \infty} |a_n| = \lim_{n \rightarrow \infty} \frac{\frac{1}{n}}{1 + \frac{1}{n^3}} = 0$ .
  - Let  $f(x) = \frac{x^2}{x^3 + 1} \Rightarrow f'(x) = \frac{x(2 - x^3)}{(x^3 + 1)^2}$ .
    - $f'(x) < 0$  if  $x > \sqrt[3]{2} \Rightarrow \{|a_n|\}_{n=2}^{\infty}$  is decreasing.
  - $\therefore \sum_{n=1}^{\infty} (-1)^{n+1} \frac{n^2}{n^3 + 1}$  is convergent.
  - However,  $\sum_{n=1}^{\infty} \frac{n^2}{n^3 + 1}$  is divergent, (compare  $\sum_{n=1}^{\infty} \frac{1}{n}$ ).
- It seems that the condition that “ $\sum |a_n|$  **converges**” is “**stronger**” than the condition that “ $\sum a_n$  **converges**”.

70 / 78

## Absolute Convergence

- Let  $\sum_{n=1}^{\infty} a_n$  be a series. We can consider a new series
  - $\sum_{n=1}^{\infty} |a_n| = |a_1| + |a_2| + \cdots + |a_n| + \cdots$ .
- Theorem.**  $\sum_{n=1}^{\infty} |a_n|$  is converges  $\Rightarrow \sum_{n=1}^{\infty} a_n$  converges.
- Examples.**
  - $\sum_{n=1}^{\infty} \frac{1}{n^2}$  converges  $\Rightarrow \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2}$  converges.
  - If  $\sum_{n=1}^{\infty} |a_n|$  is divergent, then  $\sum_{n=1}^{\infty} a_n$  is **inconclusive**.
    - $\sum_{n=1}^{\infty} 1$  diverges, and  $\sum_{n=1}^{\infty} (-1)^n$  diverges.
    - $\sum_{n=1}^{\infty} \frac{1}{n}$  diverges, but  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$  converges.

71 / 78

## Absolute Convergence

- Definition.** Let  $\sum_{n=1}^{\infty} a_n$  be a series.
  - It is **absolutely convergent** if  $\sum_{n=1}^{\infty} |a_n|$  is convergent.
  - It is **conditionally convergent** if  $\sum_{n=1}^{\infty} |a_n|$  is divergent and  $\sum_{n=1}^{\infty} a_n$  is convergent.
- Examples.**
  - $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$  is conditionally convergent.
  - $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2}$  is absolutely convergent.
  - $\sum_{n=1}^{\infty} (-1)^n$  is divergent.

72 / 78

## Proof of Absolute Convergence Theorem

- **Proof.** Separate positive and negative terms in  $\sum_{n=1}^n a_n$ .

$a_n$ :	1,	1,	-4,	-5,	1,	-3,	-1,	2,	1,	7,
$a_n^+$ :	1,	1,	0,	0,	1,	0,	0,	2,	1,	7,
$a_n^-$ :	0,	0,	4,	5,	0,	3,	1,	0,	0,	0,

$$a_n^+ = \begin{cases} a_n, & \text{if } a_n \geq 0, \\ 0, & \text{if } a_n < 0. \end{cases} \quad a_n^- = \begin{cases} 0, & \text{if } a_n \geq 0, \\ -a_n, & \text{if } a_n < 0. \end{cases}$$

- $0 \leq a_n^+ \leq |a_n|$  and  $0 \leq a_n^- \leq |a_n|$ .
- $a_n^+ + a_n^- = |a_n|$  and  $a_n^+ - a_n^- = a_n$ .

- Suppose  $\sum_{n=1}^{\infty} |a_n|$  is convergent.

$$0 \leq a_n^+, a_n^- \leq |a_n| \Rightarrow \sum_{n=1}^{\infty} a_n^+ \text{ and } \sum_{n=1}^{\infty} a_n^- \text{ are convergent.}$$

- $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} a_n^+ - \sum_{n=1}^{\infty} a_n^-$  is convergent.

73 / 78

## Examples

- **Example.** Is  $\sum_{n=1}^{\infty} \frac{\cos n}{n^2}$  convergent?

$$\circ \sum_{n=1}^{\infty} \left| \frac{\cos n}{n^2} \right| \leq \sum_{n=1}^{\infty} \frac{1}{n^2}.$$

$$\circ \sum_{n=1}^{\infty} \frac{1}{n^2} \text{ is convergent (} p\text{-series).}$$

$$\Rightarrow \sum_{n=1}^{\infty} \left| \frac{\cos n}{n^2} \right| \text{ is convergent by comparison test.}$$

$$\Rightarrow \sum_{n=1}^{\infty} \frac{\cos n}{n^2} \text{ is convergent by absolute convergence test.}$$

- **Example.**  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^p}$ .

$$\circ \text{ It is } \begin{cases} \text{absolutely convergent,} & \text{if } p > 1, \\ \text{divergent,} & \text{if } p \leq 0, \\ \text{conditionally convergent,} & \text{if } 0 < p \leq 1. \end{cases}$$

74 / 78

## Examples

- Given  $\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$ . Evaluate  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2}$ .
  - $$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2} = 1 - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \frac{1}{5^2} - \frac{1}{6^2} + \dots$$

$$= \left(1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots\right) - \left(\frac{1}{2^2} + \frac{1}{4^2} + \frac{1}{6^2} + \dots\right)$$

$$= \left(\frac{\pi^2}{6} - \frac{1}{4} \cdot \frac{\pi^2}{6}\right) - \left(\frac{1}{4} \cdot \frac{\pi^2}{6}\right) = \frac{\pi^2}{8} - \frac{\pi^2}{24} = \frac{\pi^2}{12}.$$
  - $$\sum_{n=1}^{\infty} \frac{1}{n^2} = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2} + \frac{1}{6^2} + \dots = \frac{\pi^2}{6}$$

$$= \left(1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots\right) + \left(\frac{1}{2^2} + \frac{1}{4^2} + \frac{1}{6^2} + \dots\right)$$

$$= \left(1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots\right) + \frac{1}{4} \cdot \left(1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots\right) \frac{\pi^2}{6}$$

75 / 78

## Example

- Can we evaluate  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$  similarly?
  - $$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \frac{1}{7} - \dots$$

$$= \left(1 + \frac{1}{3} + \frac{1}{5} + \frac{1}{7} + \dots\right) - \left(\frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \dots\right)$$

$$= \infty - \infty = ? \quad \leftarrow \text{indeterminate form!}$$
  - $$\lim_{n \rightarrow \infty} \frac{1/(2n-1)}{1/n} = \lim_{n \rightarrow \infty} \frac{n}{2n-1} = \frac{1}{2}.$$
  - $$\lim_{n \rightarrow \infty} \frac{1/(2n)}{1/n} = \lim_{n \rightarrow \infty} \frac{n}{2n} = \frac{1}{2}.$$
- By limit comparison test with  $\sum_{n=1}^{\infty} \frac{1}{n}$ ,
  - $$\sum_{n=1}^{\infty} \frac{1}{2n-1} = \sum_{n=1}^{\infty} \frac{1}{2n}$$
 are divergent.

76 / 78

## Conditional Convergence affects Rearrangement

- **Theorem.** Let  $\sum_{n=1}^{\infty} a_n$  be a convergent series.
  - If  $\sum_{n=1}^{\infty} a_n$  is **absolutely convergent**, then every rearrangement has the same sum.
  - If  $\sum_{n=1}^{\infty} a_n$  is **conditionally convergent**, then different rearrangements may have different sum.
    - Moreover, for any  $L$  (a real number or  $\pm\infty$ ), there is a rearrangement of  $\sum_{n=1}^{\infty} a_n$  whose sum is  $L$ .
- This theorem shows that if the series is conditionally convergent, we should not evaluate the sum by rearranging (infinitely many) terms.

77 / 78

## Example

- Find a rearrangement of  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$  whose sum is 1.
  - $1 + \frac{1}{3} + \frac{1}{5} + \frac{1}{7} + \frac{1}{9} + \frac{1}{11} + \frac{1}{13} + \frac{1}{15} + \frac{1}{17} + \dots = \infty.$
  - $\frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \frac{1}{8} + \frac{1}{10} + \frac{1}{12} + \frac{1}{14} + \frac{1}{16} + \frac{1}{18} + \dots = \infty.$
- 1. If  $S_n \geq 1$ , add the negative terms until partial sum is  $< 1$ .
- 2. If  $S_n < 1$ , add the positive terms until partial sum is  $\geq 1$ .
- $\frac{1}{1} - \frac{1}{2} + \frac{1}{3} + \frac{1}{5} - \frac{1}{4} + \frac{1}{7} + \frac{1}{9} - \frac{1}{6} + \frac{1}{11} + \frac{1}{13} - \frac{1}{8}$   
 $+ \frac{1}{15} + \frac{1}{17} - \frac{1}{10} + \frac{1}{19} + \frac{1}{21} - \frac{1}{12} + \frac{1}{23} + \dots$
- $S_1 = 1.0000. S_2 = 0.5000. S_3 = 0.8333. S_4 = 1.0333. S_5 = 0.7833. S_6 = 0.9262. S_7 = 1.0373. S_8 = 0.8706. S_9 = 0.9615. S_{10} = 1.0385. S_{11} = 0.9135. S_{12} = 0.9801. S_{13} = 1.0390. S_{14} = 0.9390. S_{15} = 0.9916. S_{16} = 1.0392. S_{17} = 0.9559. S_{18} = 0.9994.$  In fact,  
 $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} = \ln 2.$

78 / 78