

VIET NAM GENERAL CONFEDERATION OF LABOR

TON DUC THANG UNIVERSITY

FACULTY OF INFORMATION TECHNOLOGY



NGUYỄN QUANG HUY - 523H0140

NGUYỄN THANH TÙNG - 523H0192

FINAL REPORT

APPLIED CALCULUS FOR IT

HO CHI MINH CITY, 2024

VIET NAM GENERAL CONFEDERATION OF LABOR

TON DUC THANG UNIVERSITY

FACULTY OF INFORMATION TECHNOLOGY



NGUYỄN QUANG HUY - 523H0140

NGUYỄN THANH TÙNG - 523H0192

FINAL REPORT

APPLIED CALCULUS FOR IT

Instructor

Master of Science Phạm Kim Thủy

HO CHI MINH CITY, 2024

ACKNOWLEDGEMENT

I wish to show our appreciation to Ton Duc Thang University, Faculty of Information Technology, Division of Computer Science and every Lecture, Professor of the faculty or creating a positive and excellent study environment for us to indulge and prosper. We appreciate the insurmountable effort of MSc. Phạm Kim Thủy. At the end, we wish for the opportunities and experiences that have been given to us and proud to be studying at Ton Duc Thang University.

Ho Chi Minh city, January 4, 2024

Author

(Sign and write your full name)

THIS FINAL REPORT HAD BEEN COMPLETED AT TON DUC THANG UNIVERSITY

We ensure that this is our own final report with scientific guideline of MSc. Phạm Kim Thủy. The final report's contents, results is honest and never publish under any circumstances. All data in the charts serve any analytical, critique and evaluation purposes have been collected by the authors from different sources which is written down in the reference materials. Furthermore, project also uses some feedbacks, reviews as well as statistics of other authors and agencies all have citing and origins.

If anything fraud detected on the our project, we willing take responsibilities about the content. Ton Duc Thang University does not relate to any copyrights infringement cause by the authors in the process (if there any).

Ho Chi Minh city, January 4, 2024

Author

(Sign and write your full name)

TABLE OF CONTENTS

LIST OF ABBREVIATIONS.....	vii
CHAPTER 1: SOLUTIONS.....	1
1.1 Question 1	1
<i>1.1.1 Topic.....</i>	<i>1</i>
<i>1.1.2 Solution.....</i>	<i>1</i>
<i>1.1.3 Final result</i>	<i>1</i>
1.2 Question 2	3
<i>1.2.1 Topic.....</i>	<i>3</i>
<i>1.2.2 Solution.....</i>	<i>3</i>
<i>1.2.3 Final result</i>	<i>3</i>
1.3 Question 3	5
<i>1.3.1 Topic.....</i>	<i>5</i>
<i>1.3.2 Final result</i>	<i>5</i>
1.4 Question 4	6
<i>1.4.1 Topic.....</i>	<i>6</i>
<i>1.4.2 Solution.....</i>	<i>6</i>
<i>1.4.3 Final result</i>	<i>7</i>
1.5 Question 5	7
<i>1.5.1 Topic.....</i>	<i>7</i>
<i>1.5.2 Solution.....</i>	<i>7</i>
<i>1.5.3 Final result</i>	<i>8</i>
1.6 Question 6	9
<i>1.6.1 Topic.....</i>	<i>9</i>

1.6.2 Solution.....	10
1.6.3 Final result	11
1.7 Question 7	11
1.7.1 Topic.....	11
1.7.2 Final result	11
1.8 Question 8	12
1.8.1 Topic.....	12
1.8.2 Solution.....	12
1.8.3 Final result	13
1.9 Question 9	13
1.9.1 Topic.....	13
1.9.2 Solution.....	13
1.9.3 Final result	14
BIBLIOGRAPHY	15

LIST OF ABBREVIATIONS

MSc	Master of Science
M.A	Master of Art
Pcs	Pieces

CHAPTER 1: SOLUTIONS

1.1 Question 1

1.1.1 Topic

Tell whether the following functions are even, odd, or neither. Give reasons for your answer.

- $f(x) = x^2 + x$
- $f(x) = x^3 + x$
- $f(x) = \frac{4}{x^4 - 4}$
- $f(x) = \frac{x^3}{x^4 - 4}$

1.1.2. Solution

- To solve the problem of determining whether a function is even, odd or not we do the following:
- Find the domain D of the function.
- Based on the definition of even and odd functions as follows:
 - A function $f(x)$ is considered even if, for every x in its domain:

$$f(x) = -f(x)$$

- A function $f(x)$ is considered odd if, for every x in its domain

$$f(x) = -f(-x) \quad \text{or} \quad -f(x) = f(-x)$$

1.1.3. Final result

- $f(x) = x^2 + x$
 - The domain of the function $f(x): D = \mathbb{R}$
 - For every $x \in \mathbb{R} \rightarrow -x \in \mathbb{R}$, we have:

$$f(-x) = (-x)^2 + (-x) = x^2 - x \neq f(x)$$

- Because of $f(-x) \neq -f(x)$, we can conclude that the **function** $f(x) = x^2 + x$ **does not have parity**.

- $f(x) = x^3 + x$

- The domain of the function $f(x): D = \mathbb{R}$
- For every $x \in \mathbb{R} \rightarrow -x \in \mathbb{R}$, we have:

$$f(-x) = (-x)^3 + (-x) = -x^3 - x = -(x^3 + x) = -f(x)$$

- Therefore, we can conclude that the **function** $f(x) = x^3 + x$ **is odd**.

- $f(x) = \frac{4}{x^4 - 4}$

- The domain of the function $f(x)$:

$$x^4 - 4 \neq 0$$

$$\rightarrow \begin{cases} x \neq \sqrt{2} \\ x \neq -\sqrt{2} \end{cases} \rightarrow D = \mathbb{R} \setminus \{-\sqrt{2}; \sqrt{2}\}$$

- For every $x \in \mathbb{R} \setminus \{-\sqrt{2}; \sqrt{2}\} \rightarrow -x \in \mathbb{R} \setminus \{-\sqrt{2}; \sqrt{2}\}$, we have:

$$f(-x) = \frac{4}{(-x)^4 - 4} = \frac{4}{x^4 - 4} = f(x)$$

- Therefore, we can conclude that **the function** $f(x) = \frac{4}{x^4 - 4}$ **is even**.

- $f(x) = \frac{x^3}{x^4 - 4}$

- The domain of the function $f(x)$:

$$x^4 - 4 \neq 0$$

$$\rightarrow \begin{cases} x \neq \sqrt{2} \\ x \neq -\sqrt{2} \end{cases} \rightarrow D = \mathbb{R} \setminus \{-\sqrt{2}; \sqrt{2}\}$$

- For every $x \in \mathbb{R} \setminus \{-\sqrt{2}; \sqrt{2}\} \rightarrow -x \in \mathbb{R} \setminus \{-\sqrt{2}; \sqrt{2}\}$, we have:

$$f(-x) = \frac{(-x)^3}{(-x)^4 - 4} = -\frac{x^3}{x^4 - 4} = -f(x)$$

- Finally, we can conclude that **the function** $f(x) = \frac{x^3}{x^4 - 4}$ **is odd**.

1.2 Question 2

1.2.1 Topic

Find the following limit $\lim_{x \rightarrow 5} \frac{555}{x^2 - 25}$ as:

- $x \rightarrow 5^+$
- $x \rightarrow 5^-$
- $x \rightarrow -5^+$
- $x \rightarrow -5^-$

1.2.2 Solution

- Simplify the given function.
- Because when substituting the limit value of x into the function, the function becomes undefined, so consider the numerator and denominator separately to determine the limit:
 - The numerator is a constant.
 - The denominator is a polynomial that contains x substitutions for each case to find the limit of the denominator.
- Conclusion, with the numerator being a constant:
 - The denominator has a limit approaching positive tense \rightarrow given expression has $\lim = +\infty$
 - The denominator has a limit approaching negative tense \rightarrow given expression has $\lim = -\infty$

1.2.3 Final result

- $\lim_{x \rightarrow 5^+} \frac{555}{x^2 - 25}$

Consider:

- The numerator: 555 is a constant.
- The denominator: $\lim_{x \rightarrow 5^+} (x + 5)(x - 5) = 0^+$ at $x \rightarrow 5^+$.

$$\Rightarrow \lim_{x \rightarrow 5^+} \frac{555}{(x-5)(x+5)} = +\infty$$

$$\bullet \lim_{x \rightarrow 5^-} \frac{555}{x^2 - 25}$$

Consider:

- The numerator: 555 is a constant.
- The denominator: $\lim(x+5)(x-5) = 0^-$ at $x \rightarrow 5^-$.

$$\Rightarrow \lim_{x \rightarrow 5^-} \frac{555}{(x-5)(x+5)} = -\infty$$

$$\bullet \lim_{x \rightarrow -5^+} \frac{555}{x^2 - 25}$$

Consider:

- The numerator: 555 is a constant.
- The denominator: $\lim(x+5)(x-5) = 0^+$ at $x \rightarrow -5^+$.

$$\Rightarrow \lim_{x \rightarrow -5^+} \frac{555}{(x-5)(x+5)} = +\infty$$

$$\bullet \lim_{x \rightarrow -5^-} \frac{555}{x^2 - 25}$$

Consider:

- The numerator: 555 is a constant.
- The denominator: $\lim(x+5)(x-5) = 0^-$ at $x \rightarrow -5^-$.

$$\Rightarrow \lim_{x \rightarrow -5^-} \frac{555}{(x-5)(x+5)} = -\infty$$

1.3 Question 3

1.3.1 Topic

Find the derivatives $\frac{dy}{dx}$ of the following functions:

- $y = \frac{\sqrt{x} - 4}{\sqrt{x} + 4}$

- $y = \left(\frac{\sqrt{x}}{10} - 1 \right)^{-10}$

1.3.2 Final result.

- $y = \frac{\sqrt{x} - 4}{\sqrt{x} + 4}$

$$\begin{aligned}
 \frac{dy}{dx} &= \left(\frac{\sqrt{x} - 4}{\sqrt{x} + 4} \right)' \\
 &= \frac{(\sqrt{x} - 4)'(\sqrt{x} + 4) - (\sqrt{x} - 4)(\sqrt{x} + 4)'}{(\sqrt{x} + 4)^2} \\
 &= \frac{\frac{\sqrt{x} + 4}{2\sqrt{x}} - \frac{\sqrt{x} - 4}{2\sqrt{x}}}{(\sqrt{x} + 4)^2} \\
 &= \frac{\frac{8}{2\sqrt{x}}}{(\sqrt{x} + 4)^2} \\
 &= \frac{4}{\sqrt{x}(\sqrt{x} + 4)^2}
 \end{aligned}$$

- $y = \left(\frac{\sqrt{x}}{10} - 1 \right)^{-10}$

$$\begin{aligned}
 \frac{dy}{dx} &= \left[\left(\frac{\sqrt{x}}{10} - 1 \right)^{-10} \right]' = \left[\left(\frac{\sqrt{x} - 10}{10} \right)^{-10} \right]' \\
 &= -10 \cdot \left(\frac{\sqrt{x} - 10}{10} \right)^{-11} \cdot \left(\frac{\sqrt{x} - 10}{10} \right)' \\
 &= -10 \cdot \left(\frac{\sqrt{x} - 10}{10} \right)^{-11} \cdot \left(\frac{10}{2\sqrt{x}} \right) \\
 &= -10 \cdot \left(\frac{\sqrt{x} - 10}{10} \right)^{-11} \cdot \left(\frac{1}{20\sqrt{x}} \right) \\
 &= \left(\frac{-1}{2\sqrt{x}} \right) \cdot \left(\frac{\sqrt{x} - 10}{10} \right)^{-11}
 \end{aligned}$$

1.4 Question 4

1.4.1 Topic

Find an equation of the tangent line to the graph of $y = 1 + 2e^x$ at the point where $x = 0$.

1.4.2 Solution

- Based on the tangent equation: $y - y_0 = f'(x_0)(x - x_0)$ to do exercise 4.
- Find the coordinates of the point with $x = 0$ by substituting x into the function to find y .
- Apply the derivative formula to calculate the derivative y' .
- Calculate the value of y' at $x = 0$.
- Set to $y - y_0 = f'(x_0)(x - x_0)$ the formula to find the tangent equation.

1.4.3 Final result

- Because the tangent line to the graph of $y = 1 + 2e^x$ goes through the graph at the point $A(0, y_0)$, so the point A belongs to the graph.

$$y(0) = 1 + 2e^0 = 1 + 2 = 3$$

- So the coordinates of point A : $A(0,3)$.
- We have: $y'(x) = (1 + 2e^x)' = 2e^x$

$$\rightarrow y'(x_0) = y'(0) = 2e^0 = 2 \cdot 1 = 2$$

- Tangent equation of the graph of a function $y = 1 + 2e^x$:

$$y - y_0 = y'(x_0)(x - x_0)$$

$$\rightarrow y - 3 = 2(x - 0)$$

$$\rightarrow y - 3 = 2x$$

$$\rightarrow y = 2x + 3$$

- So we can conclude that the equation of the tangent to the function $y = 1 + 2e^x$ at the point $x = 0$ is: $y = 2x + 3$.

1.5 Question 5

1.5.1 Topic

Given the derivative $f'(x) = (\sin x + \cos x)(\sin x - \cos x)$, $0 \leq x \leq 2\pi$.

- What are the critical numbers of f ?
- On what open intervals is f increasing or decreasing?
- At what points, if any, does f assume local maximum and minimum values?

1.5.2 Solution

- Give the function $f'(x) = 0$ to find the critical number.
- Draw a table of variations to determine whether the function increases or decreases at the intervals found when given: $f'(x) = 0$
- Identify Local Maxima and Minima Points:

- Local Maximum: Points where the function transitions from decreasing to increasing. This occurs when the derivative changes sign from positive to negative.
- Local Minimum: Points where the function transitions from increasing to decreasing. This occurs when the derivative changes sign from negative to positive.

1.5.3 Final result

- What are the critical numbers of f ?

$$\rightarrow (\sin x + \cos x)(\sin x - \cos x) = 0$$

$$\rightarrow \begin{cases} \sin x + \cos x = 0 \\ \sin x - \cos x = 0 \end{cases}$$

$$\rightarrow \begin{cases} \sin x = -\cos x \\ \sin x = \cos x \end{cases}$$

$$\rightarrow \begin{cases} \tan x = -1 \\ \tan x = 1 \end{cases}$$

$$\rightarrow \begin{cases} \left\{ \begin{array}{l} x = \frac{3\pi}{4} \\ x = \frac{7\pi}{4} \end{array} \right. & (0 \leq x \leq 2\pi) \\ \left\{ \begin{array}{l} x = \frac{\pi}{4} \\ x = \frac{5\pi}{4} \end{array} \right. & (0 \leq x \leq 2\pi) \end{cases}$$

- So the critical number of the function $f(x)$ is $\left(\frac{3\pi}{4}; \frac{7\pi}{4}; \frac{\pi}{4}; \frac{5\pi}{4}\right)$ suitable for the given condition $0 \leq x \leq 2\pi$.
- And we have the following variation table:

x	$+\infty$	0	$\frac{\pi}{4}$	$\frac{3\pi}{4}$	$\frac{5\pi}{4}$	$\frac{7\pi}{4}$	2π	$+\infty$
$f'(x)$	<div>- 0 + 0 - 0 + 0 -</div>							
$f(x)$	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>							

- On what open intervals is f increasing or decreasing?
 - According to the variation table above, we can see that:
 - The function decreases at $\left(0, \frac{\pi}{4}\right)$, $\left(\frac{3\pi}{4}, \frac{5\pi}{4}\right)$, $\left(\frac{7\pi}{4}, 2\pi\right)$
 - The function increases at $\left(\frac{\pi}{4}, \frac{3\pi}{4}\right)$ and $\left(\frac{5\pi}{4}, \frac{7\pi}{4}\right)$
- At what points, if any, does f assume local maximum and minimum values?
 - At $x = \frac{\pi}{4}$, $x = \frac{5\pi}{4}$, We see the value of $f(x)$ from increasing to decreasing or the derivative value changing sign from negative to positive.
 - The value of $f(x)$ reaches its local minimum value at $x = \frac{\pi}{4}$, $x = \frac{5\pi}{4}$
 - At $x = \frac{3\pi}{4}$, $x = \frac{7\pi}{4}$, We see the value $f(x)$ from decreasing to increasing or the derivative value $f'(x)$ changing sign from positive to negative.
 - The value of $f(x)$ reaches its local maximum value at $x = \frac{3\pi}{4}$, $x = \frac{7\pi}{4}$

1.6 Question 6

1.6.1 Topic

Find all curves through a point where $x = 1$ whose arc length is the following L value:

$$L = \int_1^5 \sqrt{1 + \frac{1}{x^2}} dx$$

1.6.2 Solution

- According to the bibliography Applied Calculus for IT, wrote by M.A. Phạm Kim Thủy, Division of Computer Science, TON DUC THANG University, we have an equation to measure the arc length of the curve $y = f(x)$ from $x = a$ to $x = b$ is

$$L = \int_a^b \sqrt{1 + (f'(x))^2} dx$$

- So that we have

$$L = \int_a^b \sqrt{1 + (f'(x))^2} dx = \int_1^5 \sqrt{1 + \frac{1}{x^2}} dx$$

$$\rightarrow \sqrt{1 + (f'(x))^2} = \sqrt{1 + \frac{1}{x^2}} \quad \left((f'(x))^2 \geq 0, \frac{1}{x^2} \geq 0 \right) \quad \text{Square both sides}$$

$$\rightarrow 1 + (f'(x))^2 = 1 + \frac{1}{x^2} \quad \text{Subtract both side 1}$$

$$\rightarrow (f'(x))^2 = \frac{1}{x^2}$$

$$\rightarrow \begin{cases} f'(x) = \frac{1}{x} \\ f'(x) = \frac{-1}{x} \end{cases}$$

$$\rightarrow \begin{cases} \int f'(x) dx = \int \frac{1}{x} dx \\ \int f'(x) dx = -\int \frac{1}{x} dx \end{cases} \quad \text{Implication both sides}$$

$$\rightarrow \begin{cases} f(x) = \ln|x| + C_1 \\ f(x) = -\ln|x| + C_2 \end{cases}$$

- Given that the curves pass through the point where $x = 1$, we can find the constants C_1 and C_2 by substituting $x = 1$ and $y = y_0$:

$$y_0 = \ln|1| + C_1 \rightarrow C_1 = y_0$$

$$y_0 = -\ln|1| + C_2 \rightarrow C_2 = y_0$$

1.6.3 Final result:

So all the curves that pass through the point $(1, y_0)$ and satisfy the given arc length:

$$y = \ln|x| + y_0 \text{ and } y = -\ln|x| + y_0$$

1.7 Question 7

1.7.1 Topic

Given that $a_1, a_2, a_3, \dots, a_n, \dots$ are real numbers fulfilling the following conditions:

- $a_n > 0, n \in \mathbb{Z}^+$
- $a_1 \geq a_2 \geq a_3 \geq \dots \geq a_n \geq \dots$
- The series $a_2 + a_4 + a_8 + a_{16} + \dots + a_{2^n} + \dots$ diverges.

Determine the convergence or divergence of the following series. Explain.

$$a_1 + \frac{a_2}{2} + \frac{a_3}{3} + \frac{a_4}{4} \dots + \frac{a_n}{n} + \dots$$

1.7.2 Final result

- We have the divergence series S:

$$\sum_{n=1}^{\infty} a_{2^n} = a_2 + a_4 + a_8 + \dots + a_{2^n} = S$$

- Suppose that we have the T series following:

$$a_1 + \frac{a_2}{2} + \frac{a_3}{3} + \frac{a_4}{4} \dots + \frac{a_n}{n} + \dots$$

- Apply the comparison test:

$$0 < a_{2^n} \leq a_n$$

- Also we know that $\sum_{n=1}^{\infty} a_{2^n}$ is a divergence series.
- So $\sum_{n=1}^{\infty} a_n$ is a divergence series either.
- At the same time, with $a_n > 0$ so that $\sum_{n=1}^{\infty} \frac{a_n}{n}$ is also a divergence series.

1.8 Question 8

1.8.1 Topic

Find all values of x such that the following series is absolutely convergent:

$$\sum_{n=1}^{\infty} \frac{nx^n}{(n+1)(2x+1)^n}$$

1.8.2 Solution

- *Apply the ratio test:*

$$\begin{aligned} \lim_{x \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \lim_{x \rightarrow \infty} \left| \frac{(n+1)x^{n+1}}{(n+2)(2x+1)} \cdot \frac{(n+1)(2x+1)}{nx^n} \right| = \lim_{x \rightarrow \infty} \left| \frac{(n+1)^2 x}{(n+2)n} \right| \\ &= \lim_{x \rightarrow \infty} \left| \frac{n^2 + 2nx + x}{n^2 + 2n} \right| \stackrel{L'H}{=} \lim_{x \rightarrow \infty} \left| \frac{2n + 2x}{2n + 2} \right| \stackrel{L'H}{=} \lim_{x \rightarrow \infty} \left| \frac{2x}{2} \right| = \lim_{x \rightarrow \infty} |x| = L \end{aligned}$$

- So the above series is absolutely convergent, if:

$$0 \leq L < 1 \rightarrow 0 \leq |x| < 1 \rightarrow -1 < x < 1$$

1.8.3 Final result

So with every $x \in (-1, 1)$, $\sum_{n=1}^{\infty} \frac{nx^n}{(n+1)(2x+1)^n}$ will be absolutely convergent.

1.9 Question 9

1.9.1 Topic

One thousand earphones sell for \$55 each, resulting in a revenue of $(1000) (\$55) = \$55,000$. For each \$5 increase in the price, 20 fewer earphones are sold. For ex., if the price of each earphone is \$60, there will be 980 $(1000 - 20)$ earphones sold; if the price of each earphone is \$65, there will be 960 $(1000 - 20 - 20)$ earphones sold; so on. Find the revenue in case the price of each earphone is \$255

1.9.2 Solution

- We call T is the revenue (units \$).
- We call S is the total earphones sold (units pcs).
- We call G is the original cost of each earphones (units \$).
- We call N is the topic requires cost of each earphones (units \$).
- From the topic, we know that: If we increase \$5 in price for each earphone, 20 fewer earphones are sold, so we have the following equation:

$$S = 1000 - \frac{N - G}{5} \times 20 \quad (pcs)$$

- So we can find the total earphones sold by the above equation:

$$S = 1000 - \frac{225 - 55}{5} \times 20 = 200 \quad (pcs)$$

- Finally, find the revenue if we sell \$225 each earphones:

$$T = S \times N = 200 \times 255 = \$51000$$

1.9.3 Final result

If we sold 200 piece earphones by \$225 for each, we can find the revenue is \$51000.

BIBLIOGRAPHY

English

[1]. Wang Fei, MA1521 Calculus for computing.

[2]. M.A. Phạm Kim Thủy, Division of Computer Science, TON DUC
THANG University, Applied Calculus for IT - Course