

Introductory Course on Non-smooth Optimisation

Lecture 04

Backward-Backward splitting

Outline

- 1 Forward–Backward splitting revisit
- 2 MAP continue
- 3 Backward–Backward splitting
- 4 Numerical experiments

Monotone inclusion problem

Problem

Let $B : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be β -cocoercive for some $\beta > 0$, $s > 1$ be a positive integer, such that for each $i \in \{1, \dots, s\}$: $A_i : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is maximal monotone. Consider the problem

$$\text{Find } x \in \mathbb{R}^n \text{ such that } 0 \in B(x) + \sum_{i=1}^s A_i(x).$$

- A_i can be composed with linear mapping, e.g. $L^* \circ A \circ L$
- Even if the resolvent of B and each A_i is simple, the resolvent of $B + \sum_i A_i$ in most cases is not solvable
- Use the properties of operators and structure of problem to derive operator splitting schemes

Outline

- 1 Forward-Backward splitting revisit
- 2 MAP continue
- 3 Backward-Backward splitting
- 4 Numerical experiments

Monotone inclusion problem

Monotone inclusion

Find $x \in \mathbb{R}^n$ such that $0 \in A(x) + B(x)$

Assumptions:

- $A : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is maximal monotone
- $B : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is β -cocoersive
- $\text{zer}(A + B) \neq \emptyset$

Characterisation of solution: $\gamma > 0$

$$x^* - \gamma B(x^*) \in x^* + \gamma A(x^*) \quad \Leftrightarrow \quad x^* = \mathcal{J}_{\gamma A} \circ (\text{Id} - \gamma B)(x^*).$$

Example:

$$\min_{x \in \mathbb{R}^n} R(x) + F(x),$$

with $R \in \Gamma_0$ and $F \in C_L^1$.

Forward-Backward splitting

Fixed-point operator: $\gamma \in]0, 2\beta[$

$$\mathcal{T}_{\text{FB}} = \mathcal{J}_{\gamma A} \circ (\text{Id} - \gamma B)$$

- $\mathcal{J}_{\gamma A}$ is firmly non-expansive
- $\text{Id} - \gamma B$ is $\frac{\gamma}{2\beta}$ -averaged non-expansive
- \mathcal{T}_{FB} is $\frac{2\beta}{4\beta - \gamma}$ -averaged non-expansive
- $\text{fix}(\mathcal{T}_{\text{FB}}) = \text{zer}(A + B)$

Forward-Backward splitting

Let $\gamma \in]0, 2\beta[, \lambda_k \in [0, \frac{4\beta - \gamma}{2\beta}]$:

$$x_{k+1} = (1 - \lambda_k)x_k + \lambda_k \mathcal{T}_{\text{FB}}(x_k)$$

- Special case of Krasnosel'skiĭ-Mann iteration
- Recovers proximal point algorithm when $B = 0$

Outline

- 1 Forward-Backward splitting revisit
- 2 MAP continue**
- 3 Backward-Backward splitting
- 4 Numerical experiments

Method of alternating projection

Let $\mathcal{X}, \mathcal{Y} \subseteq \mathbb{R}^n$ be closed convex and non-empty, such that $\mathcal{X} \cap \mathcal{Y} \neq \emptyset$

$$\min_{x \in \mathbb{R}^n} \iota_{\mathcal{X}}(x) + \iota_{\mathcal{Y}}(x).$$

Method of alternating projection (MAP)

Let $x_0 \in \mathcal{X}$:

$$y_{k+1} = \text{proj}_{\mathcal{Y}}(x_k)$$

$$x_{k+1} = \text{proj}_{\mathcal{X}}(y_{k+1})$$

Fixed-point operator: $x_{k+1} = \mathcal{T}_{\text{MAP}}(x_k)$,

$$\mathcal{T}_{\text{MAP}} \stackrel{\text{def}}{=} \text{proj}_{\mathcal{X}} \circ \text{proj}_{\mathcal{Y}}$$

- $\text{proj}_{\mathcal{X}}, \text{proj}_{\mathcal{Y}}$ are firmly non-expansive
- \mathcal{T}_{MAP} is $\frac{2}{3}$ -averaged non-expansive
- $\text{fix}(\mathcal{T}_{\text{MAP}}) = \mathcal{X} \cap \mathcal{Y}$

Derive MAP

- Feasibility problem is equivalent to

$$\min_{x, y \in \mathbb{R}^n} \iota_{\mathcal{X}}(x) + \frac{1}{2} \|x - y\|^2 + \iota_{\mathcal{Y}}(y).$$

- Optimality condition

$$0 \in \mathcal{N}_{\mathcal{Y}}(y^*) + y^* - x^*$$

$$0 \in \mathcal{N}_{\mathcal{X}}(x^*) + x^* - y^*.$$

- Fixed-point characterisation

$$y^* = \text{proj}_{\mathcal{Y}}(x^*)$$

$$x^* = \text{proj}_{\mathcal{X}}(y^*).$$

- Fixed-point iteration

$$y_{k+1} = \text{proj}_{\mathcal{Y}}(x_k)$$

$$x_{k+1} = \text{proj}_{\mathcal{X}}(y_{k+1}).$$

Example: SDP feasibility

SDP feasibility

Find $X \in \mathcal{S}^n$ such that

$$X \succeq 0 \quad \text{and} \quad \text{Tr}(A_i X) = b_i, \quad i = 1, \dots, m.$$

Two sets and projection:

- $\mathcal{X} = \mathcal{S}_+^n$ is the positive semidefinite cone. Let $Y_k = \sum_{i=1}^n \sigma_i u_i u_i^T$ be the eigenvalue decomposition of Y_k , then

$$\text{proj}_{\mathcal{X}}(Y_k) = \sum_{i=1}^n \max\{0, \sigma_i\} u_i u_i^T.$$

- \mathcal{Y} is the affine set in \mathcal{S}^n define by the linear inequalities,

$$\text{proj}_{\mathcal{Y}}(X_k) = X_k - \sum_{i=1}^m u_i A_i,$$

where u_i are found from the normal equations

$$Gu = (\text{Tr}(A_1 X_k) - b_1, \dots, \text{Tr}(A_m X_k) - b_m), \quad G_{ij} = \text{Tr}(A_i A_j).$$

Convergence rate

Let \mathcal{X}, \mathcal{Y} be two subspaces, and assume

$$1 \leq p \stackrel{\text{def}}{=} \dim(\mathcal{X}) \leq q \stackrel{\text{def}}{=} \dim(\mathcal{Y}) \leq n - 1.$$

Principal angles The principal angles $\theta_k \in [0, \frac{\pi}{2}]$, $k = 1, \dots, p$ between \mathcal{X} and \mathcal{Y} are defined by, with $u_0 = v_0 \stackrel{\text{def}}{=} 0$, and

$$\begin{aligned} \cos(\theta_k) \stackrel{\text{def}}{=} \langle u_k, v_k \rangle = \max \langle u, v \rangle \quad \text{s.t.} \quad u \in \mathcal{X}, v \in \mathcal{Y}, \|u\| = 1, \|v\| = 1, \\ \langle u, u_i \rangle = \langle v, v_i \rangle = 0, i = 0, \dots, k-1. \end{aligned}$$

Friedrichs angle The Friedrichs angle $\theta_F \in]0, \frac{\pi}{2}]$ between \mathcal{X} and \mathcal{Y} is

$$\begin{aligned} \cos(\theta_F(\mathcal{X}, \mathcal{Y})) \stackrel{\text{def}}{=} \max \langle u, v \rangle \quad \text{s.t.} \quad u \in \mathcal{X} \cap (\mathcal{X} \cap \mathcal{Y})^\perp, \|u\| = 1, \\ v \in \mathcal{Y} \cap (\mathcal{X} \cap \mathcal{Y})^\perp, \|v\| = 1. \end{aligned}$$

Lemma

The Friedrichs angle is θ_{d+1} where $d \stackrel{\text{def}}{=} \dim(\mathcal{X} \cap \mathcal{Y})$. Moreover, $\theta_F(\mathcal{X}, \mathcal{Y}) > 0$.


Convergence rate

Example \mathcal{X}, \mathcal{Y} are defined by

$$\mathcal{X} = \{x : Ax = 0\}, \quad \mathcal{Y} = \{x : Bx = 0\}$$

Projection onto subspace

$$\text{proj}_{\mathcal{X}}(x) = x - A^T(AA^T)^{-1}Ax$$



Convergence rate

- Define diagonal matrices

$$\mathbf{c} = \text{diag}(\cos(\theta_1), \dots, \cos(\theta_p))$$

$$\mathbf{s} = \text{diag}(\sin(\theta_1), \dots, \sin(\theta_p))$$

- Suppose $p + q < n$, then there exists orthogonal matrix U such that

$$\text{proj}_{\mathcal{X}} = U \left[\begin{array}{cc|cc} \text{Id}_p & 0 & 0 & 0 \\ 0 & 0_p & 0 & 0 \\ \hline 0 & 0 & 0_{q-p} & 0 \\ 0 & 0 & 0 & 0_{n-p-q} \end{array} \right] U^*$$

$$\text{proj}_{\mathcal{Y}} = U \left[\begin{array}{cc|cc} c^2 & cs & 0 & 0 \\ cs & c^2 & 0 & 0 \\ \hline 0 & 0 & \text{Id}_{q-p} & 0 \\ 0 & 0 & 0 & 0_{n-p-q} \end{array} \right] U^*,$$

Convergence rate

- Fixed-point operator

$$\begin{aligned}\mathcal{T}_{\text{MAP}} &= \text{proj}_{\mathcal{X}} \circ \text{proj}_{\mathcal{Y}} \\ &= U \left[\begin{array}{cc|cc} c^2 & cs & 0 & 0 \\ 0 & 0_p & 0 & 0 \\ \hline 0 & 0 & 0_{q-p} & 0 \\ 0 & 0 & 0 & 0_{n-p-q} \end{array} \right] U^*\end{aligned}$$

- Consider relaxation

$$\begin{aligned}\mathcal{T}_{\text{MAP}}^\lambda &= (1 - \lambda)\text{Id} + \lambda\mathcal{T}_{\text{MAP}} \\ &= U \left[\begin{array}{cc|cc} (1 - \lambda)\text{Id}_p + \lambda c^2 & \lambda cs & 0 & 0 \\ 0 & (1 - \lambda)\text{Id}_p & 0 & 0 \\ \hline 0 & 0 & (1 - \lambda)\text{Id}_{n-2p} & 0 \end{array} \right] U^*\end{aligned}$$

Convergence rate

- Eigenvalues

$$\sigma(\mathcal{T}_{\text{MAP}}^\lambda) = \{1 - \lambda \sin^2(\theta_i) | i = 1, \dots, p\} \cup \{1 - \lambda\}$$

- Spectral radius

$$\rho(\mathcal{T}_{\text{MAP}}^\lambda) = \max \{1 - \lambda \sin^2(\theta_F), |1 - \lambda|\}$$

- No relaxation

$$\rho(\mathcal{T}_{\text{MAP}}) = \cos^2(\theta_F)$$

- Convergence rate, $C > 0$ is some constant


$$\begin{aligned} \|x_k - x^*\| &= \|\mathcal{T}_{\text{MAP}} x_{k-1} - \mathcal{T}_{\text{MAP}} x^*\| \\ &= \dots \\ &= \|\mathcal{T}_{\text{MAP}}^k (x_0 - x^*)\| \\ &\leq C \|\mathcal{T}_{\text{MAP}}\|^k \|x_0 - x^*\| \end{aligned}$$

Outline

- 1 Forward-Backward splitting revisit
- 2 MAP continue
- 3 Backward-Backward splitting**
- 4 Numerical experiments

Best pair problem

When the problem is feasible, MAP returns $x_k, y_k \rightarrow x^* \in \mathcal{X} \cap \mathcal{Y}$.



Best pair problem

Let $\mathcal{X}, \mathcal{Y} \subseteq \mathbb{R}^n$ be closed and convex, such that

$$\mathcal{X} \cap \mathcal{Y} = \emptyset.$$

Consider finding two points in \mathcal{X} and \mathcal{Y} such that they are the closest, that is

$$\min_{x \in \mathcal{X}, y \in \mathcal{Y}} \|x - y\|.$$

- MAP can be applied and

$$(x_k, y_k) \rightarrow (x^*, y^*)$$

where (x^*, y^*) is a best pair.

- When $\mathcal{X} \cap \mathcal{Y} \neq \emptyset$, $x^* = y^*$.

Backward-Backward splitting

Consider

Find $x, y \in \mathbb{R}^n$ such that $0 \in A(x) + B(y)$,

where

- $A, B : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ are maximal monotone
- The set of solution is non-empty

There exists $x^*, y^* \in \mathbb{R}^n$ and $\gamma > 0$ such that

$$y^* - x^* \in \gamma A(x^*)$$

$$x^* - y^* \in \gamma B(y^*).$$

Backward-Backward splitting

Let $x_0 \in \mathbb{R}^n$, $\gamma > 0$:

$$y_{k+1} = \mathcal{J}_{\gamma B}(x_k)$$

$$x_{k+1} = \mathcal{J}_{\gamma A}(y_{k+1})$$

Regularised monotone inclusion

Yosida approximation

$$\gamma A = \frac{1}{\gamma}(\text{Id} - \mathcal{J}_{\gamma A})$$

which is γ -cocoercive.

Regularised monotone inclusion

Find $x \in \mathbb{R}^n$ such that $0 \in A(x) + \gamma B(x)$

Forward–Backward splitting $\tau \in]0, 2\gamma]$

$$x_{k+1} = \mathcal{J}_{\tau A} \circ (\text{Id} - \tau^\gamma B)(x_k)$$

BB as special case of FB let $\tau = \gamma$

$$\begin{aligned} x_{k+1} &= \mathcal{J}_{\gamma A} \circ (\text{Id} - \gamma^\gamma B)(x_k) \\ &= \mathcal{J}_{\gamma A} \circ (\text{Id} - \gamma \frac{1}{\gamma} (\text{Id} - \mathcal{J}_{\gamma B}))(x_k) \\ &= \mathcal{J}_{\gamma A} \circ \mathcal{J}_{\gamma B}(x_k) \end{aligned}$$

Inertial BB splitting

An inertial Backward-Backward splitting

Initial: $x_0 \in \mathbb{R}^n$, $x_{-1} = x_0$ and $\gamma > 0$, $\tau \in]0, 2\gamma]$;

$$\begin{aligned} y_k &= x_k + a_{0,k}(x_k - x_{k-1}) + a_{1,k}(x_{k-1} - x_{k-2}) + \cdots, \\ x_{k+1} &= \mathcal{J}_{\gamma A} \circ \mathcal{J}_{\gamma B}(y_k), \lambda_k \in [0, 1]. \end{aligned}$$

An inertial BB splitting based on Yosida approximation

Initial: $x_0 \in \mathbb{R}^n$, $x_{-1} = x_0$ and $\gamma > 0$;

$$\begin{aligned} y_k &= x_k + a_{0,k}(x_k - x_{k-1}) + a_{1,k}(x_{k-1} - x_{k-2}) + \cdots, \\ z_k &= x_k + b_{0,k}(x_k - x_{k-1}) + b_{1,k}(x_{k-1} - x_{k-2}) + \cdots, \\ x_{k+1} &= \mathcal{J}_{\tau A} \circ (y_k - \tau^\gamma B(z_k)), \lambda_k \in [0, 1]. \end{aligned}$$


Outline

- 1 Forward-Backward splitting revisit
- 2 MAP continue
- 3 Backward-Backward splitting
- 4 Numerical experiments**

Numerical experiment

Feasibility problem for two subspaces:

$$A = [-4/5, 1] \quad \text{and} \quad B = [-1/5, 1]$$



Reference

- S. Boyd. Alternating projection, lecture notes
- H. H. Bauschke, J. Y. Bello Cruz, T. T. A. Nghia, H. M. Pha, and X. Wang. Optimal rates of linear convergence of relaxed alternating projections and generalized Douglas–Rachford methods for two subspaces. *Numerical Algorithms*, 73(1):33–76, 2016.