
a Overall Parameter Influence (Random Forest on SSE_w)
μ_{TP53} (TP53 loss-of-function)
θ_{D2} (TUMORAL / clearance threshold)
ν (Tumoral cell division)
δ_{low} (Heterozygous accumulation)
β (BRCA1 second-hit)
δ_{high} (Biallelic accumulation)
θ_{D1} (PRIMER threshold)
d (Basal division)
κ_{BRCA1} (BRCA1 D2 weight)
A_{it} (Cell age drift)
38.1%
17.6%
13.1%
11.2%
7.8%
5.8%
2.3%
1.6%
1.4%
1.2%
0
10
20
30
40
Normalized Permutation Importance (%)
b Age-Specific Parameter Influence on Penetrance (PRCC)
Partial Rank Correlation Coefficient (PRCC)
0.8
0.6
0.4
0.2
0.0
−0.2
−0.4
−0.6
−0.8
30
40
50
60
70
80
Age (years)
β (BRCA1 second-hit)
μ_{TP53} (TP53 loss-of-function)
δ_{low} (Heterozygous accumulation)
δ_{high} (Biallelic accumulation)
θ_{D1} (PRIMER threshold)
θ_{D2} (TUMORAL / clearance threshold)
d (Basal division)
κ_{BRCA1} (BRCA1 D2 weight)
A_{it} (Cell age drift)
ν (Tumoral cell division)