

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/32969549>

# Paralongidorus iberis sp.n. and P. monegrensis sp.n. from Spain with a polytomous key to the species of the genus Paralongidorus Siddiqi, Hooper & Khan, 1963 (Nematoda : Longidorid...

Article · January 1997

Source: OAI

CITATIONS

21

READS

140

2 authors:



Miguel Escuer

Spanish National Research Council

57 PUBLICATIONS 1,294 CITATIONS

SEE PROFILE



Manuel Arias

Hospital Universitario Marques de Valdecilla

630 PUBLICATIONS 8,316 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



mTOR inhibitors in renal transplantation [View project](#)



Ecological risk of priority contaminants in an insular environment. Assessment of spatial-temporal patterns and impacts on soil biodiversity [View project](#)

## ***Paralongidorus iberis* sp.n. and *P. monegrensis* sp.n. from Spain with a polytomous key to the species of the genus *Paralongidorus* Siddiqi, Hooper & Khan, 1963 (Nematoda : Longidoridae)**

Miguel ESCUER and María ARIAS

*Departamento de Agroecología, CCMA, CSIC, Serrano 115 dpdo, 28006 Madrid, Spain.*

Accepted for publication 22 March 1996.

**Summary** – Species of the genus *Paralongidorus* were found for the first time in Spain near Serreta Negra (Huesca), in the North-West of the country. Two new species, *P. iberis* sp.n. and *P. monegrensis* sp. n., are described. *P. iberis* sp. n. is characterized by its medium size (4.4-4.9 mm), rounded set off lip region, stirrup-shaped amphidial pouch and conical dorsally convex tail with conical terminus. It is close to *P. lemoni* from which it differs in tail shape, presence of males, body length, position of guiding ring, and a and c ratios. *P. monegrensis* sp. n. is characterized by its large body size (7.5-12 mm), hemispherical expanded lip region, stirrup-shaped amphidial pouch, and conical dorsally convex tail with rounded terminus. It resembles *P. sandellus* and *P. xiphinemoides*. It differs from *P. sandellus* by its longer body and odontostyle and anterior position of guiding ring, and from *P. xiphinemoides* by presence of males, odontostyle length, position of guiding ring, and c' ratio. A polytomous key of the 70 species described in the genus is proposed and two new combinations are proposed, *P. sativus* (Soni & Nama, 1983) n. comb. and *P. wiesae* (Heyns, 1994) n. comb.

**Résumé** – *Paralongidorus iberis* sp. n. et *P. monegrensis* sp. n. provenant d'Espagne et clé polytomique des espèces du genre *Paralongidorus* Siddiqi, Hooper & Khan, 1963 (Nematoda : Longidoridae) – Des espèces du genre *Paralongidorus* ont été trouvées pour la première fois en Espagne, près de Serreta Negra (Huesca) dans la région nord-ouest du pays. Deux nouvelles espèces, *P. iberis* sp. n. et *P. monegrensis* sp. n., sont décrites. *P. iberis* sp. n. est caractérisé par une taille moyenne (4,4-4,9 mm), une région labiale arrondie et en relief, des poches amphidiennes en forme d'étrier et une queue conique, convexe dorsalement et à extrémité conique. *P. iberis* sp. n., proche de *P. lemoni*, en diffère par la forme de la queue, la présence de mâles, la longueur du corps, la position du guide du stylet, et la valeurs des coefficients a et c. *P. monegrensis* sp. n. est caractérisé par sa grande taille (7,5-12 mm), une région labiale hémisphérique très en relief, des poches amphidiennes en forme d'étrier, une queue conique, convexe dorsalement à extrémité arrondie. Cette espèce est proche de *P. sandellus* et *P. xiphinemoides*. Elle diffère de *P. sandellus* par sa taille plus grande, un odontostyle plus long, la position du guide du stylet, et de *P. xiphinemoides* par la présence de mâles, la longueur de l'odontostyle, la position du guide du stylet et le coefficient c'. Une clé polytomique des 70 espèces du genre est proposée, de même que deux nouvelles combinaisons : *P. sativus* (Soni & Nama, 1983) n. comb et *P. wiesae* (Heyns, 1994) n. comb.

**Key-words** : Longidoridae, key to species, *Paralongidorus iberis* sp. n., *P. monegrensis* sp. n. *P. lemoni*, *P. sandellus*, *P. xiphinemoides*. Nematodes.

During a nematode survey near Serreta Negra (Huesca), in the North-West of Spain, two populations of the genus *Paralongidorus* Siddiqi, Hooper & Khan, 1963, were found for the first time in Spain. Two new species, *P. iberis* sp. n. and *P. monegrensis* sp. n., were identified in the same soil sample, from a dry area with xerophytic vegetation of a *Rhahmno-Cocciferetum* association, frequent in the valleys and in the shadiest places.

Khan *et al.* (1978) erected the genera *Siddiqia* Khan, Chawla & Saha, 1978 for *Paralongidorus* species having a clear offset lip region and funnel- to stirrup-shaped amphids, and *Longidoroides* Khan, Chawla & Saha, 1978 for those having a continuous lip region, exceptionally expanded, and pouch-like amphids, which left only the species having a continuous lip region and funnel shaped amphid remaining in the genus *Paralongidorus*. Luc and Doucet (1984)\* and later Jana and Baqri

(1984)\*\* as well as Coomans (1985) regarded *Siddiqia* as a junior synonym of *Paralongidorus*. Hunt (1993) proposed *Paralongidorus* and *Siddiqia* as subgenera in *Paralongidorus* based on head and amphid pouch shapes; in the same year, but later, Siddiqi *et al.* (1993) regarded *Longidoroides* as a junior synonym of *Paralongidorus* on the basis of the variability in the shape and size of amphidial pouch in some species of both genera.

Since amphidial pouches of some species –*P. duncani*; *P. hooperi*; *P. esci*; *P. sativus*; *P. spiralis*; *P. strelitziae*; *Paralongidorus* sp. in Rashid *et al.* (1986)– do not correspond to the funnel or bilobed types, we consider also the genera *Longidoroides* and *Siddiqia* as junior synonyms of *Paralongidorus*, and do not recognize the subgenera proposed by Hunt (1993) as valid. *Longidoroides sativus* Soni & Nama, 1983 and *L. wiesae* Heyns, 1994 are transferred to *Paralongidorus* as *P. sativus* (Soni &

\* August 1984.

\*\* December, 1984.

**Table 1.** Morphometric data of *Paralongidorus iberis* n. sp. (All measurements in  $\mu\text{m}$  except L in mm).

		Holotype		Paratypes			
		Females	Males	J1	J2	J3	J4
n		18	14	5	9	6	10
L	5.3	5.8 $\pm$ 0.6 (4.4-6.9)	5.1 $\pm$ 0.7 (3.6-5.9)	1.9 $\pm$ 0.6 (1.5-3.0)	2.6 $\pm$ 0.1 (1.9-3.1)	3.6 $\pm$ 0.6 (2.5-4.3)	3.9 $\pm$ 0.6 (2.8-4.6)
a	176	191.3 $\pm$ 26.1 (138-262)	179 $\pm$ 25 (121-223)	104 $\pm$ 23 (81-141)	130 $\pm$ 18 (107-160)	155 $\pm$ 16.3 (129-178)	168 $\pm$ 25 (141-202)
b	18	15.1 $\pm$ 5.9 (12-23)	16.2 $\pm$ 2.5 (11-16)	8.6 $\pm$ 3.1 (7-13)	10.3 $\pm$ 2.4 (7-15)	11.4 $\pm$ 2.8 (8-16)	13 $\pm$ 1.7 (11-16)
c	106	124 $\pm$ 17.7 (98-178)	108 $\pm$ 13 (80-124)	43.7 $\pm$ 12.2 (36-65)	58 $\pm$ 11 (39-73)	76.4 $\pm$ 14.6 (51-91)	82 $\pm$ 9.5 (72-98)
c'	3.1	2.2 $\pm$ 0.3 (1.9-3.1)	2.1 $\pm$ 0.1 (1.9-2.4)	3.2 $\pm$ 0.4 (2.7-3.9)	3 $\pm$ 0.3 (2.4-4)	2.5 $\pm$ 0.2 (2-3)	2.6 $\pm$ 0.2 (2-3)
V	51	49 $\pm$ 2.3 (46-56)	—	—	—	—	—
Stylet	110	112 $\pm$ 7.9 (104-126)	104 $\pm$ 13 (82-119)	81 $\pm$ 5.3 (75-86)	86 $\pm$ 4.1 (81-95)	94 $\pm$ 6.3 (86-101)	95 $\pm$ 10.4 (78-107)
Odontostyle	66.5	66.7 $\pm$ 4.4 (61-80)	65.7 $\pm$ 5.1 (56-73)	49 $\pm$ 3.3 (44-53)	52 $\pm$ 2.6 (47-54)	57 $\pm$ 3.6 (54-60)	57.8 $\pm$ 6.2 (48-66.5)
Odontophore	43.7	45.7 $\pm$ 5 (38-55)	38.7 $\pm$ 8.8 (21-50)	34.4 $\pm$ 6 (27-42)	34 $\pm$ 4 (28-42)	37 $\pm$ 4.4 (32-41)	40 $\pm$ 9.6 (28-63)
Rpl. od.style	—	—	—	48.6 $\pm$ 1.2 (46-49)	56 $\pm$ 3.3 (52-62)	61 $\pm$ 2 (58-63)	68.6 $\pm$ 4.6 (64-78)
Tail	50.3	47 $\pm$ 4.3 (39-55)	47.3 $\pm$ 3 (43-52)	43.3 $\pm$ 2.7 (40-47)	46 $\pm$ 3.8 (41-55)	47.5 $\pm$ 3.3 (42-51)	47 $\pm$ 4.2 (40-56)
Guiding ring	22.8	23.7 $\pm$ 1.1 (22-26)	24.2 $\pm$ 1.6 (21-28)	17 $\pm$ 1.2 (16-19)	19 $\pm$ 1 (18-21)	22 $\pm$ 2 (20-25)	21 $\pm$ 1.5 (18-24)
Spicule	—	—	34 $\pm$ 2 (30-38)	—	—	—	—
Supplements	—	—	(6-9)	—	—	—	—

Nama, 1983) n. comb. and *P. wiesae* (Heyns, 1994) n. comb.

Nematodes were extracted by Flegg's method, fixed in hot F.G. 4/1 and processed and mounted in anhydrous glycerin by the Seinhorst's rapid method.

***Paralongidorus iberis*\* sp. n.**  
(Fig. 1)

MEASUREMENTS


See Table 1.

DESCRIPTION

**Female**: Body elongate-slender, C-shaped or ventrally arcuate when relaxed by gentle heating. Large size, maximum width 24-26  $\mu\text{m}$ . Cuticle smooth, 2-3  $\mu\text{m}$  thick at oesophageal region, 4-6  $\mu\text{m}$  at midbody and on tail. Lip

region 8-10  $\mu\text{m}$  wide, rounded, flattened anteriorly, well set off by constriction at level of amphidial aperture. Amphidial aperture distinct, slit-like, 3-4  $\mu\text{m}$  wide or about half as long as head width, fovea stirrup-shaped. Lateral body pores arranged in lateral hypodermal chords irregularly along the body. Odontostyle thin, straight, slightly arcuate in posterior half, 6-8 times head width long or about 1.5-2 times odontophore length, with smooth unsplit base. Odontophore with slightly swollen posterior region. Stylet guiding ring single, 2-3 head widths from anterior end. Oesophagus dorylaimoid, posterior enlarged part cylindroid, 77-96  $\times$  9-11  $\mu\text{m}$ . Cardia cylindroid-rounded. Nerve ring just behind stylet. Vulva a transverse slit with slightly raised lips or flush with body surface, 2.62-3.20 mm from anterior end of body, 2.1-3.2 mm from tail terminus. Vagina about 60 % body width long. Two branches of reproductive organs equally developed; a distinct sphincter present at the junction of muscular and glandular parts of the uterus. Rectum 22-29  $\mu\text{m}$  long. Conoid dorsally convex tail, with conical terminus.

\* From the Latin name (Iber) of the river Ebro, close to the type locality.



**Fig. 1.** *Paralongidorus iberis* sp. n. A : Oesophageal region; B-C : Anterior region of juvenile and female respectively; D : Anterior genital branch; E-H : Tail region of ♂1, ♀2, female and male, respectively; I : spicule.

**Male** : Body ventrally curved in a C-shape; tail end curved. Cephalic region, cuticle, stylet and oesophagus as described for female. Testes paired, dorylaimoid; sperm oval to sausage-shaped, about 3.5–4 µm long and 1.5–2 µm wide. Supplements consisting of one adanal pair opposite the middle of the spicules, and a ventromedial series of six to nine supplements starting near the proximal end of the spicules or 42–49 µm from the cloacal opening. Spicules thick, slightly arcuate ventrally; with distinct lateral guiding pieces. Pores arranged irregularly along the body. Tail dorsally convex-conoid, ventrally slightly concave, attenuated towards the conical somewhat spicate terminus with thickened hyaline inner layer of the cuticle.

**Juveniles** : Resembling female except by tail length and shape (more elongate). First stage juvenile tail with a somewhat digitate terminus.

#### TYPE HABITAT AND LOCALITY

Soil around plants of *Rhamno-Cocciferetum* association in Barranco de la Valcuerna, Peñalva, province of Huesca, Spain.

#### TYPE MATERIAL

Holotype female and eleven paratypes deposited in the collection of the Centro de Ciencias Medioambientales, CSIC, Madrid, Spain and one female and male paratypes at each of : Muséum National d'Histoire Naturelle, Paris, France; International Institute of Parasitology, St. Albans, U.K.; Nematology Laboratory, University of Wageningen, The Netherlands; Indian Agricultural Institute, New Delhi, India; South African National Collection of Nematodes, Pretoria; and USDA Nematode Collection, Beltsville, MD, USA.

#### DIAGNOSIS AND RELATIONSHIPS

*P. iberis* sp. n. is recognised by its medium size, rounded lip region well set off by constriction, stirrup-shaped amphidial pouch or fovea, about half as long as head width, and stylet guiding ring two-three head widths from anterior end; tail conical, dorsally convex, with conical terminus and about two to three times anal body width long.

*P. iberis* is close to *P. lemoni* Nasira *et. al.*, 1993 in the shape of amphidial pouch and lip region. It differs in the tail shape, conical with a conoid terminus in *P. iberis* sp. n. and conical with a round terminus in *P. lemoni*; by the presence of males, and also by body length (4.4–6.9 *vs* 2.8–3.8 mm), position of the guiding ring (22–26 *vs* 32–38 µm), “a” ratio (138–262 *vs* 99–125) and “c” ratio (98–178 *vs* 77–96).

### *Paralongidorus monegrensis*\* sp. n. (Fig. 2)

#### MEASUREMENTS

See Table 2.


#### DESCRIPTION

**Female** : Body of large size, elongate-slender, maximum width of 79 (64–95) µm, C-shaped or ventrally arcuate when relaxed by gentle heating. Cuticle apparently smooth but finely striated transversely; 2–3 µm thick at oesophageal region and 4–6 µm at midbody and on the tail. Lip region hemispherical 15–21 µm width at amphidial level, smooth, slightly offset by expansion which is slightly wider than adjacent body. Amphidial aperture distinct, slit-like, 7–9 µm wide or about as long as half head width, pouch or fovea stirrup-shaped. Lateral hypodermal chords 10–12 µm wide or 1.5 of body width. Lateral body pores arranged irregularly along the body with two pores in tail region, one ventrolateral, and another dorsolateral. Odontostyle thin, straight, slightly arcuate in posterior half, 7–9 times head width long or about 1.5–2 times odontophore length, with smooth unsplit base. Odontophore with slightly swollen posterior region. Stylet guiding ring single, 1.4–1.8 head widths from anterior end. Oesophagus dorylaimoid, posterior enlarged part cylindroid 130–136 µm long, 19–25 µm wide. Cardia cylindroid-rounded. Nerve ring just posterior to stylet base. Vulva a transverse slit with slightly raised lips or flush with body surface; 4.5–6.8 mm from anterior end of body, 3.8–5.3 mm from tail terminus. Vagina about 60 % body width long. Two branches of reproductive organs equally developed; a distinct sphincter present at the junction of muscular and glandular parts of the uterus. Prerectum 114–118 µm long. Tail conoid, dorsally convex, with round terminus.

**Male** : Body ventrally curved in a C-shape; tail end slightly more curved. Cephalic region, cuticle, stylet and oesophagus as described for female. Testes paired, dorylaimoid; sperm oval to sausage-shaped, about 6–8 µm long and 2–3 µm wide. Supplements consisting of two or three adanal pairs, the first pair near the anus, the second and third pairs opposite the middle of the spicules, and a ventromedial series of ten to eleven supplements starting nearby the opposite proximal end of the spicules. Spicules thick, slightly arcuate ventrally, with distinct lateral guiding pieces. Pores arranged irregularly along the body; three caudal papillae on tail, one ventrodorsal, one dorsolateral and the other one dorsal. Tail broadly conoid, dorsally convex, ventrally slightly concave attenuated towards the blunt terminus with thickened hyaline inner layer of the cuticle.

**Juveniles** : Resembling female except in the more elongate and differently shaped tail.

\* From the name of the type locality, Monegros.



**Fig. 2.** *Paralongidorus monegrensis* sp. n. A, B : Anterior region of J2 and female, respectively; C, D : Female tails; E : Anterior genital branch; F, G : Male tail region; H, I : J1 and J2 tails respectively.

**Table 2.** Morphometric data of *Paralongidorus monegrensis* n. sp. (All measurements in  $\mu\text{m}$  except L in mm)

	Holotype	Paratypes					
		Females	Males	J1	J2	J3	J4
n		21	22	1	18	8	15
L	9.82	9.57 $\pm$ 1.2 (7.54-12.1)	9.47 $\pm$ 0.8 (7-11)	2.1	3.2 $\pm$ 0.3 (2.6-4.1)	4.4 $\pm$ 0.4 (3.8-4.7)	6.1 $\pm$ 0.7 (4.7-7.2)
a	126	119.2 $\pm$ 12.5 (99-145)	127.3 $\pm$ 16 (105-181)	75	82 $\pm$ 6.3 (73-92)	96.3 $\pm$ 12 (77-114)	109 $\pm$ 24.6 (83-172)
b	16	17.6 $\pm$ 2.9 (12-22)	17.56 $\pm$ 4.2 (11-29)	8.9	8.6 $\pm$ 1.1 (6.8-11.7)	9.1 $\pm$ 0.6 (7.8-10)	12.8 $\pm$ 2 (9-15)
c	230	272.5 $\pm$ 41.4 (198-341)	218.7 $\pm$ 26.8 (146-257)	49	75.6 $\pm$ 12.3 (60-114)	112 $\pm$ 9.4 (100-127)	158 $\pm$ 29 (89-194)
c''	0.97	0.64 $\pm$ 0.1 (0.5-0.8)	0.82 $\pm$ 0.1 (0.7-1)	2.1	1.2 $\pm$ 0.1 (1-1.7)	0.9 $\pm$ 0.1 (0.8-1)	0.8 $\pm$ 0.1 (0.68-1.1)
V	50	53.5 $\pm$ 1.8 (50-56)	—	—	—	—	—
Stylet	216.6	203 $\pm$ 10.6 (188-223)	207.4 $\pm$ 14 (185-235)	122	140.9 $\pm$ 7.8 (123-156)	166 $\pm$ 8.7 (151-178)	178 $\pm$ 20.5 (109-196)
Odontostyle	140.6	133 $\pm$ 6 (120-140)	135.1 $\pm$ 4.9 (123-146)	63	82.9 $\pm$ 4 (72-92)	104 $\pm$ 4 (97-111)	115 $\pm$ 4.6 (104-123)
Odontophore	76	70.3 $\pm$ 9.9 (56-86)	72 $\pm$ 11.6 (54-97)	59	58 $\pm$ 6.7 (43-74)	62 $\pm$ 6 (50-70)	69.2 $\pm$ 4.3 (63-78)
Rpl.od.style	—	—	—	82	103 $\pm$ 4.9 (90-111)	115 $\pm$ 5.4 (107-121)	131.4 $\pm$ 5.5 (122-143)
Tail	42.7	35 $\pm$ 4.2 (28-42)	43 $\pm$ 4.7 (29-52)	40.8	42 $\pm$ 4.3 (37-55)	39 $\pm$ 3.4 (35-44)	39 $\pm$ 5.2 (31-53)
Guiding ring	33.2	31.8 $\pm$ 1.5 (28-34)	32.6 $\pm$ 1.8 (29-35)	19	22.7 $\pm$ 1.7 (20-27.5)	26.5 $\pm$ 1.2 (25-28)	29 $\pm$ 2.7 (25-36)
Spicule	—	—	92.6 $\pm$ 4 (81-100)	—	—	—	—
Supplements	—	—	(10-11)	—	—	—	—

## TYPE HABITAT AND LOCALITY

Same as *P. iberis*.

## TYPE MATERIAL

*Holotype* female and 69 paratypes deposited in the collection of the Centro de Ciencias Medioambientales, CSIC, Madrid, Spain. One female and male paratypes at each of: Muséum National d'Histoire Naturelle, Paris, France; International Institute of Parasitology, St. Albans, U.K.; Nematology Laboratory, University of Wageningen, The Netherlands; Indian Agricultural Institute, New Delhi, India; South African National Collection of Nematodes, Pretoria, Republic of South Africa; and USDA Nematode Collection, Beltsville, MD, USA.

## DIAGNOSIS AND RELATIONSHIPS

*P. monegrensis* sp. n. is characterised by its large body size, hemispherical lip region slightly offset by expansion, stirrup-shaped amphidial pouch or fovea, about half as long as head width, and stylet guiding ring one to one-half head width from anterior end. The tail is conoid, dorsally convex, with rounded terminus, more than a half anal body width long.

*P. monegrensis* sp. n. resembles *P. sandellus* (Heyns, 1966) Coomans, 1985 and *P. xiphinemoides* Heyns, 1965 in amphidial pouch, lip region and tail shapes. It differs from *P. sandellus* in having longer body (7.5-12 vs 2.9-3.5 mm) and odontostyle length (120-140 vs 81-89  $\mu\text{m}$ ) and in the anterior position of guiding ring (28-34 vs 50-60  $\mu\text{m}$ ); and from *P. xiphinemoides* in the presence of males.

**Genus *Paralongidorus* Siddiqi,  
Hooper & Khan, 1963**

- = *Longidoroides* Khan, Chawla & Saha, 1978
- = *Siddiqia* Khan, Chawla & Saha, 1978
- = *Inagreiis* Khan, 1982
- = *Paralongidorus* (*Paralongidorus*); in Hunt, 1993
- = *Paralongidorus* (*Siddiqia*); in Hunt, 1993

## TYPE SPECIES

- 8\*. *P. sali* Siddiqi, Hooper & Khan, 1963
- = *P. (P.) sali* Siddiqi, Hooper & Khan, 1963; in Hunt, 1993

\* Numbers before each valid species refer to the order in the key.

## VALID SPECIES

- 55\*. *P. afzali* (Khan, 1964) Siddiqi & Husain, 1965  
= *Longidorus afzali* Khan, 1964  
= *Longidoroides afzali* (Khan, 1964) Khan, Chawla & Saha, 1978
4. *P. agni* Sharma & Edward, 1985  
= *P.(P.) agni* (Sharma & Edward, 1985); *in* Hunt, 1993
10. *P. australis* Stirling & McCulloch, 1984  
= *P. (P.) australis* (Stirling & McCulloch, 1984); *in* Hunt, 1993
18. *P. beryllus* Siddiqi & Hussain, 1965  
= *Siddiqia beryllus* (Siddiqi & Hussain, 1965) Khan, Chawla & Saha, 1978  
= *Inagreiis beryllus* (Siddiqi & Hussain, 1965) Khan, 1982  
= *Longidoroides beryllus* (Siddiqi & Hussain, 1965) Luc & Doucet, 1984  
= *P.(S.) beryllus* (Siddiqi & Hussain, 1965); *in* Hunt, 1993
69. *P. bikanerensis* (Lal & Mathur, 1987) Siddiqi, Baujard & Mountport, 1993  
= *Longidoroides bikanerensis* Lal & Mathur, 1987
64. *P. boshi* Khan, Saha & Seshadri, 1972  
= *Longidoroides boshi* (Khan, Saha & Seshadri, 1972) Khan, Chawla & Saha, 1978
23. *P. buchae* Lamberti, Roca & Chinappen, 1985  
= *P.(S.) buchae* (Lamberti, Roca & Chinappen, 1985); *in* Hunt, 1993
41. *P. buckeri* Sharma & Edward, 1985  
= *P.(P.) buckeri* (Sharma & Edward, 1985); *in* Hunt, 1993
14. *P. bullatus* Sharma & Siddiqi, 1990  
= *P.(S.) bullatus* (Sharma & Siddiqi, 1990); *in* Hunt, 1993
48. *P. capensis* Heyns, 1967  
= *Siddiqia capensis* (Heyns, 1967) Khan, Chawla & Saha, 1978  
= *P.(S.) capensis* (Heyns, 1967); *in* Hunt, 1993  
= *Siddiqia natalensis* Jacobs & Heyns, 1982  
= *P.natalensis* (Jacobs & Heyns, 1982) Luc & Doucet, 1984  
= *P.(S.) natalensis* (Jacobs & Heyns, 1982); *in* Hunt, 1993
5. *P. cebensis* Heyns & Coomans, 1989  
= *P. (P.) cebensis* (Heyns & Coomans, 1989); *in* Hunt, 1993
62. *P. cedari* (Khan, Chawla & Saha, 1978) Siddiqi, Baujard & Mountport, 1993  
= *Longidoroides cedari* Khan, Chawla & Saha, 1978
32. *P. christiani* Liebenberg, Heyns & Swart, 1993
27. *P. citri* (Siddiqi, 1959) Siddiqi, Hooper & Khan, 1963  
= *Xiphinema citri* Siddiqi, 1959  
= *Longidorus citri* (Siddiqi, 1959) Thorne, 1961  
= *Siddiqia citri* (Siddiqi, 1959) Khan, Chawla & Saha, 1978  
= *P.(S.) citri* (Siddiqi, 1959); *in* Hunt, 1993  
= *Paralongidorus droseri* Sukul, 1972  
= *Longidoroides droseri* (Sukul, 1972) Khan, Chawla & Saha, 1978
47. *P. clavicaudatus* (Jacobs & Heyns, 1982) Hunt, 1993  
= *Longidoroides clavicaudatus* Jacobs & Heyns, 1982  
= *Paralongidorus (P.) clavicaudatus* (Jacob & Heyns, 1982) *in* Hunt, 1993
60. *P. costatus* (Jacobs & Heyns, 1987) Siddiqi, Baujard & Mountport, 1993  
= *Longidoroides costatus* Jacobs & Heyns, 1987
44. *P. dasturi* (Ganguly, Patil & Khan, 1981) Luc & Doucet, 1984  
= *Siddiqia dasturi* Ganguly, Patil & Khan, 1981  
= *P.(S.) dasturi* (Ganguly, Patil & Khan, 1981); *in* Hunt, 1993
36. *P. deborae* (Jacobs & Heyns, 1982) Luc & Doucet, 1984  
= *Siddiqia deborae* Jacobs & Heyns, 1982  
= *P.(S.) deborae* (Jacobs & Heyns, 1982); *in* Hunt, 1993
6. *P. distinctus* Baqri & Jairajpuri, 1981
50. *P. duncani* Siddiqi, Baujard & Mountport, 1993
17. *P. epimikis* Dalmasso, 1969  
= *Siddiqia epimikis* (Dalmasso, 1969) Khan, Chawla & Saha, 1978  
= *P.(S.) epimikis* (Dalmasso, 1969); *in* Hunt, 1993
30. *P. erriae* Heyns, 1965  
= *Siddiqia erriae* (Heyns, 1965) Khan, Chawla & Saha, 1978  
= *P.(S.) erriae* (Heyns, 1965); *in* Hunt, 1993
9. *P. esci* Khan, Chawla & Saha, 1978  
= *P.(P.) esci* (Khan, Chawla & Saha, 1978); *in* Hunt, 1993
37. *P. eucalypti* Fisher, 1964  
= *Siddiqia eucalypti* (Fisher, 1964) Khan, Chawla & Saha, 1978  
= *P. (S.) eucalypti* (Fisher, 1964); *in* Hunt, 1993
67. *P. eugeni* (Khan, 1987) Hunt, 1993  
= *Inagreiis eugeni* Khan, 1986

\* Numbers before each valid species refer to the order in the key.



- = *Longidoroides eugeni* (Khan, 1987) Jairajpuri & Ahmad, 1992
- = *P.(S.) eugeni* (Khan, 1987) Hunt, 1993
25. *P. fici* Edward, Misra & Singh, 1964
- = *P.(P.) fici* (Edward, Misra & Singh, 1964); in Hunt, 1993
45. *P. fischeri* Heyns, 1972
- = *Siddiqia fischeri* (Heyns, 1972) Khan, Chawla & Saha, 1978
- = *P.(S.) fischeri* (Heyns, 1972); in Hunt, 1993
1. *P. flexus* Khan, Seshadri, Weischer & Mathen, 1971
- = *P.(P.) flexus* (Khan, Seshadri, Weischer & Mathen, 1971); in Hunt, 1993
26. *P. georgiensis* (Tulaganov, 1937) Siddiqi, 1964
- = *Longidorus georgiensis* Tulaganov, 1937
- = *Siddiqia georgiensis* (Tulaganov, 1937) Khan, Singh & Singh, 1981
- = *Paralongidorus (S.) georgiensis* (Tulaganov, 1937); in Hunt, 1993
68. *P. gloriosus* (Khan, 1982) Hunt, 1993
- = *Inagreius gloriosus* Khan, 1982
- = *Longidoroides gloriosus* (Khan, 1982) Luc & Doucet, 1984
- = *P.(S.) gloriosus* (Khan, 1982) in Hunt, 1993
31. *P. hanliae* Liebenberg, Heyns & Swart, 1993
54. *P. hooperi* Heyns, 1966
- = *Siddiqia hooperi* (Heyns, 1966) Khan, Chawla & Saha, 1978
- = *Longidoroides hooperi* (Heyns, 1966) Jacobs & Heyns, 1982
- = *P.(S.) hooperi* (Heyns, 1966); in Hunt, 1993
15. *P. iberis* sp. n.
19. *P. inagreinus* (Chawla & Samathanam, 1981) Luc & Doucet, 1984
- = *Siddiqia inagreina* Chawla & Samathanam, 1981
- = *P.(S.) inagreinus* (Chawla & Samathanam, 1981); in Hunt, 1993
28. *P. indicus* (Phukan & Sanwal, 1983) Luc & Doucet, 1984
- = *Siddiqia indicus* Phukan & Sanwal, 1983
- = *P.(S.) indicus* (Phukan & Sanwal, 1983); in Hunt, 1993
65. *P. latilabiatatus* (Jacobs & Heyns, 1982) Siddiqi, Baujard & Mountport, 1993
- = *Longidoroides latilabiatatus* Jacobs & Heyns, 1982
13. *P. lemoni* Nasira, Shahina, Firoza & Maqbool, 1993
70. *P. longiurus* (Chawla & Samathanam, 1981) Siddiqi, Baujard & Mountport, 1993
- = *Longidoroides longiurus* Chawla & Samathanam, 1981
42. *P. lutensis* Hunt & Rahman, 1991
- = *P.(P.) lutescens* (Hunt & Rahman, 1991); in Hunt, 1993 [*lapsus calami*]
52. *P. lutosus* (Heyns, 1965) Aboul-Eid, 1970
- = *Longidorus lutosus* Heyns, 1965
- = *Longidoroides lutosus* (Heyns, 1965) Khan, Chawla & Saha, 1978
- = *P.(P.) lutosus* (Heyns, 1965) Aboul-Eid, 1970 in Hunt, 1993
24. *P. major* Verma, 1973
- = *Siddiqia major* (Verma, 1973) Khan, Singh & Singh, 1981
- = *P.(S.) major* (Verma, 1973); in Hunt, 1993
35. *P. maximus* (Bütschli, 1874) Siddiqi, 1964
- = *Dorylaimus maximus* Bütschli, 1874
- = *Dorylaimus (Longidorus) maximus* Bütschli, 1874 (Micoletzky, 1922)
- = *Longidorus maximus* (Bütschli, 1874) Thorne & Swanger, 1936
- = *Siddiqia maxima* (Bütschli, 1874) Khan, Chawla & Saha, 1978
- = *P.(S.) maximus* (Bütschli, 1874); in Hunt, 1993
21. *P. mediensis* (Ganguly, Patil & Khan, 1981) Luc & Doucet, 1984
- = *Siddiqia mediensis* Ganguly, Patil & Khan, 1981
- = *P.(S.) mediensis* (Ganguly, Patil & Khan, 1981); in Hunt, 1993
12. *P. microlaimus* Siddiqi, 1964
- = *P.(P.) microlaimus* (Siddiqi, 1964); in Hunt, 1993
7. *P. monegrensis* sp. n.
29. *P. namibiensis* Jacobs & Heyns, 1987
- = *P.(S.) namibiensis* (Jacobs & Heyns, 1987); in Hunt, 1993
20. *P. nudus* (Kirjanova, 1951) Lamberti, 1975
- = *Longidorus nudus* Kirjanova, 1951
- = *P.(P.) nudus* (Kirjanova, 1951); in Hunt, 1993
11. *P. oryzae* Verma, 1973
- = *P.(P.) oryzae* (Verma, 1973); in Hunt, 1993
39. *P. paramaximus* Heyns, 1965
- = *Siddiqia paramaximus* (Heyns, 1965) Khan, Chawla & Saha, 1978
- = *P.(S.) paramaximus* (Heyns, 1965); in Hunt, 1993
63. *P. pini* (Jacobs & Heyns, 1987) Siddiqi, Baujard & Mountport, 1993
- = *Longidoroides pini* Jacobs & Heyns, 1987
59. *P. pulcher* (Jacobs & Heyns, 1982) Siddiqi, Baujard & Mountport, 1993
- = *Longidoroides pulcher* Jacobs & Heyns, 1982
56. *P. pulcheroides* (Jacobs & Heyns, 1987) Siddiqi, Baujard & Mountport, 1993

- = *Longidoroides pulcheroides* Jacobs & Heyns, 1987
22. *P. remyi* (Altherr, 1963) Siddiqi & Husain, 1965  
= *Longidorus remyi* Altherr, 1963  
= *Siddiqia remyi* (Altherr, 1963) Khan, Chawla & Saha, 1978  
= *P.(S.) remyi* (Altherr, 1963); in Hunt, 1993
40. *P. rex* Andrassy, 1986  
= *P.(S.) rex* (Andrassy, 1986); in Hunt, 1993
46. *P. rotundatus* Khan, 1987  
= *P.(P.) rotundatus* (Khan, 1987); in Hunt, 1993
3. *P. sacchari* Siddiqi, Hooper & Khan, 1963  
= *P.(P.) sacchari* (Siddiqi, Hooper & Khan, 1963); in Hunt, 1993
51. *P. sandellus* (Heyns, 1966) Coomans, 1985  
= *Xiphinema sandellum* Heyns, 1966  
= *Longidorus sandellus* (Heyns, 1966) Khan, Chawla & Saha, 1978  
= *Brevinema sandellum* (Heyns, 1966) Chaves & Coomans, 1984  
= *P. (P.) sandellus* (Heyns, 1966); in Hunt, 1993
53. *P. sativus* (Soni & Nama, 1983) n. comb.  
= *Longidoroides sativus* Soni & Nama, 1983
34. *P. seclipsi* (Khan, Singh & Singh, 1981) Jana & Baqri, 1984  
= *Siddiqia seclipsi* Khan, Singh & Singh, 1981  
= *Longidoroides seclipsi* (Khan, Singh & Singh, 1981) Luc & Doucet, 1984  
= *P.(S.) seclipsi* (Khan, Singh & Singh, 1981); in Hunt, 1993
2. *P. similis* Khan, Chawla & Prasad, 1972  
= *P.(P.) similis* (Khan, Chawla & Prasad, 1972); in Hunt, 1993
33. *P. spasskii* Heyns, 1972  
= *Siddiqia spasskii* (Heyns, 1972) Khan, Chawla & Saha, 1978  
= *P. (S.) spasskii* (Heyns, 1972); in Hunt, 1993
38. *P. spauli* (Jacobs & Heyns, 1982) Luc & Doucet, 1984  
= *Siddiqia spauli* Jacob & Heyns, 1982  
= *P.(S.) spauli* (Jacobs & Heyns, 1982); in Hunt, 1993
57. *P. spiralis* Khan, Saha & Seshadri, 1972  
= *Longidoroides spiralis* (Khan, Saha & Seshadri, 1972) Khan, Chawla & Saha, 1978
49. *P. strelitziae* (Heyns, 1966) Aboul-Eid, 1970  
= *Longidorus strelitziae* Heyns, 1966  
= *Longidoroides strelitziae* (Heyns, 1966) Khan, Chawla & Saha, 1978
66. *P. teres* (Khan, 1987) Hunt, 1993  
= *Inagreius teres* Khan, 1987  
= *Longidoroides teres* (Khan, 1987) Jairajpuri & Ahmad, 1992  
= *P. (S.) teres* (Khan, 1987); in Hunt, 1993
58. *P. utriculoides* (Corbett, 1964) Siddiqi & Husain, 1965  
= *Longidorus utriculoides* Corbett, 1964  
= *Longidoroides utriculoides* (Corbett, 1964) Khan, Chawla & Saha, 1978
61. *P. wiesae* (Heyns, 1994) n. comb.  
= *Longidoroides wiesae* Heyns, 1994
16. *P. xiphinemoides* Heyns, 1965  
= *Siddiqia xiphinemoides* (Heyns, 1965) Khan, Chawla & Saha, 1978  
= *P.(S.) xiphinemoides* (Heyns, 1965); in Hunt, 1993
43. *P. zenobiae* Hunt & Rahman, 1991  
= *P. (P.) zenobiae* Hunt & Rahman, 1991; in Hunt, 1993

## Polytomous key

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
1. <i>P. flexus</i>	1	1	1	1	2	2	34	1	4	4	2	13	1	1	1
2. <i>P. similis</i>	1	1	1	2	1	1	1	1	1	2	1	1	1	—	—
3. <i>P. sacchari</i>	1	1	1	2	1	2	34	1	3	23	23	1	2	—	—
4. <i>P. agni</i>	1	1	1	2	2	2	12	12	1	1	1	1	13	1	1
5. <i>P. cebensis</i>	1	1	1	2	2	4	5	3	2	1	3	1	3	3	3
6. <i>P. distinctus</i>	1	1	1	2	1	4	7	2	2	1	4	1	3	—	—
7. <i>P. monegrensensis</i>	1	1	1	2	2	56	45	1	23	1	3	23	3	3	1
8. <i>P. sali</i>	1	1	1	3	1	1	3	1	1	1	2	1	1	—	—
9. <i>P. esci</i>	1	1	1	3	2	2	6	2	2	1	34	1	1	1	3
10. <i>P. australis</i>	1	1	1	3	2	56	67	3	12	1	34	13	23	4	3
11. <i>P. oryzae</i>	1	1	1(3)	2	1	1	1	1	2	3	1	1	1	—	—
12. <i>P. microlaimus</i>	1	1	1(3)	2	2	12	2	1	12	23	2	1	1	1	1
13. <i>P. lemoni</i>	1	1	2	1	1	12	1	1	2	4	1	23	1	—	—
14. <i>P. bullatus</i>	1	1	2	1	1	2	57	1	23	3	34	3	12	—	—
15. <i>P. iberis</i>	1	1	2	1	2	24	12	1	34	4	1	34	1	1	1

**Polytomous key** (continued)

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
16. <i>P. xiphinemoides</i>	1	1	2	2	1	12	3	3	12	3	2	12	1	—	—
17. <i>P. epimikis</i>	1	1	2	2	2	6	7	1	2	2	4	4	2	1	2
18. <i>P. beryllus</i>	1	1	3	1	1	2	2	1	2	3	3	2	1	—	—
19. <i>P. inagreinus</i>	1	1	3	2	1	2	3	1	12	12	3	23	2	—	—
20. <i>P. nudus</i>	1	1	3	2	1	2	3	1	3	23	2	1	1	—	—
21. <i>P. mediansis</i>	1	1	3	2	1	2	45	1	2	3	2	23	1	—	—
22. <i>P. remyi</i>	1	1	3	2	1	2	7	1	2	1	1	—	—	—	—
23. <i>P. buchae</i>	1	1	3	2	1	23	4	1	2	2	3	3	2	—	—
24. <i>P. major</i>	1	1	3	2	1	24	4	1	2	2	23	23	2	—	—
25. <i>P. fici</i>	1	1	3	2	1	34	45	1	2	23	3	3	2	—	—
26. <i>P. georgiensis</i>	1	1	3	2	1	34	5	1	2	2	3	3	2	—	—
27. <i>P. citri</i>	1	1	3	2	1	4	5	1	2	23	3	3	2	—	—
28. <i>P. indicus</i>	1	1	3	2	1	45	3	1	2	2	2	3	13	—	—
29. <i>P. namibiensis</i>	1	1	3	2	1	45	67	1	12	13	4	34	12	—	—
30. <i>P. erriae</i>	1	1	3	2	2	12	1	1	12	23	2	13	1	1	2
31. <i>P. hanliae</i>	1	1	3	2	2	23	23	1	12	13	23	13	—	1	13
32. <i>P. christiani</i>	1	1	3	2	2	3	2	2	2	1	2	1	1	2	2
33. <i>P. spasskii</i>	1	1	3	2	2	34	2	1	2	2	34	23	12	12	2
34. <i>P. seclipsi</i>	1	1	3	2	2	34	56	1	2	1	4	12	3	2	2
35. <i>P. maximus</i>	1	1	3	2	2	56	7	2	2	1	6	13	3	4	2
36. <i>P. deborae</i>	1	1	3	2	2	6	7	12	2	2	45	4	23	2	23
37. <i>P. eucalypti</i>	1	1	3	23	1	23	56	2	3	3	3	23	12	—	—
38. <i>P. spauli</i>	1	1	3	3	2	23	12	1	12	3	1	23	1	1	13
39. <i>P. paramaximus</i>	1	1	4	2	2	4	4-7	12	12	12	45	34	13	13	23
40. <i>P. rex</i>	1	1	4	3	1	6	7	2	3	1	6	23	3	—	—
41. <i>P. buckeri</i>	1	2	1	2	1	1	1	1	2	2	1	1	1	—	—
42. <i>P. lutensis</i>	1	2	1	3	1	35	57	3	1	1	12	1	—	—	—
43. <i>P. zenobiae</i>	1	2	1	3	1	46	7	3	24	1	1	1	1	—	—
44. <i>P. dasturi</i>	1	2	3	2	1	23	35	1	1	1	3	12	23	—	—
45. <i>P. fischeri</i>	1	2	3	2	2	2	2	12	23	3	2	12	12	1	1
46. <i>P. rotundatus</i>	1	2	3	3	1	2	67	2	1	1	3	1	1	—	—
47. <i>P. clavicaudatus</i>	1(3)	1	1	3	2	2	2	2	12	12	1	12	1	1	1
48. <i>P. capensis</i>	1(3)	1	3	2	2	26	45	2	2-4	3	3	34	1-3	12	2
49. <i>P. strelitziae</i>	1(3)	2	1	23	2	26	25	3	12	1	34	2	3	23	23
50. <i>P. duncani</i>	2	1	2	1	2	1	1	1	34	4	1	1	1	1	12
51. <i>P. sandellus</i>	2	1	2	2	2	1	2	3	1	23	2	1	1	1	1
52. <i>P. lutosus</i>	2(3)	1	2(3)	2	2	4	23	12	2	3	2	34	13	1	1
53. <i>P. sativus</i>	2	2	1	2	1	2	1	2	—	2	2	1	1	—	—
54. <i>P. hooperi</i>	2	2	2	2	2	56	7	3	23	12	45	4	2	23	12
55. <i>P. afzali</i>	3	1	1	1	1	12	4	1	4	4	2	23	1	—	—
56. <i>P. pulcheroides</i>	3	1	1	2	1	12	1	1	2	3	12	23	1	—	—
57. <i>P. spiralis</i>	3	1	1	2	1	12	2	2	2	3	2	1	1	—	—
58. <i>P. utriculoides</i>	3	1	1	2	1	2	1	1	2	3	2	3	1	—	—
59. <i>P. pulcher</i>	3	1	1	2	1	2	2	1	12	12	2	13	13	—	—
60. <i>P. costatus</i>	3	1	1	2	1	2	34	1	24	3	2	23	12	—	—
61. <i>P. wiesae</i>	3	1	1	2	1	2	34	2	23	12	23	1	23	—	—
62. <i>P. cedari</i>	3	1	1	2	2	2	56	3	1	1	2	1	2	3	1
63. <i>P. pini</i>	3	1	1	2	2	3-6	34	23	23	3	3	23	23	2	12
64. <i>P. boshi</i>	3	1	1	3	2	2	4	3	1	1	2	1	1	1	1
65. <i>P. latilabiatius</i>	3	1	2	2	1	4	2	3	2	3	3	4	1	—	—
66. <i>P. teres</i>	3	1	3	2	1	24	45	1	2	2	3	13	2	—	—
67. <i>P. eugeni</i>	3	1	3	2	1	2-5	46	2	1	2	2	13	2	—	—
68. <i>P. gloriosus</i>	3	1	3	2	1	34	45	1	2	3	2	3	2	—	—
69. <i>P. bikanerensis</i>	3	1	3	2	1	34	5	1	2	2	3	3	1	—	—
70. <i>P. longiurus</i>	3	2	1(2)	1	1	1	2	1	4	4	2	1	1	—	—

**Polytomous key for *Paralongidorus* species identification****CHARACTERS USED FOR THE POLYTOMOUS KEY**

## The code

- A. Shape of amphidial pouches :
  - 1. Funnel- or stirrup-shaped (Fig. 3 F, J, M)
  - 2. Wine glass-shaped (Fig. 3 C, E)
  - 3. Pouch-shaped, bilobed or not (Fig. 3 A, B)
- B. Width of amphid aperture :
  - 1. Aperture of amphid half or more as wide lip region
  - 2. Aperture less than half of lip region width
- C. Shape of lip region :
  - 1. Continuous with outline of the body (Fig. 3 A, B)
  - 2. Expanded (Fig. 3 C, E, F)
  - 3. Set off by constriction without neck (Fig. 3 H, J)
  - 4. Clearly set off by a deep constriction with neck (Fig. 3 M)
- D. Tail shape of female :
  - 1. Conoid (Fig. 3 D, G, I)
  - 2. Conoid with a terminus broadly rounded to sub-hemispherical (Fig. 3 K, L, O, P)
  - 3. Hemispherical (Fig. 3 Q, N)
- E. Males :
  - 1. Unknown
  - 2. Known
- F. Body length of females :
  - 1. < 3.6 mm
  - 2. 3.6-5.6 mm
  - 3. 5.7-6.2 mm
  - 4. 6.3-7.6 mm
  - 5. 7.7-8.2 mm
  - 6. > 8.2 mm
- G. Length of odontostyle :
  - 1. < 70  $\mu\text{m}$
  - 2. 70-94  $\mu\text{m}$
  - 3. 95-110  $\mu\text{m}$
  - 4. 111-124  $\mu\text{m}$
  - 5. 125-140  $\mu\text{m}$
  - 6. 141-152  $\mu\text{m}$
  - 7. > 152  $\mu\text{m}$
- H. Distance from anterior end to guiding ring :
  - 1. < 37  $\mu\text{m}$
  - 2. 37-47  $\mu\text{m}$
  - 3. > 47  $\mu\text{m}$
- I. Tail length :
  - 1. < 27  $\mu\text{m}$
  - 2. 27-38  $\mu\text{m}$
  - 3. 39-50  $\mu\text{m}$
  - 4. > 50  $\mu\text{m}$
- J. Ratio c' :
  - 1. < 0.8
  - 2. 0.8-1.0
  - 3. 1.1-1.8
  - 4. > 1.8


- K. Lip region width :
  - 1. < 10  $\mu\text{m}$
  - 2. 10-15  $\mu\text{m}$
  - 3. 16-20  $\mu\text{m}$
  - 4. 21-25  $\mu\text{m}$
  - 5. 26-30  $\mu\text{m}$
  - 6. > 30  $\mu\text{m}$
- L. Ratio a :
  - 1. < 95
  - 2. 95-108
  - 3. 109-145
  - 4. > 145
- M. Length of basal bulb of oesophagus :
  - 1. < 108  $\mu\text{m}$
  - 2. 108-128  $\mu\text{m}$
  - 3. > 128  $\mu\text{m}$
- N. Length of spicules :
  - 1. < 60  $\mu\text{m}$
  - 2. 60-80  $\mu\text{m}$
  - 3. 81-100  $\mu\text{m}$
  - 4. > 100  $\mu\text{m}$
- O. Number of supplements in male tail :
  - 1. < 11
  - 2. 11-16
  - 3. > 16

**Acknowledgements**

Authors thank Prof. A. Bello, Drs. P. Baujard, P. González, D.J. Hunt, P.A.A. Loof and M. Luc for their valuable suggestions and Dr.A.M. Golden for lending material.

**References**

- ABOUL-El, H. Z. (1970). Systematic notes on *Longidorus* and *Paralongidorus*. *Nematologica*, 16 : 159-179.
- ALTHERR, E. (1963). Contribution à la connaissance de la faune des sables submergés en Lorraine. Nématodes. *Annls Spéol.*, 18 : 53-98.
- ANDRÁSSY, I. (1986). Egy új Tüfönálféreg faj magyarországról: *Paralongidorus rex* sp. n. (Nematoda : Longidoridae). *Allattani Közlemények*, 73 : 115-118.
- BAQRI, Q. H. & JAIRAJPURI, J. S. (1981). A new species of the genus *Paralongidorus* (Longidoridae : Nematoda). *Bull. zool. Surv. India*, 4 : 37-39.
- BÜTSCHLI, O. (1874). Zur Kenntnis der freilebenden Nematoden insbesondere der des Kieler Hafens. *Abh. Seskenb. Naturf. Ges.*, 9 : 237-292.
- CHAVES, E & COOMANS, A. (1984). Three new species of *Xiphidius* from Argentina with comments on *Xiphinema sandellum* Heyns, 1966. *Revue Nématol.*, 7 : 3-12.
- CHAWLA, M. L. & SAMATHANAM, G. J. (1981). Two new species of the superfamily Longidoroidea (Dorylaimida : Nematoda) from India. *Indian J. Nematol.*, 10 (1980) : 205-210.
- COOMANS, A. (1985). A phylogenetic approach to the classification of the Longidoridae (Nematoda : Dorylaimida). *Agron., Ecosyst. Envir.*, 12 : 335-354.



**Fig. 3.** A-C, E, F, H, J, M : Lip region and amphidial pouch shapes (Codes A and C). A (codes A3, C1) *P. pini*; B (codes A3, C1) *P. pulcheroides*; C (codes A2, C2) *P. duncani*; E (codes A2, C2) *P. lutosus*; F (codes A1, C2) *P. bullatus*; H (code C3) *P. beryllus*; J (codes A1, C3) *P. buchae*; M (codes A1, C4) *P. rex*. D, G, I, K, L, N-Q : Caudal region (Code D). D, G, I (code D1) *P. duncani*, *P. bullatus* and *P. beryllus* respectively; K, L, O, P (code D2) *P. buchae*, *P. deborae*, *P. fischeri* and *P. cebensis*; N, Q (code D3) *P. rex*, *P. australis* respectively. (Redrawn from the original descriptions).

- CORBETT, D. C. M. (1964). *Longidorus utriculoides* n. sp. (Nematoda : Dorylaimidae) from Nyasaland. *Nematologica*, 10 : 496-499.
- DALMASSO, A. (1969). Étude anatomique et taxonomique des genres *Xiphinema*, *Longidorus* et *Paralongidorus* (Nematoda : Dorylaimida). *Mém. Mus. natn. Hist. nat., Paris, Sér. A, Zool.*, 61 : 33-82.
- EDWARD, J. C., MISRA, S. L. & SINGH, G. R. (1964). A new species of *Paralongidorus* (Nematoda, Dorylaimoidea) from Allahabad, Uttar Pradesh, India. *Jap. J. appl. Ent. Zool.*, 8 : 313-316.
- FISHER, J. M. (1964). *Dolichodorus adelaidensis* n. sp. and *Paralongidorus eucalypti* n. sp. from S. Australia. *Nematologica*, 10 : 464-470.
- GANGULY, S., PATIL, K. J. & KHAN, E. (1981). Two new species of *Siddiqia* from India. *Indian J. Nematol.*, 10 (1980) : 175-181.
- HEYNS, J. (1965). New species of the genera *Paralongidorus* and *Longidorus* (Nematoda : Dorylaimoidea) from South Africa. *S. Afr. J. agric. Sci.*, 9 : 863-874.
- HEYNS, J. (1966). Further studies on South African Longidoridae (Nematoda). *S. Afr. J. agric. Sci.*, 9 : 927-944.
- HEYNS, J. (1967). *Paralongidorus capensis* n. sp. and *Longidorus belondiroides* n. sp., with a note on *L. taniwha* Clark, 1963 (Nematoda : Longidoridae). *Nematologica*, 12 (1966) : 568-574.
- HEYNS, J. (1972). Observations on South African *Paralongidorus* species (Nematoda : Longidoridae) with descriptions of two new species. *Phytophylactica*, 4 : 127-132.
- HEYNS, J. (1975). *Paralongidorus maximus* C.I.H. Descriptions of Plant-parasitic Nematodes. Set 5, n° 75, 4 p.
- HEYNS, J. (1994). Description of *Longidoroides wiesae* spec. nov. from the Kruger National Park (Nematoda : Dorylaimida). *Koedoe*, 37 : 97-103.
- HEYNS, J. & COOMANS, A. (1989). *Paralongidorus cebensis* n. sp. from the Transkei (Nematoda : Longidoridae). *Phytophylactica*, 21 : 91-94.
- HUNT, D. J. (1993). *Aphelenchida, Longidoridae and Trichodoridae : their systematics and bionomics*. Wallingford, UK, CAB International, 352 p.
- HUNT, D. J. & RAHMAN, L. (1991). Two new species of *Paralongidorus* (Nematoda : Dorylaimida) from deep water rice in Bangladesh. *Afro-Asian J. Nematol.*, 1 : 83-85.
- JACOBS, P. J. F. & HEYNS, J. (1982). *Siddiqia* species from sugar cane in Natal (Nematoda : Longidoridae). *Phytophylactica*, 14 : 169-178.
- JACOBS, P. J. F. & HEYNS, J. (1982). *Longidoroides* species from sugar cane in Natal (Nematoda : Longidoridae). *Phytophylactica*, 14 : 179-193.
- JACOBS, P. J. F. & HEYNS, J. (1987). New and known species of *Longidoroides* and *Paralongidorus* from South Africa (Nematoda : Longidoridae). *Phytophylactica*, 19 : 7-14.
- JAIRAJPURI, M. S. & AHMAD, W. (1992). *Dorylaimida. Free-living, predaceous and plant-parasitic nematodes*. New Delhi, India, Oxford & IBH Publishing Co., xiv + 458 p.
- JANA, A. & BAQRI, Q. H. (1984). Nematodes from West Bengal (India). XIV. On the occurrence of ectoparasitic nematodes of Longidoroidea and Trichodoroidea (Dorylaimida), with remarks on the validity of the genus *Siddiqia* (Longidoridae). *Bull. zool. Surv. India*, 6 : 75-79.
- KHAN, E. (1964). *Longidorus afzali* n. sp. and *Xiphinema arcum* n. sp. (Nematoda : Longidoridae) from India. *Nematologica*, 10 : 313-318.
- KHAN, E. (1982). *Inagreiis gloriosus* n. gen., n. sp. n. and descriptions of three new species of *Xiphinema* Cobb, 1913 along with report on *X. radiculicola* T. Goodey, 1936 and *X. elongatum*. Sch. Stek. & Teun., 1938 (Nematoda : Longidoroidea) from India. *Indian J. Nematol.*, 11 (1981) : 189-204.
- KHAN, E. (1987 a). Two new species of *Inagreiis* Khan, 1982 (Nematoda : Longidoroidea) from India. *Indian J. Nematol.*, 16 (1986) : 163-167.
- KHAN, E. (1987 b). One new genus and four new species in the superfamily Longidoroidea (Nematoda). *Indian J. Nematol.*, 16 (1986) : 185-193.
- KHAN, E., CHAWLA, M. L. & PRASAD, S. K. (1972). *Paralongidorus similis* sp. nov. (Nematoda : Longidoridae) from Poona. *Indian Bull. Ent.*, 12 : 52-54.
- KHAN, E., CHAWLA, M. L. & SAHA, M. (1978). Comments on the classification of the Longidoroidea (Nematoda) with description of three new species. *Indian J. Nematol.*, 6 (1976) : 47-62.
- KHAN, E., SAHA, M. P. & SESHADRI, A. R. (1972). Plant parasitic nematodes from Kumaon Hills, India. I. Two new species of *Paralongidorus* (Nematoda : Longidoridae). *Nematologica*, 18 : 38-43.
- KHAN, E., SESHADRI, A. R., WEISCHER, B. & MATHEN, K. (1971). Five new nematode species associated with coconut in Kerala, India. *Indian J. Nematol.*, 1 : 116-127.
- KHAN, E., SINGH, M. & SINGH, L. K. (1981). *Siddiqia seclipsi* sp. n. (Nematoda : Longidoroidea) from India with a key to the species of *Siddiqia* Khan, Chawla & Saha, 1978. *Indian J. Nematol.*, 10 (1980) : 211-215.
- KIRJANOVA, E. S. (1951). [Soil nematodes found in cotton fields and in virgin soil of Golodnaya Steppe (Uzbekistan)]. *Trudy Zool. Inst. Akad. Nauk. SSSR*, 9 : 625-657.
- LAL, A. & MATHUR, V. K. (1987). Nematodes in the rhizosphere of date palm with description of *Longidoroides bikaneensis* sp. n. (Nematoda : Dorylaimoidea). *Date Palm J.*, 5 : 208-214.
- LAMBERTI, F. (1975). Taxonomy of *Longidorus* (Micoletzky) Filipjev and *Paralongidorus* Siddiqi, Hooper and Khan. In : Lamberti, F., Taylor, C. E. & Seinhorst, J. W. (Eds). *Nematode vector of plant viruses*. London & New York, Plenum Press : 71-90.
- LAMBERTI, F., ROCA, F. & CHINAPPEN, M. (1985). *Paralongidorus buchae*, a new Longidoridae species associated with declining chilli in Mauritius. *Nematol. med.*, 13 : 213-216.
- LIEBENBERG, W., HEYNS, J. & SWART, A. (1993). Studies on *Paralongidorus* species from Southern Africa. I. Introduction and descriptions of *P. capensis* Heyns, 1967 and *P. hanliae*

- spec. nov. (Nematoda : Longidoridae). *Phytophylactica*, 25 : 91-100.
- LIEBENBERG, W., HEYNS, J. & SWART, A. (1993). Studies on *Paralongidorus* species from Southern Africa. II. Data on new collections, with descriptions on one new and two known species (Nematoda : Longidoridae). *Phytophylactica*, 25 : 137-148.
- LIEBENBERG, W., HEYNS, J. & SWART, A. (1993). Studies on *Paralongidorus* species from Southern Africa. III. Redescriptions of two species and new data on the morphology of several others (Nematoda : Longidoridae). *Phytophylactica*, 25 : 231-242.
- LUC, M. & DOUCET, M. E. (1984). Description of *Xiphidorus achalae* n. sp. and proposal for a classification of Longidorids (Nematoda : Dorylaimoidea), *Revue Nématol.*, 7 : 103-112.
- MICOLETZKY, H. (1922) Die freilebenden Erd-Nematoden. *Arch. Naturg.*, 87 (1921) : 1-650.
- NASIRA, K., SHAHINA, F., FIROZA, K. & MAQBOOL, M. A. (1993). Description of *Paralongidorus lemoni* n. s., and redescription of *P. major* Verma, 1973 (Nematoda : Longidoridae) from Pakistan with SEM observation. *Pakistan J. Nematol.*, 11 : 79-91.
- PHUKAN, P. N. & SANWAL, K. C. (1983). *Siddiqia indicus* sp. n. (Nematoda : Longidoroidea) from Assam, India. *Indian J. Nematol.*, 12 (1982) : 188-191.
- RASHID, F., COOMANS, A. & SHARMA, D. (1986). Longidoridae (Nematoda : Dorylaimida) from Bahia State, Brazil. *Nematol. med.*, 14 : 235-250.
- ROBBINS, R. T. (1978). Descriptions of females (emended), a male, and juveniles of *Paralongidorus microlaimus* (Nematoda : Longidoridae). *J. Nematol.*, 10 : 28-34.
- SHARMA, N. N. & EDWARD, J. C. (1985). Two new species of *Paralongidorus* (Nematoda : Longidoridae) from Maharashtra, India. *J. Soil Biol. Ecol.*, 5 : 14-19.
- SHARMA, S. R. & SIDDIQI, M. R. (1990). *Paralongidorus bullatus* n. sp. from groundnut soils in Niger and comments on *Xiphinema parasetariae* Luc. *J. Nematol.*, 22 : 579-584.
- SIDDIQI, M. R. (1959). Studies on *Xiphinema* spp. (Nematoda : Dorylaimoidea) from Aligarh (North India), with comments on the genus *Longidorus* Micoletzky, 1922. *Proc. helminth. Soc. Wash.*, 26 : 151-163.
- SIDDIQI, M. R. (1964). *Xiphinema conurum* n. sp. and *Paralongidorus microlaimus* n. sp. with a key to the species of *Paralongidorus* (Nematoda : Longidoridae). *Proc. helminth. Soc. Wash.*, 31 : 133-137.
- SIDDIQI, M. R., BAUJARD, P. & MOUNPORT, D. (1993). Description of *Paratylenchus pernoxius* sp. n. and *Paralongidorus duncani* sp. n. from Senegal, and the synonymization of *Longidoroides* with *Paralongidorus*. *Afro-Asian J. Nematol.*, 3 : 81-89.
- SIDDIQI, M. R., HOOPER, D. J. & KHAN, E. (1963). A new nematode genus *Paralongidorus* (Nematoda : Dorylaimoidea) with description of two new species and observations on *Paralongidorus citri* (Siddiqi, 1959) n. comb. *Nematologica*, 9 : 7-14.
- SIDDIQI, M. R. & HUSAIN, Z. (1965). *Paralongidorus beryllus* n. sp. (Nematoda : Dorylaimoidea) from India. *Proc. helminth. Soc. Wash.*, 32 : 243-245.
- SONI, G. R. & NAMA, H. S. (1983). On a new species of *Longidoroides* Khan, Chawla and Saha, 1978 and with a key to its species. *Current Sci.*, 52 : 935-936.
- STIRLING, G. R. & MCCULLOCH, J. S. (1984). *Paralongidorus australis* n. sp. (Nematoda : Longidoridae), causing poor growth of rice in Australia. *Nematologica*, 30 : 387-394.
- SUKUL, N. C. (1972). *Paralongidorus droseri* sp. nov. (Nematoda : Longidoridae) associated with an insectivorous plant from West Bengal. *Bull. Ent.*, 12 (1971) : 85-88.
- THORNE G. (1961). *Principles of nematology*. New York, London & Toronto, McGraw-Hill Book Co., 533 p.
- THORNE, G. & SWANGER, H. H. (1936). A monograph of the nematodes genera *Dorylaimus* Dujardin, *Aporcelaimus* n. g., *Dorylaimoides* n. g. and *Pungentus* n. g. *Capita zool.*, 6 : 1-223.
- TULAGANOV, D. A. (1937) Nematoden der Tomate und des sie umgebenden Bodens. *Zool. Anz.*, 118 : 283-285.
- VERMA, R. S. (1973). Two new species in the subfamily Longidorinae (Nematoda) from Uttar Pradesh, India, with a key to species of *Paralongidorus* Siddiqi et al., 1963. *Zool. Anz.*, 190 : 170-174.