

MH4501 Multivariate Analysis

Assignment

Name: Honda Naoki

Matriculation: N1804369J

March 19, 2019

Question 1.

(a) Solution.

Source of variation	SSCP	df
Treatments	$B = \begin{pmatrix} 1.051 & 2.173 & -1.376 & -0.760 \\ 2.173 & 4.880 & -2.373 & -1.257 \\ -1.376 & -2.373 & 2.382 & 1.384 \\ -0.760 & -1.257 & 1.384 & 0.811 \end{pmatrix}$	$G - 1 = 2$
Residuals	$W = \begin{pmatrix} 13.408 & 7.723 & 8.675 & 5.864 \\ 7.723 & 8.480 & 7.526 & 6.213 \\ 8.675 & 7.527 & 11.608 & 7.038 \\ 5.864 & 6.213 & 7.038 & 10.566 \end{pmatrix}$	$\sum_{g=1}^G n_g - G = 33$
Total	$B + W = \begin{pmatrix} 14.459 & 9.897 & 7.299 & 5.104 \\ 9.897 & 13.360 & 5.153 & 4.957 \\ 7.299 & 5.153 & 13.990 & 8.422 \\ 5.104 & 4.957 & 8.422 & 11.376 \end{pmatrix}$	$\sum_{g=1}^G n_g - 1 = 35$

Listing 1: R code for Q1 (a)

```
ds <- read.csv("~/Rscripts/MH4501_data_fish.csv")

#Q1 (a)
xbar <- colMeans(ds)[3:6]

ds.group <- split(ds[,3:6], ds$method)

ds.means <- sapply(ds.group, function(x) {
  apply(x, 2, mean)
}, simplify = 'data.frame')
```

```

# B matrix
B <- (12*(ds.means[,1]-xbar)%*%t(ds.means[,1]-xbar)
+ 12*(ds.means[,2]-xbar)%*%t(ds.means[,2]-xbar)
+ 12*(ds.means[,3]-xbar)%*%t(ds.means[,3]-xbar))

# W matrix
W <- matrix(0,4,4)
for(g in 1:3){
  for(i in 1:12){
    xi <- as.matrix(ds[ds$method == g,][i,][3:6])
    W <- W + t(xi-ds.means[,g]) %*% (xi-ds.means[,g])
  }
}

```

(b) Solution.

$$\Lambda^* = \frac{\det(W)}{\det(W+B)} = \frac{1315.113}{5858.295} = 0.224$$

Assume $n = \sum_{g=1}^G n_g = 36$ is large enough, by Bartlett's approximation we have

$$\begin{aligned}
-\left(n-1-\frac{p+G}{2}\right) \ln \Lambda^* &= -\left(36-1-\frac{2+3}{2}\right) \ln(0.224) \\
&= 48.553 \\
&\sim \chi^2(2(3-1)) = \chi^2(4)
\end{aligned}$$

Since $\chi_{0.05}^2[4] = 9.488 < 48.553$, we reject H_0

Listing 2: R code for Q1 (b)

```

#Q1 (b)
ds.manova <- manova(as.matrix(ds[,3:6]) ~ as.factor(ds$method))
ds.summary <- summary(ds.manova)
ds.summary

det(W)
det(B+W)
det(W)/det(B+W)
summary(ds.manova, 'Wilks')$stats[,2][1]

-32.5*log(det(W)/det(B+W))
qchisq(.95, df=4)

```

Question 2.

Solution. For the standardized data set we have the sample covariance matrix of

$$S = \begin{pmatrix} 1 & & & \\ 0.712 & 1 & & \\ 0.513 & 0.377 & 1 & \\ 0.398 & 0.402 & 0.668 & 1 \end{pmatrix}$$

Its eigenvalues and eigenvectors can be calculated that

$$\begin{aligned} \lambda_1 &= 2.537, & e_1 &= (-0.521, -0.491, -0.504, -0.483)^T; \\ \lambda_2 &= 0.850, & e_2 &= (0.437, 0.547, -0.468, -0.539)^T; \\ \lambda_3 &= 0.382, & e_3 &= (0.405, -0.425, 0.558, -0.587)^T; \\ \lambda_4 &= 0.231, & e_4 &= (0.611, -0.528, -0.464, 0.363)^T. \end{aligned}$$

Therefore, the principal components are

$$\begin{aligned} Y_1 &= e_1^T X = -0.521X_1 - 0.491X_2 - 0.504X_3 - 0.483X_4; \\ Y_2 &= e_2^T X = 0.437X_1 + 0.547X_2 - 0.468X_3 - 0.539X_4; \\ Y_3 &= e_3^T X = 0.405X_1 - 0.425X_2 + 0.558X_3 - 0.587X_4; \\ Y_4 &= e_4^T X = 0.611X_1 - 0.528X_2 - 0.464X_3 + 0.363X_4. \end{aligned}$$

The proportion of total variance experienced by each of the sample principle components is following.

$$\begin{aligned} (\% \text{ of variance explained by } Y_1) &= \frac{\lambda_1}{\sum_{j=1}^4 \lambda_j} = 63.42\% \\ (\% \text{ of variance explained by } Y_2) &= \frac{\lambda_2}{\sum_{j=1}^4 \lambda_j} = 21.24\% \\ (\% \text{ of variance explained by } Y_3) &= \frac{\lambda_3}{\sum_{j=1}^4 \lambda_j} = 9.56\% \\ (\% \text{ of variance explained by } Y_4) &= \frac{\lambda_4}{\sum_{j=1}^4 \lambda_j} = 5.78\% \end{aligned}$$

Listing 3: R code for Q2

```
#Q2
standard.df <- scale(ds[,3:6])
pca <- prcomp(standard.df, scale=T)

sample.cov <- (t(standard.df)%*%standard.df)/35
sample.cov
cov(standard.df,standard.df)
ev <- eigen(sample.cov)

# check
pca
(ev$values)^(1/2)
ev$vectors

100*ev$values[1]/sum(ev$values)
100*ev$values[2]/sum(ev$values)
100*ev$values[3]/sum(ev$values)
100*ev$values[4]/sum(ev$values)
```

Question 3.

Solution. By hierarchical clustering on the first 2 sample principal components with euclidean distance and complete linkage, with $k = 3$ we can construct tree and divide it into 3 cluster as depicted in Fig. 1.




Figure 1: Hierarchical Clustering with $k = 3$

By the 3 clusters we obtain, we can construct its confusion matrix as

# observations	Method 1	Method 2	Method 3	Total
Cluster 1	7	5	7	19
Cluster 2	4	5	0	9
Cluster 3	1	2	5	8
Total	12	12	12	36

Listing 4: R code for Q3

```
#Q3
X <- ds[3:6]
pc1 <- c(t(ev$vectors[,1]))%*%t(X)
pc2 <- c(t(ev$vectors[,2]))%*%t(X)

clusters <- hclust(dist(cbind(pc1,pc2),method = "euclidean"))
plot(clusters, cex=0.6)
rect.hclust(clusters, k=3, border=2:4)
clusterCut <- cutree(clusters, 3)
table(clusterCut, ds$method)
```