

Module 3b. Hypothesis testing

Data Science & AI

Sabine De Vreese Stijn Lievens Lieven Smits Bert Van Vreckem
2022–2023

**HO
GENT**

Contents

Testing procedure

Probability Value

Critical Region

Examples

Student's t-test

Errors in Hypothesis Tests

**HO
GENT**

Learning Goals

- Statistical hypothesis testing concepts
- Hypothesis testing procedure
- Apply the z - and t -test

Testing procedure

**HO
GENT**

Statistical Hypothesis Testing

Hypothesis Idea that has yet to be proven: statement regarding the numeric value of a population parameter

Hypothesis Test verification of a statement about the values of one or multiple population parameters

Null Hypothesis (H_0) Base hypothesis, we start with assuming it is true

Alternative Hypothesis (H_1, H_a) Conclusion if the null hypothesis is unlikely to be true

**HO
GENT**

Elements of a testing procedure

Test Statistic The value that is calculated from the sample

Region of Acceptance The region of values supporting the null hypothesis

Critical Region / Region of Rejection The region of values rejecting the null hypothesis

Significance Level The probability of rejecting a true null hypothesis H_0

**HO
GENT**

Testing procedure

1. Formulate both hypotheses (H_0 and H_1)
2. Determine the significance level (α)
3. Calculate the test statistic
4. Determine the critical region or the probability value
5. Draw conclusions

Hypotheses about superheroes



A superhero rescues 3.3 persons a day



Source: <http://www.cracked.com/quick-fixes/4-people-who-saved-day-while-dressed-as-superheroes/>

**HO
GENT**

Assume that, over a period of 30 days, on average 3.483 people were saved each day ($\bar{x} = 3.483$, $n = 30$)

1. Hypothesis: $H_0 : \mu = 3.3$; $H_1 : \mu > 3.3$
2. Significance level: $\alpha = 0.05$
3. Test statistic: $\bar{x} = 3.483$


Standard deviation of the population (assumed to be known): $\sigma = 0.55$

**HO
GENT**

Calculate test statistic

Based on the central limit theorem, we know that:

$$M \sim \text{Nor}(\mu = 3.3; \sigma = \frac{0.55}{\sqrt{30}} = 0.1)$$



**HO
GENT**

Probability Value

**HO
GENT**

Probability Value


p -value

The p -value is the probability, if the null hypothesis is true, to obtain a value for the test statistic that is at least as extreme as the observed value.

- $p\text{-value} < \alpha \Rightarrow$ reject H_0 : the discovered value of \bar{x} is too extreme;
- $p\text{-value} \geq \alpha \Rightarrow$ do not reject H_0 : the discovered value of \bar{x} can still be explained by coincidence.

**HO
GENT**

Probability Value



$$P(M > 3.483) = P\left(Z > \frac{3.483 - 3.3}{\frac{\sigma}{\sqrt{n}}}\right) = P(Z > 1.822) = 0.0344$$

**HO
GENT**

Critical Region

**HO
GENT**

Critical Region

Critical region

The **critical region** is the collection of all values of the test statistic for which we can reject the null hypothesis.

We look for a critical value g for which:

$$P(M > g) = \alpha$$


Determine z_α for which:

$$P(Z > z_\alpha) = \alpha \Rightarrow g = \mu + z_\alpha \cdot \frac{\sigma}{\sqrt{n}}$$

- Left of g : region of acceptance (do not reject H_0)
- Right of g : critical region (reject H_0)

**HO
GENT**

Critical Region



significance level $\alpha = 0.05 \Rightarrow$ critical value $z_{\alpha} = 1.645$
($z_{\alpha} = \text{stats.norm.isf}(1 - 0.95)$)

**HO
GENT**

Left-tailed testing

What would you have to change in the equation in order to calculate the correct critical value?

Left-tailed testing

What would you have to change in the equation in order to calculate the correct critical value? Answer:

$$g = \mu - z \times \frac{\sigma}{\sqrt{n}}$$

because

$$P(M < g) = P\left(Z < \frac{g - \mu}{\frac{\sigma}{\sqrt{n}}}\right) = 0.05$$

Left-tailed testing

Because of symmetry:

$$P\left(Z > -\left(\frac{g - \mu}{\frac{\sigma}{\sqrt{n}}}\right)\right) = 0.05$$

The corresponding z-value is 1.645, and therefore:

$$\begin{aligned} z = \frac{-g + \mu}{\frac{\sigma}{\sqrt{n}}} &\Leftrightarrow -g = \frac{\sigma}{\sqrt{n}} z_{\alpha} - \mu \\ &\Leftrightarrow g = \mu - z_{\alpha} \frac{\sigma}{\sqrt{n}} \end{aligned}$$

Two-tailed testing

Sometimes it can be necessary to perform a two-tailed test. In this case, two critical values need to be calculated, namely the left and right critical value.

$$g = \mu \pm z_{\alpha/2} \times \frac{\sigma}{\sqrt{n}} \quad (1)$$

Summary

Goal	Test regarding the value of the population mean μ using a sample of n independent values		
Prerequisite	De population has a random distribution, n is sufficiently large		
Test Type	Two-tailed	Left-tailed	Right-tailed
H_0	$\mu = \mu_0$	$\mu = \mu_0$	$\mu = \mu_0$
H_1	$\mu \neq \mu_0$	$\mu < \mu_0$	$\mu > \mu_0$
Critical Region	$ \bar{x} > g$	$\bar{x} < -g$	$\bar{x} > g$
Test statistic	$z = \frac{\bar{x} - \mu_0}{\frac{\sigma}{\sqrt{n}}}$		

Table: Summary of Testing Procedures

Requirements for z-test

- The sample needs to be random
- The sample size needs to be sufficiently large ($n \geq 30$)
- The test statistic needs to have a normal distribution
- The standard deviation of the population, σ , is known

Sometimes these assumptions will not hold and in this case we can *not* use the Z-test!

Examples

**HO
GENT**

Example 1

When drawing a random sample consisting of 50 observations from a population with variance $\sigma^2 = 55$ we obtain as sample mean $\bar{x} = 25$. We now want to find out if there is a reason to assume that the population mean is smaller than 27.

Example 1

- 1 Formulate both hypotheses
 $H_0 : \mu = 27$ en $H_1 : \mu < 27$.
- 2 Determine significance level α and sample size n
 $\alpha = 0.05$ en $n = 50$
- 3 Test statistic & value: sample mean $\bar{x} = 25$

Example 1 (cont.)

4a Probability Value

According to the central limit theorem:

$$M \sim \text{Nor}(\mu = 27, \frac{\sigma^2}{\sqrt{n}})$$

$$z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}} = \frac{25 - 27}{\sqrt{\frac{55}{50}}} \approx -1.91$$

We therefore have a probability value of 0.0281.

Using a significance level of 0.05, we can reject H_0 .

**HO
GENT**

Example 1 (cont.)


4b Calculate and plot the critical region

$$\begin{aligned}g &= \mu - z \times \frac{\sigma}{\sqrt{n}} \\&= 27 - 1.645 \times \sqrt{\frac{\sigma}{n}} \\&= 25.27470944\end{aligned}$$

We have that $\bar{x} < g$ and therefore we can reject H_0 .

HO
GENT

Example 1 (cont.)



**HO
GENT**

Example 1 (cont.)

5 Conclusion

We can conclude, based on the sample, that $\mu < 27$ for a significance level of 0.05

**HO
GENT**

Example 2

In a research about the amount of change in the pockets of our superheroes, researchers state that on average a superhero carries €25 of cash. We assume that the standard deviation of the population $\sigma = 7$. For a random sample of size $n = 64$, the average amount of money a superhero carries is $\bar{x} = €23$. For the significance level, $\alpha = 0.05$ is selected.

Example 2

- 1 Formulate both hypotheses
 $H_0 : \mu = 25$ en $H_1 : \mu \neq 25$
- 2 Determine significance level α and sample size n
 $\alpha = 0.05$ en $n = 64$.
- 3 Test statistic & value: $\bar{x} = 23$

Example 2 (cont.)

4b Calculate the critical region

$$g_1 = \mu - z \times \frac{\sigma}{\sqrt{n}} = 23.28$$

$$g_2 = \mu + z \times \frac{\sigma}{\sqrt{n}} = 26.72$$

We have that \bar{x} is inside the critical region (because $23 < 23.28$) so we can reject H_0 .

5 Based on this sample we can conclude that superheroes carry on average *less* than 25 €, using a significance level of 5%

**HO
GENT**

Student's t-test

**HO
GENT**

Student's t-test

What if the requirements for a z-test are not met? E.g.

- Sample size too small
- Population stdev (σ) unknown

If the variable is normally distributed, you can use the *t*-test

The *t*-test

Determine critical value:

$$g = \mu \pm t \times \frac{s}{\sqrt{n}}$$

- *t*-value is derived from the Student's *t*-distribution, based on the number of *degrees of freedom*, $n - 1$
- Look for value using the function `t.isf()` in Python
- Apart from this, the procedure is identical to the procedure of the *z*-test

**HO
GENT**

Errors in Hypothesis Tests

HO
GENT

Errors in Hypothesis Tests

Conclusion	Reality	
	H_0 True	H_1 True
H_0 not rejected	Correct inference	Type II error (false negative)
H_0 rejected	Type I error (false positive)	Correct inference

P(type I error) = α (= significance level)

P(type II error) = β

Calculating β is **not** trivial, but if $\alpha \searrow$ then $\beta \nearrow$

HO
GENT