

Consider the optimization problem:

$$\begin{aligned} & \min_{\mathbf{x}, \mathbf{y}, \mathbf{z}} \frac{1}{2} \|\mathbf{y} - \mathbf{D}\mathbf{x}\|_2^2 + \lambda \|\mathbf{z}\|_1 \\ & \text{subject to } \mathbf{R}^{-1}\mathbf{z} - \mathbf{R}^{-1}\mathbf{x} = 0 \\ & \mathbf{W}(\mathbf{s} - \mathbf{y}) = 0 \end{aligned} \quad (1)$$

This optimization problem has the Lagrangian function:

$$\mathcal{L}(\mathbf{x}, \mathbf{y}, \mathbf{z}, \gamma, \eta) = \frac{1}{2} \|\mathbf{y} - \mathbf{D}\mathbf{x}\|_2^2 + \lambda \|\mathbf{z}\|_1 + \gamma^H \mathbf{R}^{-1}(\mathbf{z} - \mathbf{x}) + \eta^H \mathbf{W}(\mathbf{s} - \mathbf{y}) \quad (2)$$

Now, we can augment the Lagrangian.

$$\mathcal{L}_\rho(\mathbf{x}, \mathbf{y}, \mathbf{z}, \eta, \gamma) = \frac{1}{2} \|\mathbf{y} - \mathbf{D}\mathbf{x}\|_2^2 + \lambda \|\mathbf{z}\|_1 + \gamma^H \mathbf{R}^{-1}(\mathbf{z} - \mathbf{x}) + \eta^H \mathbf{W}(\mathbf{s} - \mathbf{y}) + \frac{\rho}{2} \|\mathbf{R}^{-1}(\mathbf{z} - \mathbf{x})\|_2^2 + \frac{\rho}{2} \|\mathbf{W}(\mathbf{s} - \mathbf{y})\|_2^2 \quad (3)$$

$$\nabla_{\mathbf{x}} \mathcal{L}_\rho(\mathbf{x}, \mathbf{y}, \mathbf{z}, \eta, \gamma) = -\mathbf{R}^{-1}\gamma - \mathbf{D}^H \mathbf{y} + \mathbf{D}^H \mathbf{D}\mathbf{x} + \rho \mathbf{R}^{-2}\mathbf{x} - \rho \mathbf{R}^{-2}\mathbf{z} \quad (4)$$

For $\mathbf{x}, \mathbf{y}, \mathbf{z}, \eta, \gamma$ such that $\nabla_{\mathbf{x}} \mathcal{L}_\rho(\mathbf{x}, \mathbf{y}, \mathbf{z}, \eta, \gamma) = 0$:

$$(\mathbf{R}^{-2} + \frac{\mathbf{D}^H \mathbf{D}}{\rho})\mathbf{x} = \mathbf{R}^{-2}\mathbf{z} + \frac{\mathbf{R}^{-1}\gamma}{\rho} + \frac{\mathbf{D}^H \mathbf{y}}{\rho} \quad (5)$$

$$\mathbf{R}^{-1}(\rho \mathbf{I} + (\mathbf{D}\mathbf{R})^H (\mathbf{D}\mathbf{R}))\mathbf{R}^{-1}\mathbf{x} = \rho \mathbf{R}^{-2}\mathbf{z} + \mathbf{R}^{-1}\gamma + \mathbf{D}^H \mathbf{y} \quad (6)$$

$$(\rho \mathbf{I} + (\mathbf{D}\mathbf{R})^H (\mathbf{D}\mathbf{R}))\mathbf{R}^{-1}\mathbf{x} = \rho \mathbf{R}^{-1}\mathbf{z} + \gamma + (\mathbf{D}\mathbf{R})^H \mathbf{y} \quad (7)$$

$$\mathbf{R}^{-1}\mathbf{x} = (\rho \mathbf{I} + (\mathbf{D}\mathbf{R})^H (\mathbf{D}\mathbf{R}))^{-1}(\rho \mathbf{R}^{-1}\mathbf{z} + \gamma + (\mathbf{D}\mathbf{R})^H \mathbf{y}) \quad (8)$$

$$\min_{\mathbf{x}} \mathcal{L}_\rho(\mathbf{x}, \mathbf{y}, \mathbf{z}, \eta, \gamma) = \mathbf{R}(\rho \mathbf{I} + (\mathbf{D}\mathbf{R})^H (\mathbf{D}\mathbf{R}))^{-1}(\rho \mathbf{R}^{-1}\mathbf{z} + \gamma + (\mathbf{D}\mathbf{R})^H \mathbf{y}) \quad (9)$$

$$\nabla_{\mathbf{y}} \mathcal{L}_\rho(\mathbf{x}, \mathbf{y}, \mathbf{z}, \eta, \gamma) = \mathbf{y} - \mathbf{D}\mathbf{x} - \mathbf{W}^H \eta + \rho \mathbf{W}^H \mathbf{W}\mathbf{y} - \rho \mathbf{W}^H \mathbf{W}\mathbf{s} \quad (10)$$

$$\min_{\mathbf{y}} \mathcal{L}_\rho(\mathbf{x}, \mathbf{y}, \mathbf{z}, \eta, \gamma) = (\mathbf{I} + \rho \mathbf{W}^T \mathbf{W})^{-1}(\rho \mathbf{W}^H \mathbf{W}\mathbf{s} + \mathbf{W}^H \eta + \mathbf{D}\mathbf{x}) \quad (11)$$

$$\min_{\mathbf{y}} \mathcal{L}_{\rho, \Lambda}(\mathbf{x}, \mathbf{y}, \mathbf{z}, \eta, \gamma) = \begin{cases} \frac{1}{1+\rho}(\rho \mathbf{s} + \mathbf{D}\mathbf{x} + \eta) & \text{within signal domain} \\ \mathbf{D}\mathbf{x} & \text{outside signal domain} \end{cases} \quad (12)$$

$$\min_{\mathbf{z}} L_{\rho}(\mathbf{x}, \mathbf{y}, \mathbf{z}, \boldsymbol{\eta}, \boldsymbol{\gamma}) = S_{\frac{\lambda \mathbf{R}^2}{\rho}}(\mathbf{x} - \frac{\mathbf{R}\boldsymbol{\gamma}}{\rho}) \quad (13)$$

From this, we can get the update equations:

$$\mathbf{x}^{(k+1)} = \mathbf{R}(\rho \mathbf{I} + (\mathbf{D}\mathbf{R})^H(\mathbf{D}\mathbf{R}))^{-1}(\rho \mathbf{R}^{-1} \mathbf{z}^{(k)} + \boldsymbol{\gamma}^{(k)} + (\mathbf{D}\mathbf{R})^H \mathbf{y}^{(k)}) \quad (14)$$

$$\mathbf{y}^{(k+1)} = \begin{cases} \frac{1}{1+\rho}(\rho \mathbf{s} + \mathbf{D}\mathbf{x}^{(k+1)} + \boldsymbol{\eta}^{(k)}) & \text{within signal domain} \\ \mathbf{D}\mathbf{x}^{(k+1)} & \text{outside signal domain} \end{cases} \quad (15)$$

$$\mathbf{z}^{(k+1)} = S_{\frac{\lambda \mathbf{R}^2}{\rho}}(\mathbf{x}^{(k+1)} - \frac{\mathbf{R}\boldsymbol{\gamma}^{(k)}}{\rho}) \quad (16)$$

$$\boldsymbol{\gamma}^{(k+1)} = \boldsymbol{\gamma}^{(k)} + \rho \mathbf{R}^{-1}(\mathbf{z}^{(k+1)} - \mathbf{x}^{(k+1)}) \quad (17)$$

$$\boldsymbol{\eta}^{(k+1)} = \boldsymbol{\eta}^{(k)} + \rho(\mathbf{W}(\mathbf{s} - \mathbf{y}^{(k+1)})) \quad (18)$$

However, these equations will be simpler using slightly different updates.

$$\mathbf{R}^{-1} \mathbf{x}^{(k+1)} = (\rho \mathbf{I} + (\mathbf{D}\mathbf{R})^H(\mathbf{D}\mathbf{R}))^{-1}(\rho \mathbf{R}^{-1} \mathbf{z}^{(k)} + \boldsymbol{\gamma}^{(k)} + (\mathbf{D}\mathbf{R})^H \mathbf{y}^{(k)}) \quad (19)$$

$$\mathbf{y}^{(k+1)} = \begin{cases} \frac{1}{1+\rho}(\rho \mathbf{s} + \mathbf{D}\mathbf{R}\mathbf{R}^{-1} \mathbf{x}^{(k+1)} + \boldsymbol{\eta}^{(k)}) & \text{within signal domain} \\ \mathbf{D}\mathbf{R}\mathbf{R}^{-1} \mathbf{x}^{(k+1)} & \text{outside signal domain} \end{cases} \quad (20)$$

$$\mathbf{R}^{-1} \mathbf{z}^{(k+1)} = S_{\frac{\lambda \mathbf{R}}{\rho}}(\mathbf{R}^{-1} \mathbf{x}^{(k+1)} - \frac{\boldsymbol{\gamma}^{(k)}}{\rho}) \quad (21)$$

$$\boldsymbol{\gamma}^{(k+1)} = \boldsymbol{\gamma}^{(k)} + \rho(\mathbf{R}^{-1} \mathbf{z}^{(k+1)} - \mathbf{R}^{-1} \mathbf{x}^{(k+1)}) \quad (22)$$

$$\boldsymbol{\eta}^{(k+1)} = \boldsymbol{\eta}^{(k)} + \rho(\mathbf{W}(\mathbf{s} - \mathbf{y}^{(k+1)})) \quad (23)$$

In the above equations, the dictionary never appears unscaled (without \mathbf{R}), and the coefficients are always scaled by \mathbf{R}^{-1} . Unfortunately, \mathbf{R} still must be calculated for the \mathbf{z} -update.

For reference while coding, I would like to adjust into SPORCO package notation. I can add an S subscript to prevent confusion.

$$\mathbf{x}_S^{(k)} = \mathbf{R}^{-1} \mathbf{x}^{(k)} \quad (24)$$

$$\mathbf{y}_S^{(k)} = \begin{bmatrix} \mathbf{W}^H \mathbf{y}^{(k)} \\ \mathbf{R}^{-1} \mathbf{z}^{(k)} \end{bmatrix} \quad (25)$$

$$\mathbf{u}_S^{(k)} = \begin{bmatrix} \mathbf{W}^H \boldsymbol{\eta}^{(k)} \\ \boldsymbol{\gamma}^{(k)} \end{bmatrix} \quad (26)$$

$$\mathbf{A}_S = \begin{bmatrix} \mathbf{0} \\ -\mathbf{I} \end{bmatrix} \quad (27)$$

$$\mathbf{B}_S = \begin{bmatrix} -\mathbf{I} & \mathbf{0} \\ \mathbf{0} & \mathbf{I} \end{bmatrix} = \begin{bmatrix} -\mathbf{P}_0 & \mathbf{P}_1 \end{bmatrix} \quad (28)$$

$$\mathbf{C}_S = \begin{bmatrix} \mathbf{s} \\ \mathbf{0} \end{bmatrix} \quad (29)$$

$$\mathbf{Q}_S = \rho \mathbf{I} + (\mathbf{D}\mathbf{R})^H (\mathbf{D}\mathbf{R}) \quad (30)$$

$$\mathbf{D}_S = \mathbf{D}\mathbf{R} \quad (31)$$

$$\mathbf{x}_S^{(k+1)} = \mathbf{Q}_S^{-1} (\rho \mathbf{P}_1 \mathbf{y}_S^{(k)} + \mathbf{P}_1 \mathbf{u}_S^{(k)} + \mathbf{D}_S^H \mathbf{P}_0 \mathbf{y}_S^{(k)}) \quad (32)$$

$$\mathbf{y}_S^{(k+1)} = \begin{bmatrix} \begin{cases} \frac{1}{1+\rho} (\rho \mathbf{s} + \mathbf{D}_S \mathbf{x}_S^{(k+1)} + \mathbf{P}_0 \mathbf{u}_S^{(k)}) & \text{within signal domain} \\ \mathbf{D}_S \mathbf{x}_S^{(k+1)} & \text{outside signal domain} \end{cases} \\ \mathbf{S}_{\frac{\lambda \mathbf{R}}{\rho}}(\mathbf{x}_S^{(k+1)} - \frac{\mathbf{P}_1 \mathbf{u}_S^{(k)}}{\rho}) \end{bmatrix} \quad (33)$$

$$\mathbf{u}_S^{(k+1)} = \mathbf{u}_S^{(k)} + \rho (\mathbf{A}_S \mathbf{x}_S^{(k+1)} + \mathbf{B}_S \mathbf{y}_S^{(k+1)} + \mathbf{C}_S) \quad (34)$$