

Class 14: RNA-Seq analysis mini-project

Lena (A16420052)

Differential Expression Analysis

The authors report on differential analysis of lung fibroblasts in response to loss of the developmental transcription factor HOXA1

Data Import

Read our counts and metadata CSV files

```
library(DESeq2)
```

```
Loading required package: S4Vectors
```

```
Loading required package: stats4
```

```
Loading required package: BiocGenerics
```

```
Attaching package: 'BiocGenerics'
```

```
The following objects are masked from 'package:stats':
```

```
IQR, mad, sd, var, xtabs
```

The following objects are masked from 'package:base':

```
anyDuplicated, aperm, append, as.data.frame, basename, cbind,  
colnames, dirname, do.call, duplicated, eval, evalq, Filter, Find,  
get, grep, grepl, intersect, is.unsorted, lapply, Map, mapply,  
match, mget, order, paste, pmax, pmax.int, pmin, pmin.int,  
Position, rank, rbind, Reduce, rownames, sapply, setdiff, sort,  
table, tapply, union, unique, unsplit, which.max, which.min
```

Attaching package: 'S4Vectors'

The following object is masked from 'package:utils':

```
findMatches
```

The following objects are masked from 'package:base':

```
expand.grid, I, unname
```

Loading required package: IRanges

Loading required package: GenomicRanges

Loading required package: GenomeInfoDb

Loading required package: SummarizedExperiment

Loading required package: MatrixGenerics

Loading required package: matrixStats

Attaching package: 'MatrixGenerics'

The following objects are masked from 'package:matrixStats':

```
colAlls, colAnyNAs, colAnys, colAvgsPerRowSet, colCollapse,
colCounts, colCummaxs, colCummins, colCumprods, colCumsums,
colDiffs, colIQRDiffs, colIQRs, colLogSumExps, colMadDiffs,
colMads, colMaxs, colMeans2, colMedians, colMins, colOrderStats,
colProds, colQuantiles, colRanges, colRanks, colSdDiffs, colSds,
colSums2, colTabulates, colVarDiffs, colVars, colWeightedMads,
colWeightedMeans, colWeightedMedians, colWeightedSds,
colWeightedVars, rowAlls, rowAnyNAs, rowAnys, rowAvgsPerColSet,
rowCollapse, rowCounts, rowCummaxs, rowCummins, rowCumprods,
rowCumsums, rowDiffs, rowIQRDiffs, rowIQRs, rowLogSumExps,
rowMadDiffs, rowMads, rowMaxs, rowMeans2, rowMedians, rowMins,
rowOrderStats, rowProds, rowQuantiles, rowRanges, rowRanks,
rowSdDiffs, rowSds, rowSums2, rowTabulates, rowVarDiffs, rowVars,
rowWeightedMads, rowWeightedMeans, rowWeightedMedians,
rowWeightedSds, rowWeightedVars
```

Loading required package: Biobase

Welcome to Bioconductor

```
Vignettes contain introductory material; view with
'browseVignettes()'. To cite Bioconductor, see
'citation("Biobase")', and for packages 'citation("pkgname")'.
```

Attaching package: 'Biobase'

The following object is masked from 'package:MatrixGenerics':

```
rowMedians
```

The following objects are masked from 'package:matrixStats':

```
anyMissing, rowMedians
```

```
metaFile <- "GSE37704_metadata.csv"
countFile <- "GSE37704_featurecounts.csv"
```

```
# Import metadata and take a peak
colData = read.csv(metaFile, row.names=1)
head(colData)
```

```
          condition
SRR493366 control_sirna
SRR493367 control_sirna
SRR493368 control_sirna
SRR493369      hoxa1_kd
SRR493370      hoxa1_kd
SRR493371      hoxa1_kd
```

```
# Import countdata
countData = read.csv(countFile, row.names=1)
head(countData)
```

```
          length SRR493366 SRR493367 SRR493368 SRR493369 SRR493370
ENSG00000186092    918         0         0         0         0         0
ENSG00000279928    718         0         0         0         0         0
ENSG00000279457   1982        23        28        29        29        28
ENSG00000278566    939         0         0         0         0         0
ENSG00000273547    939         0         0         0         0         0
ENSG00000187634   3214       124       123       205       207       212
          SRR493371
ENSG00000186092         0
ENSG00000279928         0
ENSG00000279457        46
ENSG00000278566         0
ENSG00000273547         0
ENSG00000187634       258
```

Q1. Complete the code below to remove the troublesome first column from count-Data

```
# Note we need to remove the odd first $length col
countData <- as.matrix(countData[,-1])
head(countData)
```

```
          SRR493366 SRR493367 SRR493368 SRR493369 SRR493370 SRR493371
ENSG00000186092         0         0         0         0         0         0
```

ENSG00000279928	0	0	0	0	0	0
ENSG00000279457	23	28	29	29	28	46
ENSG00000278566	0	0	0	0	0	0
ENSG00000273547	0	0	0	0	0	0
ENSG00000187634	124	123	205	207	212	258

Q2. Complete the code below to filter countData to exclude genes (i.e. rows) where we have 0 read count across all samples (i.e. columns).

```
# Filter count data where you have 0 read count across all samples.
countData <- countData[rowSums(countData)!= 0, ]
head(countData)
```

	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370	SRR493371
ENSG00000279457	23	28	29	29	28	46
ENSG00000187634	124	123	205	207	212	258
ENSG00000188976	1637	1831	2383	1226	1326	1504
ENSG00000187961	120	153	180	236	255	357
ENSG00000187583	24	48	65	44	48	64
ENSG00000187642	4	9	16	14	16	16

```
nrow(countData)
```

```
[1] 15975
```

DEseq setup and analysis

Running DESeq2

```
dds = DESeqDataSetFromMatrix(countData=countData,
                              colData=colData,
                              design=~condition)
```

Warning in DESeqDataSet(se, design = design, ignoreRank): some variables in design formula are characters, converting to factors

```
dds = DESeq(dds)
```

estimating size factors

estimating dispersions

gene-wise dispersion estimates

mean-dispersion relationship

final dispersion estimates

fitting model and testing

```
res = results(dds, contrast=c("condition", "hoxa1_kd", "control_sirna"))
```

Q3. Call the `summary()` function on your results to get a sense of how many genes are up or down-regulated at the default 0.1 p-value cutoff.


```
summary(res)
```

```
out of 15975 with nonzero total read count
adjusted p-value < 0.1
LFC > 0 (up)      : 4349, 27%
LFC < 0 (down)    : 4396, 28%
outliers [1]      : 0, 0%
low counts [2]    : 1237, 7.7%
(mean count < 0)
[1] see 'cooksCutoff' argument of ?results
[2] see 'independentFiltering' argument of ?results
```

Volcano plot

Result visualization

```
plot( res$log2FoldChange, -log(res$padj) )
```



Q4. Improve this plot by completing the below code, which adds color and axis labels

```
# Make a color vector for all genes
mycols <- rep("gray", nrow(res))

# Color red the genes with absolute fold change above 2
mycols[ abs(res$log2FoldChange) > 2 ] <- "red"

# Color blue those with adjusted p-value less than 0.01
# and absolute fold change more than 2
inds <- (res$padj<0.01) & (abs(res$log2FoldChange) > 2 )
mycols[ inds ] <- "blue"

plot( res$log2FoldChange, -log(res$padj), col=mycols, xlab="Log2(FoldChange)", ylab="-Log(
```



Adding gene annotation

Geneset Enrichment

Q5. Use the `mapIDs()` function multiple times to add SYMBOL, ENTREZID and GENENAME annotation to our results by completing the code below.

```
library("AnnotationDbi")
```

Warning: package 'AnnotationDbi' was built under R version 4.3.2

```
library("org.Hs.eg.db")
```

```
columns(org.Hs.eg.db)
```


[1]	"ACCNUM"	"ALIAS"	"ENSEMBL"	"ENSEMBLPROT"	"ENSEMBLTRANS"
[6]	"ENTREZID"	"ENZYME"	"EVIDENCE"	"EVIDENCEALL"	"GENENAME"
[11]	"GENETYPE"	"GO"	"GOALL"	"IPI"	"MAP"
[16]	"OMIM"	"ONTOLOGY"	"ONTOLOGYALL"	"PATH"	"PFAM"
[21]	"PMID"	"PROSITE"	"REFSEQ"	"SYMBOL"	"UCSCKG"
[26]	"UNIPROT"				

```
res$symbol = mapIds(org.Hs.eg.db,
  keys= row.names(res),
  keytype="ENSEMBL",
  column= "SYMBOL",
  multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
res$entrez = mapIds(org.Hs.eg.db,
  keys=row.names(res),
  keytype="ENSEMBL",
  column="ENTREZID",
  multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
res$name = mapIds(org.Hs.eg.db,
  keys=row.names(res),
  keytype="ENSEMBL",
  column="GENENAME",
  multiVals="first")
```

'select()' returned 1:many mapping between keys and columns

```
head(res, 10)
```

log2 fold change (MLE): condition hoxa1_kd vs control_sirna

Wald test p-value: condition hoxa1 kd vs control sirna

DataFrame with 10 rows and 9 columns

	baseMean	log2FoldChange	lfcSE	stat	pvalue
--	----------	----------------	-------	------	--------

	<numeric>	<numeric>	<numeric>	<numeric>	<numeric>
ENSG00000279457	29.913579	0.1792571	0.3248216	0.551863	5.81042e-01
ENSG00000187634	183.229650	0.4264571	0.1402658	3.040350	2.36304e-03
ENSG00000188976	1651.188076	-0.6927205	0.0548465	-12.630158	1.43990e-36
ENSG00000187961	209.637938	0.7297556	0.1318599	5.534326	3.12428e-08
ENSG00000187583	47.255123	0.0405765	0.2718928	0.149237	8.81366e-01
ENSG00000187642	11.979750	0.5428105	0.5215598	1.040744	2.97994e-01
ENSG00000188290	108.922128	2.0570638	0.1969053	10.446970	1.51282e-25
ENSG00000187608	350.716868	0.2573837	0.1027266	2.505522	1.22271e-02
ENSG00000188157	9128.439422	0.3899088	0.0467163	8.346304	7.04321e-17
ENSG00000237330	0.158192	0.7859552	4.0804729	0.192614	8.47261e-01
	padj	symbol	entrez	name	
	<numeric>	<character>	<character>	<character>	
ENSG00000279457	6.86555e-01	NA	NA	NA	
ENSG00000187634	5.15718e-03	SAMD11	148398	sterile alpha motif ..	
ENSG00000188976	1.76549e-35	NOC2L	26155	NOC2 like nucleolar ..	
ENSG00000187961	1.13413e-07	KLHL17	339451	kelch like family me..	
ENSG00000187583	9.19031e-01	PLEKHN1	84069	pleckstrin homology ..	
ENSG00000187642	4.03379e-01	PERM1	84808	PPARGC1 and ESRR ind..	
ENSG00000188290	1.30538e-24	HES4	57801	hes family bHLH tran..	
ENSG00000187608	2.37452e-02	ISG15	9636	ISG15 ubiquitin like..	
ENSG00000188157	4.21963e-16	AGRN	375790	agrin	
ENSG00000237330	NA	RNF223	401934	ring finger protein ..	

Q6. Finally for this section let's reorder these results by adjusted p-value and save them to a CSV file in your current project directory.

Save results

```
res = res[order(res$pvalue),]
write.csv(res, file = "deseq_results.csv")
```

Pathway Analysis

Use the gage package for pathway analysis and the pathview package to draw pathway diagrams

```
library(pathview)
library(gage)
library(gageData)
```

```

data(kegg.sets.hs)
data(sigmet.idx.hs)

# Focus on signaling and metabolic pathways only
kegg.sets.hs = kegg.sets.hs[sigmet.idx.hs]

# Examine the first 3 pathways
head(kegg.sets.hs, 3)

```

```
$`hsa00232 Caffeine metabolism`
```

```
[1] "10" "1544" "1548" "1549" "1553" "7498" "9"
```

```
$`hsa00983 Drug metabolism - other enzymes`
```

```

[1] "10" "1066" "10720" "10941" "151531" "1548" "1549" "1551"
[9] "1553" "1576" "1577" "1806" "1807" "1890" "221223" "2990"
[17] "3251" "3614" "3615" "3704" "51733" "54490" "54575" "54576"
[25] "54577" "54578" "54579" "54600" "54657" "54658" "54659" "54963"
[33] "574537" "64816" "7083" "7084" "7172" "7363" "7364" "7365"
[41] "7366" "7367" "7371" "7372" "7378" "7498" "79799" "83549"
[49] "8824" "8833" "9" "978"

```

```
$`hsa00230 Purine metabolism`
```

```

[1] "100" "10201" "10606" "10621" "10622" "10623" "107" "10714"
[9] "108" "10846" "109" "111" "11128" "11164" "112" "113"
[17] "114" "115" "122481" "122622" "124583" "132" "158" "159"
[25] "1633" "171568" "1716" "196883" "203" "204" "205" "221823"
[33] "2272" "22978" "23649" "246721" "25885" "2618" "26289" "270"
[41] "271" "27115" "272" "2766" "2977" "2982" "2983" "2984"
[49] "2986" "2987" "29922" "3000" "30833" "30834" "318" "3251"
[57] "353" "3614" "3615" "3704" "377841" "471" "4830" "4831"
[65] "4832" "4833" "4860" "4881" "4882" "4907" "50484" "50940"
[73] "51082" "51251" "51292" "5136" "5137" "5138" "5139" "5140"
[81] "5141" "5142" "5143" "5144" "5145" "5146" "5147" "5148"
[89] "5149" "5150" "5151" "5152" "5153" "5158" "5167" "5169"
[97] "51728" "5198" "5236" "5313" "5315" "53343" "54107" "5422"
[105] "5424" "5425" "5426" "5427" "5430" "5431" "5432" "5433"
[113] "5434" "5435" "5436" "5437" "5438" "5439" "5440" "5441"
[121] "5471" "548644" "55276" "5557" "5558" "55703" "55811" "55821"
[129] "5631" "5634" "56655" "56953" "56985" "57804" "58497" "6240"
[137] "6241" "64425" "646625" "654364" "661" "7498" "8382" "84172"
[145] "84265" "84284" "84618" "8622" "8654" "87178" "8833" "9060"

```

```
[153] "9061"    "93034"   "953"     "9533"    "954"     "955"     "956"     "957"
[161] "9583"    "9615"
```

`gage()` function requires a named vector of fold changes, where the names of the values are the Entrez gene IDs.

```
foldchanges = res$log2FoldChange
names(foldchanges) = res$entrez
head(foldchanges)
```

```
      1266      54855      1465      51232      2034      2317
-2.422719  3.201955 -2.313738 -2.059631 -1.888019 -1.649792
```

Running gage pathway analysis...

```
# Get the results
keggres = gage(foldchanges, gsets=kegg.sets.hs)
```

The value of `same.dir` determines whether to test for changes in a gene set toward a single direction (all genes up or down regulated) or changes towards both directions simultaneously (i.e. any genes in the pathway dysregulated)

```
attributes(keggres)
```

```
$names
[1] "greater" "less"    "stats"
```

Data matrix with gene sets as rows sorted by p-value

```
# Look at the first few down (less) pathways
head(keggres$less)
```

	p.geomean	stat.mean	p.val
hsa04110 Cell cycle	8.995727e-06	-4.378644	8.995727e-06
hsa03030 DNA replication	9.424076e-05	-3.951803	9.424076e-05
hsa03013 RNA transport	1.375901e-03	-3.028500	1.375901e-03
hsa03440 Homologous recombination	3.066756e-03	-2.852899	3.066756e-03
hsa04114 Oocyte meiosis	3.784520e-03	-2.698128	3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis	8.961413e-03	-2.405398	8.961413e-03

	q.val	set.size	exp1
hsa04110 Cell cycle	0.001448312	121	8.995727e-06
hsa03030 DNA replication	0.007586381	36	9.424076e-05
hsa03013 RNA transport	0.073840037	144	1.375901e-03
hsa03440 Homologous recombination	0.121861535	28	3.066756e-03
hsa04114 Oocyte meiosis	0.121861535	102	3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis	0.212222694	53	8.961413e-03

```
#look at up (greater) pathways
head(keggres$greater)
```

	p.geomean	stat.mean	p.val
hsa04640 Hematopoietic cell lineage	0.002822776	2.833362	0.002822776
hsa04630 Jak-STAT signaling pathway	0.005202070	2.585673	0.005202070
hsa00140 Steroid hormone biosynthesis	0.007255099	2.526744	0.007255099
hsa04142 Lysosome	0.010107392	2.338364	0.010107392
hsa04330 Notch signaling pathway	0.018747253	2.111725	0.018747253
hsa04916 Melanogenesis	0.019399766	2.081927	0.019399766

	q.val	set.size	exp1
hsa04640 Hematopoietic cell lineage	0.3893570	55	0.002822776
hsa04630 Jak-STAT signaling pathway	0.3893570	109	0.005202070
hsa00140 Steroid hormone biosynthesis	0.3893570	31	0.007255099
hsa04142 Lysosome	0.4068225	118	0.010107392
hsa04330 Notch signaling pathway	0.4391731	46	0.018747253
hsa04916 Melanogenesis	0.4391731	90	0.019399766


Pathway plot with our RNA-Seq expression results shown in color with pathview()

```
pathview(gene.data=foldchanges, pathway.id="hsa04110")
```

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/lenayang/Downloads/bimm 143/Class 14

Info: Writing image file hsa04110.pathview.png



```
# A different PDF based output of the same data
pathview(gene.data=foldchanges, pathway.id="hsa04110", kegg.native=FALSE)
```

'select()' returned 1:1 mapping between keys and columns

Warning: reconcile groups sharing member nodes!

```
[,1] [,2]
[1,] "9"  "300"
[2,] "9"  "306"
```

Info: Working in directory /Users/lenayang/Downloads/bimm 143/Class 14

Info: Writing image file hsa04110.pathview.pdf

```
## Focus on top 5 upregulated pathways here for demo purposes only  
keggrespathways <- rownames(keggres$greater)[1:5]
```

```
# Extract the 8 character long IDs part of each string  
keggresids = substr(keggrespathways, start=1, stop=8)  
keggresids
```

```
[1] "hsa04640" "hsa04630" "hsa00140" "hsa04142" "hsa04330"
```

```
pathview(gene.data=foldchanges, pathway.id=keggresids, species="hsa")
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/lenayang/Downloads/bimm 143/Class 14
```

```
Info: Writing image file hsa04640.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/lenayang/Downloads/bimm 143/Class 14
```

```
Info: Writing image file hsa04630.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/lenayang/Downloads/bimm 143/Class 14
```

```
Info: Writing image file hsa00140.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/lenayang/Downloads/bimm 143/Class 14
```


```
Info: Writing image file hsa04142.pathview.png
```


```
Info: some node width is different from others, and hence adjusted!
```


'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/lenayang/Downloads/bimm 143/Class 14

Info: Writing image file hsa04330.pathview.png







Q7. Can you do the same procedure as above to plot the pathview figures for the top 5 down-regulated pathways?

```
keggrespathways <- rownames(keggres$less)[1:5]
```

```
# Extract the 8 character long IDs part of each string
keggresids = substr(keggrespathways, start=1, stop=8)
keggresids
```

```
[1] "hsa04110" "hsa03030" "hsa03013" "hsa03440" "hsa04114"
```

```
pathview(gene.data=foldchanges, pathway.id=keggresids, species="hsa")
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/lenayang/Downloads/bimm 143/Class 14
```

```
Info: Writing image file hsa04110.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/lenayang/Downloads/bimm 143/Class 14
```

```
Info: Writing image file hsa03030.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/lenayang/Downloads/bimm 143/Class 14
```

```
Info: Writing image file hsa03013.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns
```


```
Info: Working in directory /Users/lenayang/Downloads/bimm 143/Class 14
```

```
Info: Writing image file hsa03440.pathview.png
```

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/lenayang/Downloads/bimm 143/Class 14

Info: Writing image file hsa04114.pathview.png



DNA polymerase III holoenzyme

			θ	Pol III core
			ε	
			α	
			γ δ δ'	
Clamp	ψ		σ	y6 complex
β	χ		δ	

Helicase
DnaB


Primase
DnaG

SSB


RNaseH
RNaseIII

DNA polymerase I
Dpol

DNA ligase
Lig



	DNA polymerase B	DNA polymerase D
	PoIB	PolD1 PolD2
	Helicase MCM	Primase Pri1 Pri2
		RPA/SSB RPA
3'	Clamp PCNA	Clamp loader Rfc5 Rfc6
5'		RNaseH RNaseH1 RNaseH2
	Helicase Dna2	DNA ligase Lig



DNA polymerase α -primase complex

$\alpha 1$	$\alpha 2$	PriI	PriII
------------	------------	------	-------

DNA polymerase δ complex

$\delta 1$	$\delta 2$	$\delta 3$	$\delta 4$
------------	------------	------------	------------

DNA polymerase ϵ complex

$\epsilon 1$	$\epsilon 2$	$\epsilon 3$	$\epsilon 4$
--------------	--------------	--------------	--------------

MCM complex (helicase)

Mcm2	Mcm3	
Mcm4	Mcm5	
Mcm6	Mcm7	

RPA

RP11
RP24
RP32

Clamp

PCNA

Clamp loader


RFC1	RFC24	RFC36
------	-------	-------

RNaseHII


RNaseHI	RNaseHIIA	RNaseHII	RNaseHIC
---------	-----------	----------	----------


DNA ligase


Dna2	Fen1	Lig1
------	------	------



23







Gene Ontology (GO)

```
data(go.sets.hs)
data(go.subs.hs)
```

```
# Focus on Biological Process subset of GO
gobpsets = go.sets.hs[go.subs.hs$BP]
```

```
gobpres = gage(foldchanges, gsets=gobpsets, same.dir=TRUE)
```

```
lapply(gobpres, head)
```

\$greater

	p.geomean	stat.mean	p.val
G0:0007156 homophilic cell adhesion	8.519724e-05	3.824205	8.519724e-05
G0:0002009 morphogenesis of an epithelium	1.396681e-04	3.653886	1.396681e-04
G0:0048729 tissue morphogenesis	1.432451e-04	3.643242	1.432451e-04
G0:0007610 behavior	1.925222e-04	3.565432	1.925222e-04
G0:0060562 epithelial tube morphogenesis	5.932837e-04	3.261376	5.932837e-04
G0:0035295 tube development	5.953254e-04	3.253665	5.953254e-04

	q.val	set.size	exp1
G0:0007156 homophilic cell adhesion	0.1952430	113	8.519724e-05
G0:0002009 morphogenesis of an epithelium	0.1952430	339	1.396681e-04
G0:0048729 tissue morphogenesis	0.1952430	424	1.432451e-04
G0:0007610 behavior	0.1968058	426	1.925222e-04
G0:0060562 epithelial tube morphogenesis	0.3566193	257	5.932837e-04
G0:0035295 tube development	0.3566193	391	5.953254e-04

\$less

	p.geomean	stat.mean	p.val
G0:0048285 organelle fission	1.536227e-15	-8.063910	1.536227e-15
G0:0000280 nuclear division	4.286961e-15	-7.939217	4.286961e-15
G0:0007067 mitosis	4.286961e-15	-7.939217	4.286961e-15
G0:0000087 M phase of mitotic cell cycle	1.169934e-14	-7.797496	1.169934e-14
G0:0007059 chromosome segregation	2.028624e-11	-6.878340	2.028624e-11
G0:0000236 mitotic prometaphase	1.729553e-10	-6.695966	1.729553e-10

	q.val	set.size	exp1
G0:0048285 organelle fission	5.843127e-12	376	1.536227e-15
G0:0000280 nuclear division	5.843127e-12	352	4.286961e-15
G0:0007067 mitosis	5.843127e-12	352	4.286961e-15
G0:0000087 M phase of mitotic cell cycle	1.195965e-11	362	1.169934e-14
G0:0007059 chromosome segregation	1.659009e-08	142	2.028624e-11
G0:0000236 mitotic prometaphase	1.178690e-07	84	1.729553e-10

\$stats

	stat.mean	exp1
G0:0007156 homophilic cell adhesion	3.824205	3.824205
G0:0002009 morphogenesis of an epithelium	3.653886	3.653886
G0:0048729 tissue morphogenesis	3.643242	3.643242
G0:0007610 behavior	3.565432	3.565432
G0:0060562 epithelial tube morphogenesis	3.261376	3.261376
G0:0035295 tube development	3.253665	3.253665

Reactome Analysis

```
sig_genes <- res[res$padj <= 0.05 & !is.na(res$padj), "symbol"]  
print(paste("Total number of significant genes:", length(sig_genes)))
```

```
[1] "Total number of significant genes: 8147"
```

```
write.table(sig_genes, file="significant_genes.txt", row.names=FALSE, col.names=FALSE, quo
```

Q8: What pathway has the most significant “Entities p-value”? Do the most significant pathways listed match your previous KEGG results? What factors could cause differences between the two methods?

Cell cycle, mitotic has the most significant “Entities p-value). The most significant pathways listed do not match my previous KEGG results for both up and down regulated genes. The only pathway matched is cell cycle for down regulated pathway. The differences between the two methods could be due to the different databases they are drawing information from.