

1 Analysis of critical transitions at the Global Forest

2 the idea is to do a global analysis using early warning signals of ecological transitions [1]

3 Thus the question is how near/far is the global forest from a catastrophic transition?

4 We will use the MODIS vegetation continuous field, so we can analyse temporal changes.

- 5 • Hypothesis: two power laws, small patches related to deforestation dynamics, large patches related to
6 forest inner dynamics.

7 Methods

- 8 • The United Nations' International Geosphere-Biosphere Programme definition of forest (Belward 1996)
9 defined forest as pixels with tree cover equal or greater than 30%
- 10 • We should define areas with different levels of degradation to apply the spatial indicators [Very difficult
11 because is not possible to establish reliable controls]
- 12 • We should use 2D DFT and multifractals in continuous data and fit patch size distributions in discretized
13 data.[Not implemented]
- 14 • Rates of growth and shrink of patches [2]
- 15 • Portfolio concept relating [2] and [4]

1 Results

2 South America

Table 1: Model selection using Akaike criterion, and goodness of fit calculated by bootstrap. The models were fitted using maximum likelihood with three different data sets: *Complete*, the full data set; $\geq X_{min}$, values greater than or equal than X_{min} threshold; $< X_{min}$, values less than X_{min} .

							GOFp
year	Data_Set	xmin	model_name	par1	par2	delta_AICc	
2000	Complete	1	Power	1.918	NA	0	0
			PowerExp	1.918	3.36e-11	2.019	NA
			LogNorm	1.151	1.631	683753	NA
			Exp	0.003986	NA	7511320	NA
2010			Power	1.833	NA	0	0
			PowerExp	1.831	2.233e-10	6.217	NA
			LogNorm	1.266	1.653	577578	NA
			Exp	0.003998	NA	6827242	NA
2000	$\geq X_{min}$	265	Power	2.013	NA	0	1
			LogNorm	-1486	38.46	2.288	NA
			PowerExp	2.003	1.38e-13	2.688	NA
			Exp	0.0005124	NA	139893	NA
2010		216	Power	2.021	NA	0	1
			PowerExp	2.015	6.11e-12	2.312	NA
			LogNorm	-1213	34.66	2.562	NA
			Exp	0.0005397	NA	150593	NA
2000	$< X_{min}$	265	PowerExp	1.936	0.0006666	0	NA
			Power	1.956	NA	1522	0
			LogNorm	1.389	1.338	818292	NA
			Exp	0.08521	NA	1822635	NA
2010		216	PowerExp	1.829	0.001763	0	NA

							GOFp
year	Data_Set	xmin	model_name	par1	par2	delta_AICc	
			Power	1.877	NA	5615	0
			LogNorm	1.499	1.319	713859	NA
			Exp	0.08257	NA	1525334	NA




Figure 1: Power law exponent of best models by year and with different data set: $<Xmin$ the data is less than the estimated minimum patch size, *Estimated Xmin* the minimum patch size was estimated from data, and only patch sizes greater than or equal to $Xmin$ was used.

- The α with Estimated $Xmin$ correspond to big forest patches and natural forest dynamics, and there is no variation in these. The α with $xmin=1$ correspond to small patches probably influenced by deforestation.
- Besides the power law distribution is the best model is not a valid model for the complete data set. For the patch sizes greater than $Xmin$ the best model is the power law and it is not rejected by the goodness of fit test (GOF). For patches less than $Xmin$, the best model is power law with exponential cutoff, the second best is power law but it was rejected by GOF.

Related papers

- About fitting power laws [5] [6]
- About global maps [7] [8] [9]
- About cluster statistics [10] [2]

Bibliography

1. Kéfi S, Guttal V, Brock WA, Carpenter SR, Ellison AM, et al. (2014) Early Warning Signals of Ecological Transitions: Methods for Spatial Patterns. PLoS ONE 9: e92097. Available: <http://dx.doi.org/10.1371/journal.pone.0092097>.
2. Manor A, Shnerb NM (2008) Origin of Pareto-like Spatial Distributions in Ecosystems. Physical Review Letters 101: 268104. Available: <http://link.aps.org/doi/10.1103/PhysRevLett.101.268104>.
3. Manor A, Shnerb NM (2008) Facilitation, competition, and vegetation patchiness: From scale free distribution to patterns. Journal of Theoretical Biology 253: 838–842. Available: <http://www.sciencedirect.com/science/article/pii/S0022519308001914>.
4. Schindler DE, Armstrong JB, Reed TE (2015) The portfolio concept in ecology and evolution. Frontiers in Ecology and the Environment 13: 257–263. Available: <http://www.esajournals.org/doi/abs/10.1890/140275>.
5. Clauset A, Shalizi C, Newman M (2009) Power-Law Distributions in Empirical Data. SIAM Review 51: 661–703. Available: <http://epubs.siam.org/doi/abs/10.1137/070710111>.
6. White EP, Enquist BJ, Green JL (2008) On Estimating The Exponent Of Power-law Frequency Distributions. Ecology 89: 905–912. Available: <http://www.esajournals.org/doi/abs/10.1890/07-1288.1>.
7. Haddad NM, Brudvig LA, Clobert J, Davies KF, Gonzalez A, et al. (2015) Habitat fragmentation and its lasting impact on Earth’s ecosystems. Science Advances 1: e1500052—e11500052. Available: <http://advances.sciencemag.org/cgi/doi/10.1126/sciadv.1500052>.
8. Sexton JO, Song X-P, Feng M, Noojipady P, Anand A, et al. (2013) Global, 30-m resolution continuous fields of tree cover: {Landsat}-based rescaling of {MODIS} vegetation continuous fields with lidar-based estimates of error. International Journal of Digital Earth 6: 427–448. Available: <http://dx.doi.org/10.1080/17538947.2013.786146>.

- 1 9. Hansen MC, Potapov PV, Moore R, Hancher M, Turubanova SA, et al. (2013) High-Resolution Global
2 Maps of 21st-Century Forest Cover Change. *Science* 342: 850–853. Available: [http://www.sciencemag.org/
3 content/342/6160/850](http://www.sciencemag.org/content/342/6160/850).
- 4 10. Seri E, Maruvka, Shnerb NM (2012) Neutral Dynamics and Cluster Statistics in a Tropical Forest. *The
5 American Naturalist* 180: E161—E1173. Available: <http://www.jstor.org/stable/10.1086/668125>.