Analysis of critical transitions at the Global Forest

- ² The idea is to do a global analysis using insights from percolation theory to detect signals of ecological
- 3 transitions [1]
- 4 Thus the question is how near/far is the global forest from a catastrophic transition?
- 5 We will use the MODIS vegetation continuous field, so we can analyse temporal changes.
- Most probably multiple process influence the distributions of patch size at a continental scale, so we are trying to extract the main generic ones.
- How scaling laws are related to ecosystem function?
- Hypothesis: two power laws, small patches related to deforestation dynamics, large patches related to
 forest inner dynamics.

₁ Methods

- The United Nations' International Geosphere-Biosphere Programme definition of forest (Belward 1996)
 defined forest as pixels with tree cover equal or greater than 30%
- We should define areas with different levels of degradation to apply the spatial indicators [Very difficult because is not possible to establish reliable controls]
- The distribution of patches is continuous but the data is discretized so we discard the lowest values and start fitting patch sizes greater than 10.
- MODIS VCF
- Fitting four models
- Consecuences of each model
- Rates of growth an shrink of patches [2]
- Portfolio concept relating [2] and [4]

1 Results




Figure 1: Power law exponent of best models by year and with different data sets: **<Xmin* the data is less than the extimated minimum patch size, *Estimated Xmin* the minimum patch size was estimated from data, and only patch sizes greater than or equal to Xmin was used.

2 Related papers

- About fitting power laws [5] [6]
- About global maps [7] [8] [9]
- About cluster statistics [10] [2]

6 Bibliography

- 1. Kéfi S, Guttal V, Brock WA, Carpenter SR, Ellison AM, et al. (2014) Early Warning Signals of Ecological
- Transitions: Methods for Spatial Patterns. PLoS ONE 9: e92097. Available: http://dx.doi.org/10.1371/
- 9 journal.pone.0092097.




Figure 2: Estimated X_min by year.




Figure 3: X_min vs power exponent




Figure 4: Biggest patch size by year

- Manor A, Shnerb NM (2008) Origin of Pareto-like Spatial Distributions in Ecosystems. Physical Review
- Letters 101: 268104. Available: http://link.aps.org/doi/10.1103/PhysRevLett.101.268104.
- 3 Manor A, Shnerb NM (2008) Facilitation, competition, and vegetation patchiness: From scale free
- 4 distribution to patterns. Journal of Theoretical Biology 253: 838–842. Available: http://www.sciencedirect.
- 5 com/science/article/pii/S0022519308001914.
- ⁶ 4. Schindler DE, Armstrong JB, Reed TE (2015) The portfolio concept in ecology and evolution. Frontiers in
- Ecology and the Environment 13: 257–263. Available: http://www.esajournals.org/doi/abs/10.1890/140275.
- 5. Clauset A, Shalizi C, Newman M (2009) Power-Law Distributions in Empirical Data. SIAM Review 51:
- 9 661–703. Available: http://epubs.siam.org/doi/abs/10.1137/070710111.
- 6. White EP, Enquist BJ, Green JL (2008) On Estimating The Exponent Of Power-law Frequency Distributions.
- 11 Ecology 89: 905–912. Available: http://www.esajournals.org/doi/abs/10.1890/07-1288.1.
- 12 7. Haddad NM, Brudvig LA, Clobert J, Davies KF, Gonzalez A, et al. (2015) Habitat fragmentation
- and its lasting impact on Earth's ecosystems. Science Advances 1: e1500052—e11500052. Available:
- http://advances.sciencemag.org/cgi/doi/10.1126/sciadv.1500052.

- 8. Sexton JO, Song X-P, Feng M, Noojipady P, Anand A, et al. (2013) Global, 30-m resolution continuous
- 2 fields of tree cover: Landsat-based rescaling of MODIS vegetation continuous fields with lidar-based estimates
- $_3$ of error. International Journal of Digital Earth 6: 427–448. Available: http://dx.doi.org/10.1080/17538947.
- 4 2013.786146.
- 9. Hansen MC, Potapov PV, Moore R, Hancher M, Turubanova SA, et al. (2013) High-Resolution Global
- 6 Maps of 21st-Century Forest Cover Change. Science 342: 850–853. Available: http://www.sciencemag.org/
- 7 content/342/6160/850.
- 8 10. Seri E, Maruvka, Shnerb NM (2012) Neutral Dynamics and Cluster Statistics in a Tropical Forest. The
- ⁹ American Naturalist 180: E161—E1173. Available: http://www.jstor.org/stable/10.1086/668125.