

Analysis of critical transitions at the Global Forest

The idea is to do a global analysis using insights from percolation theory to detect signals of ecological transitions [1]

Thus the question is how near/far is the global forest from a catastrophic transition?

We will use the MODIS vegetation continuous field, so we can analyze temporal changes.

- Most probably multiple process influence the distributions of patch size at a continental scale, so we are trying to extract the main generic ones.
- How scaling laws are related to ecosystem function?
- Hypothesis: the patch distribution is a power law.
- We use percolation theory to calculate the distance to a critical point, the exact value is dependent on details we can not determine but some exponents are invariant.
- Two assumptions:
 - 1) if we view the forest as a static landscape the isotropic percolation universality class is plausible
 - 2) If we view the dynamic of forest the directed percolation universality class is plausible

Methods

- The United Nations' International Geosphere-Biosphere Programme definition of forest (Belward 1996) defined forest as pixels with tree cover equal or greater than 30%
- We defined broad regions based in connectivity
- The distribution of patches is continuous but the data is discretized so we discard the lowest values and start fitting patch sizes greater than 9.
- MODIS VCF
- Fitting four models
- Distance to the critical point is a two step procedure:

- 1) Correlation length is close to the linear length (average linear length) then we can assume the critical point is close and we can apply the universal exponents
- 2) Correlation length is lower than the linear length we are far from the critical point.
- 3) Correlation length is greater than the linear length \rightarrow something is wrong!

Results




Figure 1: Power law exponent of best models by year and with different data sets: $** < X_{min}$ the data is less than the estimated minimum patch size, *Estimated X_{min}* the minimum patch size was estimated from data, and only patch sizes greater than or equal to X_{min} was used.

Related papers

- About fitting power laws [2] [3]
- About global maps [4] [5] [6]
- About cluster statistics [7] [8]

Bibliography

1. Kéfi S, Guttal V, Brock WA, Carpenter SR, Ellison AM, et al. (2014) Early Warning Signals of Ecological Transitions: Methods for Spatial Patterns. PLoS ONE 9: e92097. Available: <http://dx.doi.org/10.1371/journal.pone.0092097>.




Figure 2: Estimated X_{min} by year.




Figure 3: X_{min} vs power exponent




Figure 4: Biggest patch size by year

- 1 2. Clauset A, Shalizi C, Newman M (2009) Power-Law Distributions in Empirical Data. SIAM Review 51:
- 2 661–703. Available: <http://epubs.siam.org/doi/abs/10.1137/070710111>.
- 3 3. White EP, Enquist BJ, Green JL (2008) On Estimating The Exponent Of Power-law Frequency Distributions.
- 4 Ecology 89: 905–912. Available: <http://www.esajournals.org/doi/abs/10.1890/07-1288.1>.
- 5 4. Haddad NM, Brudvig LA, Clobert J, Davies KF, Gonzalez A, et al. (2015) Habitat fragmentation
- 6 and its lasting impact on Earth’s ecosystems. Science Advances 1: e1500052—e11500052. Available:
- 7 <http://advances.sciencemag.org/cgi/doi/10.1126/sciadv.1500052>.
- 8 5. Sexton JO, Song X-P, Feng M, Noojipady P, Anand A, et al. (2013) Global, 30-m resolution continuous
- 9 fields of tree cover: Landsat-based rescaling of MODIS vegetation continuous fields with lidar-based estimates
- 10 of error. International Journal of Digital Earth 6: 427–448. Available: [http://dx.doi.org/10.1080/17538947.](http://dx.doi.org/10.1080/17538947.2013.786146)
- 11 [2013.786146](http://dx.doi.org/10.1080/17538947.2013.786146).
- 12 6. Hansen MC, Potapov PV, Moore R, Hancher M, Turubanova SA, et al. (2013) High-Resolution Global
- 13 Maps of 21st-Century Forest Cover Change. Science 342: 850–853. Available: [http://www.sciencemag.org/](http://www.sciencemag.org/content/342/6160/850)
- 14 [content/342/6160/850](http://www.sciencemag.org/content/342/6160/850).
- 15 7. Seri E, Maruvka, Shnerb NM (2012) Neutral Dynamics and Cluster Statistics in a Tropical Forest. The

- ¹ American Naturalist 180: E161—E1173. Available: <http://www.jstor.org/stable/10.1086/668125>.
- ² 8. Manor A, Shnerb NM (2008) Origin of Pareto-like Spatial Distributions in Ecosystems. Physical Review
- ³ Letters 101: 268104. Available: <http://link.aps.org/doi/10.1103/PhysRevLett.101.268104>.