

Analysis of critical transitions at the Global Forest

Power laws are ubiquitous in nature [1] and much of them are related to vegetation, like patch patterns in semi-arid regions [2] photosynthetic biofilms [3], savanna forest transitions [4] and tree size structure in tropical forest [5]. Several of these power laws are related to patch sizes which define fragmentation [6]. Fragmentation is a global issue and have multiple impacts on ecosystems from reducing biodiversity to changes on nutrient cycling [7]. The concept of percolation shows that there is a critical threshold where suitable habitat patches are essentially isolated [8]. Here we analyzed forest at a global scale to determine if some regions are close to the fragmentation threshold. Following percolation theory, if a system is near a threshold its patch structure must follow a power-law, beside this is not a proof is a necessary condition []. If we are near the fragmentation threshold the biggest patch will have major fluctuations, because the addition or loose of a small patches will change its size the fluctuations must be also be power-law and that if we are approaching the threshold fluctuations will increase with time. We observed that the forest patch structure follows power laws in four major continental biomes: Africa, North America, South America, South East Asia; and it follows a log-normal distribution in Eurasia. Fluctuation of biggest patch are all power law reveals that Africa, and South East Asia are near a fragmentation threshold, North America and South America are not approaching it thus the power laws could be the results of fire and facilitation. Eurasia is also in a pattern of increasing fluctuations.

Results

Related papers

- About fitting power laws [9] [10]
- About global maps [7] [11] [12]
- About cluster statistics [13] [14]

Bibliography

1. Newman MEJ (2005) Power laws, Pareto distributions and Zipf's law. *Contemporary Physics* 46: 323–351. Available: <http://www.tandfonline.com/doi/abs/10.1080/00107510500052444>.
2. Solé R (2007) Scaling laws in the drier. *Nature* 449: 151–153. Available: <http://www.nature.com/nature/journal/v449/n7159/full/449151a.html>.
3. Weerman EJ, Van Belzen J, Rietkerk M, Temmerman S, Kéfi S, et al. (2012) Changes in diatom patch-size distribution and degradation in a spatially self-organized intertidal mudflat ecosystem. *Ecology* 93: 608–618. Available: <http://www.esajournals.org/doi/abs/10.1890/11-0625.1>.
4. Abades SR, Gaxiola A, Marquet PA (2014) Fire, percolation thresholds and the savanna forest transition: a neutral model approach. *Journal of Ecology* 102: 1386–1393. Available: <http://onlinelibrary.wiley.com/doi/10.1111/1365-2745.12321/abstract>.
5. Farrior CE, Bohlman SA, Hubbell S, Pacala SW (2016) Dominance of the suppressed: Power-law size structure in tropical forests. *Science* 351: 155–157. Available: <http://science.sciencemag.org/content/351/6269/155.abstract>.
6. Bascompte J, Solé RV, Sole RV (1996) Habitat fragmentation and extinction thresholds in spatially explicit models. *Journal of Animal Ecology* 65: 465–473. Available: <http://www.jstor.org/discover/10.2307/5781?uid=3737512&uid=>
7. Haddad NM, Brudvig LA, Clobert J, Davies KF, Gonzalez A, et al. (2015) Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances* 1: e1500052. Available: <http://advances.sciencemag.org/cgi/doi/10.1126/sciadv.1500052>.
8. Solé RV, Bascompte J (2006) *Self-organization in complex ecosystems*. New Jersey, USA.: Princeton

- University Press. Available: <http://books.google.com.ar/books?id=v4gpGH6Gv68C>.
9. Clauset A, Shalizi C, Newman M (2009) Power-Law Distributions in Empirical Data. *SIAM Review* 51: 661–703. Available: <http://epubs.siam.org/doi/abs/10.1137/070710111>.
10. White EP, Enquist BJ, Green JL (2008) On Estimating The Exponent Of Power-law Frequency Distributions. *Ecology* 89: 905–912. Available: <http://www.esajournals.org/doi/abs/10.1890/07-1288.1>.
11. Sexton JO, Song X-P, Feng M, Noojipady P, Anand A, et al. (2013) Global, 30-m resolution continuous fields of tree cover: Landsat-based rescaling of MODIS vegetation continuous fields with lidar-based estimates of error. *International Journal of Digital Earth* 6: 427–448. Available: <http://dx.doi.org/10.1080/17538947.2013.786146>.
12. Hansen MC, Potapov PV, Moore R, Hancher M, Turubanova SA, et al. (2013) High-Resolution Global Maps of 21st-Century Forest Cover Change. *Science* 342: 850–853. Available: <http://www.sciencemag.org/content/342/6160/850>.
13. Seri E, Maruvka, Shnerb NM (2012) Neutral Dynamics and Cluster Statistics in a Tropical Forest. *The American Naturalist* 180: E161—E173. Available: <http://www.jstor.org/stable/10.1086/668125>.
14. Manor A, Shnerb NM (2008) Origin of pareto-like spatial distributions in ecosystems. *Physical Review Letters* 101: 268104. Available: <http://link.aps.org/doi/10.1103/PhysRevLett.101.268104>.