

Analysis of critical transitions at the Global Forest

Abstract

Global forests provide critical habitat for many species, essential ecosystem services, and are coupled to atmospheric dynamics through exchanges of energy, water and gases. One of the most important changes produced in the biosphere is the replacement of forest areas with human dominated landscapes. These areas contain many fewer species than the natural ecosystem they replaced, and this change is one of the main drivers of extinctions in the Anthropocene. Another effect is that habitat replacement leads to fragmentation, altering the sizes of patches, the structure and function of the forest. In different kinds of vegetation, including forests, the patch size distributions follow power laws. This have been used as a signal of critical phase transitions, where the system state changes suddenly at a critical point; in the case of forest patches, this implies an abrupt change in connectivity that causes a increased fragmentation level and this poses an additional risk to forest function as an habitat and ecosystem. All these patterns have been observed at relatively small scales, however. Here we studied the distribution of forest patch sizes at a global level, its changes over the last fourteen years, and different signals of a critical transition. We defined wide regions of connected forest at continental level and also studied big islands. The conditions that indicate that a region is near a critical fragmentation threshold are: a patch size distributions following a power law; temporal fluctuations of the largest patch following a power law; variance of largest patch fluctuations increasing in time; and a negative skewness of the largest patch fluctuations. Most regions, except the Eurasian mainland, followed a power-law distribution. We found that only the tropical forest of Africa and South America met the five criteria and thus seemed to be near a critical fragmentation threshold. This implies that the combined influence of human pressures and climate forcings might trigger undesired effects of fragmentation, such as species loss and degradation of ecosystems services, in these regions. The simple criteria proposed here could be used as early warningto estimate the distance to a fragmentation threshold in forest around the globe, and provide a guide to direct conservation efforts at a continental level.

Introduction

Forests are one of the most important ecosystems on earth, providing habitat for a large proportion of species and contributing extensively to global biodiversity (Crowther *et al.*, 2015). In the previous century human activities have influenced global bio-geochemical cycles (Bonan, 2008; Canfield *et al.*, 2010), with one of the most dramatic changes being the replacement of 40% of Earth's formerly biodiverse land areas with

landscapes that contain only a few species of crop plants, domestic animals and humans (Foley *et al.*, 2011). These local changes have accumulated over time and now constitute a global forcing (Barnosky *et al.*, 2012). Another global scale forcing that is tied to habitat destruction is fragmentation, which is defined as the division of a continuous habitat into separated portions that are smaller and more isolated. Fragmentation produces multiple interwoven effects: reductions of biodiversity between 13% and 75%, decreasing forest biomass and changes in nutrient cycling (Haddad *et al.*, 2015). The effects of fragmentation are not only important from an ecological point of view but also human activities and ecosystem services supply and flow are deeply influenced by the level of landscape fragmentation (Mitchell *et al.*, 2015).

Ecosystems have complex interactions between species and present feedbacks at different levels of organization (Gilman *et al.*, 2010), the external forcings can produce abrupt changes from one state to another, called critical transitions (Scheffer *et al.*, 2009). These abrupt state shifts cannot be linearly forecasted from past changes, and are thus difficult to predict and manage (Scheffer *et al.*, 2009). Critical transitions had been detected mostly at local scales (Drake & Griffen, 2010; Carpenter *et al.*, 2011), but the accumulation of changes in local communities that overlap geographically can propagate and cause an abrupt change of the entire system at larger scales (Barnosky *et al.*, 2012). Coupled with the existence of global scale forcings, this implies the possibility that a critical transition could occur at a global scale (Rockstrom *et al.*, 2009; Folke *et al.*, 2011).

Complex systems can experience two general classes of critical transitions (Solé, 2011). In the so called first order transitions, a catastrophic regime shift that is mostly irreversible occurs because of the existence of alternative stable states (Scheffer *et al.*, 2001). This class of transitions is suspected to be present in a variety of ecosystems like: lakes, woodlands, coral reefs (Scheffer *et al.*, 2001), semi-arid grasslands (Bestelmeyer *et al.*, 2011), and fish populations (Vasilakopoulos & Marshall, 2015) and they could be the result of positive feedback mechanisms (Martín *et al.*, 2015). For example: fires in forest ecosystems were more likely to occur in previously burned areas than in unburned places (Kitzberger *et al.*, 2012).

The other class of critical transitions are continuous or second order transitions (Solé & Bascompte, 2006). In these cases, there is a narrow region where the system suddenly changes from one domain to another, with the change being continuous and in theory reversible. This kind of transitions were suggested to be present in tropical forest (Pueyo *et al.*, 2010), semi-arid mountain ecosystems (McKenzie & Kennedy, 2012), tundra shrublands (Naito & Cairns, 2015). The transition happens at critical point where we can observe a distinctive spatial pattern: scale invariant fractal structures characterized by power law patch distributions (Stauffer & Aharony, 1994).

1 The spatial phenomena observed in continuous critical transitions deal with connectivity, a fundamental
2 property of general systems and ecosystems from forests (Ochoa-Quintero *et al.*, 2015) to marine ecosystems
3 (Leibold & Norberg, 2004) and the whole biosphere (Lenton & Williams, 2013). When a system goes from
4 a fragmented to a connected state we say that it percolates (Solé, 2011). Percolation implies that there is a
5 path of connections that involves the whole system. Thus we can characterize two domains or phases: one
6 dominated by short-range interactions where information cannot spread, and another in which long range
7 interactions are possible and information can spread over the whole area. (The term “information” is used
8 in a broad sense and can represent species dispersal or movement.)

9 Thus, there is a critical “percolation threshold” between the two phases, and the system could be driven
10 close to or beyond this point by an external force; climate change and deforestation are the main forces that
11 could be the drivers of such a phase change in contemporary forests (Bonan, 2008; Haddad *et al.*, 2015).
12 Species’ dispersal strategies are influenced by percolation thresholds in three-dimensional forest structure
13 (Solé *et al.*, 2005), and it has been shown that species distributions also have percolation thresholds (He
14 & Hubbell, 2003). This implies that pushing the system below the percolation threshold could produce a
15 biodiversity collapse (Pardini2010; Bascompte *et al.*, 1996; Solé *et al.*, 2004); conversely, being in a connected
16 state (above the threshold) could accelerate the invasion of forest into prairie (Loehle *et al.*, 1996; Naito &
17 Cairns, 2015).

18 One of the main challenges with systems that can experience critical transitions—of any kind—is that the
19 value of the critical threshold is not known in advance. In addition, because near the critical point a small
20 change can precipitate a state shift of the system, they are difficult to predict. Several methods have been
21 developed to detect if a system is close to the critical point, e.g. a deceleration in recovery from perturbations,
22 or an increase in variance in the spatial or temporal pattern (Hastings & Wysham, 2010; Carpenter *et al.*,
23 2011; Boettiger & Hastings, 2012; Dai *et al.*, 2012).

24 There are several processes that can convert a catastrophic transition to a second order transitions (Martín
25 *et al.*, 2015). These include stochasticity, such as demographic fluctuations, spatial heterogeneities, and/or
26 dispersal limitation. All these components are present in forest around the globe (Seidler & Plotkin, 2006;
27 Filotas *et al.*, 2014; Fung *et al.*, 2016), thus continuous transitions might be more probable than catastrophic
28 transitions. Moreover there is some evidence of recovery in some systems that supposedly suffered an irre-
29 versible transition produced by overgrazing (Zhang *et al.*, 2005) and desertification (Allington & Valone,
30 2010).

31 In this study, our objective is to look for evidence that forests around the globe are near continuous critical
32 point that represent a fragmentation thresholds. We use the framework of percolation to first evaluate if

forest patch distribution at a continental scale is described by a power law distribution and then examined the fluctuations of the largest patch. The advantage of using data at a continental scale is that for very large systems the transitions are very sharp (Solé, 2011) and thus much easier to detect than at smaller scales, where noise can mask the signals of the transition.

Methods

Study areas definition

We analyzed mainland forests at a continental scale, covering the whole globe, by delimiting land areas with a near-contiguous forest cover, separated with each other by large non-forested areas. Using this criterion, we delimited the following forest regions. In America, three regions were defined: South America temperate forest (SAT), subtropical and tropical forest up to Mexico (SAST), and USA and Canada forest (NA). Europe and North Asia were treated as one region (EUAS). The rest of the delimited regions were South-east Asia (SEAS), Africa (AF), and Australia (OC). We also included in the analysis islands larger than 10^5 km². The mainland region has the number 1 e.g. OC1, and the nearby islands have consecutive numeration (Supplementary figure S1-S6).

Forest patch distribution

We studied forests patch distribution in each defined area from 2000 to 2014 using the MODerate-resolution Imaging Spectroradiometer (MODIS) Vegetation Continuous Fields (VCF) Tree Cover dataset version 051 (DiMiceli *et al.*, 2015). This dataset is produced at a global level with a 231-m resolution, from 2000 onwards on an annual basis. There are several definition of forest based on percent tree cover (Sexton *et al.*, 2015), we choose a 30% threshold to convert the percentage tree cover to a binary image of forest and non-forest pixels. This was the definition used by the United Nations' International Geosphere-Biosphere Programme (Belward, 1996), and studies of global fragmentation (Haddad *et al.*, 2015). This definition avoids the errors produced by low discrimination of MODIS VCF between forest and dense herbaceous vegetation at low forest cover (Sexton *et al.*, 2015). Patches of contiguous forest were determined in the binary image by grouping connected forest pixels using a neighborhood of 8 forest units (Moore neighborhood).

1 Percolation theory

2 A more in-depth introduction to percolation theory can be found elsewhere (Stauffer & Aharony, 1994) and a
3 review from an ecological point of view is available (Oborny *et al.*, 2007). Here to explain the basic elements
4 of percolation theory we formulate a simple model: we represent our area of interest by a square lattice
5 and each site of the lattice can be occupied—e.g. by forest—with a probability p . The lattice will be more
6 occupied when p is greater, but the sites are randomly distributed. We are interested in the connection
7 between sites, so we define a neighborhood as the eight adjacent sites surrounding any particular site. The
8 sites that are neighbors of other occupied sites define a patch. When there is a patch that connects the
9 lattice from opposite sides, it is said that the system percolates. When p is increased from low values, a
10 percolating patch suddenly appears at some value of p called the critical point p_c .

11 Thus percolation is characterized by two well defined phases: the unconnected phase when $p < p_c$ (called
12 subcritical in physics), in which species cannot travel far inside the forest, as it is fragmented; in a general
13 sense, information cannot spread. The second is the connected phase when $p > p_c$ (supercritical), species
14 can move inside a forest patch from side to side of the area (lattice), i.e. information can spread over the
15 whole area. Near the critical point several scaling laws arise: the structure of the patch that spans the area
16 is fractal, the size distribution of the patches is power-law, and other quantities also follow power-law scaling
17 (Stauffer & Aharony, 1994).

18 The value of the critical point p_c depends on the geometry of the lattice and on the definition of the
19 neighborhood, but other power-law exponents only depend on the lattice dimension. Close to the critical
20 point, the distribution of patch sizes is:

$$21 \quad (1) \quad n_s(p_c) \propto s^{-\alpha}$$

22 where $n_s(p)$ is the number of patches of size s . The exponent α does not depend on the details of the
23 model and it is called universal (Stauffer & Aharony, 1994). These scaling laws can be applied for landscape
24 structures that are approximately random, with correlations if the forest sites are correlated over short
25 distances (Gastner *et al.*, 2009). In physics this is called “isotropic percolation universality class”, and
26 corresponds to an exponent $\alpha = 2.05495$. If we observe that the patch size distribution has another exponent
27 it will not belong to this universality class and some other mechanism should be invoked to explain it.
28 Percolation thresholds can also be generated by models that have some kind of memory (Hinrichsen, 2000;
29 Ódor, 2004): for example, a patch that has been exploited for many years will recover different than a
30 recently deforested forest patch. In this case, the system could belong to a different universality class, or in
31 some cases there is no universality, in which case the value of α will depend on the parameters and details

of the model (Corrado *et al.*, 2014).

Patch size distributions

We fitted the empirical distribution of forest patch areas, based on the MODIS-derived data described above, to four distributions using maximum likelihood estimation (Goldstein *et al.*, 2004; Clauset *et al.*, 2009). The distributions were: power-law, power-law with exponential cut-off, log-normal, and exponential. We assumed that the patch size distribution is a continuous variable that was discretized by remote sensing data acquisition procedure.

We set a minimal patch size (X_{min}) at nine pixels to fit the patch size distributions to avoid artifacts at patch edges due to discretization (Weerman *et al.*, 2012). Besides this hard X_{min} limit we set due to discretization, the power-law distribution needs a lower bound for its scaling behaviour. This lower bound is also estimated from the data by maximizing the Kolmogorov-Smirnov (KS) statistic, computed by comparing the empirical and fitted cumulative distribution functions (Clauset *et al.*, 2009). We also calculated the uncertainty of the parameters using a non-parametric bootstrap method (Efron & Tibshirani, 1994), and computed corrected Akaike Information Criteria (AIC_c) and Akaike weights for each model (Burnham & Anderson, 2002). Akaike weights (w_i) are the weight of evidence in favor of model i being the actual best model given that one of the N models must be the best model for that set of N models.

Additionally, we computed the goodness of fit of the power-law model following the bootstrap approach described by Clauset *et al.* (2009), where simulated data sets following the fitted model are generated, and a p -value computed as the proportion of simulated data sets that has a KS statistic less extreme than empirical data. The criterion to reject the power law model suggested by Clauset *et al.* (2009) was $p \leq 0.1$, but as we have a very large n , meaning that negligible small deviations could produce a rejection (Klaus *et al.*, 2011), we chose a $p \leq 0.05$ to reject the power law model.

To test for differences between the fitted power law exponent for each study area we used a generalized least squares linear model (Zuur *et al.*, 2009) with weights and a residual auto-correlation structure. Weights were the bootstrapped 95% confidence intervals and we added an auto-regressive model of order 1 to the residuals to account for temporal autocorrelation.

Largest patch dynamics

The largest patch is the one that connects the highest number of sites in the area. This has been used extensively to indicate fragmentation (Gardner & Urban, 2007; Ochoa-Quintero *et al.*, 2015). The relation

of the size of the largest patch S_{max} to critical transitions has been extensively studied in relation to percolation phenomena (Bazant2000; Botet2004; Stauffer & Aharony, 1994), but seldom used in ecological studies (but see (2009)). When the system is in a connected state ($p > p_c$) the landscape is almost insensitive to the loss of a small fraction of forest, but close to the critical point a minor loss can have important effects (Solé & Bascompte, 2006; Oborny *et al.*, 2007), because at this point the largest patch will have a filamentary structure, i.e. extended forest areas will be connected by thin threads. Small losses can thus produce large fluctuations.

One way to evaluate the fragmentation of the forest is to calculate the proportion of the largest patch against the total area (Keitt *et al.*, 1997). The total area of the regions we are considering (Supplementary figure S1-S6) may not be the same than the total area that the forest could potentially occupy, thus a more accurate way to evaluate the weight of S_{max} is to use the total forest area, that can be easily calculated summing all the forest pixels. We calculate the proportion of the largest patch for each year, dividing S_{max} by the total forest area of the same year: $RS_{max} = S_{max} / \sum_i S_i$. This has the effect of reducing the S_{max} fluctuations produced due to environmental or climatic changes influences in total forest area. When the proportion RS_{max} is large (more than 60%) the largest patch contains most of the forest so there are fewer small forest patches and the system is probably in a connected phase. Conversely, when it is low (less than 20%), the system is probably in a fragmented phase (Saravia & Momo, 2015).

The RS_{max} is a useful qualitative index that does not if the system is near or far from the critical transition; this can be evaluated using the temporal fluctuations. We calculate the fluctuations around the mean with the absolute values $\Delta S_{max} = S_{max}(t) - \langle S_{max} \rangle$, using the proportions of RS_{max} . To characterize fluctuations we fitted three empirical distributions: power-law, log-normal, and exponential, using the same methods described previously. We expect that large fluctuation near a critical point have heavy tails (log-normal or power-law) and that fluctuations far from a critical point have exponential tails, corresponding to Gaussian processes (Rooij *et al.*, 2013). As the data set spans 15 years, we do not have enough power to reliably detect which distribution is better (Clauset *et al.*, 2009). To improve this we performed the goodness of fit test described above for all the distributions. We generated animated maps showing the fluctuations of the two largest patches to aid in the interpretations of the results.

A robust way to detect if the system is near a critical transition is to analyze the increase in variance of the density (Benedetti-Cecchi *et al.*, 2015). It has been demonstrated that the variance increase in density appears when the system is very close to the transition (Corrado *et al.*, 2014), thus practically it does not constitute an early warning indicator. An alternative is to analyze the variance of the fluctuations of the largest patch ΔS_{max} : the maximum is attained at the critical point but a significant increase occurs

well before the system reaches the critical point (Corrado *et al.*, 2014). In addition, before the critical fragmentation the skewness of the distribution of ΔS_{max} should be negative, implying that fluctuations below the average are more frequent. We characterized the increase in the variance using quantile regression: if variance is increasing the slopes of upper or/and lower quartiles should be positive or negative.

All statistical analyses were performed using the GNU R version 3.3.0 (R Core Team, 2015), using code provided by Cosma R. Shalizi for fitting the power law with exponential cutoff model and the `powerLaw` package (Gillespie, 2015) for fitting the other distributions. For the generalized least squares linear model we used the R function `gls` from package `nlme` (Pinheiro *et al.*, 2016); and we fitted quantile regressions using the R package `quantreg` (Koenker, 2016). Image processing was done in MATLAB r2015b (The Mathworks Inc.). The complete source code for image processing and statistical analysis, and the patch size data files are available at figshare <http://dx.doi.org/10.6084/m9.figshare.XXXXX>.

Results

The power law distribution was selected as the best model in 92% of the cases (Supplementary Figure S7). In a small number of cases (1%) the power law with exponential cutoff was selected, but the value of the parameter α was similar by ± 0.02 to the pure power law. Additionally the patch size where the exponential tail begins is very large, thus we used the power law parameters for this cases (See Supplementary data, region EUAS3). In finite-size systems the favored model should be the power law with exponential cutoff, because the power-law tails are truncated to the size of the system (Stauffer & Aharony, 1994). Here the regions are so large that the cutoff is practically not observed.

There is only one region that did not follow a power law: Eurasia mainland, which showed a log-normal distribution, representing 7% of the cases. The log-normal and power law are both heavy tailed distributions and difficult to distinguish, but in this case Akaike weights have very high values for log-normal (near 1), meaning that this is the only possible model. In addition, the goodness of fit tests clearly rejected the power law model in all cases for this region (Supplementary table S1, region EUAS1). In general the goodness of fit test rejected the power law model in fewer than 10% of cases. In large forest areas like Africa mainland (AF1) or South America tropical-subtropical (SAST1), larger deviations are expected and the rejections rates are higher so the proportion is 30% or less (Supplementary Table S1).

Taking into account the bootstrapped confidence intervals of each power law exponent (α) and the temporal autocorrelation, there were no significant differences between α for the regions with the biggest (greater than 10^7 km²) forest areas (Figure 1 and Supplementary figure S8). There were also no differences between these

regions and smaller ones (Supplementary Table S2 & S3), and all the slopes of α were not different from 0 (Supplementary Table S3). This implies a global average $\alpha = 1.908$, with a bootstrapped 95% confidence interval between 1.898 and 1.920.

The proportion of the largest patch relative to total forest area RS_{max} for regions with more than 10^7 km² of forest is shown in figure 2. South America tropical and subtropical (SAST1) and North America (NA1) have a higher RS_{max} of more than 60%, and other big regions 40% or less. For regions with less total forest area (Supplementary figure S9 & Table 1), the United Kingdom (EUAS3) has a very low proportion near 1%, while other regions such as New Guinea (OC2) and Malaysia/Kalimantan (OC3) have a very high proportion. SEAS2 (Philippines) is a very interesting case because it seems to be under 30% until the year 2005, fluctuates in the range 30-60%, and then stays over 60% (supplementary figure S9). This seems to be an example of a transition from an unconnected to connected state, and deserves further in-depth study [TODO: I'd move previous sentence to discussion].

We analyzed the distributions of fluctuations of the largest patch relative to total forest area ΔRS_{max} and the fluctuations of the largest patch ΔS_{max} . The model selection for ΔS_{max} resulted in power law distributions for all regions (Supplementary table S6). For ΔRS_{max} instead some regions showed exponential distributions: Eurasia mainland (EUAS1), New Guinea (OC2), Malaysia (OC3), New Zealand (OC6, OC8) and Java (OC7), all others were power laws (Supplementary Table S7). The goodness of fit test (GOF) did not reject power laws in any case, but neither rejected the other models except in a few cases; this was due to the small number of observations. We only considered fluctuations to follow a power law when this distribution was selected for both absolute and relative fluctuations.

The animations of the two largest patches (Supplementary data) qualitatively shows the nature of fluctuations and if the state of the forest is connected or not. If the largest patch is always the same patch over time, the forest is probably not fragmented; this happens for regions with RS_{max} of more than 40% such as AF2 (Madagascar), NA1 (North America), and OC3 (Malaysia). Moreover these regions maintain always the same largest patch, which could represent a core area for conservation. [TODO: I'd move previous sentence to discussion] In the regions with RS_{max} between 40% and 30% the identity of the largest patch could change or stay the same in time. For OC7 (Java) the largest patch changes and for AF1 (Africa mainland) it stays the same. Only for EUAS1 (Eurasia mainland) we observed that the two largest patches are always the same, implying that this region is probably composed of two independent domains and should be divided in further studies. The regions with RS_{max} less than 25%: SAST2 (Cuba) and EUAS3 (United Kingdom), the largest patch always changes reflecting their fragmented state. In the case of SEAS2 (Philippines) a transition is observed, with the identity of the largest patch first variable, and then constant after 2010.




Figure 1: Power law exponents (α) of forest patch distributions for regions with total forest area $> 10^7$ km². Dashed horizontal lines are the fitted generalized least squares linear model, with 95% confidence interval error bars estimated by bootstrap resampling. The regions are AF1: Africa mainland, EUAS1: Eurasia mainland, NA1: North America mainland, SAST1: South America subtropical and tropical, SEAS1: Southeast Asia mainland. For EUAS1 the best model is log-normal but the exponents are included here for comparison.




Figure 2: Largest patch proportion relative to total forest area RS_{max} , for regions with total forest area $> 10^7$ km². Dashed lines are averages across time. The regions are AF1: Africa mainland, EUAS1: Eurasia mainland, NA1: North America mainland, SAST1: South America tropical and subtropical, SEAS1: Southeast Asia mainland.

1 The results of quantile regressions are identical for ΔRS_{max} and ΔS_{max} (supplementary table S4). Among
 2 the biggest regions, Africa (AF1) has upper and lower quantiles with significantly negative slopes, but the
 3 lower quantile slope is lower, implying that negative fluctuations and variance are increasing (Figure 3).
 4 Eurasia mainland (EUAS1) has only the upper quantile significant with a positive slope, suggesting an
 5 increase in the variance. North America mainland (NA1) exhibits a significant lower quantile with positive
 6 slope, implying decreasing variance. Finally, for mainland Australia, all quantiles are significant, and the
 7 slope of the lower quantiles is greater than the upper ones, showing that variance is decreasing (Supplementary
 8 figure S10). These results are summarized in Table 1.
 9 The conditions that indicate that a region is near a critical fragmentation threshold are that patch size
 10 distributions follow a power law; temporal ΔRS_{max} fluctuations follow a power law; variance of ΔRS_{max} is
 11 increasing in time; and skewness is negative. All these conditions were true only for Africa mainland (AF1)
 12 and South America tropical & subtropical (SAST1).




Figure 3: Largest patch fluctuations for regions with total forest area $> 10^7 \text{km}^2$. The patch sizes are relative to the total forest area of the same year. Dashed lines are 90%, 50% and 10% quantile regressions, to show if fluctuations were increasing. The regions are AF1: Africa mainland, EUAS1: Eurasia mainland, NA1: North America mainland, SAST1: South America tropical and subtropical, SEAS1: Southeast Asia mainland.

Table 1: Regions and indicators of closeness to a critical fragmentation threshold. Where, RS_{max} is the largest patch divided by the total forest area, ΔRS_{max} are the fluctuations of RS_{max} around the mean, skewness was calculated for RS_{max} and the increase or decrease in the variance was estimated using quantile regressions, NS means the results were non-significant.

Region			Average	Patch Size	ΔRS_{max}	Skewness	Variance
Description			RS_{max}	Distrib	Distrib.		
AF	1	Africa mainland	0.36	Power	Power	-1.8630	Increase
	2	Madagascar	0.65	Power	Power	-0.2478	NS
EUAS	1	Eurasia, mainland	0.36	LogNormal	Exp	0.4016	Increase
	2	Japan	0.94	Power	Power	0.0255	NS
	3	United Kingdom	0.07	Power	Power	2.1330	NS
NA	1	North America, mainland		0.71 Power	er Power		-1.5690 De- crease
	5	Newfoundland	0.87	Power	Power	-0.7411	NS
OC	1	Australia, Mainland	0.28	Power	Power	0.0685	Decrease
	2	New Guinea	0.97	Power	Exp	0.1321	Decrease
	3	Malaysia/Kalimantan	0.97	Power	Exp	-0.9633	NS
	4	Sumatra	0.92	Power	Power	1.3150	Increase
	5	Sulawesi	0.87	Power	Power	-0.3863	NS
	6	New Zealand South Island	0.76	Power	Exp	-0.6683	NS

Region			Average	Patch Size	ΔRS_{max}	Skewness	Variance
		Description	RS_{max}	Distrib	Distrib.		
	7	Java	0.38	Power	Exp	-0.1948	NS
	8	New Zealand North Island	0.75	Power	Exp	0.2940	NS
SAST	1	South America, Tropical and subtropical forest up to Mexico	0.68	Power	Power	-2.7760	Increase
	2	Cuba	0.21	Power	Power	0.2751	NS
SAT	1	South America, Temperate forest	0.60	Power	Power	-1.5070	Decrease
SEAS	1	Southeast Asia, Mainland	0.40	Power	Power	3.0030	NS
	2	Philippines	0.54	Power	Power	0.3113	Increase

1 Discussion

2 We found that the forest patch distribution of most regions of the world followed power laws spanning
3 seven orders of magnitude. These include tropical rainforest, boreal and temperate forest. Power laws have
4 previously been found for several kinds of vegetation, but never at global scales as in this study. Interestingly,
5 Eurasia does not follow a power law, but it is a geographically extended region, consisting of different domains
6 (as we observed in the Supplementary Information animations). It is known that the union of two independent
7 power law distributions produces a lognormal distribution (Rooij *et al.*, 2013). Future studies should split
8 this region into two or more new regions, and test if the underlying distributions are power laws.

9 Several mechanisms have been proposed for the emergence of power laws in forest: the first is related self
10 organized criticality (SOC), when the system is driven by its internal dynamics to a critical state; this has
11 been suggested mainly for fire-driven forests (Zinck & Grimm, 2009; Hantson *et al.*, 2015). Real ecosystems
12 do not seem to meet the requirements of SOC dynamics: their dynamics are influenced by external forces,
13 and interactions are non-homogeneous (i.e. vary from place to place) (Solé *et al.*, 2002). Moreover, SOC
14 requires a memory effect: fire scars in a site should accumulate and interfere with the propagation of a new

1 fire. Pueyo et al. (2010) did not find such effect for tropical forests, and suggest that other mechanisms
2 might produce the observed power laws. Other studies also found that SOC models did not reproduce the
3 patterns of observed fires (McKenzie & Kennedy, 2012). Thus a mechanism which resembled SOC i.e. with
4 a double separation of scales, seems not a plausible explanation for the global forest dynamics.

5 The mechanism suggested by Pueyo et al. (2010) is isotropic percolation, when a system is near the critical
6 point power law structures arise. This is equivalent to the random forest model that we explained previously,
7 and requires the tuning of an external environmental condition to carry the system to this point. We did not
8 expect forest growth to be a random process at local scales, but it is possible that combinations of factors
9 cancel out to produce seemingly random forest dynamics at large scales. If this is the case the power law
10 exponent should be theoretically near $\alpha = 2.055$, but this value is outside the confidence interval we observed,
11 and thus other explanations are needed.

12 The third mechanism suggested as the cause of pervasive power laws in patch size distribution is facilitation
13 (Manor & Shnerb, 2008; Irvine *et al.*, 2016): a patch surrounded by forest will have a smaller probability of
14 been deforested or degraded than an isolated patch. We hypothesize that models that include facilitation
15 could explain the patterns observed here. The model of Scanlon et al. (2007) represented the dynamics
16 of savanna trees includes a global constraint (mean rainfall), a local facilitation mechanism, and two states
17 (tree/non-tree), producing power laws across a rainfall gradient without the need for tuning an external
18 parameter. The results for this model showed an $\alpha = 1.34$ which is also different from our results. Another
19 model but with three states (tree/non-tree/degraded), including local facilitation and grazing, was also used
20 to obtain power laws patch distributions without external tuning, and exhibited deviations from power laws
21 at high grazing pressures (Kéfi *et al.*, 2007). The values of the power law exponent α obtained for this model
22 are dependent on the intensity of facilitation, when facilitation is more intense the exponent is higher, but
23 the maximal values they obtained are still lower than the ones we observed. The interesting point is that
24 the value of the exponent is dependent on the parameters, thus the observed α might be obtained with some
25 parameter combination. Kéfi et al. (2007) proposed that a deviation in power law behavior with the form
26 of an exponential decay or cut-off could be a signal of a critical transition. At the continental scales studied
27 here, we did not observe exponential cut-offs, but we observed other signals of a transition, this confirms
28 previous results (Weerman *et al.*, 2012; Kéfi *et al.*, 2014) showing that different mechanisms could produce
29 seemingly different spatial patterns near the transition and that early warnings based only on spatial patterns
30 are not universal for all the systems.

31 It has been suggested that a combination of spatial and temporal indicators could more reliably detect
32 critical transitions (Kéfi *et al.*, 2014). In this study, we combined five criteria to detect the closeness to a

1 fragmentation threshold. Two of them were spatial: the forest patch size distribution, and the proportion
2 of the largest patch relative to total forest area (RS_{max}). The other three were the distribution of temporal
3 fluctuations in the largest patch size, the trend in the variance, and the skewness of the fluctuations. Each
4 one of these is not a strong individual predictor, but their combination gives us an increased degree of
5 confidence about the system being close to a critical transition.

6 We found that only the tropical forest of Africa and South America met the five criteria, and thus seem
7 to be near a critical fragmentation threshold. This means that the combined influence of human pressures
8 and climate forcings might trigger all the undesired effects of fragmentation in these extended areas. A
9 small but continuous increase in forest loss could produce a biodiversity collapse (Solé *et al.*, 2004), and
10 this threshold effect has been observed in different kind of models, experimental microcosms (Starzomski &
11 Srivastava, 2007), field studies (Pardini *et al.*, 2010; Martensen *et al.*, 2012) and food webs (Martinson *et*
12 *al.*, 2012). Of these two areas, Africa seems to be more affected, because the proportion of the largest patch
13 relative to total forest area (RS_{max}) is near 30%, which could indicate that the transition is already started.
14 Moreover, this region was estimated to be potentially bistable, with the possibility to completely transform
15 into a savanna (Staver *et al.*, 2011). The region of South America tropical forest has a RS_{max} of more than
16 60% suggesting that the fragmentation transition is approaching but not yet started.

17 At low levels of habitat reduction species population will decline proportionally; this can happen even when
18 the habitat fragments retain connectivity. As habitat reduction continues, the critical threshold is approached
19 and connectivity will have large fluctuations (Brook *et al.*, 2013). This could trigger several synergistic
20 effects: populations fluctuations and the possibility of extinctions will rise, increasing patch isolation and
21 decreasing connectivity (Brook *et al.*, 2013). This positive feedback mechanism will be enhanced when the
22 fragmentation threshold is reached, resulting in the loss of most habitat specialist species at a landscape scale
23 (Pardini *et al.*, 2010). Some authors argue that since species have heterogeneous responses to habitat loss
24 and fragmentation, and biotic dispersal is limited, the importance of thresholds is restricted to local scales
25 or even that its existence is questionable (Brook *et al.*, 2013). Fragmentation is by definition a local process
26 that at some point produces emergent phenomena over the entire landscape, even if the area considered is
27 infinite (Oborny *et al.*, 2005). In addition, after a region's fragmentation threshold connectivity decreases,
28 there is still a large and internally well connected patch that can maintain sensitive species (Martensen *et*
29 *al.*, 2012). What is the time needed for these large patches to become fragmented, and pose a real danger
30 of extinction to a myriad of sensitive species? If a forest is already in a fragmented state, a second critical
31 transition from forest to non-forest could happen, this was called the desertification transition (Corrado *et*
32 *al.*, 2014). Considering the actual trends of habitat loss, and studying the dynamics of non-forest patches—

1 instead of the forest patches as we did here—the risk of this kind of transition could be estimated. To
2 improve the estimation of non-forest patches other data set as the MODIS cropland probability should be
3 incorporated (Sexton *et al.*, 2015). The simple models proposed previously could also be used to estimate if
4 these thresholds are likely to be continuous and reversible or discontinuous and irreversible, and the degree
5 of protection (e.g. using the set-asides strategy (Banks-Leite *et al.*, 2014)) than would be necessary to stop
6 this trend.

7 The effectiveness of landscape management is related to the degree of fragmentation, and the criteria to
8 direct the efforts could be focusing on the regions that are near a transition (Oborny *et al.*, 2007). Regions
9 that are in an unconnected state require large efforts to recover a connected state, but regions that are near
10 a transition could be easily pushed to a connected state; feedbacks due to facilitation mechanisms might
11 help to maintain this state.

12 Crossing the fragmentation critical point in forests could have negative effects on biodiversity and ecosystem
13 services (Haddad *et al.*, 2015), but it could also produce feedback loops at different levels of the biological
14 hierarchy. This means that a critical transition produced at a continental scale could have effects at the level
15 of communities, food webs, populations, phenotypes and genotypes (Barnosky *et al.*, 2012). All these effects
16 interact with climate change, thus there is a potential production of cascading effects that could lead to a
17 global collapse. Therefore, even if critical thresholds are reached only in some forest regions at a continental
18 scale, a cascading effect with global consequences could still be produced, and may contribute to reach a
19 planetary tipping point (Reyer *et al.*, 2015). The risk of such event will be higher if the dynamics of separate
20 continental regions are coupled (Lenton & Williams, 2013). Using the time series obtained in this work the
21 coupling of the continental could be further investigated. It has been proposed that to assess the probability
22 of a global scale shift, different small scale ecosystems should be studied in parallel (Barnosky *et al.*, 2012).
23 As forest comprises a major proportion of such ecosystems, we think that the transition of forests could be
24 used as a proxy for all the underling changes and as a successful predictor of a planetary tipping point.

25 References

- 26 Allington GRH, Valone TJ (2010) Reversal of desertification: The role of physical and chemical soil properties.
27 *Journal of Arid Environments*, **74**, 973–977.
- 28 Banks-Leite C, Pardini R, Tambosi LR et al. (2014) Using ecological thresholds to evaluate the costs and
29 benefits of set-asides in a biodiversity hotspot. *Science*, **345**, 1041–1045.
- 30 Barnosky AD, Hadly EA, Bascompte J et al. (2012) Approaching a state shift in Earth’s biosphere. *Nature*,

1 **486**, 52–58.

2 Bascompte J, Solé RV, Sole RV (1996) Habitat fragmentation and extinction thresholds in spatially explicit
3 models. *Journal of Animal Ecology*, **65**, 465–473.

4 Belward AS (1996) The IGBP-DIS Global 1 Km Land Cover Data Set “DISCover”: Proposal and Imple-
5 mentation Plans : Report of the Land Recover Working Group of IGBP-DIS. IGBP-DIS Office, pp.

6 Benedetti-Cecchi L, Tamburello L, Maggi E, Bulleri F (2015) Experimental Perturbations Modify the Per-
7 formance of Early Warning Indicators of Regime Shift. *Current biology*, **25**, 1867–1872.

8 Bestelmeyer BT, Ellison AM, Fraser WR et al. (2011) Analysis of abrupt transitions in ecological systems.
9 *Ecosphere*, **2**, 129.

10 Boettiger C, Hastings A (2012) Quantifying limits to detection of early warning for critical transitions.
11 *Journal of The Royal Society Interface*, **9**, 2527–2539.

12 Bonan GB (2008) Forests and Climate Change: Forcings, Feedbacks, and the Climate Benefits of Forests.
13 *Science*, **320**, 1444–1449.

14 Brook BW, Ellis EC, Perring MP, Mackay AW, Blomqvist L (2013) Does the terrestrial biosphere have
15 planetary tipping points? *Trends in Ecology & Evolution*.

16 Burnham K, Anderson DR (2002) Model selection and multi-model inference: A practical information-
17 theoretic approach, 2nd. edn. Springer-Verlag, New York, pp.

18 Canfield DE, Glazer AN, Falkowski PG (2010) The Evolution and Future of Earth’s Nitrogen Cycle. *Science*,
19 **330**, 192–196.

20 Carpenter SR, Cole JJ, Pace ML et al. (2011) Early Warnings of Regime Shifts: A Whole-Ecosystem
21 Experiment. *Science*, **332**, 1079–1082.

22 Clauset A, Shalizi C, Newman M (2009) Power-Law Distributions in Empirical Data. *SIAM Review*, **51**,
23 661–703.

24 Corrado R, Cherubini AM, Pennetta C (2014) Early warning signals of desertification transitions in semiarid
25 ecosystems. *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics*, **90**, 62705.

26 Crowther TW, Glick HB, Covey KR et al. (2015) Mapping tree density at a global scale. *Nature*, **525**,
27 201–205.

28 Dai L, Vorselen D, Korolev KS, Gore J (2012) Generic Indicators for Loss of Resilience Before a Tipping

1 Point Leading to Population Collapse. *Science*, **336**, 1175–1177.

2 DiMiceli C, Carroll M, Sohlberg R, Huang C, Hansen M, Townshend J (2015) Annual Global Automated
3 MODIS Vegetation Continuous Fields (MOD44B) at 250 m Spatial Resolution for Data Years Beginning
4 Day 65, 2000 - 2014, Collection 051 Percent Tree Cover, University of Maryland, College Park, MD, USA.

5 Drake JM, Griffen BD (2010) Early warning signals of extinction in deteriorating environments. *Nature*,
6 **467**, 456–459.

7 Efron B, Tibshirani RJ (1994) An Introduction to the Bootstrap. Taylor & Francis, New York, pp.

8 Filotas E, Parrott L, Burton PJ et al. (2014) Viewing forests through the lens of complex systems science.
9 *Ecosphere*, **5**, 1–23.

10 Foley JA, Ramankutty N, Brauman KA et al. (2011) Solutions for a cultivated planet. *Nature*, **478**, 337–342.

11 Folke C, Jansson Å, Rockström J et al. (2011) Reconnecting to the Biosphere. *AMBIO*, **40**, 719–738.

12 Fung T, O’Dwyer JP, Rahman KA, Fletcher CD, Chisholm RA (2016) Reproducing static and dynamic
13 biodiversity patterns in tropical forests: the critical role of environmental variance. *Ecology*.

14 Gardner RH, Urban DL (2007) Neutral models for testing landscape hypotheses. *Landscape Ecology*, **22**,
15 15–29.

16 Gastner MT, Oborny B, Zimmermann DK, Pruessner G (2009) Transition from Connected to Fragmented
17 Vegetation across an Environmental Gradient: Scaling Laws in Ecotone Geometry. *The American Naturalist*,
18 **174**, E23–E39.

19 Gillespie CS (2015) Fitting Heavy Tailed Distributions: The powerLaw Package. *Journal of Statistical*
20 *Software*, **64**, 1–16.

21 Gilman SE, Urban MC, Tewksbury J, Gilchrist GW, Holt RD (2010) A framework for community interactions
22 under climate change. *Trends in Ecology & Evolution*, **25**, 325–331.

23 Goldstein ML, Morris SA, Yen GG (2004) Problems with fitting to the power-law distribution. *The European*
24 *Physical Journal B - Condensed Matter and Complex Systems*, **41**, 255–258.

25 Haddad NM, Brudvig La, Clobert J et al. (2015) Habitat fragmentation and its lasting impact on Earth’s
26 ecosystems. *Science Advances*, **1**, 1–9.

27 Hantson S, Pueyo S, Chuvieco E (2015) Global fire size distribution is driven by human impact and climate.
28 *Global Ecology and Biogeography*, **24**, 77–86.

29 Hastings A, Wysham DB (2010) Regime shifts in ecological systems can occur with no warning. *Ecology*

1 *Letters*, **13**, 464–472.

2 He F, Hubbell S (2003) Percolation Theory for the Distribution and Abundance of Species. *Physical Review*
3 *Letters*, **91**, 198103.

4 Hinrichsen H (2000) Non-equilibrium critical phenomena and phase transitions into absorbing states. *Ad-*
5 *vances in Physics*, **49**, 815–958.

6 Irvine MA, Bull JC, Keeling MJ (2016) Aggregation dynamics explain vegetation patch-size distributions.
7 *Theoretical Population Biology*, **108**, 70–74.

8 Keitt TH, Urban DL, Milne BT (1997) Detecting critical scales in fragmented landscapes. *Conservation*
9 *Ecology*, **1**, 4.

10 Kéfi S, Rietkerk M, Alados CL, Pueyo Y, Papanastasis VP, ElAich A, Ruiter PC de (2007) Spatial vegetation
11 patterns and imminent desertification in Mediterranean arid ecosystems. *Nature*, **449**, 213–217.

12 Kéfi S, Guttal V, Brock WA et al. (2014) Early Warning Signals of Ecological Transitions: Methods for
13 Spatial Patterns. *PLoS ONE*, **9**, e92097.

14 Kitzeberger T, Aráoz E, Gowda JH, Mermoz M, Morales JM (2012) Decreases in Fire Spread Probability
15 with Forest Age Promotes Alternative Community States, Reduced Resilience to Climate Variability and
16 Large Fire Regime Shifts. *Ecosystems*, **15**, 97–112.

17 Klaus A, Yu S, Plenz D (2011) Statistical analyses support power law distributions found in neuronal
18 avalanches. (ed Zochowski M). *PloS one*, **6**, e19779.

19 Koenker R (2016) quantreg: Quantile Regression.

20 Leibold MA, Norberg J (2004) Biodiversity in metacommunities: Plankton as complex adaptive systems?
21 *Limnology and Oceanography*, **49**, 1278–1289.

22 Lenton TM, Williams HTP (2013) On the origin of planetary-scale tipping points. *Trends in Ecology &*
23 *Evolution*, **28**, 380–382.

24 Loehle C, Li B-L, Sundell RC (1996) Forest spread and phase transitions at forest-prairie ecotones in Kansas,
25 U.S.A. *Landscape Ecology*, **11**, 225–235.

26 Manor A, Shnerb NM (2008) Origin of pareto-like spatial distributions in ecosystems. *Physical Review*
27 *Letters*, **101**, 268104.

28 Martensen AC, Ribeiro MC, Banks-Leite C, Prado PI, Metzger JP (2012) Associations of Forest Cover,
29 Fragment Area, and Connectivity with Neotropical Understory Bird Species Richness and Abundance. *Con-*

1 *ervation Biology*, **26**, 1100–1111.

2 Martinson HM, Fagan WF, Denno RF (2012) Critical patch sizes for food-web modules. *Ecology*, **93**, 1779–
3 1786.

4 Martín PV, Bonachela JA, Levin SA, Muñoz MA (2015) Eluding catastrophic shifts. *Proceedings of the*
5 *National Academy of Sciences*, **112**, E1828–E1836.

6 McKenzie D, Kennedy MC (2012) Power laws reveal phase transitions in landscape controls of fire regimes.
7 *Nat Commun*, **3**, 726.

8 Mitchell MGE, Suarez-Castro AF, Martinez-Harms M et al. (2015) Reframing landscape fragmentation’s
9 effects on ecosystem services. *Trends in Ecology & Evolution*, **30**, 190–198.

10 Naito AT, Cairns DM (2015) Patterns of shrub expansion in Alaskan arctic river corridors suggest phase
11 transition. *Ecology and Evolution*, **5**, 87–101.

12 Oborny B, Meszéna G, Szabó G (2005) Dynamics of Populations on the Verge of Extinction. *Oikos*, **109**,
13 291–296.

14 Oborny B, Szabó G, Meszéna G (2007) Survival of species in patchy landscapes: percolation in space and
15 time. In: *Scaling biodiversity*, pp. 409–440. Cambridge University Press.

16 Ochoa-Quintero JM, Gardner TA, Rosa I, de Barros Ferraz SF, Sutherland WJ (2015) Thresholds of species
17 loss in Amazonian deforestation frontier landscapes. *Conservation Biology*, **29**, 440–451.

18 Ódor G (2004) Universality classes in nonequilibrium lattice systems. *Reviews of Modern Physics*, **76**, 663–
19 724.

20 Pardini R, Bueno A de A, Gardner TA, Prado PI, Metzger JP (2010) Beyond the Fragmentation Threshold
21 Hypothesis: Regime Shifts in Biodiversity Across Fragmented Landscapes. *PLoS ONE*, **5**, e13666.

22 Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team (2016) nlme: Linear and Nonlinear Mixed Effects
23 Models. pp.

24 Pueyo S, de Alencastro Graça PML, Barbosa RI, Cots R, Cardona E, Fearnside PM (2010) Testing for
25 criticality in ecosystem dynamics: the case of Amazonian rainforest and savanna fire. *Ecology Letters*, **13**,
26 793–802.

27 R Core Team (2015) R: A Language and Environment for Statistical Computing.

28 Reyer CPO, Rammig A, Brouwers N, Langerwisch F (2015) Forest resilience, tipping points and global

change processes. *Journal of Ecology*, **103**, 1–4.

Rockstrom J, Steffen W, Noone K et al. (2009) A safe operating space for humanity. *Nature*, **461**, 472–475.

Rooij MMJW van, Nash B, Rajaraman S, Holden JG (2013) A Fractal Approach to Dynamic Inference and Distribution Analysis. *Frontiers in Physiology*, **4**.

Saravia LA, Momo FR (2015) Biodiversity collapse in a phase transition between neutral and niche communities. *PeerJ PrePrints*, **3**, e1589v2.

Scanlon TM, Caylor KK, Levin SA, Rodriguez-iturbe I (2007) Positive feedbacks promote power-law clustering of Kalahari vegetation. *Nature*, **449**, 209–212.

Scheffer M, Walkerk B, Carpenter S, Foley J a, Folke C, Walker B (2001) Catastrophic shifts in ecosystems. *Nature*, **413**, 591–596.

Scheffer M, Bascompte J, Brock WA et al. (2009) Early-warning signals for critical transitions. *Nature*, **461**, 53–59.

Seidler TG, Plotkin JB (2006) Seed Dispersal and Spatial Pattern in Tropical Trees. *PLoS Biology*, **4**, e344.

Sexton JO, Noojipady P, Song X-P et al. (2015) Conservation policy and the measurement of forests. *Nature Climate Change*, **advance on**, 192–196.

Solé RV (2011) Phase Transitions. Princeton University Press, pp.

Solé RV, Bascompte J (2006) Self-organization in complex ecosystems. Princeton University Press, New Jersey, USA., pp.

Solé RV, Alonso D, McKane A (2002) Self-organized instability in complex ecosystems. *Philosophical transactions of the Royal Society of London. Series B*, **357**, 667–681.

Solé RV, Alonso D, Saldaña J (2004) Habitat fragmentation and biodiversity collapse in neutral communities. *Ecological Complexity*, **1**, 65–75.

Solé RV, Bartumeus F, Gamarra JGP (2005) Gap percolation in rainforests. *Oikos*, **110**, 177–185.

Starzomski BM, Srivastava DS (2007) Landscape geometry determines community response to disturbance. *Oikos*, **116**, 690–699.

Stauffer D, Aharony A (1994) Introduction To Percolation Theory. Taylor & Francis, London, pp.

Staver AC, Archibald S, Levin SA (2011) The Global Extent and Determinants of Savanna and Forest as

- 1 Alternative Biome States. *Science*, **334**, 230–232.
- 2 Vasilakopoulos P, Marshall CT (2015) Resilience and tipping points of an exploited fish population over six
3 decades. *Global Change Biology*, **21**, 1834–1847.
- 4 Weerman EJ, Van Belzen J, Rietkerk M, Temmerman S, Kéfi S, Herman PMJ, Koppel JV de (2012) Changes
5 in diatom patch-size distribution and degradation in a spatially self-organized intertidal mudflat ecosystem.
6 *Ecology*, **93**, 608–618.
- 7 Zhang JY, Wang Y, Zhao X, Xie G, Zhang T (2005) Grassland recovery by protection from grazing in a
8 semi-arid sandy region of northern China. *New Zealand Journal of Agricultural Research*, **48**, 277–284.
- 9 Zinck RD, Grimm V (2009) Unifying wildfire models from ecology and statistical physics. *The American*
10 *naturalist*, **174**, E170–85.
- 11 Zuur AF, Ieno EN, Walker N, Saveliev AA, Smith GM (2009) Mixed effects models and extensions in ecology
12 with R. Springer New York, New York, NY, pp.