

# MA359 Glossary

James Palmer and Luca Seemungal

April 22, 2021

## 1 Week 1: Outer Measures

**Outer Measure** An Outer measure is a function  $\mu^* : P(X) \rightarrow [0, \infty]$  such that the following three axioms are satisfied:

- (i)  $\mu^*(\emptyset) = 0$
- (ii)  $A \subseteq B \Rightarrow \mu^*(A) \leq \mu^*(B)$
- (iii) if  $\{A_n\}_{n=1}^{\infty}$  is a sequence of subsets of  $X$ , then we have

$$\mu^*\left(\bigcup_{n=1}^{\infty} A_n\right) \leq \sum_{n=1}^{\infty} \mu^*(A_n)$$

## 2 Week 2: Sigma Algebra

**Measurability** Let  $\mu^*$  be an outer measure on  $X$ . We say  $B$  such that  $B \subset X$  is measurable if  $\mu^*(A) = \mu^*(A \cap B) + \mu^*(A \cap B^c)$  for all  $A$  such that  $A \subseteq X$ .

**$\sigma$ -algebra** A collection of sets  $\mathcal{A} \subseteq P(X)$  is a  $\sigma$ -algebra if it has the following properties:

- (i)  $X \in \mathcal{A}$
- (ii)  $A \in \mathcal{A} \Rightarrow A^c \in \mathcal{A}$
- (iii) For every countable sequence of sets  $\{A_i\}_{i=1}^{\infty}$ , the union of these sets is also in  $\mathcal{A}$ . i.e.  $\mu^*\left(\bigcup_{n=1}^{\infty} A_n\right) \in \mathcal{A}$

## 3 Week 3: Measures

**Measure** In general a measure  $\mu$  is a function of a  $\sigma$ -algebra  $\mathcal{A}$  such that  $\mu(\emptyset) = 0$  and  $\mu$  is countably additive, i.e.  $\mu\left(\bigcup_{n=1}^{\infty} A_n\right) = \sum_{n=1}^{\infty} \mu(A_n)$  when  $A_n$  are all disjoint sets.

**Measure Space and Measurable Space**  $(X, \mathcal{A}, \mu)$  is a measure space,  $(X, \mathcal{A})$  is a measurable space. (Domain set  $X$ ,  $\sigma$ -algebra  $\mathcal{A}$  and measure  $\mu$ ).

**Measure Continuity** If  $A_k$  is an increasing sequence of sets in  $\mathcal{A}$  then  $\mu(\bigcup_{k=1}^{\infty} A_k) = \lim_{k \rightarrow \infty} \mu(A_k)$

If  $A_k$  is a decreasing sequence of sets in  $\mathcal{A}$  then  $\mu(\bigcap_{k=1}^{\infty} A_k) = \lim_{k \rightarrow \infty} \mu(A_k)$

**Translation Invariance** A measure is translation invariant:  $\mu(A + h) = \mu(A)$

## 4 Week 4: Measureable Functions

**Measurable Function** If  $A \subset X$  and  $A \in \mathcal{A}$  function  $f : A \rightarrow [-\infty, +\infty]$ , then the conditions

- (a) for all  $t \in \mathbb{R}$  the set  $\{x \in A : f(x) \leq t\}$
  - (b) for all  $t \in \mathbb{R}$  the set  $\{x \in A : f(x) \geq t\}$
  - (c) for all  $t \in \mathbb{R}$  the set  $\{x \in A : f(x) \leq t\}$
  - (d) for all  $t \in \mathbb{R}$  the set  $\{x \in A : f(x) \geq t\}$
- are equivalent

**Simple Functions** A simple function is a function of the form  $f = \sum_{i=0}^N a_i \chi_{A_i}$  where  $a_i$  are non-negative real numbers and  $A_i$  are disjoint subsets of  $X$ .

**Integral of a simple Functions** The integral of a simple function with respect to measure  $\mu$  is  $\int f d\mu = \sum_{i=0}^N a_i \mu(A_i)$

**For any non-negative function, there is a sequence of simple functions such that the limit of the sequence is our original function**

## 5 Week 5: Integrable Functions

**Integral of a non-negative function**  $\int f d\mu = \sup\{\int g d\mu : g \text{ is simple and } g \leq f\}$

**Integral of a function**  $\int f d\mu = \int f^+ d\mu + \int f^- d\mu$  where  $f^+ = \max\{0, f\}$  and  $f^- = \max\{0, -f\}$

**The integral of simple functions, the integral of a non-negative function and the integral of a function are all linear functionals**

**BTEC Monotone convergence theorem** if we have a monotone sequence of simple functions  $f_i$  that converge pointwise to a measurable function  $f$ , then the sequence of the integrals of  $f_i$  converges to the integral of  $f$  (integral with respect to  $\mu$ )

## 6 Week 6: Integration Theorems

**Monotone Convergence Theorem** Let  $f_i$  be a monotone (increasing) sequence of measurable functions that converge to measurable function  $f$ . Then  $\int f d\mu = \lim_{i \rightarrow \infty} \int f_i d\mu$

**Beppo Levi's Theorem** Let  $(X, \mathcal{A}, \mu)$  be a measure space, then

$$\int \sum_{i=1}^{\infty} f_i d\mu = \sum_{i=1}^{\infty} \int f_i d\mu$$

**Fatou's Lemma** Let  $(X, \mathcal{A}, \mu)$  be a measure space and let  $\{f_i\}$  be a sequence of  $[-\infty, \infty]$  valued  $\mathcal{A}$ -measurable functions on  $X$ . Then

$$\int \liminf_{i \rightarrow \infty} f_i(x) \leq \liminf_{i \rightarrow \infty} \int f_i(x)$$

**Dominated Convergence Theorem** Let  $(X, \mathcal{A}, \mu)$  be a measure space. Let  $g$  be an integrable function on  $X$  and let  $f_i$  be a sequence of  $[-\infty, \infty]$  valued  $\mathcal{A}$ -measurable functions such that the sequence converges to  $f$  and  $|f_i(x)| \leq g(x)$  for all  $n \in \mathbb{N}$ . Then  $f$  and  $f_i$  are integrable and  $\int f d\mu = \lim_{i \rightarrow \infty} \int f_i d\mu$ .

**Markow's inequality** If we have the set  $A_t = \{x \in X : f(x) \geq t\}$  then  $\mu(A_k) \leq \frac{1}{t} \int_{A_t} f d\mu \leq \frac{1}{t} \int f d\mu$ .

**Link to Riemann Integrals** Let  $[a, b]$  be a closed and bounded interval and let  $f$  be a bounded real valued function on  $[a, b]$ . Then

(a)  $f$  is Riemann integrable if and only if it is continuous almost everywhere on  $[a, b]$ .

(b) if  $f$  is Riemann integrable then  $f$  is Lebesgue integrable and the integrals coincide.

## 7 Week 7: Modes of Convergence

**Convergence in Measure** Sequence of  $\mathcal{A}$ -measurable functions valued on  $[-\infty, \infty]$   $f_i$  converge to  $f$  in measure if for any  $\epsilon > 0$ ,  $\lim_{i \rightarrow \infty} \mu(\{x \in X : |f_i(x) - f(x)| > \epsilon\}) = 0$

**Pointwise Convergence** A sequence of  $\mathcal{A}$ -measurable functions valued on  $[-\infty, \infty]$   $f_i$  converge to  $f$  pointwise if for every  $x \in X$ ,  $\lim_{i \rightarrow \infty} f_i(x) = f(x)$

**Pointwise Convergence almost everywhere** Same as above but the set of points that fail make up a set of measure zero.

**Convergence in mean** A sequence of  $\mathcal{A}$ -measurable functions valued on  $[-\infty, \infty]$   $f_i$  converge to  $f$  in mean if  $\lim_{i \rightarrow \infty} \int |f_i - f| d\mu = 0$

**Egoroff's Theorem** If  $f_i$  converges pointwise to  $f$  then for any  $\epsilon > 0$  there is a subset  $B$  of  $X$  such that  $f_i$  restricted to  $B$  converges uniformly to  $f$  restricted to  $B$  and  $\mu(B^c) < \epsilon$

## 8 Week 8: $L^p$ spaces

**$L^p$  spaces** The vector space  $L^p(X, \mathcal{A}, \mu)$  is the space of functions such that  $(\int_X |f|^p d\mu)^{\frac{1}{p}}$ , factored by the equivalence classes of functions that are equal a.e. i.e.  $f \sim g$  iff  $f(x) = g(x)$  almost everywhere.

(NB, if  $|f - g|_p = 0$ , this only implies  $f = g$  almost everywhere due to the definition of this norm.  $L^p$  is also complete, so any Cauchy sequence of functions will have a limit in  $L^p$ )

**Young's Inequality** If  $\frac{1}{p} + \frac{1}{q} = 1$   $x^{\frac{1}{p}} y^{\frac{1}{q}} \leq \frac{x}{p} + \frac{y}{q}$

**Holder's Inequality** Assume  $\frac{1}{p} + \frac{1}{q} = 1$  and  $p > 1$ , If  $f \in L^p$  and  $g \in L^q$ , then  $\int fg d\mu \leq |f|_p |g|_q$

**Minkowski's Inequality** Basically triangle rule for the  $L^p$  norm

## 9 Week 9: Product Sigma Algebras

**Dynkin Systems** A family  $\mathcal{D}$  of sets is a d-system if it satisfies axioms (i) and (ii) of a  $\sigma$ -algebra and (iii) if  $A_n$  is an increasing sequence of sets in  $\mathcal{D}$ , then  $\bigcup_{n=1}^{\infty} A_n$  is in  $\mathcal{D}$  as well.

**$\pi$ -system** Same as a  $\sigma$ -algebra but with finite union instead of countable union for axiom (iii).

**Dynkin's Theorem** Let  $X$  be a set and  $\mathcal{C}$  be a  $\pi$ -system on  $X$ . Then the  $\sigma$ -algebra generated by  $\mathcal{C}$  coincides with the d-system generated by  $\mathcal{C}$ .

**Agreement on Generators** Let  $(X, \mathcal{A})$  be a measurable space and let  $\mathcal{C}$  be a  $\pi$ -system such that  $\mathcal{A} = \sigma(\mathcal{C})$ . If  $\mu$  and  $\nu$  are measures on  $(X, \mathcal{A})$  that satisfy  $\mu(X) = \nu(X) < \infty$  and  $\mu(C) = \nu(C)$  for  $C \in \mathcal{C}$  then  $\mu = \nu$ .

**Product Sigma Algebra** Let  $(X_1, \mathcal{A}_1)$  and  $(X_2, \mathcal{A}_2)$  be two measurable spaces. Then, the family of sets  $\mathcal{A}_1 \otimes \mathcal{A}_2 := \sigma(\{A_1 \times A_2 : A_1 \in \mathcal{A}_1 \text{ and } A_2 \in \mathcal{A}_2\})$ . is called the product  $\sigma$ -algebra.

**Product Measure (Note - this is a theorem and not a definition)** Given two measure spaces  $(X_1, \mathcal{A}_1, \mu_1)$  and  $(X_2, \mathcal{A}_2, \mu_2)$  where the measure spaces are  $\sigma$ -finite. Then there exists a unique measure  $\mu_1 \otimes \mu_2: (\mathcal{A}_1, \mathcal{A}_2) \rightarrow [0, \infty]$  such that  $(\mu_1 \otimes \mu_2)(A_1 \times A_2) = \mu_1(A_1)\mu_2(A_2)$  for all sets  $A_1, A_2$  in  $\mathcal{A}_\infty, \mathcal{A}_\infty$  respectively. This is called the product measure.

Futhermore, for each  $E \in \mathcal{A}_1 \otimes \mathcal{A}_2$ , the function  $x \rightarrow \mu_2(E_x)$  where  $E_x := \{y \in X_2 : (x, y) \in E\}$  is measurable and the function  $y \rightarrow \mu_1(E_y)$  where  $E_y := \{x \in X_1 : (x, y) \in E\}$  is measurable. We also have:  $(\mu_1 \otimes \mu_2)(E) = \int_{X_1} \mu_2(E_x) d\mu_1(x) = \int_{X_2} \mu_1(E_y) d\mu_2(x)$  for all  $E \in \mathcal{A}_1 \times \mathcal{A}_2$ .

## 10 Week 10: Fubini's Theorem

**Fubini's Theorem Part 1** Let  $(X_1, \mathcal{A}_1, \mu_1)$  and  $(X_2, \mathcal{A}_2, \mu_2)$  be measure spaces which are  $\sigma$ -finite.

let  $f : X_1 \times X_2 \rightarrow [0, \infty]$  be an  $\mathcal{A}_1 \times \mathcal{A}_2$  measurable function. Then  $x \rightarrow \int f_x d\mu_2$  is  $\mathcal{A}_1$  measurable and  $y \rightarrow \int f_y d\mu_1$  is  $\mathcal{A}_2$  measurable.

Furthermore,  $\int_{X \times Y} f d(\mu_1 \times \mu_2) = \int_X (\int_Y f_X d\mu_2) d\mu_1$  and  $\int_{X \times Y} f d(\mu_1 \times \mu_2) = \int_Y (\int_X f_Y d\mu_1) d\mu_2$

**Fubini's Theorem Part 2 - what is listed as Fubini in Cohn** Same hypotheses in part 1.

(a)  $\mu_1$ -a.e.  $x \in X$  the section  $f_X$  is  $\mu_2$ -integrable and for  $\mu_2$ -a.e. the section  $f_Y$  is  $\mu_1$ -integrable

(b) The fuctions  $I_f(x) = \begin{cases} \int_Y f_X d\mu_2 & \text{if } f_X \text{ is } \mu_2\text{-integrable} \\ 0 & \text{else} \end{cases}$

and  $J_f(x) = \begin{cases} \int_X f_Y d\mu_1 & \text{if } f_Y \text{ is } \mu_1\text{-integrable} \\ 0 & \text{else} \end{cases}$

then  $I_f \in \mathcal{L}^1(X, \mu_1)$  and  $J_f \in \mathcal{L}^1(Y, \mu_2)$

(c)  $\int_{X \times Y} f d(\mu_1 \times \mu_2) = \int_Y J_f d\mu_2 = \int_X I_f d\mu_1$