Profile of studies by WHO region

1.0
evaluations per 100,000 DALYs

47%
report QALYs or DALYs

56%
use some modelling


Communicable · Non-communicable · Injuries

AFR

African Region

n = 1,379


0.23 per 100,000 DALYs



Top three diseases

HIV	25%
Maternal & newborn health	19%
Malaria & dengue	10%

46% report QALYs or DALYs




AMR

Region of the Americas

n = 9,580


2.83 per 100,000 DALYs



Top three diseases

Breast cancer	4%
Diabetes	3%
HIV	3%

50% report QALYs or DALYs




SEAR

South-East Asia Region

n = 1,146


0.19 per 100,000 DALYs



Top three diseases

Maternal & newborn health	9%
Tuberculosis	6%
HIV	4%

52% report QALYs or DALYs




EUR

European Region

n = 9,825


3.06 per 100,000 DALYs



Top three diseases

Depression & mental disorders	6%
Diabetes	4%
Psoriasis & rheumatoid arthritis	3%

60% report QALYs or DALYs




EMR

Eastern Mediterranean Region

n = 694


0.28 per 100,000 DALYs



Top three diseases

Maternal & newborn health	7%
Viral hepatitis	6%
Diabetes	5%

51% report QALYs or DALYs




WPR

Western Pacific Region

n = 6,492


0.99 per 100,000 DALYs



Top three diseases

Lung cancer	5%
Viral hepatitis	3%
Breast cancer	3%

56% report QALYs or DALYs



Full economic evaluations only; studies naming multiple regions contribute to each. Disease dots show fractional shares across three GBD classes among disease-coded evaluations; one dot is about 1%. Utility bar: QALY or DALY reporting; orange marker: all linked evaluations. Trends: cumulative share of countries with ≥1 evaluation (solid), ≥2 (dashed), all regions ≥1 (grey); 2026 excluded as incomplete.