

Studies by main type of intervention, and the share set in low- and middle-income countries

A | What is studied: each dot is a study, placed near studies with similar abstracts



B | Where it is studied: share of each intervention type in low- and middle-income settings



Primary intervention

- Behavioural & preventive (24% L&MICs)
- Care models & organisation (23% L&MICs)
- Devices & supplies (18% L&MICs)
- Financing & payment policy (33% L&MICs)
- Other (29% L&MICs)
- Pharmaceuticals (31% L&MICs)
- Screening & diagnosis (25% L&MICs)
- Surgery & procedures (23% L&MICs)
- Vaccines (43% L&MICs)

A: positions come from abstract wording (text embeddings, UMAP; frozen topic coordinates from the September run, filtered to the current studies), so studies group by topic, mainly disease area; the axes have no meaning. Grey: no topic cluster or no intervention label. 39,598 studies (27 have no coordinates). B: One dot = 20 single-country studies; darker = high-income settings, lighter (filling from the bottom) = low- and middle-income settings. Circle area = number of studies. Ordered by L&MICs share. Base: 27,417 single-country studies (reviews only if they include an original evaluation). Income: World Bank group in force on 1 January of the publication year. Intervention: the primary class assigned to each study's abstract by a language model (secondary classes not counted).