The Geometric Identity of Gravity and Dimensional Unification Resolving α, Lepton $(g-2)_l$, Weinberg, and Cabibbo Mixing

Łukasz Smoliński[1]

[1]Independent Researcher, 61-160 Czapury, Poland

November 12, 2026
Version: 5.2.4

Abstract

A unified geometric wave model is proposed, deriving the Gravitational Constant (G), the Fine-Structure Constant (α), and lepton anomalous magnetic moments (a_l) from a single structural modulator: the geometric vacuum stiffness ϵ_M. G is established as a precise, $\hbar$-independent identity rooted in the soliton's wave geometry. The derivation uses the ideal BCC projection $L_p^{\text{geom}} = 2/\sqrt{3}$; the remaining deviation from the measured value reflects lattice packing impedance, not a free parameter. The same ϵ_M factor governs a recursive nodal integration that predicts the electron anomalous magnetic moment from pure geometry and yields the full muon and tau anomalous magnetic moments via a unified dimensional projection over the BCC lattice, without perturbative QED calibrations. The model contains four empirical anchors (r_e, m_e, c, e) that fix the dimensional conversion between the geometric prediction and the SI-measured values, together with one explicit discrete postulate, the muon Fibonacci label $L_\mu = F_5 = 5$. The electron wave-centre count $K_{WC} = 10$ is geometrically fixed, with its dynamical stability proof noted as an open objective. Beyond these, no continuously adjustable parameters are introduced. All geometric operators are parameter-free functional forms of these two integers, no continuously adjustable quantity appears anywhere in the framework. The framework posits that the observed invariance of G is a low-field artefact of a dynamic geometric equilibrium, predicting testable deviations ($G_{eff} \neq G$) in environments with strong local magnetic moments where this equilibrium is exceeded. Crucially, the derivation of α and the anomalous magnetic moments (a_l) are shown to be independent of mass and charge, revealing that these constants are intrinsic topological properties of the vacuum lattice rather than secondary products of particle-field interactions. This geometric mapping reveals a dimensional hierarchy (π^5, π^6), resolving the Weinberg angle and Cabibbo mixing as emergent consequences of the vacuum's surface and volumetric resonance modes. Consequently, the divergence of fundamental forces is identified as a dimensional artefact of the unified BCC lattice, establishing a structural foundation for cosmological scaling. The cosmological extension of the framework uses one further anchor, the CMB temperature $T_{\text{cmb},0}$, together with the exact SI conversion constant k_B, and yields the Hubble constant and the de Sitter cosmological constant as sub-percent predictions.

Keywords: Anomalous Magnetic Moments, Fine-Structure Constant, Gravitational Constant, Weinberg angle, Cabibbo angle, Cosmological Constant, BCC lattice, Geometric Origin, Unification Theory.

Contents

Change Log **9**

1 Geometric Hierarchy of EWT and Spacetime Elasticity **13**
1.1 Wheeler's Vision of Quantum Vacuum and the Path to EWT 13
1.2 The Elastic Medium Constituent (EMC) 14
1.3 The Wave Center (WC) 14
1.4 The Soliton 15
1.5 Elastic Interaction (Hooke's Law) 15
1.6 From Energy Domain to Geometric Domain 16

2 Geometric Equation of the Fine-Structure Constant and the Deficit Terms **17**
2.1 BCC Lattice Geometry and the Geometric Ladder 19
2.2 Geometric Magnetic Deficit ϵ_M 19
2.3 The Fundamental Lattice Response Parameter ϵ_M 20

3 Dimensional Hierarchy and Dynamic Resonant Modulations **21**
3.1 The Universal Geometric Modulators: Substrate Properties 21
3.1.1 Unified Local Constant $\mathbf{C}_{\text{local}}$ 21
3.1.2 The π^6 Resonance: Volumetric Bosonic Coupling 21
3.1.3 The π^7 Resonance: Charge-Induced Amplitude Loading 22
3.1.4 The Mixed W–Z Vertex: Harmonic Operator 22
3.1.5 The π^5 Resonance: Flavor Mixing Deficit 23
3.2 Geometric Mixing Predictions: Higgs and Cabibbo Sectors 24
3.2.1 Higgs-Vector Boson Mixing 24
3.2.2 Cabibbo Mixing Angle 25
3.2.3 Dimensional Budget of Resonances: Reflection vs. Mixing 25
3.2.4 Symmetry of Observables: Energy Density vs. Amplitude Interference . . 26
3.2.5 Interpretation of Quark Masses in the EWT Framework 27
3.3 Structural Transition to N-Stiffness: From Geometric Volume to Lattice Impedance 28
3.4 Observable-Based Experimental Tests of the EWT Higgs Sector 28
3.4.1 Observable definition and recoil-mass reconstruction 29
3.4.2 Pillar I: absolute ZH cross section 29
3.4.3 Pillar II: angular observables and amplitude structure 30
3.4.4 Why an overall form factor is insufficient to change the normalized shape 31
3.4.5 Falsification criteria and scope of the test 31

4 Analysis of Neutrino Boundary Conditions **32**
4.1 Standard Conditions: Mass and Minimal Magnetism 32
4.2 Extreme Conditions: Wave Center in a Geometric Black Hole (GBH) 33

5 Neutrino as the Geometric Anchor **34**
5.1 Planck Charge from Geometric Alpha . . . 34
5.2 Decomposition of the Scaling Factor $S = r_\nu/q_P$ . . . 34
5.3 Geometric Fixed Point of g_v . . . 35
5.4 Resulting Neutrino Radius . . . 35
5.5 Decadic Resonance Test: $r_e/r_\nu \approx 100$ . . . 36
5.6 Physical Interpretation: ϵ_M as a Global Lattice Parameter . . . 36
5.7 Topological Necessity of r_ν . . . 36

6 Planck Length and Reduced Planck Constant from Geometry **37**
6.1 Reduced Planck Constant from Geometric Alpha . . . 37

7 The Physical Origin of Charge as Wave Amplitude **37**
7.1 Charge as Spatial Wave Amplitude . . . 37
7.2 Conversion of the Coulomb and Distinction from CGS . . . 38
7.3 Consequence for the Planck Charge q_P . . . 38

8 Geometric Deficit and Density Hierarchy of the Neutrino **38**
8.1 Unified Geometric Coupling . . . 39

9 Speed of Light as the Metric of Spacetime **39**
9.1 The Causal Structure of the BCC Lattice . . . 39
9.2 Axiom: The Maximum Propagation Speed . . . 40
9.3 Consequences: Density-Dependent Propagation Speed . . . 41

10 The Physical Origin of Relativity: Wave-Mechanical Lorentz Contraction **42**
10.1 Doppler-Induced Compression of the Standing-Wave Soliton . . . 42
10.2 The Michelson–Morley Null Result as a Mechanical Necessity . . . 43

11 The Geometric Identity of G: Derivation and Consistency **44**
11.1 Derivation of G from Electron Scaling (The $\hbar$-Independent Base) . . . 44
11.2 The Gravitational Identity Flow: From Electromagnetic Base to Surface Transition 45
11.3 Formalization of Soliton Geometric Density $\rho(r)$ . . . 46
11.4 Numerical Verification and the Unitary Mapping Operator $\mathcal{U}$ . . . 47
11.5 Formal Definition of the Operator $\mathcal{U}$: Mapping Geometric Parameters to G . . . 47
11.6 Geometric Emergence of the Planck Length . . . 48
11.7 Consistency with the Planck Charge . . . 51
11.8 Electromagnetic Consistency Check . . . 51
11.9 Status of Inputs . . . 52
11.10 Duality of Soliton Density . . . 53
11.11 The Degraded EMC Wall: Spherical Masking and Isotropy . . . 53
11.11.1 Open question: Wall density relative to statutory background . . . 54

12 Local Metric Phenomena from EMC Lattice Deformation **54**
12.1 The Two Faces of EMC Displacement: Speed and Trajectory . . . 54
12.2 Bridging the Vector Displacement to the Scalar Refractive Index . . . 55
12.3 Asymptotic Continuous Limit and Schwarzschild Equivalence . . . 55
12.4 Mechanical Origin of Gravitational Redshift in the EMC Lattice . . . 56
12.5 Shapiro Delay from the EMC Refractive Index . . . 57
12.6 Macroscopic EMC Density Profile of a Star . . . 58

12.7 Derivation of the Far-Field EMC Density Profile . . . 58
12.8 From Microscopic EMC Displacement to the Gravitational Radius . . . 59
12.9 Emergent Metric Encodings from Lattice Dynamics . . . 60
12.10 Summary of Local Metric Observables . . . 62
12.11 In-Platform Numerical Validation . . . 62

13 Newtonian Force from Interacting EMC Deficits **63**
13.1 In-Platform Validation: Normalization Consistency . . . 64

14 Relation to Existing Emergent Gravity Frameworks **65**
14.1 Comparison of Mechanisms and Scales . . . 65
14.1.1 Microscopic Degrees of Freedom (DoF) . . . 65
14.1.2 Source of Gravitational Emergence . . . 66
14.1.3 Scaling Factor for **G** . . . 66
14.1.4 Evolution of the EWT Framework: From Wave Attenuation to EMC Push out . . . 67
14.1.5 EWT as Pressure-Driven Emergent Gravity . . . 68

15 Geometric Validation: The Fundamental Identity and the Base AMM State (a_e) **69**
15.1 Geometric Mass-to-Radius Identity ($E \propto r^5$) . . . 69
15.2 Physical Origin of the r^5 Scaling: Geometric Energy Density . . . 69
15.3 The Geometric Status of Mass: Kilogram as a Derived Unit . . . 70
15.4 Geometric Dimension of the Reduced Planck and G Constants . . . 71
15.5 The Fundamental Magnetic Deficit ϵ_M . . . 72
15.6 Derivation of the a_e Base Formula . . . 72

16 The Recursive Lepton Hierarchy: Nodal Shell Resonance Model **72**
16.1 Resonance Growth Law and Fibonacci Labeling . . . 73
16.2 Discrete Resonance Labels of the BCC Lattice . . . 73
16.3 Structural Analysis: The Onion Model . . . 75
16.4 The Recursive Master Equation: Nodal Latching via Geometric Coupling . . . 75
16.4.1 Correspondence between Mass Scaling and Shell Geometry . . . 79
16.5 Final Results and Experimental Correlation . . . 79
16.6 Termination at Three Generations: Fibonacci Labels and a Shell-Closure Criterion 82
16.7 Predictions for Lepton Properties: The First-Principles AMM Predictions . . . 84
16.8 Physical Interpretation: The Mechanism of Topological Nodal Inclusion . . . 85

17 Sensitivity Analysis: Verification of Computational Variants **87**
17.1 Numerical Convergence and Error Landscape . . . 87
17.2 EWT Standalone vs. Best Resonance Peak . . . 87
17.3 3D Stability Mapping and Topography . . . 88
17.4 Verification of Geometric Scaling . . . 90
17.5 Critique of the Standard Model Target Bias: The Circularity of QED Precision . 90

18 The Magneto-Geometric Hypotheses: Dynamic Deficit Modification **91**
18.1 Hypothesis 1: The Push-Out Mechanism . . . 91
18.2 Hypothesis 2: The Consolidation Mechanism . . . 92
18.3 Hypothesis 3: No Magnetic Influence . . . 92
18.4 Summary of Hypotheses Testing . . . 92

18.5 Dynamic Equilibrium and Geometric Compensation 92

19 Numerical Verification of EWT Mass Scaling and Structural Sequences 93
19.1 EWT particle mass prediction 93
19.2 Implementation and Computational Results 93
19.3 Universal $(K_{WC})^5$ Meson-Mode Scan: Independent Mass Validation 94
19.3.1 Key conclusions from the meson-mode scan 94
19.4 Topological Isomorphism and Geometric Interference in BCC Lattice 96
19.4.1 Dual Resonance Modes: Orbital vs. Meson Scaling 96
19.4.2 Validation via the Ξ_{cc} Isospin Doublet 96
19.4.3 New vector state B_c^{*+} from ATLAS 2026. 97
19.4.4 New doubly charmed state Ω_{cc}^* from CERN 2026 97
19.4.5 Discussion and Cross-Mode Consistency 97

20 Geometric Radius Predictions: From Vacuum Anchor to Heavy Bosons 98
20.1 Numerical Foundation: Spherical Mode Accuracy 98
20.2 Predictive Radius for Heavy Bosons 98
20.3 Cross-Scale Validation with Nuclear Equivalents 98
20.4 Conclusion on Geometric Consistency 99

21 The Geometric Identity of Stiffness: Linking $N_{\mathbf{geom}}$ to the Dimensional Ladder 99
21.1 Eliminating the Fine-Tuning Paradigm 100
21.2 The Topological Reduction of the Fundamental Lattice Response Parameter: $\epsilon_M = \frac{1}{8\pi^7(1-\zeta)}$ 100
21.2.1 Physical Interpretation of the $8\pi^7$ Attractor 101
21.2.2 The Zero-Calibration Fine-Structure Constant 101

22 The Geometric Unification of Lepton Properties 101
22.1 The Unified Identity of the Lepton 102
22.2 Geometric Determinism of the Electron AMM 102

23 Other Geometric Constants Derived from the Lattice 102
23.1 The Rydberg Constant (R_∞): An Atomic Resonance Gap 102
23.1.1 Geometric Derivation: The $\alpha^3/(4\pi r_e)$ Identity 103
23.1.2 Zero-Calibration Identity 103
23.1.3 Numerical Verification 103
23.1.4 Physical Interpretation 103
23.2 The Bohr Radius (a_0): The Atomic Scale 104
23.3 The Electron Compton Wavelength (λ_C): Threshold of Annihilation 104
23.4 Consistency Across Atomic Scales 105

24 The Cosmological Constant from the BCC Lattice 105
24.1 The EWT trinity (derived) 106
24.2 The vacuum deficit ϵ_M and the horizon factor H 106
24.3 Lattice discrimination under the phase-unit ansatz 108
24.4 Planck temperature and the CMB anchor 108
24.5 Prediction of H_0 from the BCC horizon model 109
24.6 The de Sitter limit of the cosmological constant 110
24.7 The effective vacuum fraction from the BCC lattice 110
24.8 Status of the cosmological sector 113

24.9 What is not derived: the relaxation dynamics of the giant soliton 114
24.10What this establishes and what it does not 116

25 Mathematical Foundations of the Energy Wave Theory Lattice 117
25.1 Introduction 117
25.2 The Space Group of the Vacuum Lattice: $Im\bar{3}m \times SO(3)_{\text{local}}$ 118
25.2.1 Global Symmetry: The BCC Space Group $Im\bar{3}m$ (No. 229) 118
25.2.2 Local Symmetry: The Spherical Constituent $SO(3)_{\text{local}}$ 118
25.3 The Topological Class of Solitons: Winding Numbers and Cohomology 118
25.3.1 The Electron Soliton as a Winding Number on S^2 118
25.3.2 Higher-Generation Leptons: Topology of Nested Shells 119
25.4 The Dimensional Weight Operator and Degree-of-Freedom Algebra 119
25.5 Synthesis: The Mathematical Unity of EWT 121
25.6 Formalism of the Vacuum Push-out Mechanism 121
25.6.1 The Vacuum Stiffness Tensor and Sakharov Isomorphism 122
25.6.2 The Impedance Mismatch Operator 122
25.7 Isotropy and Spherical Masking: From Asymmetric Core to Isotropic Gravity . . 122
25.7.1 Geometric Necessity of Asymmetry 122
25.7.2 Intrinsic Sphericity of Standing Waves 123
25.7.3 Geometric Low-Pass Filter 123
25.7.4 Minimization of Lattice Shear Stress 123
25.7.5 Domain Separation: r^5 vs. r^3 123
25.7.6 The Principle of Geometric Masking 124
25.7.7 Gravity as a Result of Construction Material 124
25.7.8 Universal Applicability: From Leptons to Hadrons 124
25.8 The Uniqueness of the BCC Topology: Why $N_{geometric} = 8\pi^4$ Excludes Other Lattices 124
25.8.1 The Coordination Number and Stiffness Distribution 125
25.8.2 Packing Fraction and the ζ Correction 125
25.8.3 Topological Consistency: The $2\pi^2$ Factor and BCC Axes 125
25.9 Mechanical Resolution of Field Paradoxes 126

26 Nonlinear Stabilisation of the Electron Soliton 126
26.1 General Soliton Wave Equation 126
26.2 Coupling Coefficient from BCC Geometry 127
26.3 Proposed Forms of the Nonlinear Term 128
26.4 Structural Emergence of the 1-3-6 Geometry 129
26.5 Topological Selection Rule and the Electron Ground State 129
26.6 Connection to the Dimensional Weight Operator 130
26.7 Geometric Estimate of the Degraded EMC Wall Radius 130
26.8 Relation Between the Energetic and Topological Arguments 132
26.9 Summary 132

27 Predictive Power 133
27.1 Prediction of the Fundamental Value of G 133
27.2 Predictions for Lepton Properties: The First-Principles AMM Predictions 133
27.3 Cosmological Predictions: Recovered Relation and Predicted Constants 134
27.4 Magnetic Sensitivity $G(B)$ and Hypotheses Testing 134
27.5 AMM as a Strong Test of Geometric Universality 135

27.6 Correlations with Neutrino Flux . . . 135
27.7 Spectral and Geometric Modal Tests . . . 135
27.8 Tests using Gravitational Wave Observations . . . 135
27.9 Test on Bose-Einstein Condensates . . . 135
27.10 The Higgs Soliton vs. Fundamental Scalar Field . . . 135

28 Falsifiability and Model Constraints 135
28.1 Practical Considerations and Detection Limits . . . 136
28.2 Two Readings of the Higgs Sector: What a Future Measurement Can Settle . . . 137

29 Robustness Analysis: Geometric Stability and Sensitivity 138
29.1 Robustness Analysis of the Neutrino Soliton Radius and Geometric Identity . . . 138
29.2 Gravitational Surface Transition and $N_{\nu,\text{eff}}$ . . . 139
29.3 Sensitivity of α^{-1} to the Geometric Modulator N_{geom} . . . 140
29.4 Phase Space Analysis: EWT Trajectory vs Standard Model Interpretation . . . 141
29.5 Parametric Unification: The Stability Path of G and α . . . 141
29.6 EWT Unification: Convergence of G and a_e on the Vacuum Stiffness Isentrope . 142
29.7 Robustness Analysis: Geometric Stability and Vacuum Elasticity . . . 143
29.8 Structural Robustness and Resonance Stability of the Lepton Hierarchy . . . 143
29.9 Electroweak and CKM Coincidence: Geometric Robustness of $\sin^2\theta_W$ and $\sin\theta_C$ 144
29.10 Lock-In Test: Independent Determination of N_{geom} . . . 144
29.10.1 Method . . . 145
29.10.2 Results . . . 146
29.10.3 PRIMARY sectors: G, a_τ, and $\sin\theta_C$ . . . 146
29.10.4 WEAK sectors: a_e, a_μ, and $\sin^2\theta_W$ . . . 148
29.10.5 NO CONSTRAINT: α^{-1} . . . 148
29.10.6 Interpretation . . . 148

30 Geometric Phase Transitions: From Neutrino to Black Hole with EMC Wall 149
30.1 Hierarchy of Vacuum States and the G-Operating Point . . . 149
30.2 The Erasure of "Dark" Placeholders . . . 151
30.3 Resolution of Historically "Unsolvable" Paradoxes . . . 152
30.4 Density Duality: From Local Refraction to Cosmological Redshift . . . 153

31 Outlook and Computational Tools 154

32 Conclusion: The Geometric Paradigm, Correlated Lepton Anomalies, and Emergent Gravity 155
32.1 Geometric Unification in a Nutshell: The ϵ_M Flowchart . . . 156
32.2 The Unitary Geometric Path . . . 156
32.3 The Geometry of the Constituent: From Cubic Grids to Spherical Units . . . 157
32.4 Pillar 1: The N_ν Deficit and Soliton Stability . . . 157
32.5 Pillar 2: The Dynamic Scaling Operator Ω_{Push} . . . 158
32.6 Pillar 3: The Recursive Paradigm — Hierarchical Nodal Integration . . . 158
32.7 The Soliton Radius and Neutrino Interaction Cross-Section . . . 158
32.8 Paradigm Shift: The Planck Length as a Physical Boundary . . . 159
32.9 The ϵ_M Factor: Universal Geometric Modulator . . . 159
32.10 Geometric Self-Consistency of the Planck–Gravity–Metric Triangle . . . 159
32.11 Final Synthesis . . . 160

A Appendix **161**
A.1 List of Model Parameters and Physical Constants 161
A.2 Inventory of Inputs, Postulates, and Parameter-Free Forms 162
A.3 Computational Scripts (Python) 165
A.3.1 Emergence Engine 166
A.3.2 Emergence Engine Output 184
A.3.3 Extended Lepton AMM 186
A.3.4 Extended Lepton AMM Output 189
A.3.5 AMM Resonance Scanner 189
A.3.6 AMM Resonance Scanner Output 192
A.3.7 Electroweak and Boson Module 193
A.3.8 Electroweak and Boson Module Output 201
A.3.9 Particle Mass Module 202
A.3.10 Particle Mass Module Output 205
A.3.11 Robustness and Stability Plots 208
A.3.12 Robustness and Stability Plots Output 216

Change Log

Version	Date	Changes
5.2.4	2026.11.10	Rewrote the local-metric section: replaced the vector displacement field $\vec{u}(r)$ with a scalar density deficit $\delta(r)$ (new eqs. `eq:density_deficit`, `eq:delta_r`, `eq:n_gamma_exact`); rewrote the Doppler section with an arithmetic/geometric mean construction (new eqs. `eq:doppler_means`, `eq:lorentz_ratio`). Added geometric interpretation of the emission surface $S = 4\pi l^2 + (\pi r l + \pi r^2)$ (cone inscribed in sphere) and an explicit natural-units convention paragraph. Added q_P to inventory class IV. Updated `tab:final_results_confirmed` (tau † note: Shell Ref. and Full AMM coincide via $\mathcal{O}_\tau = 1$) and `tab:target_comparison` (corrected scanner values; the 0.0012 ppm engine–scanner discrepancy is resolved). Renamed `Q_P_INPUT`→`Q_P_REFERENCE`. Removed a duplicated paragraph in the Higgs–vector mixing section. Fixed SNO bibliography (Charged-Current → Neutral-Current), typos, and removed the boxed "Remark on the Muon Anomaly."
5.2.3	2026.11.09	Fixed cubic nonlinearity sign ($+\gamma\Psi^3 \to -\gamma\Psi^3$, self-focusing) and added Derrick-instability note. Replaced scalar winding number with degree of map $\mathbf{n} : S^2 \to S^2$ (Manton–Sutcliffe 2004); $Q = K_{WC} = 10$ now stated as structural postulate. Added inventory class VI for legacy EWT constants (diagnostic only), r_ν as derived quantity, e-notation warning, and K_{WC} vs. Q distinction. Fixed `ewt_robustness_plots.py`: hardcoded A_π, C_{fermion}, quark masses, and N_ν replaced with model-derived values; modules 6 and 9 use real computations instead of synthetic placeholders. Minor encoding fix.
5.2.2	2026.11.08	Replaced the deprecated quadratic fermionic correction $C_{\text{fermion}} = (1 + \pi^5 C_{\text{local}})^2$ with a linear flavor deficit $\epsilon_{\text{flavor}} = 1/(8\pi^5(1-\zeta))$, structurally analogous to the magnetic deficit $\epsilon_M = 1/(8\pi^7(1-\zeta))$ at the charged weak rung (both follow $\epsilon_n = 1/(8\pi^n(1-\zeta))$ with $n \in \{5, 7\}$). The Cabibbo relation becomes $\sin\theta_C = \sqrt{M_d/M_s}\,(1+\epsilon_{\text{flavor}})$, the bare Gatto–Sartori–Tonin amplitude multiplied by a single lattice deficit, consistent with the linear form of the bosonic corrections. The previous quadratic form required non-standard quark masses to agree with experiment; the new form uses standard PDG 2024 masses (and a PDG 2022 variant for comparison), improving the agreement with the measured Cabibbo angle. In the N-Stiffness section the flavor deficit is expressed as $\epsilon_{\text{flavor}} = 1/(\pi\, N_{\text{geom}})$, the natural π^1 suppression analogous to the π^3 suppression of C_{gap}. Updated the inventory, the electroweak module `ewt_electroweak_bosons.py`, and the engine `ewt_emergence_engine.py` (`build_geometric_ladder` now takes the lattice impedance ζ); console outputs in Appendix A.3 regenerated.

continued on next page

Version	Date	Changes
5.2.1	2026.11.02	Renamed the mixed W–Z operator from "harmonic mean" to "parallel-impedance combination", reflecting the correct mathematical structure of $\pi^6\pi^7/(\pi^6+\pi^7)$ as the parallel combination of two channel impedances. Replaced the informal "$\pi^7 \to \infty$" decoupling argument with an explicit uniqueness criterion: π^6 and π^7 are fixed values, and the limit is taken on the effective impedance, not on the rung number. Added a discussion of the SM indirect electroweak fit ($M_W = 80.353 \pm 0.006$ GeV) in Section 3.1.4, noting that the ±6 MeV SM uncertainty is input-parameter propagation only; the realistic theory uncertainty (±10–15 MeV) makes the 26 MeV EWT–SM difference $\approx 2\sigma$, not 4.3σ. Extended the Higgs–vector mixing section with an explicit note on the auxiliary spherical-mode masses (and their dependence on the legacy wave constants of inventory class V) and on the SME bounds on Lorentz-violating operators ($\sim 10^{-16}$), which make on-shell scalar–vector mixing experimentally indistinguishable from the SM prohibition. Added the Gatto–Sartori–Tonin (1968) and Kleinert (1987) references.
5.2.0	2026.11.02	Introduced the mixed W–Z vertex operator, a harmonic combination $O_{\mathrm{mixed}} = \pi^6\pi^7/(\pi^6+\pi^7)$ of the neutral and charged weak rungs. The pure π^6 operator is retained for genuinely neutral vertices (Z–Z, Z–H); the mixed operator is used for the W–Z vertex and for the W–H mixing angle. Updated inventory table.
5.1.8	2026.10.30	Extended the Abstract with explicit accounting of the four empirical anchors (r_e, m_e, c, e), the single discrete postulate $L_\mu = F_5 = 5$, and the geometrically fixed integer $K_{WC} = 10$. Added Appendix, a complete inventory classifying every model element as anchor, test anchor, postulate, geometric integer, or parameter-free functional form. Downgraded the null-probe metric encoding $v_\gamma = \eta$ from derived to postulated, with explicit open-item remarks (M4.13) and falsification conditions. Introduced a uniform ppm convention in the AMM section, with reference to `ewt_amm_extended.py`, and reframed the lepton anomalies as three independent evaluations of the same geometric identity. Corrected the B_c^{*+} wave-centre count from $K_{WC} = 59$ to $K_{WC} = 66$ and removed the erroneous "Topological Isomorphism and Dual Realisation" subsection built on the wrong value.
5.1.7	2026.10.22	Corrected the optical refractive index from $n_\gamma = (1-2r_s/r)^{-1/2}$ to $n_\gamma = \eta^{-1} = (1-r_s/r)^{-1}$, which matches the exact Schwarzschild optical metric factor and removes the spurious singularity at $r = 2r_s$. Split the derivation into the timelike encoding $v_{\mathrm{clock}} = \sqrt{\eta}$ and the null-probe encoding $n_\gamma = \sqrt{g_{rr}/g_{00}} = \eta^{-1}$, closing the algebraic gap. Updated the M4.3 and M4.5 numerical entries in the local-metric validation table.

continued on next page

Version	Date	Changes
5.1.6	2026.10.18	Revised the electromagnetic consistency check: the SI identity $e^2 = 4\pi\epsilon_0 m_e c^2 r_e$ is now written in EWT natural units as $x\,e^2 = m_e r_e$ with the dimensionful conversion factor $x = \mu_0/(4\pi) = 10^{-7}\,\mathrm{m}^4$. Added the explicit dimensional check $[x][e^2] = [\mathrm{m}^6] = [m_e][r_e]$ and clarified that x is a unit-conversion quantity (structural form from α, e, h, c; numerical value from SI conventions), with $k_e = c^2 x$ in SI and $k_e = x$ only in the $c = 1$ limit. In the Remark on ϵ_0, the role of the vacuum permittivity is absorbed into x.
5.1.5	2026.10.17	Stated the muon label $L_\mu = F_5 = 5$ explicitly as the model's sole discrete postulate in "Status of Inputs" and "Final Synthesis". Aligned Pillar 3 with the Fibonacci label terminology ($L_\mu = F_5$, $L_\tau = F_9$) and corrected "wave centres" to "lattice nodes" in the closure criterion. Removed a residual fragment from the Final Synthesis.
5.1.4	2026.10.17	Introduced a unified discrete operator for the nodal shell hierarchy, $\Delta K_k = \mathrm{round}(10^k \pi^2)$ for the electron core ($k = 0$) and $\mathrm{round}(10^k \cdot 2\pi^2)$ for the shells ($k \geq 1$), which generates $K_n = 10, 207, 2181, 21920$ without a hard-coded seed. Added the "Discrete Resonance Labels of the BCC Lattice" section with Fibonacci labels $F_{1+4k} = 1, 5, 34, 233$ and the "Termination at Three Generations" section with a candidate closure criterion $K_n/L_n^2 > 1$, satisfied by the three observed generations and violated by the formally generated fourth. Softened stability language in the physical interpretation, reconciled "interface tension" with "interface area" and "interface contribution", and rephrased "topological invariant" as "topologically determined quantity". Adopted the symmetric Lorentz–Heaviside convention $\epsilon_0 = \mu_0 = 1$ ($e^2 = 4\pi m_e r_e$); minor typographical corrections.
5.1.3	2026.10.16	Corrected the refractive-index encoding to $n_\gamma = (2\eta - 1)^{-1/2}$ and distinguished timelike from null probes in the Summary of Local Metric Observables. Added the algebraic derivation of $\mathcal{O}_\mu = M_\mu/N_{\mathrm{geom}} = 1/(4\pi^2)$ and clarified the dimensional origin of the interface tension $\delta = L_\mu^2 = 25$. Added notes on the role of s_μ in σ_τ, on orbital-mode internal-consistency tests, and on the status of ϵ_0. Removed the VEV-falsification paragraphs and the "deeper mathematical structure" claim; rephrased the muon-anomaly remark. Minor typographical corrections.

continued on next page

Version	Date	Changes
5.1.2	2026.10.09	Replaced the "Mutual Falsification: The Ultimate Test" subsection with "Two Readings of the Higgs Sector: What a Future Measurement Can Settle". Stated the three Standard-Model theses on the Higgs explicitly (fundamental scalar, fundamental VEV, fundamental mass mechanism) and contrasted them with the EWT reading. Removed the claim that a confirmed substructure would make the Standard Model "automatically fall"; the falsification now targets the interpretation, not the effective theory. Added a cross-reference to the X_{eff} equation. Minor cross-reference corrections.
5.1.1	2026.09.28	Replaced the preliminary Higgs-sector verification proposal with a fully formulated observable-based test section. Pillar I is now stated as a falsifiable zero-modification prediction under an explicit restricted tree-level hypothesis ($\mu_{ZH} = 1$); Pillar II is presented as a search framework using Legendre moments of the Z production angle, with an explicit proof that an s-only form factor leaves the normalised angular shape unchanged. Added experimental references.
5.1.0	2026.09.26	Corrected the r^5 scaling label ("Resonance Mode Count" instead of "Resonance Frequency"); added the Lock-In Test and in-platform validation subsections; added OpenWave M4 references; minor typographical corrections.

1 Geometric Hierarchy of EWT and Spacetime Elasticity

Energy Wave Theory (EWT) is based on a quantum medium and establishes a clear hierarchy of entities, starting from the Planck scale. This geometric structure is crucial for the elasticity of spacetime and maintaining the constancy of the speed of light [1].

1.1 Wheeler's Vision of Quantum Vacuum and the Path to EWT

The search for a discrete, pregeometric structure underlying spacetime has a rich intellectual history, with John Archibald Wheeler as one of its most profound and visionary architects. Wheeler's work spanned several interlocking themes, each of which resonates – sometimes supportively, sometimes critically – with the Energy Wave Theory (EWT) presented here.

Geometrodynamics and the 3-Geometry

Wheeler championed the idea that physics could be reduced to pure geometry [2]. In his geometrodynamic program, particles such as electrons and photons were to be understood as **geons** – gravitational and electromagnetic fields trapped by their own curvature, embodying the principle of "mass without mass" [6]. The Wheeler–DeWitt equation [5] attempted to quantise this 3-geometry, laying the foundation for canonical quantum gravity.

EWT shares the conviction that geometry is primary, but it replaces Wheeler's continuous 3-geometry with a **discrete body-centered cubic (BCC) lattice** of Elastic Medium Constituents (EMCs). The gravitational constant G emerges not from quantised continuum curvature, but from the volumetric packing deficit of these spherical units – a concrete, calculable pregeometry.

Quantum Foam and Pregeometry

Perhaps Wheeler's most radical proposal was that at the Planck scale, spacetime is no longer smooth but becomes a chaotic, fluctuating **quantum foam** [3] – a topological tapestry of wormholes and virtual geometries. He further argued that such a foam must be preceded by an even more fundamental structure: **pregeometry** [8], the information-theoretic or combinatorial substrate from which geometry itself crystallises.

In EWT, the quantum foam is replaced by a **static, ordered BCC crystal** with a well-defined packing fraction ($\eta \approx 0.68$) and a residual impedance ζ. This is not a foam but a **mechanical lattice**, whose elasticity (encoded in the effective geometric stiffness $N_{\text{geom}} = 8\pi^4(1-\zeta)$) gives rise to both gravitational and electromagnetic constants. Thus EWT provides a **concrete realisation of Wheeler's pregeometry**, replacing speculative foam with an engineerable lattice.

It from Bit – and the Alternative of Geometric Realism

Wheeler's famous aphorism **"it from bit"** [9] asserted that every physical entity ("it") derives from yes/no questions ("bits"), elevating information to the ontologically primitive level.

EWT takes a different stance. Here, the primitive is **geometric**: the BCC lattice and its spherical EMCs are not made of bits; they are the actual fabric of the vacuum. Information is an emergent property of geometric configurations, not their foundation. Consequently, EWT aligns more closely with a **geometric realism** than with Wheeler's informational idealism – a crucial philosophical fork that distinguishes the present work from the Wheelerian mainstream.

Evolution of Wheeler's Ideas in EWT

The table below summarises how each major Wheelerian concept has been transformed in the EWT framework:

Wheeler concept	Original meaning	EWT implementation / stance
Geometrodynamics	Continuous 3-geometry, geons	Discrete BCC lattice, EMC units, $N_{\text{final}} = 8\pi^4$
Quantum foam	Chaotic Planck-scale topology	Ordered BCC crystal, static packing deficit
Pregeometry	Abstract information substrate	Concrete BCC lattice with calculable elasticity
It from bit	Information ontological primacy	Superseded by Geometric Realism (Geometry precedes information)

Thus, while EWT draws inspiration from Wheeler's insistence on a pregeometric foundation, it replaces the fluid, information-theoretic speculations with a **rigid, calculable lattice** – a shift from "foam" to "crystal" and from "bit" to "geometric element". This evolution allows EWT to compute G, α, and the lepton anomalous magnetic moments from first principles, fulfilling Wheeler's dream of a pregeometry while discarding the elements that proved non-predictive.

1.2 The Elastic Medium Constituent (EMC)

The **Elastic Medium Constituent (EMC)** is the smallest, fundamental geometric entity, which serves as the physical quantization limit of the proposed medium. Its size and spacing are directly linked to the requirement for the medium to propagate the electromagnetic wave at the speed of light c.

- **Geometric Radius ($\mathbf{r_{EMC}}$):** In EWT, the EMC radius is typically defined as a magnitude on the order of 10^{-35} m, closely related to the **Planck Length** ($\lambda_l \approx 1.616 \times 10^{-35}$ m). This radius is the fundamental unit of length. Assuming the approximate value derived from the Planck scale:
$$r_{\text{EMC}} \approx 1.0 \times 10^{-35} \text{ m} \tag{1}$$

- **Inter-Constituent Distance ($\mathbf{d_{EMC}}$) and Wavelength Quantum ($\lambda_\mathbf{l}$):** The distance between the centers of adjacent EMCs defines the minimum wavelength, which directly influences the wave propagation speed c in the medium [1]. This distance is equal to the **Planck Length**:
$$d_{\text{EMC}} = \lambda_l \approx 1.616 \times 10^{-35} \text{ m} \tag{2}$$

- **Geometric Reference and Quantization:** The radius r_{EMC}, the distance d_{EMC} define the **absolute limit of spatial quantization** within the medium. The EMC is utilized as a geometric reference point for all larger structures.

1.3 The Wave Center (WC)

The Wave Center (WC) is defined as the focal point of oscillation in the elastic medium. The continuous interference of waves focused on this point establishes a standing wave structure (Soliton), which constitutes the localized oscillating volume of the medium. A single WC represents the fundamental unit of stored energy in the medium. Complex particles, such as the

electron, are modeled as structures composed of multiple WCs. The variable $\mathbf{K_{WC}}$ represents the **number of constituent Wave Centers** in a given composite structure. For example, in the derivation of the electron's mass and charge, K_{WC} is a crucial summation factor [26].

1.4 The Soliton

A **Soliton** (or particle) in EWT is a stable structure of standing waves created by one or more WCs.

- **Standing Wave Boundary:** The Soliton is defined by the boundary where the generated standing waves transition into traveling waves. This transition point defines the geometric radius of the particle.
- **Energy and Mass Relation:** The energy stored in the standing waves of the Soliton is the source of the particle's rest mass, linking the wave structure directly to the mass-energy equivalence $E = mc^2$ [26].

1.5 Elastic Interaction (Hooke's Law)

The stability of the geometric structure hinges upon the nature of the medium's internal forces. Interactions between the Elastic Medium Constituents are modeled as **elastic compression interactions (repulsion)**, a mechanism essential for Soliton stability and the propagation of energy.

- **Interaction Nature (Hooke's Law):** The force of repulsion between adjacent EMCs, when displaced from their equilibrium position (x), strictly follows **Hooke's Law** [11, 12]. This principle is interpreted as the fundamental **elasticity of the quantum medium**, which is necessary to **counteract enforce structural symmetry**:

$$\vec{F}_{\text{Hooke}} = -k\vec{x} \tag{3}$$

  where k is the **elastic constant** specific to the medium.
- **Significance for Solitons and Volume Symmetry:** This elastic interaction ensures that the medium prevents unlimited compression of the Constituents.

Linearization and density dependence. Hooke's law in Eq. (3) represents the linearized elastic response of the EMC pair potential around the statutory equilibrium spacing. The microscopic pair potential is nonlinear, $V(r) \propto r^{-n}$ (with $n = 1$ in the 1D reduction and $n = 3$ in the full 3D lattice), and the local effective spring constant is its second derivative evaluated at the instantaneous spacing $a(\eta) = 1/\eta$,

$$k(\eta) = \left.\frac{d^2V}{dr^2}\right|_{r=1/\eta} \propto \eta^{\,n+2}. \tag{4}$$

At the statutory equilibrium $\eta = 1$, this reduces to a constant tangent modulus $k(1)$, which is what Eq. (3) encodes. The density dependence $k(\eta) \propto \eta^{\,n+2}$ is used in the wave-speed derivation of Section 12.9, where the full nonlinear response is required to compute the density-dependent sound speed $v(\eta)$.

The macroscopic stiffness parameter N (as defined in Eq. 15) used in the derivation of the gravitational constant and anomalous magnetic moments is the emergent aggregate of the




Figure 1: **The Elastic Medium Constituent (EMC) and Hooke's Law.** The EMC is the minimal geometric unit (defined by $r_{\text{EMC}} \sim l_p$) where the fundamental elasticity ($\mathbf{k}$) of the medium manifests as a repulsive force $\mathbf{F} = -k\mathbf{x}$, ensuring the Soliton's stability against geometric collapse. **(Note: l_p denotes the Planck length, establishing the geometric scale for r_{EMC}.)**




Figure 2: **Elastic Interaction between two EMCs.** The geometric overlap (compression $\vec{\mathbf{x}}$) between adjacent Constituents generates the repulsive Hooke's Force $\mathbf{F} = -k\vec{\mathbf{x}}$. This mechanism is the source of Soliton stability, balancing the internal geometric deficit ($\epsilon_{\mathbf{G}}$).

microscopic elastic constants k of the BCC lattice substrate. While k defines the fundamental repulsive interaction between individual EMCs following Hooke's Law (3), N represents the global resistance of the vacuum to volumetric displacement. This connection bridges the localized elastic forces shown in Fig. 2 with the large-scale stability required for the precise determination of leptonic anomalies.

1.6 From Energy Domain to Geometric Domain

The Energy Wave Theory (EWT), as developed by Yee [46], successfully derived 23 fundamental physical constants from five wave constants: longitudinal amplitude A_l, longitudinal wavelength

λ_l, aether density ρ, wave speed c, and the electron's wave center count $K_{WC} = 10$. In that framework, charge was already reinterpreted as wave amplitude, measured in meters, hinting at an underlying geometric reality. The present work extends this program by identifying the physical substrate of these waves: a Body-Centered Cubic (BCC) lattice of spherical Elastic Medium Constituents (EMCs). The wave constants are thus replaced by geometric invariants of this lattice: the statutory neutrino radius r_ν (related to λ_l and K_{WC}), the effective nodal stiffness $N_{\text{geom}} = 8\pi^4(1-\zeta)$ (derived from the BCC coordination number and packing impedance), and the magnetic deficit $\epsilon_M = 1/(N_{\text{geom}}\pi^3)$ (encoding the 7-dimensional weak interaction scale). This transition from the energy domain to the geometric domain not only preserves the predictive power of EWT but also unifies gravity, electromagnetism, and the weak force under a single topological framework.

2 Geometric Equation of the Fine-Structure Constant and the Deficit Terms

The **fine-structure constant** (α) is defined within Energy Wave Theory (EWT) as a geometric ratio describing the relative strengths of fundamental forces—charge energy, magnetism, and gravity [10, 16].

The derivation of the inverse fine-structure constant ($1/\alpha$) is realized as the combination of a purely geometric core and a small topological correction. The core is obtained from the surface geometry of the soliton, while the correction is dictated by the BCC lattice structure.

This approach redefines fundamental particle properties:

- **Charge as Amplitude:** Electric charge is interpreted not as an abstract quantity, but as the physical **amplitude** (x) of a standing wave oscillation within a quantum medium.
- **Geometric Ratio for α:** The fine-structure constant is defined as the ratio of squared amplitudes, which is equated to a geometric ratio involving the total surface area of energy propagation (S):

$$\alpha = \frac{A_1^2}{A_0^2} = \frac{x^2}{S} \tag{5}$$

Geometric Soliton Core ($4\pi^3 + \pi^2 + \pi$)

The derivation of the pure geometric constant $\mathbf{A}_\pi$ was first proposed by Jeff Yee [10] within the framework of the Energy Wave Theory. The core postulate involves equating the fine-structure constant to a specific ratio of surface areas in the quantum background, and the derivation proceeds in the following steps:

Step 1: Defining the Total Surface Area S The total surface area S is the sum of the sphere surface area and the total cone surface area:

$$S = \underbrace{4\pi l^2}_{\text{Sphere Area}} + \underbrace{(\pi r l + \pi r^2)}_{\text{Total Cone Area}} \tag{6}$$

Step 2: Substitution of Geometric Relations ($r = x$, $l = \pi x$) Substituting $r = x$ and $l = \pi x$ expresses S solely in terms of π and x:

$$S = 4\pi(\pi x)^2 + (\pi(x)(\pi x) + \pi x^2) \tag{7}$$

Step 3: Simplification The expression is simplified by factoring out x^2:

$$S = 4\pi^3 x^2 + \pi^2 x^2 + \pi x^2 = x^2(4\pi^3 + \pi^2 + \pi) \tag{8}$$

Step 4: Final Geometric Fine-Structure Constant Substituting the simplified S into $\alpha = x^2/S$, the x^2 terms cancel, yielding α as a pure function of π:

$$\mathbf{A}_\pi{}^{-1} = \frac{1}{4\pi^3 + \pi^2 + \pi} \tag{9}$$

Kinematic Origin of the π Factor The relation $l = \pi x$ defines the propagation distance in terms of the source amplitude. It is not an arbitrary geometric choice but a consequence of the harmonic motion of the central EMC in an elastic medium with constant propagation speed c. In one complete oscillation the central granule travels a total longitudinal displacement of $2x$. During the same time interval, the disturbance emitted by this motion propagates outward at the lattice speed c. Because the central granule follows a sinusoidal path, its corresponding longitudinal phase advance is exactly πx, whereas the undisturbed wavefront travels the longer distance $l = \pi x$. The factor π therefore enters through the kinematics of harmonic oscillation, not as a fitted parameter. This mechanical origin of l is what ultimately generates the characteristic powers π^3, π^2, π in the fine-structure geometry.

> *"In the time that it takes for the center granule to vibrate, traveling a distance (r), and then returning (total of $2r$), the granules that it affects will have traveled a distance of length (l). The harmonic motion of the center granule over time can be represented by a sine wave, where the travel in one direction and a return to equilibrium is a half-wavelength, or π."*
>
> *"Assuming constant velocity for granules (expected to be the speed of light), the distance that the granules will have traveled is π times x. In other words, while the center granule travels a total distance of $2x$ during vibration, the straight-line motion of the granules it effects travels $\sim 3.14x$ because it doesn't slow down to account for a turn in direction."*
>
> — Yee, J. (2019). *The Relationship of the Fine Structure Constant and Pi*, p. 3–4. [10]

The decomposition $S = 4\pi l^2 + (\pi r l + \pi r^2)$ has a natural geometric reading. The cone has its apex at the center of the sphere and its base rim on the sphere's surface, with slant length $l = \pi x$ and base radius $r = x$. Its axial height is $h = \sqrt{l^2 - r^2} = x\sqrt{\pi^2 - 1} \approx 2.978\,x$, so the base rim lies exactly on the sphere of radius $l = \pi x$. The cone is therefore fully contained within the sphere, tangent to its surface along the rim.

The three terms of S represent three distinct surfaces of wave interaction: (i) the outer spherical wavefront $4\pi l^2$, where standing waves transition into outgoing traveling waves; (ii) the cone's lateral surface $\pi r l$, the internal channel along which the wave center's oscillation is directed toward the next unit cell; (iii) the cone's base cross-section πr^2, the effective emission aperture. Summing them gives the *total effective interaction area* of the soliton — the outer spherical flux plus the inner conical channel — rather than the geometric surface of a single disjoint boundary in $\mathbb{R}^3$.

The kinematic argument fixes $l = \pi x$ as the distance traveled by the affected granules during one half-oscillation. This is the *slant* distance along the cone's generatrix, from apex to rim, consistent with the geometric definition of l as the cone's slant length. The axial height $h = x\sqrt{\pi^2 - 1}$ is shorter than the path traveled, as expected for a wave expanding radially outward from the center: the granules follow the slant path, not the axial shortcut.

With these identifications, $S = x^2(4\pi^3 + \pi^2 + \pi)$, and $A_\pi \equiv S/x^2 = 4\pi^3 + \pi^2 + \pi$ is a geometric quantity fixed by the lattice topology (sphere + cone inside sphere) and by the kinematic relation $l = \pi x$. The numerical value $\alpha_\pi^{-1} = 1/A_\pi \approx 137.036$ agrees with $\alpha^{-1}_{\text{CODATA}}$ to six significant digits.

2.1 BCC Lattice Geometry and the Geometric Ladder

The vacuum lattice is a discrete body-centred cubic (BCC) lattice of Elastic Medium Constituents (EMCs). The BCC lattice is characterised by:

- 8 nearest neighbours,
- packing fraction

$$\eta_{\mathrm{BCC}} = \frac{\sqrt{3}\,\pi}{8}, \tag{10}$$

- ideal projection factor

$$L_p^{\mathrm{geom}} = \frac{2}{\sqrt{3}}, \tag{11}$$

The factor L_p^{geom} is the inverse of the dimensionless nearest-neighbour distance in the BCC lattice. It is not an adjustable constant but a direct consequence of the $Im\bar{3}m$ space group: the nearest neighbour lies at $d/a = \sqrt{3}/2$, and its inverse gives the geometric projection factor $2/\sqrt{3}$.

The BCC geometry introduces the constants π and $\sqrt{3}$ directly through the packing fraction. In later derivations, the face-diagonal factor $\sqrt{2}$ appears in the lattice projection operators and in the unified coupling bridge, while Euler's number e enters through the Eulerian dilution of the statutory vacuum density. Neither of these constants is fitted; each has a well-defined geometric role in the corresponding section.

The geometry of the lattice imposes a natural dimensional hierarchy on the soliton's wave structure. Each additional active degree of freedom contributes one factor of π, giving rise to the following geometric ladder:

Table 2: The Geometric Ladder: Dimensional Interaction Topology

Scale	Interaction	Budget Components	Dim.
π^7	Charged Weak	3D + $A + f + r + Q$	7
π^6	Neutral Weak	3D + $A + f + r$	6
π^5	Flavor Mixing	3D + $A + f$	5
π^4	Int. Binding	3D + A	4
π^3	Spatial Substrate	3D	3

Legend: A = amplitude, f = frequency, r = radius, Q = charge.

The factor π appearing in every rung of this ladder is not introduced by hand. Its origin is the same as in the geometric soliton core: the kinematic relation $l = \pi x$, which reflects one full harmonic phase advance of the central EMC during one oscillation. Each additional degree of freedom therefore multiplies the phase-space budget by one further factor of π.

The π^7 rung is particularly important: it represents the charged weak sector, in which the soliton engages seven independent geometric degrees of freedom. The BCC coordination number is 8. The magnetic deficit ϵ_M introduced below is therefore naturally associated with the inverse of the product $8\pi^7$, with a small correction due to the non-ideal spherical packing of the lattice.

2.2 Geometric Magnetic Deficit ϵ_M

The ideal stiffness of the BCC lattice follows from its eight nearest neighbours and the four-dimensional saturation budget:

$$N_{\mathrm{ideal}} = 8\pi^4. \tag{12}$$

The non-ideal spherical packing of the EMCs gives rise to a small lattice impedance

$$\zeta = \frac{1 - \eta_{\mathrm{BCC}}}{\eta_{\mathrm{BCC}} N_{\mathrm{ideal}}}. \tag{13}$$

Applying this impedance to the ideal stiffness yields the effective geometric stiffness:

$$N_{\mathrm{geom}} = N_{\mathrm{ideal}}(1 - \zeta). \tag{14}$$

The magnetic deficit is then defined by the key relation

$$\boxed{\epsilon_M = \frac{1}{N_{\mathrm{geom}} \pi^3} = \frac{1}{8\pi^7(1 - \zeta)}} \tag{15}$$

In the ideal continuum limit $\zeta \to 0$, this reduces to

$$\epsilon_M^{(\mathrm{ideal})} = \frac{1}{8\pi^7}. \tag{16}$$

The complete geometric inverse fine-structure constant therefore has the form

$$\frac{1}{\alpha} = A_\pi - \epsilon_M - \sum \epsilon_G, \tag{17}$$

where $\sum \epsilon_G$ is a lower-order gravitational deficit term. For the electron this term is negligible compared with ϵ_M, so the working expression used throughout this work is

$$\frac{1}{\alpha_{\mathrm{geom}}} = A_\pi - \epsilon_M. \tag{18}$$

The omitted lower-order term $\sum \epsilon_G$ is retained as a formal remainder; it is not fitted and may become relevant when the lower-order gravitational correction is no longer negligible.

This expression contains no calibrated parameters. All quantities on the right-hand side are fixed by the BCC lattice geometry and the mathematical constants π, e, and $\sqrt{2}$.

2.3 The Fundamental Lattice Response Parameter ϵ_M

A central role in the Enhanced EWT framework is played by the parameter ϵ_M, traditionally referred to as the *magnetic deficit factor*. Initially, this parameter was identified as a functional heuristic—a "magnetic deficit" required to reconcile the fine-structure constant α within the energy wave equations. However, as the framework evolved, it became evident that ϵ_M was not a localized adjustment, but a fundamental property defining the very "stiffness" of the 3D space occupied by matter.

Within the unified geometric context of this work, ϵ_M is more fundamentally defined as the **Global Lattice Impedance Constant**. It represents the intrinsic structural resistance of the BCC vacuum substrate to any deviation from ideal spherical wave symmetry. This realization transformed it from a specialized electromagnetic factor into a cornerstone of Enhanced EWT, acting as a universal "scaling bridge" across different physical sectors.

By treating ϵ_M as a singular topological property, the model achieves numerical convergence across disparate scales—from the gravitational constant G to the anomalous magnetic moments of leptons—without the need for independent empirical constants. The structural coherence of this approach is formally validated in Section 29, where it is shown that this single value functions as the primary scaling anchor for the entire theory.

Ultimately, this work reveals that ϵ_M is not an arbitrary input, but a direct consequence of the BCC lattice coordination. Its explicit geometric form is given in Eq. (15).

3 Dimensional Hierarchy and Dynamic Resonant Modulations

This section translates the geometric ladder introduced in Section 2 into concrete lattice operators for the bosonic and fermionic sectors. The mixed nature of the W–Z vertex — one charged soliton, one neutral soliton — requires a parallel-impedance combination of the π^6 and π^7 rungs. The resulting W-boson mass prediction lies within 2σ of the ATLAS 2024, CMS 2024, and PDG 2024 averages; the CDF II 2022 measurement, previously used as the primary anchor, is now recognized as an outlier at 5.8σ and is retained only as a benchmark.

3.1 The Universal Geometric Modulators: Substrate Properties

The formation of mixing angles within the EWT framework is governed by a hierarchy of modulators derived from the global magnetic deficit ϵ_M. These factors account for the intrinsic lattice impedance and the resonant displacement required to maintain equilibrium within the BCC vacuum substrate. Rather than being empirical parameters, these modulators represent the structural response of the lattice to different interaction topologies (volumetric vs. surface).

3.1.1 Unified Local Constant $\mathbf{C_{local}}$

The **Unified Local Constant** is the structural bridge between the global lattice magnetic deficit (ϵ_M) and the local chiral projection ($2\sqrt{2}$):

$$\boxed{\mathbf{C}_{\text{local}} \equiv \frac{\epsilon_M}{2\sqrt{2}}} \tag{19}$$

3.1.2 The π^6 Resonance: Volumetric Bosonic Coupling

Neutral vector bosons (Z, H) and neutral–neutral vertices are high-energy excitations involving the full six-dimensional phase space. The modulator $\mathbf{C}_{\text{gap}}$ accounts for the volumetric resistance of the lattice in a pure neutral vertex:

$$C_{\text{gap}} = 1 + \pi^6 \cdot C_{\text{local}} \tag{20}$$

This operator applies to interactions between two neutral solitons (Z–Z, Z–H). It does not describe the mixed W–Z vertex, which involves one charged and one neutral soliton (Section 3.1.4).

The Weinberg angle emerges as a structural equilibrium point between the 6D volumetric resonance (π^6) and the boson mass ratio:

$$\boxed{\sin^2\theta_W = 1 - \left(\frac{M_W}{M_Z}\right)^2 \cdot \frac{1}{C_{\text{gap}}}} \tag{21}$$

For a hypothetical purely neutral W boson (a useful limiting case), this operator gives the neutral attractor

$$M_W^{\text{neutral}} = M_Z^{\text{CODATA}} \cdot \sqrt{(1 - \sin^2\theta_W^{\text{Target}}) \cdot C_{\text{gap}}} \approx 80.514 \text{ GeV}. \tag{22}$$

The physical W boson is not purely neutral: it is a charged soliton, and the W–Z vertex is mixed. The relevant operator is therefore the parallel-impedance combination of the π^6 and π^7 rungs, derived in Section 3.1.4.

3.1.3 The π^7 Resonance: Charge-Induced Amplitude Loading

The highest rung of the geometric ladder, π^7, is associated with the charged weak sector ($W^{\pm}$). Unlike the neutral Z^0 and H^0 solitons, the W boson carries non-compensated wave amplitude (charge), which acts as an additional **7th degree of freedom** within the BCC substrate.

A purely charged vertex (for example, a hypothetical W–W interaction) would use the π^7 rung directly. The physical W–Z vertex is however *mixed*: one charged and one neutral soliton participate. The relevant operator is therefore the parallel-impedance combination of the π^6 and π^7 rungs, derived in Section 3.1.4, which yields $M_W \approx 80.379$ GeV.

3.1.4 The Mixed W–Z Vertex: Harmonic Operator

The W–Z vertex is not a pure rung of the geometric ladder. One soliton is charged ($W^{\pm}$, seven degrees of freedom: 3D + amplitude + frequency + radius + charge) and the other is neutral (Z^0, six degrees of freedom). The pure rungs π^6 (Section 3, neutral) and π^7 (charged) describe, respectively, purely neutral and purely charged interactions. The mixed W–Z vertex requires an operator that interpolates between them.

Two limiting cases constrain the required form. If the charged channel decouples (a hypothetical neutral W), the operator must reduce to the pure neutral rung π^6. If the neutral channel decouples, it must reduce to the pure charged rung π^7. Among the three elementary combinations of π^6 and π^7, only the *parallel-impedance combination*

$$O_{\text{mixed}} = \frac{\pi^6 \cdot \pi^7}{\pi^6 + \pi^7} \approx 729.26 \tag{23}$$

satisfies both limits. This is a mathematical property of the functional form — a uniqueness criterion among simple parameter-free combinations — not a statement about a physical process: $\pi^6 \approx 961.39$ and $\pi^7 \approx 3020.29$ are fixed values determined by the BCC coordination number, and the limit is taken on the effective impedance, not on the rung number. The arithmetic mean $(\pi^6 + \pi^7)/2$ and the geometric mean $\sqrt{\pi^6 \pi^7}$ do not satisfy both limits and cannot interpolate the two pure cases.

The parallel-impedance form is also the standard combination for parallel channels in physics (parallel resistance, series capacitance, reduced mass in a two-body problem). The physical picture is that the charged and neutral channels of the W–Z exchange add as parallel conductances rather than as serial resistances.

The resulting C_{gap} operator for the mixed vertex is

$$C_{\text{gap}}^{\text{mixed}} = 1 + O_{\text{mixed}}\, C_{\text{local}} \approx 1.010677, \tag{24}$$

and the W-boson mass prediction becomes

$$\boxed{M_W^{\text{EWT}} = M_Z^{\text{CODATA}} \cdot \sqrt{\left(1 - \sin^2\theta_W^{\text{MS-bar}}\right) \cdot C_{\text{gap}}^{\text{mixed}}} \approx 80.379 \text{ GeV}.} \tag{25}$$

Table 3 compares this prediction with current experimental benchmarks. The prediction lies within 2σ of the ATLAS 2024, CMS 2024, and PDG 2024 averages. The earlier pure π^6 value ($M_W \approx 80.514$ GeV) was 10–16σ above these averages; the improvement is due to the mixed-operator correction, not to any adjustment of the input electroweak anchors. The only remaining tension is with the CDF II 2022 measurement (80.4335 GeV), which is 5.8σ *above* the EWT prediction — consistent with the CDF II result now being recognized as an outlier by the ATLAS and CMS 2024 measurements.

Status. The parallel-impedance combination is a *methodologically motivated ansatz*, not a first-principles derivation. The two limiting cases fix its form uniquely among simple parameter-free combinations, but the detailed dynamics of wave interference between the neutral and charged channels — from which the parallel-impedance form should ultimately follow — is not derived in this work.

Table 3: Comparison of the EWT W-boson mass prediction with current experimental benchmarks. The prediction uses the mixed W–Z operator of Eq. (23), with M_Z and $\sin^2\theta_W$ (MS-bar) fixed to their CODATA values. Δ is in MeV relative to the PDG 2024 average. The last column gives the deviation of the EWT prediction from each benchmark in units of the experimental uncertainty.

Source	M_W **[GeV]**	Δ **vs PDG 2024 [MeV]**	σ **(EWT)**
Experimental benchmarks			
CDF II 2022	80.4335 ± 0.0094	$+64.3$	-5.8σ
PDG 2022 average	80.3770 ± 0.0120	$+7.8$	$+0.2\sigma$
ATLAS 2024	80.3600 ± 0.0160	-9.2	$+1.2\sigma$
CMS 2024	80.3602 ± 0.0099	-9.0	$+1.9\sigma$
PDG 2024 average	80.3692 ± 0.0133	0.0	$+0.7\sigma$
Standard Model reference			
SM tree (MS-bar, no rad. corr.)	79.9535	-415.7	—
SM full (radiative-corrected)	80.3600	-9.2	—

The EWT mixed-operator prediction is $M_W = 80.379$ GeV, and the SM indirect electroweak fit predicts $M_W^{\mathrm{SM}} = 80.353 \pm 0.006$ GeV (PDG 2024). The two predictions differ by 26 MeV.

This difference should be interpreted with care. The SM uncertainty of ± 6 MeV reflects only the propagation of input-parameter uncertainties (M_Z, m_t, m_H, α_s, HVP); it does not include the theoretical uncertainty from truncating the perturbative series at two loops, the scheme-conversion uncertainty between different renormalization schemes, or the model-dependence of the hadronic vacuum polarization. Published estimates of the full theoretical uncertainty on the SM indirect fit are in the range ± 10–15 MeV. At this level, the EWT-SM difference of 26 MeV corresponds to approximately 2σ, not 4.3σ.

Both predictions are ansatz-based. The SM calculation uses a truncated perturbative expansion in a chosen renormalization scheme, with the hadronic vacuum polarization imported from external data or lattice QCD. The EWT calculation uses the parallel-impedance combination $\pi^6\pi^7/(\pi^6 + \pi^7)$, which is a methodologically motivated ansatz whose algebraic limits are exact but whose full derivation from BCC lattice dynamics is an open objective.

3.1.5 The π^5 Resonance: Flavor Mixing Deficit

For the quark sector and flavour mixing, the interaction topology shifts from the volume to the **soliton boundary**. The active budget is five-dimensional (3D space + amplitude + frequency), excluding the volumetric radius r, reflecting the surface character of the two-quark mixing vertex. The corresponding geometric deficit at the π^5 rung is

$$\boxed{\epsilon_{\mathrm{flavor}} = \frac{1}{8\pi^5(1-\zeta)} \approx 4.087 \times 10^{-4}} \tag{26}$$

analogous in structure to the magnetic deficit $\epsilon_M = 1/(8\pi^7(1-\zeta))$ at the charged weak rung (Eq. 15). Both follow the pattern $\epsilon_n = 1/(8\pi^n(1-\zeta))$ with $n \in \{5, 7\}$, where 8 is the BCC coordination number and $(1-\zeta)$ is the packing-impedance correction of the discrete lattice.

The flavor correction enters the Cabibbo relation as a single additive lattice deficit,

$$\sin\theta_C = \sqrt{\frac{M_d}{M_s}} \cdot (1 + \epsilon_{\text{flavor}}), \tag{27}$$

consistent with the linear form of the bosonic corrections $C_{\text{gap}} = 1 + \pi^6 C_{\text{local}}$ (pure neutral) and $C_{\text{gap}}^{\text{mixed}} = 1 + O_{\text{mixed}} C_{\text{local}}$ (mixed W–Z). The earlier quadratic operator $C_{\text{fermion}} = (1+\pi^5 C_{\text{local}})^2$ used in previous versions of this work is replaced by this single-deficit form: it systematically increased $\sin\theta_C$ and was only consistent with experiment when paired with non-standard quark masses. The physical picture is that the bare Gatto–Sartori–Tonin amplitude $\sqrt{M_d/M_s}$ captures the bi-local interference of the two quark states at the soliton boundary, while ϵ_{flavor} represents the *single* vacuum deficit of the pair, not a per-quark rescaling.

A detailed derivation of the Cabibbo angle and its comparison with PDG quark masses is given in Section 3.4 (see also Section 3.2.4 for the linear form of the fermionic observable).

3.2 Geometric Mixing Predictions: Higgs and Cabibbo Sectors

3.2.1 Higgs-Vector Boson Mixing

The mixing operators differ by the nature of the vertex. The Z–H mixing is a pure neutral–neutral vertex (both solitons are neutral), so it uses the pure neutral rung $C_{\text{gap}} = 1 + \pi^6 C_{\text{local}}$. The W–H mixing is a mixed charged–neutral vertex (one charged, one neutral soliton), so it uses the parallel-impedance combination $C_{\text{gap}}^{\text{mixed}} = 1 + O_{\text{mixed}} C_{\text{local}}$ defined in Eq. (24):

Table 4: Geometric Mixing Angles for Higgs-Vector Boson Pairs

Mixing Interaction	Calculation Formula	EWT Prediction
$\sin^2\theta_{ZH}$	$1 - \left(\frac{M_Z^{\text{EWT}}}{M_H^{\text{EWT}}}\right)^2 \cdot \frac{1}{C_{\text{gap}}}$	**0.4686**
$\sin^2\theta_{WH}$	$1 - \left(\frac{M_W^{\text{EWT}}}{M_H^{\text{EWT}}}\right)^2 \cdot \frac{1}{C_{\text{gap}}^{\text{mixed}}}$	**0.5906**

$$\boxed{\sin^2\theta_{ZH}^{\text{EWT}} = \mathbf{0.4686}} \tag{28}$$

The Higgs boson interactions are computed as a consistency check of the auxiliary spherical-mode masses, not as a first-principles prediction from the four primary anchors. The calculation uses $M_Z^{\text{EWT}} = 91.731$ GeV (from `mass_spherical(110)`), $M_H^{\text{EWT}} = 124.961$ GeV (from `mass_spherical(117)`), and the mixed W-mass prediction $M_W^{\text{EWT}} = 80.379$ GeV of Eq. (25). The spherical mode is used here rather than the meson mode because the spherical-mode masses reproduce M_Z and M_H at the 0.2–0.6% level, whereas the meson mode gives $M_Z \approx 82.3$ GeV (9.7% off); the spherical-mode calculation inherits the legacy wave constants ρ_a, A_l, λ_l (inventory class V, Section A.2), which is why the mixing angles are classified as auxiliary-mode outputs.

Status. The angles $\sin^2\theta_{ZH}$ and $\sin^2\theta_{WH}$ are not on-shell scalar–vector mixing angles. Within the Standard Model, on-shell mixing between a spin-0 Higgs boson and a spin-1 gauge boson is forbidden by Lorentz invariance and electric charge conservation. This is a symmetry-based statement, not a direct experimental measurement. Experimental searches for such mixing are necessarily indirect and are interpreted within the Standard-Model Extension (SME), an effective

field theory that parametrizes all Lorentz-violating operators consistent with gauge invariance and power-counting renormalizability. Within the minimal SME, loop corrections and other indirect data yield upper bounds on the relevant Higgs-sector coefficients of order 10^{-16} or better, so that any deviation from the SM prohibition would be unobservably small. The angles $\sin^2\theta_{ZH}$ and $\sin^2\theta_{WH}$ are therefore topological projection parameters of the soliton overlap region, not experimentally accessible on-shell mixing angles, as discussed in Section 3.4.

3.2.2 Cabibbo Mixing Angle

The leading-order form $\sin\theta_C \approx \sqrt{M_d/M_s}$ is the classic Gatto–Sartori–Tonin relation [64], well established in chiral perturbation theory. In the EWT framework this relation is reinterpreted geometrically: the square-root form is a signature of surface amplitude interference at the π^5 phase-surface (Section 3.2.4).

The flavor mixing operator is a geometric deficit at the π^5 rung, analogous to the magnetic deficit $\epsilon_M = 1/(8\pi^7(1-\zeta))$ at the charged weak rung:

$$\epsilon_{\text{flavor}} = \frac{1}{8\pi^5(1-\zeta)} \approx 4.087 \times 10^{-4}, \tag{29}$$

where 8 is the BCC coordination number, π^5 is the five-dimensional flavor mixing budget (Table 27), and $\zeta \approx 6.03 \times 10^{-4}$ is the lattice impedance from BCC packing. The Cabibbo angle is then

$$\boxed{\sin\theta_C = \sqrt{\frac{M_d}{M_s}} \cdot (1 + \epsilon_{\text{flavor}})} \tag{30}$$

Using the EWT-derived spherical-mode quark masses at $K_{WC} = 15$ and $K_{WC} = 28$:

$$M_d^{\text{EWT}} = 0.004034\,\text{GeV}, \tag{31}$$

$$M_s^{\text{EWT}} = 0.094885\,\text{GeV}, \tag{32}$$

the EWT prediction is $\sin\theta_C = 0.2063$ (Variant A in Table 5), a relative difference of about 8.03% from the measured value $\sin\theta_C = 0.22430$. The residual reflects the auxiliary spherical-mode quark masses, which differ from PDG by 10–14% (Section 3.2.5).

To isolate the geometric mixing mechanism from the quark-mass determination, the calculation is repeated using standard PDG light-quark masses at $\mu = 2$ GeV. With PDG 2024 values ($M_d = 0.00470$ GeV, $M_s = 0.09350$ GeV), the prediction is $\sin\theta_C = 0.22430$ (Variant B), a residual of 0.002% against the measured value. With PDG 2022 values ($M_d = 0.00467$ GeV, $M_s = 0.09340$ GeV), the prediction is $\sin\theta_C = 0.22370$ (Variant C), a residual of 0.27%. Both residuals are well within the $\sim$ 10–14% uncertainty on the light-quark masses themselves (PDG 2024). The bare Gatto–Sartori–Tonin relation, without the flavor correction, gives $\sin\theta_C = 0.22420$ (PDG 2024) and 0.22361 (PDG 2022), corresponding to residuals of 0.04% and 0.31%. The flavor correction ϵ_{flavor} therefore leaves the leading-order agreement essentially unchanged, consistent with the small size of the geometric deficit ($\epsilon_{\text{flavor}} \approx 4 \times 10^{-4}$ versus the $\sim 10\%$ quark-mass uncertainty). The π^5 operator does not distort the successful GST relation; it adds a lattice correction of the same relative order as the input uncertainties, and its value is not fitted to the Cabibbo angle.

3.2.3 Dimensional Budget of Resonances: Reflection vs. Mixing

The scaling of the modulators is determined by the "dimensional budget" of the resonance involved in the interaction with the BCC lattice:

Table 5: Cabibbo angle prediction: sensitivity to quark-mass input. Variant A uses EWT spherical-mode masses; Variants B and C use standard PDG light-quark masses at $\mu = 2$ GeV. The bare Gatto–Sartori–Tonin relation $\sqrt{M_d/M_s}$, without the flavor deficit, is shown for reference. The geometric deficit is $\epsilon_{\text{flavor}} = 1/(8\pi^5(1-\zeta)) \approx 4.087 \times 10^{-4}$.

Variant	M_d **[GeV]**	M_s **[GeV]**	$\sin\theta_C$	**Error**
A: EWT spherical mode	0.004034	0.094885	0.2063	8.03%
B: PDG 2024 + ϵ_{flavor}	0.00470	0.09350	0.22430	0.002%
C: PDG 2022 + ϵ_{flavor}	0.00467	0.09340	0.22370	0.27%
Bare GST (PDG 2024)	0.00470	0.09350	0.22420	0.04%
Bare GST (PDG 2022)	0.00467	0.09340	0.22361	0.31%
PDG 2024 (reference)	–	–	0.22430	–

- **Bosons (π^6 and π^7):** These are volumetric excitations where the action occurs in 3D space, involving 1D amplitude, 1D frequency, and 1D radius (r). For charged bosons, an additional 1D of freedom (charge) expands the budget to π^7. These processes represent a unified "reflection" of the energy packet off the lattice, resulting in a linear correction (C_{gap} for pure neutral vertices, $C_{\text{gap}}^{\text{mixed}}$ for mixed W–Z vertices).

- **Fermions (π^5):** In the fermionic sector, the interaction is localised on the phase-surface. The budget includes 3D space, 1D amplitude, and 1D frequency, but excludes the volumetric radius (r). This π^5 resonance describes the surface integrity of the soliton.

- **The Pair Deficit (Linear):** The Cabibbo angle represents the mixing of two such π^5 states. The bare Gatto–Sartori–Tonin relation $\sqrt{M_d/M_s}$ captures the *bi-local interference* between the two quarks at the amplitude level: since mass scales as $M \propto A^2$, the square root of the mass ratio directly accesses the wave amplitude A (Section 3.2.4). The lattice response to the pair is a *single* geometric deficit at the π^5 rung,

$$\epsilon_{\text{flavor}} = \frac{1}{8\pi^5(1-\zeta)} \approx 4.087 \times 10^{-4}, \tag{33}$$

analogous to the magnetic deficit $\epsilon_M = 1/(8\pi^7(1-\zeta))$ at the charged weak rung (Eq. 15). Both follow the same structural pattern

$$\epsilon_n = \frac{1}{8\pi^n(1-\zeta)}, \qquad n \in \{5, 7\}, \tag{34}$$

with $n = 7$ for the charged weak sector (ϵ_M) and $n = 5$ for flavor mixing. The flavor correction is therefore *linear*,

$$\sin\theta_C = \sqrt{\frac{M_d}{M_s}}\,(1 + \epsilon_{\text{flavor}})\,, \tag{35}$$

consistent with the linear form of the bosonic corrections C_{gap} and $C_{\text{gap}}^{\text{mixed}}$.

3.2.4 Symmetry of Observables: Energy Density vs. Amplitude Interference

The contrasting mathematical forms of the key observables—$\sin^2\theta_W$ for the bosonic sector and $\sin\theta_C \propto \sqrt{M_d/M_s}$ for the fermionic sector—are not coincidental. They reflect a fundamental

transition in the nature of the interaction within the BCC lattice, rooted in the dimensional hierarchy of the Geometric Ladder.

- **Bosonic Sector (Volumetric Energy Density):** The Weinberg angle, $\sin^2\theta_W$, emerges from the ratio of the W and Z boson masses. In the EWT framework, mass is a measure of volumetric energy density, which at the π^6 and π^7 levels involves the full set of degrees of freedom: 3D space, amplitude A, frequency f, radius r, and (for charged bosons) charge Q. The squared form of the observable directly mirrors the squared relation between mass and the underlying wave amplitude ($E \propto A^2$), making $\sin^2\theta_W$ the natural expression for a volumetric, energy-density-based coupling.

- **Fermionic Sector (Surface Amplitude Interference):** In contrast, the Cabibbo angle, $\sin\theta_C$, describes the mixing of two boundary-localised fermions (d and s quarks). This is a bi-local interference process occurring on the π^5 phase-surface, where the relevant physical quantity is the wave amplitude A itself, not the volume-integrated energy. At the π^5 level, the soliton possesses 3D space, amplitude A, and frequency f—the minimal set required for a massive surface excitation. Its mass scales as $M \propto A^2$, as energy is quadratic in amplitude. To access the amplitude A directly, one must therefore take the square root of the mass. This operation effectively projects the energy-based observable from the π^5 level down to the π^4 level of the Geometric Ladder, where the fundamental amplitude A resides in 3D space as the static structural skeleton of the particle (3D + A). The linear form $\sin\theta_C \propto \sqrt{M_d/M_s}$ is thus the direct signature of amplitude interference at the soliton boundary.

This dichotomy—$\sin^2$ for volumetric energy ratios versus sin for surface amplitude interference—is a direct and testable consequence of the dimensional hierarchy. It confirms that the EWT framework does not merely fit parameters, but derives the very structure of physical laws from the topology of the vacuum substrate.

The geometric flavor deficit $\epsilon_{\text{flavor}} = 1/(8\pi^5(1-\zeta))$ of Eq. (33) enters as a small multiplicative correction on top of the bare amplitude relation $\sqrt{M_d/M_s}$; it does not change the linear form of the fermionic observable, just as the bosonic corrections C_{gap} and $C_{\text{gap}}^{\text{mixed}}$ do not change the squared form of the bosonic observable.

3.2.5 Interpretation of Quark Masses in the EWT Framework

While the Standard Model operates on the abstraction of point-particles within a passive vacuum, the Enhanced EWT model restores the physical necessity of the medium. Matter cannot exist in space without being of space. The $\boldsymbol{\pi^4}$ Quark Stability budget formalises this necessity: the 3D spatial substrate ($\boldsymbol{\pi^3}$) is not merely a background, but the indispensable structural foundation for any dynamic amplitude ($\boldsymbol{\pi^1}$) and/or frequency ($\boldsymbol{\pi^1}$) to manifest as a stable particle.

Remark on Quark Masses and the Nature of Confinement:
In the EWT framework, quarks are not fundamental particles but topological excitations confined within hadronic solitons. Their "masses" are not intrinsic properties of free entities, but effective measures of the binding energy within the collective π^5 phase-surface. The permanent isolation of a quark is geometrically impossible: it would require severing the π^4 amplitude-space core from the π^5 surface, leaving a residual frequency without a spatial substrate.

3.3 Structural Transition to N-Stiffness: From Geometric Volume to Lattice Impedance

To unify the model, the correction operators C_{gap} and ϵ_{flavor} are expressed directly through the geometric vacuum stiffness N_{geom} introduced in Section 2. Using the identity $C_{\text{local}} = \epsilon_M/(2\sqrt{2})$ and $\epsilon_M = 1/(N_{\text{geom}}\pi^3)$, together with $N_{\text{geom}} = 8\pi^4(1-\zeta)$, the geometric structure reveals the scaling hierarchy:

1. Direct N-Representation

$$C_{\text{gap}} = 1 + \pi^6 \cdot C_{\text{local}} = 1 + \frac{\pi^3}{2\sqrt{2} \cdot N_{\text{geom}}} \tag{36}$$

$$\epsilon_{\text{flavor}} = \frac{1}{8\pi^5(1-\zeta)} = \frac{1}{\pi \cdot N_{\text{geom}}} \tag{37}$$

2. Physical Interpretation: The Pi-Power Scaling

- **C_{gap} (Volumetric Coupling):** The π^3/N_{geom} term identifies the magnetic gap as a 3D volumetric resonance. The π^3 factor represents the interaction of the spherical soliton with the BCC lattice stiffness.
- **ϵ_{flavor} (Surface Scaling):** The reduction to $1/(\pi N_{\text{geom}})$ identifies the flavor correction as a surface effect at the π^5 rung. Unlike the bosonic correction, which appears multiplicatively inside C_{gap}, the flavor deficit enters additively as $1 + \epsilon_{\text{flavor}}$. The additive form reflects the bi-local nature of the two-quark mixing at the soliton boundary: the interference is a small perturbation of the bare GST amplitude $\sqrt{M_d/M_s}$, not a rescaling of each quark mass.
- **The N_{geom}-Anchor:** Both corrections are locked to the same geometric stiffness parameter, proving that the difference between the magnetic (π^3 suppression, $n = 7$) and flavor (π^1 suppression, $n = 5$) sectors is purely a matter of the dimensionality of the active rung.

> The transition to the N_{geom}-anchor reveals a fundamental split in the vacuum's response: the bosonic sector is governed by 3D volumetric resonance, while the fermionic sector is dictated by 2D surface-coupling dynamics.

3.4 Observable-Based Experimental Tests of the EWT Higgs Sector

The EWT Higgs sector should be confronted with experimentally defined observables rather than with an unspecified "production angle." The process considered in this section is Higgsstrahlung,

$$e^+e^- \to ZH, \tag{38}$$

where the notation ZH is used throughout. The notation WZ refers to a different final state and is not used here as a synonym for Higgsstrahlung.

The parameter θ_{ZH} introduced earlier is a topological projection parameter of the soliton overlap region, not an on-shell scalar–vector mixing angle. Its physical consequences therefore do not include a uniform rescaling of the on-shell HZZ coupling; this is why the tests below are framed in terms of observables rather than in terms of a production angle. The absence of a tree-level modification is itself the EWT prediction under the restricted hypothesis stated in Pillar I.

The proposed tests are divided into two logically distinct pillars. Pillar I tests the absolute normalization of the ZH production rate under a restricted tree-level EWT hypothesis. Pillar II tests differential observables that can be sensitive to momentum-dependent or tensorial deformations of the production amplitude. The distinction is essential: agreement with the Standard Model in Pillar I does not by itself validate EWT, while a shape test requires an explicit EWT amplitude beyond a purely energy-dependent overall normalization.

3.4.1 Observable definition and recoil-mass reconstruction

A particularly clean observable is the inclusive Higgsstrahlung cross section measured with the recoil-mass technique. For a reconstructed dileptonically decaying Z boson,

$$Z \to \ell^+\ell^-, \qquad \ell = e, \mu, \tag{39}$$

the recoil four-momentum is

$$p_{\text{recoil}} = p_{e^+} + p_{e^-} - p_{\ell\ell}. \tag{40}$$

In the centre-of-mass frame of the collision, the recoil mass is therefore

$$m^2_{\text{recoil}} = s + m^2_{\ell\ell} - 2\sqrt{s}\, E_{\ell\ell}, \tag{41}$$

where $s = (p_{e^+} + p_{e^-})^2$, $m_{\ell\ell}$ is the reconstructed dilepton invariant mass, and $E_{\ell\ell}$ is the dilepton energy in the same centre-of-mass frame. This observable permits an inclusive measurement of $e^+e^- \to ZH$ without requiring reconstruction of the Higgs decay mode, and is consequently appropriate for an absolute test of the ZH production rate [57, 58, 59].

In a realistic analysis, Eq. (41) is not compared directly with an ideal parton-level distribution. The prediction and measurement must use the same treatment of initial-state radiation, beamstrahlung, beam-energy spread, finite detector resolution, event selection, backgrounds, and acceptance. The theoretical prediction must also specify the perturbative order and the input-parameter scheme. These requirements are part of the observable definition rather than optional post-processing corrections.

3.4.2 Pillar I: absolute ZH cross section

The first test is formulated through the signal-strength-like ratio

$$\mu_{ZH}(s) \equiv \frac{\sigma_{\text{EWT}}(e^+e^- \to ZH; s)}{\sigma_{\text{SM}}(e^+e^- \to ZH; s)}. \tag{42}$$

The restricted EWT hypothesis tested in this pillar is

$$\begin{aligned} g^{\text{EWT}}_{HZZ} &= g^{\text{SM}}_{HZZ}, \\ g^{\text{EWT}}_{eeZ} &= g^{\text{SM}}_{eeZ}, \\ \mathcal{M}_{\text{contact}} &= 0, \\ \mathcal{M}_{\text{new tensor}} &= 0, \end{aligned} \tag{43}$$

with the EWT and SM predictions evaluated at the same perturbative order and with identical kinematic cuts. At tree level, Eq. (43) implies

$$\mathcal{M}^{(0)}_{\text{EWT}} = \mathcal{M}^{(0)}_{\text{SM}}, \qquad \mu^{(0)}_{ZH}(s) = 1. \tag{44}$$

Thus, the statement $\mu_{ZH} = 1$ is a consequence of the explicitly stated restricted hypothesis. It is not an independent numerical prediction of the full EWT framework unless EWT specifies why all additional HZZ structures and contact interactions vanish.

If the amplitude contains a small deformation,

$$\mathcal{M}_{\rm EWT} = \mathcal{M}_{\rm SM} + \delta\mathcal{M}, \tag{45}$$

then, to first order in the deformation,

$$\mu_{ZH} \simeq 1 + 2\,\mathrm{Re}\left(\frac{\delta\mathcal{M}}{\mathcal{M}_{\rm SM}}\right), \tag{46}$$

with the understanding that the experimentally measured quantity is a phase-space integral including the relevant acceptance and radiative effects. This makes the interference contribution explicit and prevents the rate comparison from being interpreted as a comparison of amplitudes alone.

The corresponding experimental statement should be written as

$$\mu_{ZH}^{\rm meas}(s) = \mu_{ZH}^{\rm pred}(s) \pm \delta_{\rm exp} \pm \delta_{\rm th}, \tag{47}$$

where the experimental and theoretical uncertainties must be defined. A statistically significant discrepancy with Eq. (44), after the specified corrections and nuisance parameters have been included, would falsify the restricted hypothesis of Eq. (43). It would not, by itself, falsify every possible realization of EWT.

The reference normalization $\kappa_Z = 1$ is the Standard Model value. Any statement that present data are consistent with this value must be attached to a specified experimental combination or global fit; it should not be presented as an uncited universal numerical result. Future lepton-collider studies use precisely this type of inclusive ZH measurement to constrain the absolute Higgsstrahlung rate and Higgs couplings [58, 59, 60].

3.4.3 Pillar II: angular observables and amplitude structure

The second pillar concerns the shape of the production-angle distribution. Let θ_Z denote the angle between the incoming electron direction and the outgoing Z momentum in the e^+e^- centre-of-mass frame,

$$\cos\theta_Z \equiv \frac{\mathbf{p}_{e^-} \cdot \mathbf{p}_Z}{|\mathbf{p}_{e^-}|\,|\mathbf{p}_Z|}. \tag{48}$$

This production angle must be distinguished from decay or helicity angles defined in the rest frame of the Z boson. If a decay angle is used in a future analysis, it must be introduced as a separate observable with its own frame and axis convention.

The normalized production-angle distribution is

$$f(\cos\theta_Z) \equiv \frac{1}{\sigma_{ZH}}\frac{d\sigma_{ZH}}{d\cos\theta_Z}, \qquad \int_{-1}^{1} f(x)\,dx = 1. \tag{49}$$

A convention-independent set of Legendre moments is

$$\langle P_n \rangle \equiv \int_{-1}^{1} dx\, P_n(x) f(x), \qquad x = \cos\theta_Z, \tag{50}$$

where $P_n(x)$ denotes the Legendre polynomial of order n. With this definition, $\langle P_0 \rangle = 1$, and the distribution can be reconstructed formally as

$$f(x) = \frac{1}{2} \sum_{n=0}^{\infty} (2n+1) \langle P_n \rangle P_n(x). \tag{51}$$

In an experimental analysis, the reported moments must specify whether they are detector-level, efficiency-corrected, unfolded, or obtained from a likelihood fit. The covariance matrix of the moments should be retained because the moments are generally correlated.

Angular observables can probe information that is not contained in the total rate, including interference between different helicity amplitudes and momentum-dependent tensor structures. Such observables have been studied as probes of anomalous Higgs interactions and EFT operators at future e^+e^- colliders [61, 62, 63].

3.4.4 Why an overall form factor is insufficient to change the normalized shape

Suppose that the EWT modification is represented only by an energy-dependent multiplicative form factor,

$$\mathcal{M}_{\rm EWT}(s,t,u) = F(s)\, \mathcal{M}_{\rm SM}(s,t,u). \tag{52}$$

At fixed centre-of-mass energy, this gives

$$\frac{d\sigma_{\rm EWT}}{d\cos\theta_Z} = |F(s)|^2 \frac{d\sigma_{\rm SM}}{d\cos\theta_Z}, \qquad \sigma_{\rm EWT} = |F(s)|^2 \sigma_{\rm SM}. \tag{53}$$

Consequently,

$$\frac{1}{\sigma_{\rm EWT}} \frac{d\sigma_{\rm EWT}}{d\cos\theta_Z} = \frac{1}{\sigma_{\rm SM}} \frac{d\sigma_{\rm SM}}{d\cos\theta_Z}, \tag{54}$$

and all normalized angular moments are unchanged. Therefore, a form factor depending only on s is a rate deformation, not a shape deformation.

A genuine angular-shape prediction requires an amplitude of the more general form

$$\mathcal{M}_{\rm EWT} = \mathcal{M}_{\rm SM} + \sum_i C_i(s,t,u)\, \mathcal{M}_i, \tag{55}$$

where at least one coefficient or tensor structure depends on additional kinematic variables or changes the helicity content. Examples include momentum-dependent HZZ form factors, derivative interactions, contact terms, modified longitudinal/transverse amplitudes, beam-polarization dependence, or nontrivial t- and u-dependence. Only after such a structure has been specified can EWT make a quantitative prediction for a shift

$$\Delta\langle P_n \rangle \equiv \langle P_n \rangle_{\rm EWT} - \langle P_n \rangle_{\rm SM}\,. \tag{56}$$

3.4.5 Falsification criteria and scope of the test

The experimental logic should distinguish three outcomes:

1. A statistically significant deviation of the absolute rate from the prediction in Eq. (44), after all specified theoretical and experimental corrections, would falsify the restricted EWT hypothesis tested in Pillar I.

2. A statistically significant deviation in one or more angular moments, relative to the SM prediction and the explicitly specified EWT amplitude, would falsify the corresponding shape hypothesis in Pillar II.

3. Agreement with the SM prediction would constrain the tested EWT parameter space but would not prove EWT, because a model can reproduce the SM limit in the observables considered.

Accordingly, Pillar I is a falsifiable null-deformation test, whereas Pillar II is a quantitative falsification test only after EWT supplies explicit coefficients, tensor structures, parameter values, and a perturbative prescription. Before those ingredients are provided, Pillar II should be described as a well-defined search strategy rather than as a completed numerical prediction.

The most transparent implementation is therefore a joint likelihood in the absolute rate and differential observables,

$$\mathcal{L} = \mathcal{L}(\mu_{ZH}, \langle P_1 \rangle, \langle P_2 \rangle, \ldots; \boldsymbol{\eta}), \tag{57}$$

where $\boldsymbol{\eta}$ contains experimental nuisance parameters and any explicitly defined EWT or EFT coefficients. This formulation keeps the observable definition, the model assumptions, and the falsification criterion logically separate.

Summary. Pillar I is a falsifiable zero-modification prediction under a clearly stated hypothesis. Pillar II is a well-defined search framework whose quantitative prediction awaits the derivation of the soliton form factor. This separation of status is deliberate: it distinguishes what EWT currently predicts from what it currently proposes to test.

4 Analysis of Neutrino Boundary Conditions

Before proceeding to the detailed analysis of the neutrino's boundary conditions, it is crucial to place the Geometric Gravity mechanism of EWT—derived from the Gravitational Deficit ϵ_G—within the broader context of modern theoretical physics. The Energy Wave Theory's postulate that gravity arises from a deficit in the **Elastic Medium Constituent** (EMC) components conceptually aligns with the **Emergent Gravity** paradigm. This framework suggests that gravity is not a fundamental force, but rather an effective phenomenon arising from the microstructure or thermodynamics of spacetime [19]. Key proponents, such as Padmanabhan and Verlinde, develop theories where gravity emerges from spacetime thermodynamics or the information encoded on a holographic screen [20].While such models face significant theoretical and empirical challenges [21], the EWT approach offers a concrete geometric mechanism linking a specific wave-centre structure to a macroscopic gravitational effect, without invoking entropic or thermodynamic assumptions. This links a specific wave-centre structure to a macroscopic gravitational effect, providing a unique, finite, and quantifiable emergent model.

The stability of the Neutrino as a Soliton is dependent on the precise balancing of internal geometric forces. These forces are represented by the **Constant Geometric Deficit (ϵ_G)**, the **global Magnetic Deficit (ϵ_M)**, and the **Geometric Soliton Core**.

4.1 Standard Conditions: Mass and Minimal Magnetism

Under standard conditions, the Neutrino is confirmed to possess non-zero mass (empirically validated by oscillation experiments [22, 23]) and a negligible, but potentially non-zero, magnetic moment [24].

- **Stability (Hooke's Force):** Non-zero mass is understood to imply that the Neutrino possesses an internal **Constant Geometric Deficit** ϵ_G. The internal geometric pressure, caused by ϵ_G and other factors, is **balanced** by the repulsion force resulting from **Hooke's**

Law (Elastic Interaction) between the constituents, which keeps the Soliton in a stable state.

- **Charge:** The Soliton's charge is **considered effectively zero** (≈ 0).
- **Magnetism and ϵ_M:** The global magnetic deficit ϵ_M is fixed by the BCC lattice geometry and does not vanish for the neutrino. However, because the neutrino carries no charge and no spin, it does not generate the local magnetic torque that compresses charged leptons. As a result, the neutrino exhibits only a negligible local magnetic response, even though a tiny moment $\mu_\nu \neq 0$ is not excluded.

4.2 Extreme Conditions: Wave Center in a Geometric Black Hole (GBH)

In the case of a Geometric Black Hole (GBH) [17], extreme geometric compression and the dominance of the accumulated gravitational field ($\sum \epsilon_G$) impose severe conditions.

- **Dominance of ϵ_G and Stability (Hooke's):** At the Critical Distance (d_{crit}) of the GBH, geometric forces are so extreme that **all geometric terms other than gravity are suppressed**. However, the **Elastic Interaction ($F_{\mathbf{Hooke}}$)** is an invariant property and **is still active**, guaranteeing that the Soliton's radius does not fall below r_ν (the limit of minimal Soliton stability), and the Neutrino remains a stable WC.
- **Empirical Validation (Supernovae):** The observation that **99% of supernova energy is released as neutrinos** constitutes empirical confirmation that the collapse of matter under extreme gravity leads to the conversion of the dominant mass into the **minimal geometric state** of the Neutrino (r_ν) at the point d_{crit}. This fact justifies the adoption of the Neutrino as the fundamental WC for the GBH model.
- **Local Magnetic Suppression:** Under conditions of extreme compression, the soliton's local spin and magnetic degrees of freedom are geometrically frozen. This does not mean that the global lattice parameter ϵ_M changes; rather, the local contribution of the spin sector to the soliton's energy balance is suppressed.
- **Pure WC (Boundary Condition):** At the critical point (d_{crit}), the geometric boundary conditions of the GBH force the complete elimination of charge and the local magnetic response. The Neutrino is converted into a **pure Wave Center (WC)** with the single dominant characteristic (ϵ_G):

$$\text{Charge} = 0 \quad \text{and the local magnetic contribution is suppressed.} \tag{58}$$

Hypothesis of Spin Recovery Beyond the GBH Horizon

Within the EWT framework, the geometric structure of the WC (and thus its local spin/magnetic properties) may be **partially recovered or enhanced** outside the GBH's critical boundary d_{crit}. This phenomenon is driven by the reduced compression and resulting non-linear elasticity of the spacetime medium away from the core, aligning conceptually with information preservation hypotheses near the horizon.

5 Neutrino as the Geometric Anchor

The statutory neutrino radius r_ν sets the fundamental length scale of the BCC lattice. In this section, r_ν is not treated as an independent input. Instead, it emerges from the geometric fine-structure constant α_{geom}, the elementary charge amplitude e, and the topology of the BCC lattice.

5.1 Planck Charge from Geometric Alpha

The Planck charge in EWT is interpreted as the fundamental geometric amplitude. It is not introduced as a separate empirical constant; it follows from the relation between the elementary charge amplitude e and the geometric fine-structure constant:

$$q_P = \frac{e}{\sqrt{\alpha_{\text{geom}}}}, \tag{59}$$

where $e = 1.602176634 \times 10^{-19}$ m is the CODATA elementary charge amplitude, and α_{geom} is the purely geometric fine-structure constant derived in Section 2.

The resulting ratio

$$\frac{q_P}{e} = \frac{1}{\sqrt{\alpha_{\text{geom}}}} \approx 11.70624886 \tag{60}$$

is therefore a geometric prediction, not a fitted input.

5.2 Decomposition of the Scaling Factor $S = r_\nu/q_P$

The dimensionless scaling factor $S = r_\nu/q_P$ is obtained by combining three physically distinct mechanisms that act in the undisturbed BCC vacuum lattice.

1. **Static lattice projection.** The soliton potential, described by the inverse geometric fine-structure constant $\alpha_{\text{inv, geom}}$, is distributed over the eight BCC coordination nodes and the spherical wave front:
$$S_{\text{proj}} = \frac{\alpha_{\text{inv, geom}}}{8 + \pi}. \tag{61}$$

2. **Dynamic wave expansion.** The natural exponential decay of the standing-wave amplitude from the centre outward introduces Euler's number e[1] as an integrated factor:
$$S_{\text{exp}} = e. \tag{62}$$

3. **Discrete lattice impedance.** The discrete BCC lattice and wave propagation along the face diagonals add a small correction:
$$\delta_{\text{imp}} = (1 - g_v)(\sqrt{2} - 1). \tag{63}$$
Here g_v is the geometric correction factor associated with the neutrino ground state. The factor $\sqrt{2}-1$ measures the excess diagonal path length, while $1-g_v$ quantifies the deviation of the relaxed neutrino state from an ideal continuum response.

[1] Throughout this work, e denotes Euler's number ($e = 2.71828\ldots$) when it appears in geometric expressions such as $S_{\text{exp}} = e$, $r_\nu = 2q_P e^2/g_v$, or the $2e$ prefactor in Eq. (87). The elementary charge is written e_{geom} or e_{charge} wherever ambiguity could arise, and appears as such in $q_P = e/\sqrt{\alpha_{\text{geom}}}$ (Section 5.1). See Appendix A.2 for the full classification.

The total scaling factor is the sum of these components:

$$S_{\text{tot}} = S_{\text{proj}} + S_{\text{exp}} + \delta_{\text{imp}}. \tag{64}$$

The static projection of the soliton potential over the 8 BCC coordination nodes (8 discrete degrees of freedom) together with the single continuous degree of freedom of the spherically symmetric wavefront π, combined additively because the two contributions enter the projection energy additively rather than multiplicatively.

5.3 Geometric Fixed Point of g_v

The dynamic relation for the neutrino radius is

$$r_\nu = \frac{2q_P e^2}{g_v}. \tag{65}$$

In terms of the scaling factor, this becomes

$$S_{\text{tot}} = \frac{2e^2}{g_v}. \tag{66}$$

Equating this with Eq. (64) gives

$$\frac{2e^2}{g_v} = \frac{\alpha_{\text{inv, geom}}}{8+\pi} + e + (1-g_v)(\sqrt{2}-1). \tag{67}$$

Rearranging leads to the quadratic equation

$$(\sqrt{2}-1)g_v^2 - \left(\frac{\alpha_{\text{inv, geom}}}{8+\pi} + e + \sqrt{2} - 1\right) g_v + 2e^2 = 0. \tag{68}$$

This equation has two roots:

$$g_v \approx 0.983594 \qquad \text{and} \qquad g_v \approx 36.27. \tag{69}$$

Only the first root satisfies $0 < g_v < 1$, which is required for stable wave propagation and a physical soliton. Thus

$$g_v \approx 0.983594 \tag{70}$$

is the unique geometric fixed point of the BCC lattice. It is not an adjustable parameter, but a direct consequence of the lattice topology.

5.4 Resulting Neutrino Radius

Using the fixed-point value of g_v and the geometric Planck charge q_P, the neutrino radius is

$$r_\nu = q_P S_{\text{tot}} \approx 2.817935 \times 10^{-17} \text{ m}. \tag{71}$$

This value is in close agreement with the earlier dynamical expression

$$r_\nu = \frac{2q_P e^2}{g_v}, \tag{72}$$

and the two approaches differ only at the 10^{-6} level. This confirms that r_ν is derived from the same BCC geometry as α_{geom} and ϵ_M.

5.5 Decadic Resonance Test: $r_e/r_\nu \approx 100$

An immediate test of the geometric anchor is provided by the ratio of the classical electron radius to the neutrino radius:

$$\frac{r_e}{r_\nu} \approx 100.00017. \tag{73}$$

The corresponding fifth power

$$\left(\frac{r_e}{r_\nu}\right)^5 \approx 1.0000087 \times 10^{10} \tag{74}$$

reproduces the 10^{10} scaling expected from the difference between the electron wave-centre count $K_{WC} = 10$ and the neutrino count $K_{WC} = 1$. This is a direct consequence of the geometric mass-to-radius identity, Eq. (197), discussed in Section 15.1. It provides a non-trivial consistency test linking the geometric radius to the r^5 energy scaling used throughout the model.

5.6 Physical Interpretation: ϵ_M as a Global Lattice Parameter

The magnetic deficit ϵ_M is a global topological property of the BCC lattice and does not vanish for any soliton. What differs between the neutrino and charged leptons is the local magnetic response.

The neutrino carries no charge and no spin; it therefore generates no local magnetic torque. Consequently, its soliton relaxes to the natural radius r_ν given by Eq. (71).

For the electron, the local magnetic response compresses the soliton below this relaxed state. The factor g_v measures exactly this difference: it is the ratio between the relaxed neutrino radius and the compressed wavelength that would follow from the simple wave constants. The global ϵ_M remains unchanged; it is the local magnetic compression that differs between particles.

Thus the neutrino is not a "small electron", but the fundamental, torque-free ground state of the BCC lattice. The electron is a compressed and locally magnetised excitation of this same ground state.

5.7 Topological Necessity of r_ν

The decomposition of S_{tot} into

$$\frac{\alpha_{\text{inv, geom}}}{8+\pi}, \quad e, \quad (1-g_v)(\sqrt{2}-1) \tag{75}$$

shows that the neutrino radius is fixed by three simultaneous geometric constraints:

- the static soliton potential distributed over the eight BCC directions and the spherical wave front,
- the natural exponential decay encoded by Euler's number e,
- the discrete lattice correction along the face diagonals.

No free parameter is introduced. All quantities are determined by π, e, $\sqrt{2}$, the BCC coordination number, and the geometric fine-structure constant.

This completes the emergence of the neutrino as a topological anchor of the BCC lattice. No fitted parameter is introduced in this derivation.

6 Planck Length and Reduced Planck Constant from Geometry

The geometric fine-structure constant α_{geom}, the electron radius r_e, and the electron mass m_e determine the reduced Planck constant without invoking $\hbar$ as an independent input. The same geometric machinery then yields the Planck length λ_l as a derived quantity.

6.1 Reduced Planck Constant from Geometric Alpha

In EWT, the classical electron radius is not defined through $\hbar$ but is treated as a geometric scale fixed by the electron soliton. The reduced Planck constant follows from the relation

$$\hbar_{\text{geom}} = \frac{m_e c r_e}{\alpha_{\text{geom}}}, \tag{76}$$

where c is the metric conversion factor defined in Section 9.

Using the CODATA values

$$m_e = 9.1093837015 \times 10^{-31}\ \text{kg}, \qquad r_e = 2.8179403262 \times 10^{-15}\ \text{m}, \tag{77}$$

and the geometric fine-structure constant α_{geom}, Eq. (76) gives

$$\hbar_{\text{geom}} \approx 1.0545738 \times 10^{-34}\ \text{kg}\,\text{m}^2\,\text{s}^{-1}, \tag{78}$$

in agreement with the CODATA value at the level of the underlying experimental inputs. No new free parameter is introduced.

7 The Physical Origin of Charge as Wave Amplitude

The identification of charge with wave amplitude is a foundational postulate of Energy Wave Theory, established in the primary EWT literature [29]. The present section makes the dimensional reduction explicit, in the same way that the kilogram was reduced to a geometric unit in Section 15.3.

7.1 Charge as Spatial Wave Amplitude

In EWT, the elementary charge e of a single electron represents the physical wave displacement amplitude of the lattice constituents at the first wavelength. Because displacement in a spatial lattice is fundamentally a geometric length, charge carries the dimension of length ($[Q] = [L]$).

In the International System of Units (SI), where the fundamental unit of spatial length is the metre, the elementary charge amplitude is given by:

$$e_{\text{geom}} = 1.602176634 \times 10^{-19}\ \text{m}. \tag{79}$$

This is a direct dimensional mapping: the numerical value of the CODATA elementary charge is preserved, while the dimension changes from coulombs (C) to metres (m).

7.2 Conversion of the Coulomb and Distinction from CGS

From the definition 1 C = 1 A s and the geometric representation of charge as amplitude (m), the coulomb is recognized as a derived unit of spatial displacement:

$$e_{\mathrm{SI}} = 1.602176634 \times 10^{-19}\ \mathrm{C} \quad \Longleftrightarrow \quad e_{\mathrm{geom}} = 1.602176634 \times 10^{-19}\ \mathrm{m}. \tag{80}$$

Resolution of the CGS / Statcoulomb Ambiguity:
It is crucial to emphasize that e_{geom} is a spatial displacement ($[L]$), not a force-derived quantity. In Gaussian CGS units, the statcoulomb is defined via Coulomb's Law as $\mathrm{g}^{1/2}\,\mathrm{cm}^{3/2}\,\mathrm{s}^{-1}$, which assigns charge a composite mechanical dimension rather than a pure spatial length.
Transforming e_{geom} to CGS spatial units simply corresponds to expressing the displacement in centimetres ($1.602176634 \times 10^{-17}$ cm), rather than substituting the numerical value of the statcoulomb (4.8032×10^{-10}). The numerical equality $1\ \mathrm{C} \to 1\ \mathrm{m}$ is anchored specifically to the SI convention, where spatial displacement is measured in metres.

7.3 Consequence for the Planck Charge q_P

The Planck charge in EWT represents the maximum fundamental geometric amplitude of the BCC lattice:

$$q_P = \frac{e_{\mathrm{geom}}}{\sqrt{\alpha_{\mathrm{geom}}}}. \tag{81}$$

Evaluating this relation with $e_{\mathrm{geom}} = 1.602176634 \times 10^{-19}$ m yields:

$$q_P = 1.8755478 \times 10^{-18}\ \mathrm{m}. \tag{82}$$

The resulting quantity q_P is therefore an absolute spatial length of the lattice. The zero-calibration derivation of the statutory neutrino radius $r_\nu = q_P \cdot S_{\mathrm{tot}}$ is a purely geometric relation mapping this amplitude scale to the lattice step.

8 Geometric Deficit and Density Hierarchy of the Neutrino

The neutrino soliton serves as the reference wave centre for the BCC vacuum. Its geometric radius r_ν, derived in Section 5, sets the fundamental length scale from which the EMC density hierarchy follows.

At this stage the EMC spacing λ_l is treated as a formal geometric parameter. Its numerical value is fixed later by the self-consistent gravitational identity of Section 11.6, where the Planck-length condition and the geometric expression for G form a closed algebraic system. The density hierarchy below therefore uses λ_l symbolically before that value is substituted.

This geometric saturation limit and the associated density hierarchy follow from the EWT granular density model developed by Yee [11].

1. Absolute Maximum Capacity ($N_{\nu,\mathbf{max}}$)

$$N_{\nu,\max} = \left(\frac{r_\nu}{\lambda_l}\right)^3 \approx 5.3004 \times 10^{54} \tag{83}$$

2. Statutory Background Density ($N_{\nu,\mathbf{stat}}$)

$$N_{\nu,\mathrm{stat}} = \left(\frac{r_\nu}{2\lambda_l e}\right)^3 \approx 3.2986 \times 10^{52} \tag{84}$$

3. Effective Gravitational Density ($N_{\nu,\mathbf{eff}}$)

The effective density inside the electron soliton is obtained from the statutory density and the geometric dilution factor X_{eff}:

$$N_{\nu,\mathrm{eff}} = \frac{N_{\nu,\mathrm{stat}}}{X_{\mathrm{eff}}}, \tag{85}$$

where X_{eff} is the geometric dilution factor defined in Section 11.6.

8.1 Unified Geometric Coupling

The transition from the statutory background to the effective density is governed by the unified coupling operator C_{unif} and the geometric dilution factor X_{eff}, already introduced in Section 6. Both quantities are fixed by the BCC lattice geometry and the electron wave-centre count $K_{WC} = 10$.

The resulting effective gravitational density is

$$N_{\nu,\mathrm{eff}} = \frac{N_{\nu,\mathrm{stat}}}{X_{\mathrm{eff}}} \approx 6.247 \times 10^{48}. \tag{86}$$

This value follows from the geometric dilution of the statutory vacuum by the soliton wave structure. The concept of such a density deficit and its role in gravitational emergence originates from the EWT volume-deficit models of Yee [11, 1]. In the present formulation the deficit is not fitted; it is a direct consequence of the same BCC geometry that determines α_{geom}, r_ν, and λ_l.

The extreme dilution of approximately 99.98% relative to $N_{\nu,\mathrm{stat}}$ explains the weakness of gravity as a pressure deficit within the high-density medium.

9 Speed of Light as the Metric of Spacetime

Before deriving the gravitational constant, the role of the speed of light c must be addressed. In the conventional formulation, c is treated as a fundamental constant, empirically determined and independent of any underlying structure. Within the EWT framework, this status is revised: c is not an independent parameter but the intrinsic metric conversion factor between the spatial and temporal dimensions of the BCC lattice, determined entirely by the discrete geometry of the elastic medium.

9.1 The Causal Structure of the BCC Lattice

The BCC spacetime lattice is characterised by three distinct density levels that define its structural states:

- **Statutory density** $N_{\nu,\mathrm{stat}}$: the equilibrium density of the undisturbed vacuum, derived in Eq. (84).
- **Effective density** $N_{\nu,\mathrm{eff}}$: the reduced density inside a soliton (matter-occupied region), derived in Eq. (85).

- **Maximum density** $N_{\nu,\max}$: the absolute saturation limit of the lattice, given by Eq. (83).

The lattice is characterised by two fundamental geometric scales:

- **The spatial step:** the inter-constituent distance λ_l, which follows from the packing geometry of the Elastic Medium Constituents (EMCs) and is identified with the Planck length. Its value is fixed by the statutory neutrino radius r_ν and the Eulerian dilution factor $2e$ through the relation

$$\lambda_l = \frac{r_\nu}{2e\, N_{\nu,\mathrm{stat}}^{1/3}}, \tag{87}$$

  where $N_{\nu,\mathrm{stat}}$ is the statutory background density. Equivalently, it appears in the uncorrected neutrino wavelength

$$\lambda_{\mathrm{uncorr}} = 2q_P e^2, \tag{88}$$

  which, after the geometric correction g_v, yields the statutory radius $r_\nu = \lambda_{\mathrm{uncorr}}/g_v$. The spatial step λ_l is thus a derived property of the BCC lattice, not an independent input.

- **The temporal step:** the fundamental cycle time t_p, which represents the characteristic oscillation period of an EMC in the lattice. In the spring-mass representation of the BCC medium, this period is determined by the effective elastic constant of the lattice and the Planck mass, both of which are themselves consequences of the lattice geometry (Sec. 32.8). The formal derivation of t_p from first principles is a natural direction for future research; for the present purposes, it suffices to note that t_p is a structural invariant of the lattice, defined by the discrete causal structure itself.

The Planck charge q_P is not an independent parameter but the fundamental amplitude scale of the BCC lattice. It is related to the elementary charge e through the fine-structure constant:

$$e^2 = \alpha q_P^2. \tag{89}$$

This relation follows directly from the definitions of α and q_P, and it holds identically in both SI units (where $q_P = \sqrt{4\pi\epsilon_0\hbar c}$) and in the EWT amplitude representation (where both charges are measured in meters). It expresses the fact that the elementary charge is the lattice amplitude reduced by the coupling factor α.

9.2 Axiom: The Maximum Propagation Speed

The following fundamental axiom is posited:

> *In the BCC lattice at the statutory density $N_{\nu,stat}$, there exists a maximum propagation speed c for any disturbance. This speed is a structural property of the lattice, determined by its density and elastic response, and is independent of the amplitude or frequency of the disturbance.*

The BCC lattice is anisotropic at the Planck scale, where the wavelength of a propagating mode is comparable to the lattice spacing λ_l; the dispersion relation is not exactly linear, and wave speeds depend on direction in the crystal frame. The effective Lorentz invariance and isotropic speed of light c are infrared properties: at scales $\lambda \gg \lambda_l$, the dispersion relation becomes linear to leading order, and the elastic tensor averages over many unit cells to an isotropic effective continuum. The Planck-scale BCC crystal is the ultraviolet completion of the model; the observed isotropy of c is a long-wavelength property, not a Planck-scale symmetry.

This axiom is physically motivated: in any elastic medium with finite density and stiffness, there exists a speed of propagation (the speed of sound in the medium). The BCC lattice, as a discrete elastic medium, is no exception. The speed c is therefore not an empirical input but a derived property of the lattice, on the same footing as the speed of sound in a solid.

In any discrete causal manifold, the maximum propagation speed is fixed by the ratio of the spatial step to the temporal step:

$$c \equiv \frac{\lambda_l}{t_p}. \tag{90}$$

Because the speed of light is the fixed metric conversion $c \equiv \lambda_l/t_p$, specifying λ_l therefore fixes the temporal step $t_p = \lambda_l/c$. In the natural units of the lattice ($c = 1$) the spatial and temporal steps are identical.

This relation is not a postulate nor an empirical input; it is the definition of the causal structure of the lattice, expressing the unique conversion factor between the temporal and spatial coordinates in the metric:

$$ds^2 = -c^2dt^2 + dx^2 + dy^2 + dz^2. \tag{91}$$

9.3 Consequences: Density-Dependent Propagation Speed

The axiom immediately implies that the propagation speed in the lattice is a function of the local EMC density:

$$v(N_\nu) = c \cdot f\left(\frac{N_\nu}{N_{\nu,\mathrm{stat}}}\right), \tag{92}$$

where the function f encodes the elastic response of the lattice. The three density regimes yield distinct physical consequences:

- **In vacuum** ($N = N_{\nu,\mathrm{stat}}$), the speed is exactly c, explaining the constancy of the speed of light in free space. The vacuum is defined as the region where the lattice maintains its statutory density.
- **In matter** ($N = N_{\nu,\mathrm{eff}} < N_{\nu,\mathrm{stat}}$), the speed is reduced, providing a fundamental mechanism for refraction and the slowing of light in media. This is the origin of the refractive index $n = c/v > 1$.
- **In the extreme EMC packaging density limit** ($N \to N_{\nu,\mathrm{max}}$), the lattice approaches its elastic saturation limit. The propagation properties of this regime — relevant to early-universe cosmology — lie beyond the scope of the present analysis and are reserved for future work.

The ratio of the spatial step to the temporal step in the lattice gives the maximum propagation speed in the statutory vacuum as defined in eq. 90.

Remark on the status of t_p in natural units. The preceding analysis noted that the formal derivation of the temporal step t_p from the elastic constants of the BCC medium remains an open objective in SI units. Within the natural units of the lattice ($c = 1$), however, this problem does not arise: the defining relation $c \equiv \lambda_l/t_p$ immediately yields

$$t_p = \frac{\lambda_l}{c} = \lambda_l, \tag{93}$$

so the lattice possesses a *single* fundamental scale λ_l, and t_p is not an independent quantity but an alternative labelling of the same geometric step in the temporal direction. The open objective

of deriving the *numerical value* of c in SI units (metres per second) is therefore reframed as a question about the external convention that separates the metre from the second — not a question about the internal structure of the lattice. Within EWT, the lattice is characterised by one geometric primitive: λ_l.

Remark on the Dimensional Status of c:

The speed of light c does not appear in any of the dimensionless predictions of the EWT framework (the fine-structure constant α, the anomalous magnetic moments a_e, a_μ, a_τ, or the lepton mass ratios). It enters only when expressing dimensional SI quantities such as G or m_e, where it plays the role of a *metric conversion factor* between the spatial step λ_l and the temporal step t_p of the BCC lattice ($c \equiv \lambda_l/t_p$).

In natural units ($c = 1$), the EWT framework is parameter-free: all physical predictions reduce to dimensionless ratios determined by BCC lattice topology alone. The appearance of c in SI expressions reflects the unit system's convention for separating spatial and temporal dimensions, not an independent physical input to the theory. Its numerical value in SI units is therefore a consequence of the arbitrary definitions of the metre and the second, not a fundamental property of the vacuum lattice. This status is consistent with the SI 2019 definition, in which c is an exact definitional constant rather than a measured quantity.

With this geometric interpretation of the speed of light, the base gravitational scaling may now be derived.

10 The Physical Origin of Relativity: Wave-Mechanical Lorentz Contraction

As established in Sec. 9, the speed of light c is the structural conversion factor between the spatial and temporal steps of the BCC lattice. The present section demonstrates that this same value is measured by all inertial observers, not as a primary postulate, but as a mechanical consequence of the wave-structure of matter. The derivation follows the Doppler-based approach developed by Gabriel LaFrenière [51] and historically proposed by H.A. Lorentz [52] and G.F. FitzGerald [53] as a physical explanation of the Michelson–Morley null result.

10.1 Doppler-Induced Compression of the Standing-Wave Soliton

In EWT, a particle is a standing wave (soliton) formed by the interference of incoming and outgoing waves in the BCC medium. When such a soliton moves relative to the lattice with velocity $v = \beta c$, the underlying waves undergo an anisotropic Doppler shift.

Let λ_0 be the wavelength of the soliton at rest. For a soliton moving at speed v, the forward-propagating wave is compressed to λ_f and the backward wave is expanded to λ_b:

$$\lambda_f = \lambda_0(1 - \beta), \qquad \lambda_b = \lambda_0(1 + \beta). \tag{94}$$

Two mean wavelengths can be constructed from the Doppler-shifted pair (λ_f, λ_b). The *arithmetic mean* represents the laboratory-frame average of the two counter-propagating waves; the *geometric mean* represents the wavelength after the phase self-consistency condition of the standing wave is imposed (the wave must return to itself after one round trip across the soliton boundary, which in the multiplicative phase space of Fourier amplitudes corresponds to the geometric

mean of the two wavelengths):

$$\lambda_{\rm am} = \frac{\lambda_f + \lambda_b}{2} = \lambda_0, \qquad \lambda_{\rm gm} = \sqrt{\lambda_f \cdot \lambda_b} = \lambda_0\sqrt{1-\beta^2} = \frac{\lambda_0}{\gamma}, \tag{95}$$

with $\gamma = (1-\beta^2)^{-1/2}$ the Lorentz factor. The contraction factor of the soliton is the *ratio* of these two means,

$$g \;\equiv\; \frac{\lambda_{\rm gm}}{\lambda_{\rm am}} = \sqrt{1-\beta^2} = \frac{1}{\gamma}. \tag{96}$$

The physical distance between the wave-center nodes — the "size" of the particle as seen in the laboratory frame — therefore contracts by $g = 1/\gamma$. This is the form of the wave-mechanical Lorentz contraction derived by LaFrenière [51] from the Doppler-shifted standing-wave structure of a moving soliton in his analysis of the compressibility of standing waves in a material medium. The key point is that neither of the two individual Doppler wavelengths follows the Lorentz factor; the contraction emerges only as the ratio of the geometric mean (which encodes the phase self-consistency of the standing wave) to the arithmetic mean (the laboratory-frame reference scale).

10.2 The Michelson–Morley Null Result as a Mechanical Necessity

The invariance of the measured speed of light in the Michelson–Morley experiment is usually interpreted as a property of spacetime. In EWT, it is a property of the **measuring apparatus**. Because all matter (including the interferometer arms) consists of waves in the BCC lattice, the apparatus itself undergoes a physical contraction:

1. **Longitudinal arm ($L_\parallel$):** Contracts by exactly $1/\gamma$ owing to the wave-node compression derived above.

2. **Transverse arm ($L_\perp$):** Remains unchanged in length, but the wave path is elongated by γ because of the diagonal trajectory through the medium.

The total round-trip times for the longitudinal path ($t_\parallel$) and transverse path ($t_\perp$) become identical:

$$t_\parallel = \frac{L_0/\gamma}{c-v} + \frac{L_0/\gamma}{c+v} = \frac{2L_0}{c}\,\gamma, \qquad t_\perp = \frac{2L_0\sqrt{1+(v\gamma/c)^2}}{c} = \frac{2L_0}{c}\,\gamma. \tag{97}$$

Thus $t_\parallel = t_\perp$. The null result is not due to the absence of a medium, but because the wave-mechanical contraction of the soliton perfectly masks the motion through the lattice.

Remark on time dilation: Substituting $c = 1$ (the natural units of the BCC lattice, Sec. 9) into the Michelson–Morley round-trip times yields $t_\parallel = t_\perp = 2L_0\gamma$ for both arms. In the lattice rest frame the round-trip time is therefore $t = 2L_0\gamma$. A clock carried by the moving interferometer, whose rate is dilated by $1/\gamma$, measures

$$t' = \frac{t}{\gamma} = 2L_0,$$

the same elapsed time as an identical apparatus at rest in the BCC lattice. The oscillation period of the soliton therefore dilates to $T' = \gamma T_0$, with the observed tick rate scaling as $f' = f_0/\gamma$ — the temporal counterpart of the spatial contraction derived above.

11 The Geometric Identity of G: Derivation and Consistency

The core hypothesis of the Energy Wave Theory (EWT) is that the gravitational constant $\mathbf{G}$ is not a fundamental constant but an **emergent, calculated geometric identity** derived from the electron's geometric parameters and the BCC lattice topology. This section derives a precise $\hbar$-independent formula for $\mathbf{G}$ and explains the physical meaning of its scaling terms.

11.1 Derivation of G from Electron Scaling (The $\hbar$-Independent Base)

Traditional definitions of $\mathbf{G}$ rely on Planck units, which inherently contain the reduced Planck constant ($\hbar$). EWT posits that the gravitational constant is derived from the scaling of the electron's properties ($\mathbf{m_e}, \mathbf{r_e}$), which are geometric wave-based quantities.

The Base Gravitational Scaling ($\mathbf{G_{Base}}$)

The fundamental scaling constant for gravity is established by the ratio of energy density to mass, using the speed of light (c), the classical electron radius ($\mathbf{r_e}$), and the electron mass ($\mathbf{m_e}$):

$$\mathbf{G}_{\text{Base}} = \frac{c^2 \mathbf{r_e}}{\mathbf{m_e}} \approx 2.780252 \times 10^{32} \text{ m}^3\text{kg}^{-1}\text{s}^{-2} \tag{98}$$

This term is the sole source of the required physical units for the gravitational constant:

$$\left[\frac{c^2 \mathbf{r_e}}{\mathbf{m_e}}\right] = \frac{\left(\text{m} \cdot \text{s}^{-1}\right)^2 \cdot \text{m}}{\text{kg}} = \text{m}^3 \cdot \text{kg}^{-1} \cdot \text{s}^{-2} \tag{99}$$

All subsequent geometric factors in the final identity for $\mathbf{G}$ are therefore dimensionless.

The $\hbar$-Independence of $\mathbf{G_{Base}}$

The form of $\mathbf{G}_{\text{Base}}$ is inherently independent of $\hbar$. This is demonstrated by substituting the standard definition of the classical electron radius ($\mathbf{r_e} = \alpha \frac{\hbar}{\mathbf{m_e} c}$) into the conventional expression for the gravitational constant based on particle properties ($\mathbf{G}_{\text{Base}} = \alpha \frac{\hbar c}{\mathbf{m_e}^2}$):

$$\mathbf{G}_{\text{Base}} = \alpha \frac{\hbar c}{\mathbf{m_e}^2} = \alpha \frac{\left(\frac{\mathbf{r_e} \mathbf{m_e} c}{\alpha}\right) c}{\mathbf{m_e}^2} = \frac{\mathbf{r_e} \mathbf{m_e} c^2}{\mathbf{m_e}^2} = \frac{c^2 \mathbf{r_e}}{\mathbf{m_e}} \tag{100}$$

The cancellation of $\hbar$ confirms that the base gravitational scale is a classical geometric property of the electron soliton ($\mathbf{m_e}, \mathbf{r_e}, c$), rather than a purely quantum effect.

Status of m_e and r_e in the Present Work

In the current formulation, m_e, r_e, and c are the metric and structural inputs used to anchor the electron soliton. The radius r_e is related to the geometric neutrino anchor by

$$r_e = 100\, r_\nu, \tag{101}$$

while m_e is constrained by the $E \propto r^5$ scaling law discussed in Section 15.1. Their full derivation from the BCC lattice alone is a central open objective.

The subsequent derivation of G therefore contains no free dimensional parameters: the only inputs are r_e, m_e, c, and the elementary charge amplitude e. In natural units ($c = 1$), the number of independent inputs reduces further, and the gravitational constant becomes a purely geometric prediction.

The Final Geometric Identity for G

The observed gravitational constant $\mathbf{G}$ is obtained by scaling $\mathbf{G}_{\text{Base}}$ with the total dimensionless geometric weakness factor. In the zero-calibration formulation, this factor is built solely from the geometric soliton core A_π, the effective BCC stiffness N_{geom}, the electron wave-centre count $K_{WC} = 10$, and the effective volume deficit $N_{\nu,\text{eff}}$.

The geometric identity for $\mathbf{G}$ is therefore:

$$\mathbf{G} \equiv \mathbf{G}_{\text{Base}} \cdot \frac{1}{A_\pi} \cdot \left(\frac{1}{N_{\text{geom}} A_\pi} \right)^3 \cdot \frac{1}{K_{WC} \sqrt{N_{\nu,\text{eff}}}} \tag{102}$$

The only dimensional factor in this expression is $\mathbf{G}_{\text{Base}}$. Every other factor is a pure geometric number determined by the BCC lattice and the electron soliton.

Geometric Dependence on λ_l

The effective volume deficit $N_{\nu,\text{eff}}$ depends on the EMC spacing λ_l through the statutory density:

$$N_{\nu,\text{eff}} = \frac{N_{\nu,\text{stat}}}{X_{\text{eff}}}, \qquad N_{\nu,\text{stat}} = \left(\frac{r_\nu}{2e\lambda_l} \right)^3. \tag{103}$$

Thus Eq. (102) defines G as a function of the formal parameter λ_l. This is not a fault of the model; it is the expected self-consistent coupling between gravity and the fundamental lattice spacing. The value of λ_l will be fixed in Section 11.6 by combining the geometric G with the Planck-length definition.

Compact ϵ_M Form

Using the geometric magnetic deficit

$$\epsilon_M = \frac{1}{N_{\text{geom}} \pi^3}, \tag{104}$$

Eq. (102) can also be written as

$$\mathbf{G} = \frac{\mathbf{G}_{\text{Base}}}{A_\pi} \left(\frac{\epsilon_M \pi^3}{A_\pi} \right)^3 \frac{1}{K_{WC} \sqrt{N_{\nu,\text{eff}}}}. \tag{105}$$

This form shows that gravity and the magnetic deficit share the same geometric origin. It also makes clear that the weakness of gravity is a consequence of the cubic suppression by ϵ_M / A_π, followed by the surface projection $N_{\nu,\text{eff}}^{-1/2}$.

11.2 The Gravitational Identity Flow: From Electromagnetic Base to Surface Transition

The derivation of the Gravitational Constant $\mathbf{G}$ follows a systematic, multi-step geometric flow. This process resolves the scale disparity between electromagnetic and gravitational interactions by accounting for the displacement of the medium's constituents within the Soliton structure.

Step 1: Establishment of the EM Base ($\mathbf{G}_{\text{Base}}$): The process is initiated by defining $\mathbf{G}_{\text{Base}}$, which links the classical electron radius (r_e) and its mass (m_e). This term represents the maximum potential coupling scale before geometric dilution is applied.

Step 2: Geometric Stability Factor ($\mathbf{A_\pi}^{-1}$): The first scaling step introduces the normalization to the static wavelength-to-radius ratio of the Soliton.

Step 3: Nodal Push-Out Potential ($\mathbf{\Omega_{Push}}$): This phase incorporates the volumetric redistribution of energy. By applying the cubic factor $(N_{\text{geom}} A_\pi)^{-3}$, the model quantifies the structural dilution of the soliton in the BCC lattice.

Step 4: Geometric Surface Transition ($\mathbf{T_{Surface}}$): The conversion of the internal displacement into the observable gravitational force is governed by the factor $(K_{WC} \cdot \sqrt{N_{\nu,\text{eff}}})^{-1}$. Here, $K_{WC} = 10$ Wave Centers act as the primary operators of the Push-Out mechanism, evacuating the medium from its statutory density to the effective density of the matter-occupied region.

The square root $(N_{\nu,\text{eff}}^{-1/2})$ serves as the surface transition operator, projecting the three-dimensional internal packing deficit onto the two-dimensional effective surface of the Soliton. Gravity is thus revealed as an emergent, pressure-driven phenomenon acting on this surface.

Step 5: Final Gravitational Identity: The integration of these modular scales yields the geometric identity for $\mathbf{G}$ given in Eq. (102).

Physical Significance

The extreme weakness of gravity is a direct consequence of this hierarchical dilution. While the Soliton's mass is defined by high-frequency energy localization, the gravitational constant $\mathbf{G}$ is a measure of the effective displacement deficit. The transition from the statutory vacuum to the diluted interior creates a pressure gradient, whereby the universal medium strives for equilibrium by filling the geometric void.

The Holographic Nature and Experimental Implications

The presence of the square root $(\sqrt{N_{\nu,\text{eff}}})$ in the final identity reinforces the principle that gravity is an emergent surface phenomenon. By projecting the three-dimensional volumetric deficit onto a two-dimensional boundary, the model aligns with holographic and thermodynamic interpretations of gravitational flux.

11.3 Formalization of Soliton Geometric Density $\rho(r)$

To avoid ambiguity between energy density ρ_E and the packing density, the internal density function $\rho(r)$ is introduced to represent the local distribution of Elastic Medium Constituents (EMC) within the Soliton. This profile is physically important, but the numerical value of the effective deficit is fixed algebraically by the unified coupling described in Section 8:

$$N_{\nu,\text{eff}} = \frac{N_{\nu,\text{stat}}}{X_{\text{eff}}}, \tag{106}$$

where X_{eff} is the geometric dilution factor from Section 11.6 and Eq. (116). The radial profile $\rho(r)$ is compatible with this value, and the structural parameters of any assumed profile do not enter as free inputs into the gravitational identity.

11.4 Numerical Verification and the Unitary Mapping Operator $\mathcal{U}$

Numerical Consistency

Using the geometric value N_{geom} and the self-consistent λ_l from Eq. (121), the geometric identity Eq. (102) gives

$$G_{\text{EWT}} \approx 6.67752 \times 10^{-11}\ \text{m}^3\,\text{kg}^{-1}\,\text{s}^{-2}. \tag{107}$$

The corresponding CODATA 2022 value is

$$G_{\text{CODATA}} = 6.674305 \times 10^{-11}\ \text{m}^3\,\text{kg}^{-1}\,\text{s}^{-2}, \tag{108}$$

so the relative deviation is approximately 0.048%.

This is a genuine geometric prediction.

The Unitary Mapping Operator $\mathcal{U}_{\mathbf{Geom}\to G}$

The final step in the geometric formalization is the introduction of the **Unitary Mapping Operator** $\mathcal{U}$, which symbolically represents the non-linear transformation of the Soliton's internal geometry into the macroscopic gravitational constant:

$$\mathcal{U}_{\text{Geom}\to G}:\ A_\pi,\ N_{\text{geom}},\ N_{\nu,\text{eff}} \longmapsto G \equiv \frac{c^2 r_e}{m_e}\cdot\Omega_G. \tag{109}$$

This operator formalizes that G is not a standalone fundamental constant but a direct manifestation of the geometric scaling applied to the electron's $\hbar$-independent base.

11.5 Formal Definition of the Operator $\mathcal{U}$: Mapping Geometric Parameters to G

To codify the micro–macro transformation in a rigorous mathematical manner, a nonlinear functional operator $\mathcal{U}$ is defined:

$$\mathcal{U}:\ \mathcal{D}\subset\mathbb{R}^n \to \mathbb{R}, \qquad \mathcal{U}(\mathbf{P}) \mapsto G \tag{110}$$

where the parameter vector $\mathbf{P}$ contains all relevant microscopic variables of the model. This functional approach expresses the gravitational constant G as a product of the **Base Soliton Identity** ($\mathbf{G}_{\text{Base}}$) and a **Geometric Scaling Functional** (F_{geom}):

$$\mathcal{U}(\mathbf{P}) = \mathbf{G}_{\text{Base}}\cdot F_{\text{geom}}(\mathbf{P}), \qquad \mathbf{G}_{\text{Base}} = \frac{c^2\mathbf{r_e}}{\mathbf{m_e}}. \tag{111}$$

The explicit geometric form used in this work is

$$\boxed{\mathcal{U}(\mathbf{P}) = \frac{c^2 r_e}{m_e}\cdot\frac{1}{A_\pi}\cdot\left(\frac{1}{N_{\text{geom}}A_\pi}\right)^3\cdot\frac{1}{K_{WC}\sqrt{N_{\nu,\text{eff}}}}} \tag{112}$$

where A_π is the core geometric factor and $N_{\nu,\text{eff}}$ is the effective volume deficit derived from the unified coupling C_{unif}.

Properties of the Operator $\mathcal{U}$

- **Non-linearity:** The mapping is characterized by cubic scaling of the stiffness factor and an inverse square-root dependency on the effective constituent count.
- **Monotonicity:** For a fixed statutory background, any increase in $N_{\nu,\text{eff}}$ results in a deterministic decrease in the emergent value of G.
- **Zero calibration:** The operator contains only geometric and structural inputs: A_π, N_{geom}, K_{WC}, $N_{\nu,\text{eff}}$.

the G implementation result showed in table 6.

Table 6: Numerical Verification of the Gravitational Constant G within the Geometric EWT Framework

Parameter / Result	Numerical Value	Source and Physical Context
I. Base Soliton Identity	$2.7802522591 \times 10^{32}$	$G_{\text{Base}} = \frac{c^2 r_e}{m_e}$. Fundamental Soliton base.
II. Raw Geometry Value	$6.6804369615 \times 10^{-11}$	$G_{\text{EWT, raw}}$. Pure $K+1$ projection; error 0.09187% relative to CODATA.
III. Geometric Result	$6.6775199755 \times 10^{-11}$	Zero-calibration geometric G. Uses L_p^{geom} and N_{geom}; error 0.04817%.
IV. CODATA Target	$6.6743050000 \times 10^{-11}$	Experimental reference value (CODATA 2022).

Note: All values are expressed in units of $m^3 kg^{-1} s^{-2}$.

11.6 Geometric Emergence of the Planck Length

The Planck length in EWT is identified with the fundamental EMC spacing λ_l. In the zero-calibration formulation, λ_l is not taken from experiment; it follows from the geometric identity for the gravitational constant and the Planck-length definition.

Before the effective dilution factor X_{eff} can be written down, the unified coupling operator C_{unif} that governs it must be derived. The operator bridges the raw geometric projection of the wave centres onto the BCC lattice with the electromagnetic response of the medium, and its three terms each have a distinct structural origin.

The Fundamental Geometric Ratio: C_{Raw}

Prior to any electromagnetic coupling, the model identifies the pure interaction ratio between the soliton's wave centres and the background lattice. This factor,

$$C_{\text{Raw}} = 1 + \frac{1}{K_{WC}}, \tag{113}$$

is the "ideal" geometric scaffold of the displacement. The term $1/K_{WC}$ is the harmonic reciprocal of the $K_{WC} = 10$ wave centres; it normalises the global volume deficit to the individual node's displacement capacity. The additive $+1$ is the statutory equilibrium of the medium, the

unperturbed vacuum state that the correction is applied on top of. For $K_{WC} = 10$ this gives $C_{\text{Raw}} = 1.1$, the baseline efficiency of the vacuum displacement before any elastic or fine-structure corrections.

The Lattice Projection Factor L_p^{geom}

The natural geometric starting point for the interfacial correction is the ideal BCC projection factor

$$L_p^{\text{geom}} = \frac{2}{\sqrt{3}} \approx 1.154700538, \tag{114}$$

which arises as the inverse of the dimensionless nearest-neighbour distance ($d_{NN}/a = \sqrt{3}/2$) in the $Im\bar{3}m$ lattice. Its combination $\pi L_p^{\text{geom}} = 2\pi/\sqrt{3}$ is the dimensionless fundamental wavevector for a standing wave along the [111] body diagonal, capturing the projection of the 3D lattice onto the 2D soliton interface.

The Unified Bridge: C_{unif}

The full operator adds the electromagnetic coupling across the lattice interface:

$$C_{\text{unif}} = \frac{1}{K_{WC}} + 1 + \frac{\alpha_{\text{geom}}}{\pi L_p^{\text{geom}}}. \tag{115}$$

The three terms correspond to the three distinct contributions of the medium's response:

- **Harmonic reciprocal** $1/K_{WC}$: the specific contribution of the $K_{WC} = 10$ wave centres, ensuring that the global deficit is normalised to the individual node's displacement capacity.
- **Unitary baseline** $+1$: the statutory equilibrium of the medium, the unperturbed vacuum state. It ensures the correction is additive to the existing background density $N_{\nu,\text{stat}}$, maintaining continuity of the elastic field.
- **Interfacial tension bridge** $\alpha_{\text{geom}}/(\pi L_p^{\text{geom}})$: the electromagnetic strain coupled through the lattice projection factor and the geometric π. This term scales the large-scale volumetric deficit down to the interfacial tension at the soliton boundary.

With C_{unif} established, the effective dilution factor follows directly:

$$X_{\text{eff}} = \frac{A_\pi \, 3 \, K_{WC} \sqrt{2}}{C_{\text{unif}}}, \tag{116}$$

where $A_\pi = 4\pi^3 + \pi^2 + \pi$ is the geometric core of the soliton. The numerator counts the active geometric budget of the soliton (the core geometry A_π, the three spatial dimensions, the K_{WC} wave centres, and the $\sqrt{2}$ face-diagonal coupling); the denominator C_{unif} is the structural impedance of the medium through which that budget is projected. The ratio is the geometric dilution of the statutory background density inside the soliton.

The self-consistent derivation of λ_l starts from the Planck-length definition,

$$\lambda_l = \sqrt{\frac{\hbar_{\text{geom}} \, G_{\text{geom}}}{c^3}}, \tag{117}$$

in which $\hbar_{\rm geom}$ is given by Eq. (76), while $G_{\rm geom}$ is the geometric gravitational constant whose explicit form is derived in Section 11. In the present section it is sufficient to use its geometric structure,

$$G_{\rm geom} = \frac{G_{\rm Base}}{A_\pi}\left(\frac{1}{N_{\rm geom}A_\pi}\right)^3 \frac{1}{K_{WC}\sqrt{N_{\nu,\rm eff}}}, \tag{118}$$

where

$$G_{\rm Base} = \frac{c^2 r_e}{m_e}. \tag{119}$$

The effective density $N_{\nu,\rm eff}$ is related to the statutory density $N_{\nu,\rm stat}$ by

$$N_{\nu,\rm eff} = \frac{N_{\nu,\rm stat}}{X_{\rm eff}}, \qquad N_{\nu,\rm stat} = \left(\frac{r_\nu}{2e\lambda_l}\right)^3. \tag{120}$$

Substituting these expressions into the Planck-length definition cancels m_e, c, and the common A_π factors. Solving the resulting algebraic equation for λ_l gives

$$\lambda_l = \left[\frac{r_e^2\sqrt{X_{\rm eff}}}{\alpha_{\rm geom}A_\pi^4 N_{\rm geom}^3 K_{WC}\left(\frac{r_\nu}{2e}\right)^{3/2}}\right]^2, \tag{121}$$

where e is Euler's number. Normalized form is:

$$\lambda_l = \left[\frac{r_e^2\sqrt{\dfrac{3\sqrt{2}\,A_\pi}{C_{\rm unif}}}}{\alpha_{\rm geom}A_\pi^4 N_{\rm geom}^3\sqrt{K_{WC}}\left(\dfrac{r_\nu}{2e}\right)^{3/2}}\right]^2. \tag{122}$$

Equation (122) substitutes the definition of $X_{\rm eff}$ directly, so that the wave-centre count enters once through $\sqrt{K_{WC}}$ rather than twice. The fine-structure constant appears twice in the expression, but in two distinct roles: the leading $\alpha_{\rm geom}$ is the electromagnetic coupling of the soliton itself, while the $\alpha_{\rm geom}$ inside $C_{\rm unif}$ is the electromagnetic strain across the soliton–vacuum interface. The two appearances are not a redundancy; they reflect the two surfaces through which the same coupling is projected.

Planck introduced this length as a formal dimensional combination of $\hbar$, G, and c, without reference to a microscopic substrate. In the present framework, the same quantity appears as the self-consistent solution of the geometric BCC equations, thereby providing a physical origin for the scale that Planck treated only as a formal unit.

Equations (121) and (122) are therefore not an independent ansatz. They are the direct consequence of combining the geometric fine-structure constant, the geometric gravitational scaling, and the Planck-length definition into a single closed algebraic constraint. Here e is Euler's number and r_ν is the geometric neutrino radius derived in Section 5.

Numerically,

$$\lambda_l \approx 1.6166464 \times 10^{-35}\ \text{m}. \tag{123}$$

The CODATA value is

$$\lambda_{l,\rm CODATA} = 1.6162 \times 10^{-35}\ \text{m}, \tag{124}$$

so the relative deviation is approximately 0.0276%. This is a genuine geometric prediction.

$X_{\rm eff}$ is not a free parameter; it is the geometric bridge that connects the statutory EMC background to the effective soliton deficit. It encapsulates how $K_{WC} = 10$, the ideal BCC projection, and the geometric core A_π jointly dilute the vacuum density inside matter.

11.7 Consistency with the Planck Charge

The geometric Planck charge was derived earlier as

$$q_P = \frac{e}{\sqrt{\alpha_{\text{geom}}}}. \tag{125}$$

With the above values,

$$q_P \approx 1.8755478 \times 10^{-18}\ \text{m}, \tag{126}$$

which matches the expected amplitude scale.

The fact that λ_l, q_P, and $\hbar_{\text{geom}}$ all emerge from the same BCC geometry confirms that the model does not treat the Planck scale as an external boundary but as an internal consequence of the lattice structure.

11.8 Electromagnetic Consistency Check

In SI units, the relation between charge, $\hbar$, and α is

$$e^2 = 4\pi\epsilon_0 \hbar c \alpha. \tag{127}$$

Substituting the geometric expression for $\hbar$, Eq. (76), cancels α_{geom} identically and yields

$$e^2 = 4\pi\epsilon_0 m_e c^2 r_e. \tag{128}$$

In the natural units of the BCC lattice ($c = 1$; see Section 9), the SI electromagnetic units are absorbed into a single dimensionful conversion factor x,

$$x\, e^2 = m_e r_e, \qquad x \equiv 10^{-7}\ \text{m}^4. \tag{129}$$

The factor x is a unit-conversion quantity, not an independent geometric parameter: its numerical value 10^{-7} m^4 is set by the SI definitions of the metre, second, and coulomb. Its structural form, $x = \mu_0/(4\pi) = \alpha h/(2\pi e^2 c)$, is a combination of α, e, h, and c, which in EWT are geometric. Thus x inherits whatever geometric content those constants carry, but its specific numerical value is a SI convention. In dimensionful EWT units it equals

$$x = \frac{\mu_0}{4\pi} = 10^{-7}\ \text{m}^4, \tag{130}$$

the pre-2019 SI value of the Coulomb constant combination $\mu_0/(4\pi)$, expressed in the EWT dimensions $[\mu_0] = [\text{H/m}] = [\text{m}^4]$. Its inverse $1/x = 4\pi/\mu_0$ does not coincide with the Coulomb constant $k_e = c^2\mu_0/(4\pi)$ when c is expressed numerically in SI units; the two differ by the factor c^2. They coincide only in the $c = 1$ limit of the BCC lattice, where $k_e = x$.

The identity (129) is dimensionally consistent,

$$[x] \cdot [e^2] = [\text{m}^4] \cdot [\text{m}^2] = [\text{m}^6] = [\text{m}^5] \cdot [\text{m}] = [m_e] \cdot [r_e], \tag{131}$$

and numerically both sides equal 2.5670×10^{-45} m^6.

This is not an independent derivation of e, since m_e, r_e, and x are inputs, but it is a powerful internal consistency test: it shows that the electromagnetic scale (e) is rigidly tied to the soliton's mass and radius through a single conversion factor x, with no residual dependence on $\hbar$ or α.

Remark on the Status of ϵ_0

The SI vacuum permittivity ϵ_0 is not an independent parameter of nature. Historically, it appeared as a "property of the vacuum" only because the SI separates electric and magnetic units. The relation

$$c^2 = \frac{1}{\mu_0 \epsilon_0} \tag{132}$$

carries no physical content in natural units: setting $c = 1$ reduces it to the unit-definition identity $\mu_0 \epsilon_0 = 1$. What appears as a "vacuum property" in SI is therefore a convention for relating electric and magnetic units, not a dynamical parameter of the medium.

After the 1983 redefinition of c and the 2019 SI revision, this is manifest: ϵ_0 is a derived unit-conversion factor,

$$\epsilon_0 = \frac{e^2}{2\alpha h c}, \tag{133}$$

and carries no geometric information beyond that already encoded in α, e, h, and c. In the EWT framework, where charge is identified with wave amplitude in metres, the vacuum permittivity is therefore not required as a dynamical parameter. Its entire role is absorbed into the single dimensionful factor $x = 10^{-7}$ m^4 of Eq. (129).

11.9 Status of Inputs

The only experimental inputs in model are

$$r_e, \quad m_e, \quad c, \quad e. \tag{134}$$

In natural units ($c = 1$; see Section 9), c is a unit conversion and m_e is geometrically tied to r_e through the r^5 scaling law.

In SI units the relation between charge, $\hbar$, and α is $e^2 = 4\pi\epsilon_0 \hbar c \alpha$. In the EWT framework ϵ_0 is not an independent input: together with μ_0 it is fixed by the metric condition $c^2 = 1/(\mu_0 \epsilon_0)$ and therefore carries no new geometric information. No calibrated parameters appear in this section. Note that the lepton AMM sector, discussed in Section 16, introduces one additional discrete postulate, the muon Fibonacci label $L_\mu = F_5 = 5$, which is absent from the gravitational and electromagnetic constants derived here.

The cosmological sector, discussed in Section 24, introduces one further empirical anchor beyond the four above: the present CMB temperature $T_{\mathrm{cmb},0} = 2.7255 \pm 0.0006$ K (COBE/-FIRAS [70]). The exact SI conversion constant k_B enters the same sector, but it carries no dynamical information: like c, it is a unit-conversion factor fixed by the SI 2019 definitions. With these two additions, the cosmological chain predicts H_0 and the de Sitter value of Λ from the same BCC geometry, as detailed in Section 24.

The quantities G_{geom}, λ_l, $\hbar_{\mathrm{geom}}$, and α_{geom} are not independent predictions. They form a single closed algebraic system fixed by the BCC lattice geometry, the electron soliton inputs r_e, m_e, c, e, and the geometric self-consistency condition

$$\lambda_l = \sqrt{\frac{\hbar_{\mathrm{geom}}\, G_{\mathrm{geom}}}{c^3}}. \tag{135}$$

Solving this system is the geometric self-consistency of the model, not a stepwise derivation of one constant from another.

In the standard framework the Planck-length relation links three independent empirical constants. In the present model the same relation emerges as an internal geometric condition,

because both $\hbar_{\text{geom}}$ and G_{geom} are already determined by the BCC lattice. The Planck scale is therefore not imported from the Standard Model; it is reconstructed as a consequence of the underlying geometry.

11.10 Duality of Soliton Density

Presented model inherently describes a fundamental duality of densities within the Soliton. The localized energy density ρ_E inside the Soliton must be higher than the surrounding medium, due to localized standing wave energy, to account for its mass $E = mc^2$. Conversely, the geometric packing density function $\rho(r)$ must be lower than the uniform density of the surrounding space, creating the measurable packing deficit $N_{\nu,\text{eff}}$.

This dichotomy means the Soliton is a region of high energy but low geometric packing efficiency. The dynamic processes described in models unifying mass and charge as wave energy [26, 29, 15] can be hypothesized to be the cause of the packing density deficit of the EMC, by continuously forcing the Soliton's wave components outward, defining the effective volume and maintaining its stability.

The accompanying constant geometric term A_π^{-1} is specifically interpreted here as the Geometric Stability Factor. This factor quantifies the fixed boundary ratio resulting from the continuous dynamic process of outward forcing on the Soliton's internal wave structure.

This definition of density introduces a central principle of the Enhanced EWT Model: gravity is an emergent, pressure-driven phenomenon that acts on the Soliton's effective surface, as the universal medium strives for equilibrium by filling this geometric deficit. This provides a basis for interpreting gravity as a push force induced by a vacuum pressure gradient [34].

11.11 The Degraded EMC Wall: Spherical Masking and Isotropy

A fundamental puzzle in any geometric theory of matter is how an asymmetric, rotating not perfect spherical soliton can generate a perfectly isotropic gravitational field. The resolution lies in a structural boundary layer that surrounds every soliton – the **Degraded EMC Wall**.

The Degraded EMC Wall is a spherical shell of finite thickness located at a radius r_{mask} where the packing density of Elastic Medium Constituents (EMCs) transitions from the diluted interior of the soliton toward the statutory background density $N_{\nu,\text{stat}}$. The wall is a region of **elevated EMC density** caused by the active push-out mechanism: as the soliton's standing waves displace EMCs from the interior, these constituents accumulate just outside the soliton boundary, creating a local density peak:

$$\rho_{\text{EMC}}(r_{\text{mask}}^-) \;<\; \rho_{\text{EMC}}(r_{\text{wall}}), \tag{136}$$

where r_{wall} denotes a point located strictly inside the Degraded EMC Wall.

The exact value of this peak relative to $N_{\nu,\text{stat}}$ depends on the balance between the internal push-out force and the external statutory pressure of the surrounding BCC lattice. In ordinary solitons this equilibrium yields a moderate peak, while in extreme gravitational environments the peak can become very large.

The Degraded EMC Wall acts as a mandatory transducer between the asymmetric core and the exterior. Because the individual EMCs are spherical units, the BCC lattice can only achieve a stable, static interface with the statutory background by adopting a spherical shape. The wall therefore enforces a spherical boundary condition on the energy density distribution emerging from the asymmetric core. The effect can be quantified by the Isotropy Operator $\mathcal{I}$ (introduced in Section 25.7).

11.11.1 Open question: Wall density relative to statutory background

The precise value of the peak EMC density within the Degraded EMC Wall, ρ_{wall}, relative to the statutory background $N_{\nu,\text{stat}}$, remains undetermined. Two scenarios are physically permissible:

- **Scenario A (asymptotic approach):** $\rho_{\text{wall}} < N_{\nu,\text{stat}}$, with the density increasing monotonically from the soliton interior toward the statutory value.
- **Scenario B (local maximum):** $\rho_{\text{wall}} > N_{\nu,\text{stat}}$, corresponding to a genuine local density peak caused by the active push-out.

Both scenarios are compatible with the integral determination of $N_{\nu,\text{eff}}$ and with the derivation of G. Future high-resolution simulations may discriminate between them. For the purpose of the present work, we treat the wall density ratio as an open parameter $\eta = \rho_{\text{wall}}/N_{\nu,\text{stat}}$.

12 Local Metric Phenomena from EMC Lattice Deformation

12.1 The Two Faces of EMC Displacement: Speed and Trajectory

In the Enhanced EWT framework, light propagation involves two distinct, though coupled, consequences of the EMC lattice strain field. They are frequently conflated in continuous medium optics, so it is necessary to state them with structural precision.

Light propagation through the strained EMC lattice is described by two distinct, though coupled, consequences of the same underlying density-deficit field. They are frequently conflated in continuous- medium optics, so it is necessary to state them with structural precision.

1. **Coordinate phase velocity from node availability.** In natural lattice units ($c \equiv 1$), the statutory speed of light is the undisturbed cell ratio $c = \lambda_l/t_p = 1$. Around a matter soliton, the local EMC density drops below the statutory value, so a wave packet traversing the strained region finds fewer nodes per unit coordinate length. This reduces the effective coordinate speed of light, $v_\gamma(r) < 1$, encoded by an isotropic scalar refractive index $n_\gamma(r) = 1/v_\gamma(r) > 1$. The reduction is parameterized by the *dimensionless fractional density deficit*

$$\delta(r) \;\equiv\; 1 - \eta(r) \;=\; 1 - \frac{N_\nu(r)}{N_{\text{stat}}} \;\geq\; 0, \tag{137}$$

   which measures how far the local density has fallen below the statutory background. The functional relation between n_γ and δ (equivalently, between v_γ and $\eta = 1-\delta$) is established by the postulate of Eq. (143) in Section 12.3: $v_\gamma(r) = \eta(r)$, hence $n_\gamma(r) = 1/\eta(r)$. The scalar deficit $\delta(r)$, and not a physical vector displacement, is the geometric object that encodes the lattice strain; a literal displacement $\vec{u}_{\text{phys}}(r)$ of dimension $[L]$ is not required.

2. **Trajectory deflection from the refractive-index gradient.** A light ray traverses the strained lattice along a curved trajectory because the scalar refractive index $n_\gamma(r)$ varies spatially. The deflection is governed by the transverse gradient $\nabla_\perp n_\gamma(r)$, which continuously reorients the wavefront toward the mass center, exactly as in classical ray optics in a medium with a spatially varying index. No separate vector restoring field is required: the same scalar $n_\gamma(r)$ that slows the phase velocity also bends the ray.

12.2 Bridging the Vector Displacement to the Scalar Refractive Index

The connection between the local density deficit and the scalar refractive index is established by the density profile of the soliton together with the postulate of Eq. (143). A 3D standing-wave soliton exhibits an asymptotic radial energy-density deficit decaying as $1/r$. In the simplest spherical weak-field model, the normalized local EMC density takes the form

$$\eta(r) \equiv \frac{N_\nu(r)}{N_{\mathrm{stat}}} = 1 - \frac{r_s}{r}, \tag{138}$$

where $r_s = 2GM$ is the gravitational radius in natural units ($c = 1$). The corresponding dimensionless fractional deficit is

$$\delta(r) = 1 - \eta(r) = \frac{r_s}{r} = 2\Phi_N(r), \tag{139}$$

where $\Phi_N(r) = GM/r = r_s/(2r)$ is the dimensionless Newtonian gravitational potential per unit energy. The deficit $\delta(r)$ appearing in Eq. (139) is a scalar, dimensionless quantity; it is not a physical displacement of dimension $[L]$, and its interpretation as a "radial displacement field" used in earlier versions of this work is superseded by the scalar-deficit description above.

For a null probe, the coordinate refractive index follows directly from the postulate $v_\gamma(r) = \eta(r)$:

$$n_\gamma(r) = \frac{1}{v_\gamma(r)} = \frac{1}{\eta(r)} = \left(1 - \frac{r_s}{r}\right)^{-1}. \tag{140}$$

In the weak-field limit $r_s/r \ll 1$, the binomial expansion gives

$$n_\gamma(r) = 1 + \frac{r_s}{r} + \left(\frac{r_s}{r}\right)^2 + \cdots \approx 1 + \frac{r_s}{r} = 1 + 2\Phi_N(r), \tag{141}$$

which reproduces the standard first-order gravitational refractive index and provides the leading-order input for the solar-deflection integral of Section 12.5.

12.3 Asymptotic Continuous Limit and Schwarzschild Equivalence

Eq. (141) corresponds to the first-order weak-field expansion of the full continuous expression:

$$n_\gamma(r) = \frac{1}{\eta(r)} = \left(1 - \frac{r_s}{r}\right)^{-1}. \tag{142}$$

For a null probe, we *postulate* the coordinate speed of light

$$v_\gamma(r) = \eta(r), \qquad v_{\mathrm{clock}}(r) = \sqrt{\eta(r)}, \tag{143}$$

on the following physical ground: a light ray traverses the strained lattice, so its coordinate speed is governed by the ratio of the local spatial strain to the local temporal strain, while a clock is a bound standing wave whose period is set only by the local node availability. In the language of an effective metric with $+---$ signature, this is

$$g_{00}(r) = \eta(r), \qquad g_{rr}(r) = -\frac{1}{g_{00}(r)} = -\eta(r)^{-1}, \tag{144}$$

with g_{rr} obtained from g_{00} and v_γ by the identity $g_{rr} = -g_{00}/v_\gamma^2$. The coordinate refractive index follows as

$$n_\gamma(r) = \frac{1}{v_\gamma(r)} = \eta(r)^{-1} = \left(1 - \frac{r_s}{r}\right)^{-1}, \tag{145}$$

which is identical to the radial Schwarzschild optical factor $n_{\rm GR} = \sqrt{-g_{rr}/g_{00}} = (1-r_s/r)^{-1}$, and reproduces the standard identity $c_{\rm coord}(r) = c\sqrt{g_{00}/(-g_{rr})}$ of a static metric with the coefficients in Eq. (144).

The expansion of Eq. (142) is

$$n_\gamma(r) = \left(1 - \frac{r_s}{r}\right)^{-1} \approx 1 + \frac{r_s}{r} + \frac{r_s^2}{r^2} + \mathcal{O}\left(\left(\frac{r_s}{r}\right)^3\right), \tag{146}$$

with $r_s/r = 2\Phi_N(r)$ in natural units, giving the factor 2 in front of $\Phi_N(r)$ and the structural foundation for the observed $1.75''$ solar limb deflection. In the undisturbed statutory vacuum ($r_s/r \to 0$), $n_\gamma(r) \to 1$, restoring the fundamental lattice ratio $c = \lambda_l/t_p = 1$.

12.4 Mechanical Origin of Gravitational Redshift in the EMC Lattice

In Enhanced EWT, both electromagnetic propagation and gravitational time dilation arise from a single microscopic mechanism: the spatial deformation of the EMC node separation $\lambda_l(r)$ around a matter soliton.

In the natural units of the BCC lattice, the speed of light is the fundamental metric conversion factor

$$c \equiv \frac{\lambda_l}{t_p} = 1.$$

Consequently, length and time share the same dimension, $[m] = [s]$, and the gravitational potential is dimensionless without the explicit c^2 factor that appears only in the SI-unit convention. In these natural units, the Newtonian potential is simply

$$\Phi_N(r) = \frac{GM}{r}, \tag{147}$$

and the gravitational radius of the source is

$$r_s = 2GM. \tag{148}$$

Define the dimensionless local EMC lattice density ratio as:

$$\eta(r) \equiv \frac{N_\nu(r)}{N_{\rm stat}} = 1 - \frac{r_s}{r} = 1 - 2\Phi_N(r), \tag{149}$$

where $\Phi_N(r) = GM/r = r_s/(2r)$ in natural units.

A physical clock is modeled as a standing-wave soliton whose fundamental period $T(r)$ is determined by the internal round-trip time of a longitudinal EMC lattice wave traversing the soliton core. Because the internal wave relies on node-to-node propagation across the same strained lattice spacing $\lambda_l(r)$ that dictates external optics, its effective propagation rate scales with the square root of the local node availability $\eta(r)$:

$$v_{\rm clock}(r) = \sqrt{\eta(r)} = \sqrt{1 - 2\Phi_N(r)} \approx 1 - \Phi_N(r). \tag{150}$$

For a rest-frame internal path length $2L_0$, the clock period in the gravitational potential field becomes:

$$T(r) = \frac{2L_0}{v_{\rm clock}(r)} = \frac{T_0}{\sqrt{1 - 2\Phi_N(r)}} \approx T_0\left(1 + \Phi_N(r)\right), \qquad T_0 = 2L_0. \tag{151}$$

The fractional frequency shift reproduces the standard gravitational redshift:

$$\frac{\Delta f}{f} = \frac{f(r) - f_0}{f_0} = -\Phi_N(r) = -\frac{GM}{r} = -\frac{r_s}{2r}. \tag{152}$$

Structural Unity of Metric Phenomena. This completes the mechanical equivalence between General Relativity metric components and EWT lattice strain:

- **Temporal metric component ($\sqrt{g_{00}}$):** Encoded by the internal clock wave speed $v_{\text{clock}}(r) = \eta(r)^{1/2}$, producing gravitational time dilation.
- **Effective optical metric factor:** Encoded by the optical refractive index $n_\gamma(r) = 1 + r_s/r = 1 + 2\Phi_N(r)$, whose weak-field form is the first-order expansion of $\eta(r)^{-1} = (1 - r_s/r)^{-1}$, producing solar light deflection.

  In standard GR language, this effective index combines the contributions of both g_{00} and g_{rr}. In EWT it arises directly from the integrated EMC displacement u_r.

Both fundamental phenomena are physically unified as dual manifestations of the same local EMC density profile $\eta(r) = 1 - r_s/r$, but they project onto different observables: the clock samples the local node availability through $\sqrt{\eta}$, while the light ray samples the combined spatial and temporal strain through η^{-1}.

12.5 Shapiro Delay from the EMC Refractive Index

The third local metric observable is the Shapiro delay: a light signal passing near a massive body takes longer to cross the region than it would in the undisturbed statutory vacuum. In the Enhanced EWT framework, this delay is not an independent postulate but a direct consequence of the same EMC lattice strain that produces light bending and gravitational redshift.

In natural lattice units ($c \equiv 1$), the coordinate speed of a photon in the strained EMC geometry is

$$v_\gamma(r) = \frac{1}{n_\gamma(r)} = 1 - \frac{r_s}{r} = \eta(r), \tag{153}$$

where $n_\gamma(r)$ is the optical refractive index introduced in Sec. 12.3. The coordinate time required for a signal to travel along a path $\mathcal{C}$ is therefore

$$t = \int_{\mathcal{C}} \frac{ds}{v_\gamma(r)} = \int_{\mathcal{C}} n_\gamma(r)\, ds. \tag{154}$$

In the weak-field limit,

$$n_\gamma(r) - 1 = \frac{r_s}{r} = 2\Phi_N(r), \qquad \Phi_N(r) = \frac{GM}{r}, \tag{155}$$

so the excess delay relative to the statutory vacuum along a one-way path is

$$\Delta t_{\text{one-way}} = \int_{\mathcal{C}} \big(n_\gamma(r) - 1\big)\, ds = 2\int_{\mathcal{C}} \Phi_N(r)\, ds. \tag{156}$$

For a round-trip radar signal grazing the solar limb, the dominant contribution comes from the near-Sun segment of the trajectory. Approximating the path as a straight line with impact parameter $b \simeq R_\odot$, the integral over the two-way path gives the logarithmic form

$$\Delta t_{\text{round-trip}} \approx 4GM\, \ln\!\left(\frac{4R_E R_P}{b^2}\right) \qquad (c = 1), \tag{157}$$

where R_E and R_P are the heliocentric distances of the emitter and receiver. Restoring the speed of light c yields the standard SI expression,

$$\Delta t_{\text{round-trip}} \approx \frac{4GM}{c^3}\, \ln\!\left(\frac{4R_E R_P}{b^2}\right) \qquad \text{(SI)}, \tag{158}$$

which matches the classic relativistic Shapiro delay.

The Shapiro delay constitutes the third independent projection of the single EMC lattice deformation surrounding a matter soliton.

12.6 Macroscopic EMC Density Profile of a Star

The EMC density profile around a macroscopic body such as the Sun is not a simple scaled copy of an individual lepton's internal profile. A star is a vast collection of overlapping matter solitons, each contributing a local displacement of the EMC medium. The resulting effective density field is therefore a collective, smoothed macroscopic structure rather than a single soliton boundary wall.

For a static, spherically symmetric body of total mass M, the normalised EMC density ratio,

$$\eta(r) \equiv \frac{N_\nu(r)}{N_{\mathrm{stat}}}, \tag{159}$$

can be naturally partitioned into three characteristic regions:

1. **Interior deficit region ($r < R_\odot$).** Inside the body, the conserved energy density of the constitutive matter solitons forces a local reduction of the EMC lattice density. This deficit is finite and, for an ordinary star, remarkably shallow compared with the extreme depletion expected in a compact remnant or black hole configuration. Its exact functional shape depends on the internal equation of state and is not required for external metric tests.

2. **Stellar transition layer ($r \approx R_\odot$).** At the physical boundary of the body, the EMC density transitions from the interior deficit to the statutory background. For a star, this layer is broad and diffuse relative to the subatomic Degraded EMC Wall of a single electron, arising from the macroscopic superposition of countless individual soliton boundaries.

3. **Far-field tail ($r \gg R_\odot$).** Far outside the body, the EMC density relaxes toward the statutory value N_{stat} according to

$$\eta(r) = 1 - \frac{r_s}{r}, \qquad r_s = 2GM \quad (c \equiv 1). \tag{160}$$

   This represents the weak-field, spherically symmetric solution consistent with the long-range lattice strain generated by the enclosed mass M.

The local metric phenomena evaluated in this work — solar light bending, gravitational redshift, and Shapiro delay — are strictly insensitive to the detailed structure of the interior and transition regions. They are entirely governed by the far-field tail $\eta(r) = 1 - r_s/r$, which is universal for any static, spherically symmetric EMC deficit of total strain magnitude r_s.

Thus, the solar EMC profile used in our analytical and numerical validations represents the *macroscopic far-field tail* of the collective density field, distinct from the microscopic Degraded EMC Wall of an isolated lepton.

12.7 Derivation of the Far-Field EMC Density Profile

The far-field density profile, $\eta(r) = 1 - r_s/r$, is derived from the static continuum equilibrium of the EMC lattice rather than selected as an ad hoc fitting ansatz.

In the long-wavelength limit, $r \gg \lambda_l$, the discrete BCC lattice is well approximated by an isotropic elastic continuum. Outside a bounded source region, where external volume forces

vanish, the static response of such a medium is governed by the Laplace equation for the scalar strain field. Writing the normalized local EMC density as

$$\eta(r) = 1 + \delta\eta(r), \qquad |\delta\eta(r)| \ll 1, \tag{161}$$

the linearised equilibrium condition takes the form

$$\nabla^2 \, \delta\eta(r) = 0, \qquad r > R, \tag{162}$$

where R is the effective radius of the central matter distribution.

Under spherical symmetry and the boundary condition that the perturbation vanishes at spatial infinity ($\delta\eta \to 0$ as $r \to \infty$), the general solution to Eq. (162) is uniquely constrained to the harmonic radial tail:

$$\delta\eta(r) = -\frac{A}{r}, \qquad A > 0. \tag{163}$$

By analogy with electrostatics, the coefficient A plays the role of a monopole charge for the strain field: it measures the total strength of the deformation induced by the central source. The integration constant A is fixed by applying Gauss's theorem to the strain flux across a closed surface surrounding the mass M. In natural units ($c \equiv 1$), this gives

$$A = r_s = 2GM. \tag{164}$$

Consequently, the far-field profile is uniquely determined as:

$$\eta(r) = 1 - \frac{r_s}{r}, \qquad r \gg R. \tag{165}$$

Thus, the profile contains no free functional parameters and no empirical tuning coefficients. The $1/r$ form is required by the harmonic equilibrium of the lattice, and its amplitude is fixed by the enclosed gravitational radius r_s.

12.8 From Microscopic EMC Displacement to the Gravitational Radius

The far-field EMC density profile

$$\eta(r) = 1 - \frac{r_s}{r} \tag{166}$$

contains a single characteristic length scale, r_s. In the preceding numerical implementations, this amplitude was identified with the gravitational radius of the source. This section demonstrates that this identification is a necessary consequence of the additive microscopic EMC displacements of constituent matter solitons, rather than an independent boundary input.

Consider a static, spherically symmetric body of total mass $M = \sum_i m_i$, composed of elementary matter solitons. In the long-wavelength, weak-field regime, the lattice continuum behaves as a linear elastic medium, ensuring that the total strain field is the linear superposition of individual soliton perturbations.

For any elementary soliton i of mass m_i, the fundamental geometric identity of EWT yields the individual monopole strain amplitude:

$$a_i = 2\, G_{\mathrm{EWT,\,geo}}\, m_i, \tag{167}$$

where $G_{\mathrm{EWT,\,geo}}$ is the uncalibrated geometric constant derived purely from the ideal BCC lattice projection parameter $L_p^{\mathrm{geom}} = 2/\sqrt{3}$. Owing to field linearity in the far zone, the total monopole amplitude A of the macroscopic body is the sum of the constituent amplitudes:

$$A = \sum_i a_i = 2\,G_{\mathrm{EWT,\,geo}} \sum_i m_i = 2\,G_{\mathrm{EWT,\,geo}}\,M. \tag{168}$$

Equation (168) establishes that the macroscopic strain amplitude is strictly proportional to the total mass enclosed.

Applying Gauss's divergence theorem to the radial displacement field $\delta\eta(r) = \eta(r) - 1$, the outward strain flux through a sphere of radius r is given by:

$$\oint_{\partial V} \nabla(\delta\eta) \cdot d\mathbf{S} = 4\pi r^2 \frac{d\,\delta\eta}{dr} = 4\pi A. \tag{169}$$

For the fundamental harmonic solution $\delta\eta(r) = -A/r$, the radial derivative evaluates to:

$$\frac{d\,\delta\eta}{dr} = \frac{A}{r^2}. \tag{170}$$

Consequently, the Neumann boundary condition enforced at the source radius R in the numerical implementation is fully derived from first principles:

$$\left.\frac{d\,\delta\eta}{dr}\right|_R = \frac{2\,G_{\mathrm{EWT,\,geo}}\,M}{R^2}. \tag{171}$$

Integration yields the unique far-field density profile:

$$\eta(r) = 1 - \frac{2\,G_{\mathrm{EWT,\,geo}}\,M}{r}, \tag{172}$$

from which the gravitational radius naturally emerges as a derived effective quantity:

$$r_s \equiv 2\,G_{\mathrm{EWT,\,geo}}\,M. \tag{173}$$

No empirical value of G is injected into this derivation. The minor numerical discrepancy (~ 5 ppm) between $G_{\mathrm{EWT,\,geo}}$ and the CODATA value lies within the experimental uncertainty of Newton's constant and is fully accounted for by the non-ideal spherical packing impedance ζ (Sec. 25.8.2). This completes the logical closure: the far-field profile and metric artifacts are generated strictly from microscopic lattice parameters and the enclosed mass.

12.9 Emergent Metric Encodings from Lattice Dynamics

The elastic constant k introduced in Sec. 1.5 can be derived from a microscopic pair potential $V(r)$ between neighbouring EMCs. In the continuum limit of a one-dimensional lattice, the equilibrium spacing is

$$a(\eta) \equiv \frac{1}{\eta}. \tag{174}$$

where η is the normalised EMC density.

In this one-dimensional chain, the natural contact interaction is modelled by the repulsive potential

$$V(r) = \frac{V_0}{r}, \tag{175}$$

whose first derivative gives the force between neighbouring sites and whose second derivative evaluated at the equilibrium spacing defines the local spring stiffness:

$$k(\eta) = \left.\frac{d^2V}{dr^2}\right|_{r=1/\eta} \propto \eta^3. \tag{176}$$

The mass per lattice site is fixed, since each site corresponds to one EMC. The mass per unit length is therefore

$$\rho = \frac{m_0}{a(\eta)} = m_0\,\eta. \tag{177}$$

The long-wavelength speed of sound in a one-dimensional lattice is not $\sqrt{k/m_0}$, but rather

$$v(\eta) = a(\eta)\sqrt{\frac{k(\eta)}{m_0}}. \tag{178}$$

Using the one-dimensional modulus

$$E(\eta) = k(\eta)\,a(\eta) \;\propto\; \eta^2, \tag{179}$$

this can be written in the standard continuum form

$$v(\eta) = \sqrt{\frac{E(\eta)}{\rho(\eta)}} \;\propto\; \sqrt{\frac{\eta^2}{\eta}} = \sqrt{\eta}. \tag{180}$$

This is exactly the clock-speed encoding

$$v_{\text{clock}} \;\propto\; \sqrt{\eta}, \tag{181}$$

now obtained as an emergent property of the microscopic lattice dynamics.

This wave-speed $\sqrt{\eta}$ is the propagation speed of a timelike probe in the strained lattice. For a null probe we *postulate* the coordinate speed $v_\gamma(\eta) = \eta$ (see Sec. 12.3, Eq. (143)): a light ray samples both the spatial and the temporal strain factors of the medium, whereas a clock samples only the local node availability. The two metric coefficients are therefore

$$g_{00}(\eta) = \eta, \qquad g_{rr}(\eta) = -\frac{1}{\eta}, \tag{182}$$

with g_{rr} obtained from g_{00} and v_γ by $g_{rr} = -g_{00}/v_\gamma^2$, and the corresponding refractive index is

$$n_\gamma(\eta) = \frac{1}{v_\gamma(\eta)} = \frac{1}{\eta} = \left(1 - \frac{r_s}{r}\right)^{-1}. \tag{183}$$

A physical clock in this medium is a standing-wave soliton of fixed linear dimension L. Its fundamental frequency is therefore

$$f(\eta) \;\propto\; \frac{v(\eta)}{L} \;\propto\; \sqrt{\eta}. \tag{184}$$

The clock-speed exponent follows directly from the 1D lattice dynamics derived above. The light-index exponent rests on the postulate of Sec. 12.3, Eq. (143); it is not derived from the lattice here.

Dimensional reduction and 3D extension: The one-dimensional chain represents the effective longitudinal propagation along a high-symmetry axis of the three-dimensional BCC lattice. In a full 3D lattice, the equilibrium spacing scales as $a(\eta) \propto \eta^{-1/3}$, while the volumetric mass density scales as $\rho_{3D} \propto \eta$. The same wave-speed scaling $v(\eta) = \sqrt{E/\rho} \propto \sqrt{\eta}$ is then recovered from a microscopic inverse-cube pair potential $V(r) \propto r^{-3}$.

Note on the separation of gravitational and electromagnetic domains: In the EWT framework, electromagnetic phenomena operate in the *soliton-energy domain*. Charge and radiation are carried by the amplitude and phase of standing and travelling waves. Consequently, two

counter-propagating electromagnetic waves superpose linearly in free space: their interference does not produce a leading-order change in the EMC packing density $\rho(r)$. Gravity, by contrast, operates in the *EMC density domain*. It is the macroscopic response of the lattice to a localized deficit in the packing density $N_{\nu,\mathrm{eff}}$. This distinction explains why gravity is universal, acting on all forms of energy that displace the medium, while remaining exceptionally weak compared to electromagnetic effects, which directly involve the energy amplitudes of the solitons.

12.10 Summary of Local Metric Observables

Thus, the single normalized EMC density profile $\eta(r) = 1 - r_s/r$ unifies all three classic local metric observables without introducing any free parameters:

- **Solar light bending** through the spatial gradient of the postulated index $n_\gamma(r) = \eta^{-1} = \left(1 - r_s/r\right)^{-1}$ (Sec. 12.3),
- **Gravitational redshift** through the internal clock speed $v_{\mathrm{clock}} = \sqrt{\eta} = \sqrt{1 - r_s/r}$,
- **Shapiro delay** through the integrated excess of $n_\gamma(r) - 1$ along the propagation path.

The two encodings—$\sqrt{\eta}$ for timelike probes and η^{-1} for null probes—reflect that a clock samples the local node availability while a light ray samples the combined spatial and temporal strain of the same lattice deformation.

12.11 In-Platform Numerical Validation

The analytical derivations presented in this section are validated numerically in the OpenWave research suite. The scripts implement the same EMC density profile $\eta(r) = 1 - r_s/r$ and the same encodings used in the analytical treatment, and compare the results against established reference values. The numerical outcomes are summarised in Table 7.

Table 7: Numerical validation of the local metric phenomena. All scripts implement the same EMC density profile and encodings derived analytically in this section.

Artifact	Quantity	EWT result	Reference	Difference
M4.3	Solar-limb light bending	1.751722 arcsec	1.7517 arcsec (GR)	0.0012%
M4.4	Gravitational redshift $\Delta f/f$	-2.123135×10^{-6}	GR exact factor	0 (identity)
M4.5	Shapiro round-trip delay	239.014 μs	239.014 μs	0.000020%
M4.6	Far-field profile $\eta(r)$	RK4 solution	analytic $1 - r_s/r$	5.7×10^{-14}
M4.8	Monopole amplitude A	2.954320×10^3 m	CODATA G	65.65 ppm
M4.9	Stiffness exponent α	3.0	expected 3.0	< 0.05
M4.9	Wave-speed exponent β	0.5	expected 0.5	< 0.05

The numerical results reproduce the analytical predictions of this section. The scripts are part of the OpenWave research repository [55].

For M4.3, M4.4, and M4.5 the encoding of the EMC density profile into the refractive index and the clock speed is taken from this manuscript; the scripts perform the resulting integrations and confirm the agreement with the general-relativistic reference values.

For M4.6, the radial Laplace equation is solved numerically with the monopole flux condition at the solar surface and the asymptotic Robin boundary condition at the outer boundary. The resulting profile matches the analytic form to 5.7×10^{-14} of η.

For M4.8, the monopole amplitude is computed from the geometric G chain of the emergence engine with the packing impedance $\zeta_{\text{est}} = (1 - \eta_{\text{BCC}})/(\eta_{\text{BCC}} N_{\text{ideal}})$, without calibrating to the measured fine-structure constant. The resulting amplitude differs from the CODATA value by 65.65 ppm. The pure geometric limit ($\zeta = 0$) gives 1743.57 ppm, establishing the packing correction as the relevant physical input.

For M4.9, the one-dimensional lattice dynamics with a repulsive pair potential $V(r) = V_0/r$ yields the exponents $\alpha = 3$ (stiffness scaling $k \propto \eta^\alpha$) and $\beta = 0.5$ (wave-speed scaling $v \propto \eta^\beta$), which are the values used in the encodings of the clock speed and the refractive index.

13 Newtonian Force from Interacting EMC Deficits

The same far-field density deficit that reproduces the local metric observables also gives rise to the Newtonian force law.

Consider two matter solitons of masses M_1 and M_2, separated by a distance R. Each soliton sources a monopole density deficit

$$\delta\eta_i(r) = -\frac{A_i}{r}, \qquad A_i = \frac{2G_{\text{EWT, geo}} M_i}{c^2}. \tag{185}$$

The static interaction energy arises from the integrated overlap of the two density gradients in the surrounding medium,

$$U_{\text{int}}(R) = -K_{\text{emc}} \int \nabla(\delta\eta_1) \cdot \nabla(\delta\eta_2)\, dV, \tag{186}$$

The minus sign follows from the negative-definite energy functional adopted for the EMC field; this functional is the geometric counterpart of the Newtonian field energy $-\frac{|\nabla\Phi|^2}{8\pi G}$, and the normalization constant is

$$K_{\text{emc}} = \frac{c^4}{16\pi G_{\text{EWT, geo}}}. \tag{187}$$

In spherical coordinates centred on the first source, the scalar product of the two gradients is

$$\nabla(\delta\eta_1) \cdot \nabla(\delta\eta_2) = A_1 A_2 \frac{r^2 - rR\cos\theta}{r^3 (r^2 + R^2 - 2rR\cos\theta)^{3/2}}. \tag{188}$$

Performing the angular integration over the spherical shell at radius r yields

$$\int_0^{2\pi} \int_0^{\pi} [\nabla(\delta\eta_1) \cdot \nabla(\delta\eta_2)] \sin\theta\, d\theta\, d\phi = \begin{cases} 0, & r < R, \\ \dfrac{4\pi A_1 A_2}{r^4}, & r \geq R, \end{cases} \tag{189}$$

which is the exact field-theoretic analogue of Newton's shell theorem.

Integrating over the volume element $dV = r^2 \sin\theta\, dr\, d\theta\, d\phi$ for the non-vanishing region $r \geq R$ gives

$$U_{\text{int}}(R) = -\frac{c^4}{16\pi G_{\text{EWT, geo}}} \cdot 4\pi A_1 A_2 \int_R^\infty \frac{dr}{r^2} = -\frac{c^4 A_1 A_2}{4 G_{\text{EWT, geo}} R}. \tag{190}$$

The attractive force directed along the line connecting the centers is

$$F = -\frac{dU_{\text{int}}}{dR} = -\frac{c^4 A_1 A_2}{4 G_{\text{EWT, geo}} R^2}. \tag{191}$$

Substituting the monopole amplitudes $A_i = 2G_{\mathrm{EWT,\ geo}}M_i/c^2$ yields precisely

$$F = -\frac{G_{\mathrm{EWT,\ geo}}M_1M_2}{R^2}. \tag{192}$$

No additional fundamental force is postulated. The inverse-square law is the static consequence of two EMC density deficits interacting through vacuum push-out pressure. The 4π factor emerges dynamically from the 3D angular integration, while the 16π factor represents the field stress-energy normalization.

13.1 In-Platform Validation: Normalization Consistency

The derivation of the inverse-square force law from EMC field overlap is validated numerically in the OpenWave research suite. The baseline gate compares the field-overlap force F_{EMC} with the Newtonian force F_{Newton}, both built from the same geometric G_{geom} derived from the BCC lattice. The result is summarised in Table 8.

Table 8: Baseline gate for the Newtonian force from EMC field overlap. Both F_{EMC} and F_{Newton} are built with $G_{\mathrm{geom}} = 6.677519975508460 \times 10^{-11}$ m^3 kg^{-1} s^{-2}.

Quantity	**Value**
F_{EMC}	$3.544205553218 \times 10^{22}$ N
F_{Newton}	$3.544205979545 \times 10^{22}$ N
Relative difference	1.203×10^{-5}%

Because G_{geom} enters both the monopole amplitude $A_i = 2G_{\mathrm{geom}}M_i/c^2$ and the coupling $K_{\mathrm{emc}} = c^4/(16\pi G_{\mathrm{geom}})$, it cancels identically in $F_{\mathrm{EMC}} \equiv F_{\mathrm{Newton}}$. The gate therefore passes for any value of G_{geom} and remains a normalization-consistency check on A, K_{emc}, and the angular factor 4π. Mutating any single normalization factor breaks the agreement, as shown in Table 9.

Table 9: Mutation test of the normalization-consistency gate. Each mutation changes a single normalization factor while keeping the rest of the derivation unchanged.

Mutation	**Observed relative difference**	**Gate result**
Baseline	1.203×10^{-5}%	PASS
Angular factor ($4\pi \to 3\pi$)	25.0%	FAIL
Monopole amplitude ($2GM/c^2 \to GM/c^2$)	75.0%	FAIL
Coupling denominator ($16\pi \to 8\pi$)	100.0%	FAIL

The amplitude can also be written without the symbol G_{geom}, directly in the chain factors of the emergence engine:

$$A = \frac{2Mr_e}{m_e} \cdot \frac{\sqrt{X_{\mathrm{eff}}}}{A_\pi^4\, N_{\mathrm{geom}}^3\, K_{WC}\sqrt{N_{\nu,\mathrm{stat}}}}, \qquad X_{\mathrm{eff}} = \frac{N_{\nu,\mathrm{stat}}}{N_{\nu,\mathrm{eff}}}. \tag{193}$$

Every factor on the right-hand side is geometric: $A_\pi = 4\pi^3 + \pi^2 + \pi$, $N_{\mathrm{geom}} = 8\pi^4(1-\zeta)$, $K_{WC} = 10$, and X_{eff}, $N_{\nu,\mathrm{stat}}$ from the geometric chain. The anchors r_e, m_e are the measured dimensional quantities. The product of Eq. (193) is the engine's G_{EWT} formula multiplied by $2M/c^2$, term for term, so $A = 2G_{\mathrm{geom}}M/c^2$ by value. The geometric form agrees with the G-based form to machine precision (1.5×10^{-16} for the solar amplitude, 2.0×10^{-16} for the

terrestrial amplitude): the two are the same expression, so the agreement is an identity, not a confirmation.

The force gate does not depend on the value of G_{geom}. Making it a test of G_{geom} requires an amplitude whose value is fixed without passing through the G_{EWT} formula; this remains an open clause in the research program.

Strength clause. The two halves of the strength clause for the gravitational sector are stated as follows.

- **Circularity: discharged.** G_{geom} enters the force test directly, derived from the BCC geometry, without re-entering through an input.
- **Accuracy: stated, not met.** The G_{geom} residual against CODATA is 0.048169%, or 21.9× the CODATA 2022 relative uncertainty on G (22 ppm). At that distance the derived value does not agree with the measured G within its uncertainty. The dimensional anchors r_e, m_e, c are the measured quantities used to build the dimensionless geometric ratio; the derivation produces G_{geom} as a dimensionless ratio against $c^2 r_e/m_e$, not G from nothing.

The scripts are part of the OpenWave research repository [56].

14 Relation to Existing Emergent Gravity Frameworks

The Geometric Identity Model (EWT) is fundamentally aligned with the concept of Emergent Gravity (EG), where gravity is viewed as a collective, induced, or thermodynamic phenomenon [31, 32, 30]. In particular, Kleinert's 1987 formulation of gravity as a theory of defects in a crystalline medium with second-gradient elasticity [65] provides a direct precedent for the present BCC-lattice picture: the EWT gravitational field emerges as a static strain field of a discrete elastic substrate, in close analogy with the defect-based mechanism that Kleinert derived from the elastic energy of a crystal. The present work differs in that the BCC lattice is not just a mathematical model, but is identified with a physical Elastic Medium Constituent lattice, and the gravitational constant is derived from the packing fraction and coordination number of that lattice rather than from an effective elastic action. The derivation of G from explicit microscopic geometry and its dependence on the vacuum's effective volume deficit places this work within the general realm of EG approaches. However, EWT offers a unique, explicit microscopic realization that is independent of purely thermodynamic or entropic principles.

14.1 Comparison of Mechanisms and Scales

The EWT model differs from established EG frameworks in three critical aspects: the nature of the microscopic degrees of freedom, the source of gravitational emergence, and the explicit scaling factor for the gravitational constant G.

14.1.1 Microscopic Degrees of Freedom (DoF)

- **Thermodynamic/Entropic (Jacobson, Verlinde):** The DoF are typically macroscopic or informational, related to the **area elements of a holographic screen** or **entanglement** between quantum fields in the vacuum.
- **Induced (Sakharov):** The DoF are the **quantum fields themselves**, with gravity arising from the zero-point energy of the vacuum.

- **EWT (Geometric Soliton):** The DoF are **explicitly geometric** and local: the internal structure and energy density profile $\rho(\mathbf{r})$ of the Soliton, and its internal magnetic coherence $\epsilon_{\mathbf{M}}$. This provides a tangible, particle-scale physical realization of the underlying DoF.

14.1.2 Source of Gravitational Emergence

The EWT model proposes a **dual mechanism** for the strength of gravity, distinguishing the source of the mass/energy from the source of its weakness, which is not present in standard EG theories:

- **Thermodynamic/Entropic:** The source is the **thermodynamic state** (e.g., temperature T) or the **information content** of spacetime.
- **Induced (Sakharov):** The source is the **bulk change in vacuum energy** caused by spacetime curvature.
- **EWT (Geometric Soliton):**
  a) **Primary Source ($\mathbf{G_{Base}}$):** The gravitational force originates from the fundamental **quantity of EMC** (energy-mass-coherence) within the Soliton ($\mathbf{m_e}$).
  b) **Emergence/Weakness:** The vast weakness of gravity (scaling from $\mathbf{G}_{\text{Base}}$ to G) is caused by the **Geometric Deficit**—the ratio between the Soliton's compact volume and the much larger effective volume it influences in the surrounding EWT medium.

14.1.3 Scaling Factor for G

The frameworks rely on fundamentally different scaling parameters to define the observed value of G:

- **Thermodynamic/Entropic:** G is typically scaled by $\hbar$ and parameters related to the **temperature of the horizon ($\mathbf{T}$)**, e.g., $\mathbf{G} \propto 1/(\mathbf{S} \cdot \mathbf{T})$.
- **Induced (Sakharov):** G is inversely proportional to the **fourth power of the quantum field theory cutoff scale ($\mathbf{\Lambda}$)**, $\mathbf{G} \propto 1/\mathbf{\Lambda}^4$.
- **EWT (Geometric Soliton):** G is scaled by the complete, derived, dimensionless scaling factor $\mathbf{\Omega_G}$:

$$\mathbf{G} = \mathbf{G}_{\text{Base}} \cdot \mathbf{\Omega_G}, \quad \text{where } \mathbf{\Omega_G} = \frac{1}{\mathbf{A}_\pi} \cdot \left(\frac{1}{\mathbf{N}_{\text{final}} \mathbf{A}_\pi} \right)^3 \cdot \frac{1}{\mathbf{K_{WC}} \cdot \sqrt{\mathbf{N}_{\nu,\text{effective}}}} \tag{194}$$

  The EWT scaling factor $\mathbf{\Omega_G}$ is **explicitly constructed** from the Soliton's geometric parameters ($\mathbf{A}_\pi$, $\mathbf{N}_{\text{final}}$) and the unified volume deficit ($\mathbf{N}_{\nu,\text{effective}}$), linking G to the Soliton's internal dynamics rather than external thermodynamic quantities.

The EWT model thus acts as a bridge, providing the explicit, geometric and non-thermodynamic identity $\mathcal{U}(\mathbf{P})$ required to link fundamental particle parameters to the macroscopic gravitational constant G, while remaining compatible with the holographic and entropic concepts that underlie the vacuum structure.

Comparison with Induced Gravity (Sakharov)

The EWT scaling factor $\mathbf{\Omega_G}$ exhibits a profound structural isomorphism with the concept of **Induced Gravity** proposed by Andrei Sakharov [30]. In Sakharov's framework, gravity is not a fundamental force but an emergent "elasticity of the vacuum," where the gravitational constant G scales inversely with the fourth power of a quantum field theory cutoff, $G \propto 1/\Lambda^4$.

In the EWT model, this "cutoff" is replaced by the geometric coupling constant $\mathbf{A}_\pi$. The total suppression factor $\mathbf{\Omega_G}$ incorporates the term $\mathbf{A}_\pi^{-4}$ (derived from the linear and volumetric saturation: $\mathbf{A}_\pi^{-1} \cdot \mathbf{A}_\pi^{-3}$), providing a **purely geometric derivation** for the quartic attenuation observed in emergent gravity theories.

While Sakharov required vacuum fluctuations to define the scale, EWT identifies this scale as the physical phase-saturation point of the BCC lattice. This suggests that the 10^{42} weakness of gravity is the result of the push-out effective mechanism modulated by "stiffness" coupled with the holographic projection of nodal energy ($\sqrt{\mathbf{N}_{\nu,\text{eff}}}$).

Gravity is scaled by the total energy density of the space occupied by matter, where the attenuation follows $\mathbf{G} \propto 1/\mathbf{A}_\pi^4$, further diluted by the holographic transition to the soliton's surface.

This perspective implies that within a certain range, the gravitational strength is determined by the lattice's volumetric resistance, while the final 10^{42} weakness is dominated by the geometric projection:

$$\mathbf{G} = \underbrace{\frac{\mathbf{G}_{\text{Base}}}{\mathbf{A}_\pi{}^4 \cdot \mathbf{N}_{\text{final}}{}^3}}_{\text{Volumetric Energy Density (push-out)}} \cdot \underbrace{\frac{1}{K_{WC} \cdot \sqrt{\mathbf{N}_{\nu,\text{eff}}}}}_{\text{Surface Dilution}} \tag{195}$$

14.1.4 Evolution of the EWT Framework: From Wave Attenuation to EMC Push out

The introduction of the Push out Mechanism marks a fundamental shift in the interpretation of gravitational forces within the Energy Wave Theory (EWT). While the baseline EWT model [11] treats the BCC lattice primarily as a passive *wave carrier*—where gravity is attributed to a minute loss of wave energy (shadow effect)—the current framework redefines the BCC lattice as an **active actor**.

In this refined model, the gravitational constant is no longer dependent on an arbitrary wave attenuation coefficient. Instead, the extreme weakness of gravity ($\sim 10^{-42}$) is a direct consequence of **holographic energy dilution**. The energy density, concentrated within the volumetric saturation of the lattice ($A_\pi{}^4$), is projected onto the 3D surface of the matter soliton and further corrected by the effective EMC density $N_{\nu,\text{eff}}$.

This transition is governed by the fundamental geometric derivative of the vacuum state, where the volumetric saturation A_π^4 yields the **push-out force density**:

$$y = A_\pi{}^4 \implies y' = 4A_\pi{}^3 \tag{196}$$

- **Baseline EWT:** Relies on a nearly infinitesimal energy loss during wave propagation to account for the weakness of gravity, which lacks a direct structural justification.
- **Push-out Model (Enhanced EWT):** Identifies gravity as a "blurred image" of an extremely strong lattice interaction. The 10^{-42} factor emerges naturally from the geometric ratio between the 4D saturation base ($A_\pi{}^4$) and the holographic surface projection of the soliton.

By replacing "leaking wave energy" with the **structural stiffness** and **statutory density** ($N_{\nu,\mathbf{stat}}$) of the lattice, the model provides a deterministic explanation for gravitational emergence. Gravity is thus revealed as the mechanical reaction of the BCC medium to a localized EMC deficit, governed by the derivative of the vacuum's volumetric impedance.

Structural Range: Continuity vs. Wave Dissipation

A significant challenge for the baseline EWT model is explaining the immense, theoretically infinite range of gravity. In a framework based on infinitesimal wave attenuation, a signal as weak as 10^{-42} would realistically be dissipated by the vacuum's stochastic noise over long distances.

The **EMC Push-out Mechanism** provides a structural solution to this problem:

- **Geometric Continuity:** Unlike a wave signal that requires constant energy propagation, the push-out mechanism creates a **static structural deformation** of the BCC lattice. Since the medium is a continuous mechanical entity, the gradient between $N_{\nu,\text{stat}}$ and $N_{\nu,\text{eff}}$ must propagate throughout the entire network to maintain topological equilibrium.
- **Infinite Reach:** The $1/r^2$ scaling is revealed as a geometric requirement of 3D projection from the 4D saturation base (A_π^4), not a result of "fading waves." This explains why gravity remains stable across cosmological scales—it is not a "message" being sent, but a permanent "tilt" in the lattice geometry.

By shifting the focus from "energy loss" to **lattice elasticity**, the Enhanced EWT accounts for both the extreme weakness of the force (via holographic dilution) and its universal reach (via structural integrity), identifying gravity as the deterministic mechanical response of the vacuum's impedance.

14.1.5 EWT as Pressure-Driven Emergent Gravity

The fundamental mechanism proposed by EWT—where mass creates a **Geometric Deficit** ($\mathbf{N}_{\nu,\text{effective}}$) which is then compensated by the surrounding EWT medium—aligns the model with modern interpretations of gravity as a **Push Force** or a vacuum pressure effect.

This interpretation contrasts sharply with the classic Newtonian view (attraction) and even with General Relativity (spacetime curvature), instead focusing on local density gradients:

- **Density Deficit and Energy Conservation:** The Soliton structure (the particle) constitutes a localized region of **lower packing** (a geometric deficit $\mathbf{N}_\nu$) compared to the undisturbed background of the EWT medium (the vacuum). Simultaneously, due to continuous internal wave interference, the Soliton **conserves** a large, localized amount of **energy** (mass equivalent).
- **Equilibrium Drive (Gravity):** The resulting force (gravity) is the expression of the **EWT medium's fundamental drive to restore equilibrium** by moving into the region of lower packing. This motion generates a net **pressure gradient**, which pushes all matter towards the center of the deficit.

This conceptualization places EWT in close affinity with models advocating that gravity is generated by the **pressure of a Quantum Vacuum** (or: *Medium*) flowing from higher to lower energy density areas [35], offering a coherent, picture of the gravitational interaction that is both emergent and pressure-driven.

This interpretation provides direct justification for the appearance of the square root term ($\mathbf{N}_{\nu,\text{effective}}{}^{-1/2}$) in the scaling factor $\mathbf{\Omega_G}$. This term represents the **geometric transition** from

the underlying three-dimensional volume deficit ($\mathbf{N}_{\nu,\text{effective}}$) to the **two-dimensional surface effect** (pressure) exerted by the EWT medium, aligning EWT with the surface-based scaling principles of holographic gravity models.

15 Geometric Validation: The Fundamental Identity and the Base AMM State (a_e)

The internal consistency of the Energy Wave Theory (EWT) is verified through the geometric relationship between a soliton's energy and its spatial extent. In this framework, the anomalous magnetic moment (AMM) is treated as a deterministic consequence of the vacuum's elastic properties.

15.1 Geometric Mass-to-Radius Identity ($E \propto r^5$)

The core principle of EWT [37, 29] dictates that the energy E of a soliton scales with the fifth power of its geometric radius r ($E \propto r^5$). This identity provides the deterministic link between a particle's measured mass and its geometric size:

$$\frac{E_x}{E_e} = \left(\frac{r_x}{r_e}\right)^5 \quad \Longrightarrow \quad r_x = r_e \left(\frac{E_x}{E_e}\right)^{1/5} \tag{197}$$

This relationship ensures that the geometric radius r_f is not an arbitrary parameter but is directly derivable from the particle's measured energy. This identity confirms that the soliton's scale is governed by the Wave Count hierarchy, establishing a unified set of parameters for both mass and magnetic anomaly. Radius scaling is directly related to the mass scaling model presented in section 19.3.

15.2 Physical Origin of the r^5 Scaling: Geometric Energy Density

The quintic relationship between energy and radius ($E \propto r^5$), as identified in the foundational EWT framework [37, 29], is here demonstrated to be a rigorous requirement of energy conservation within the vacuum lattice. While the r^5 scaling provides a powerful predictive tool, its physical origin lies in the convergence of three fundamental geometric and dynamic factors of the soliton:

1. **Volumetric Occupancy (r^3):** The three-dimensional spatial extent of the wave center within the BCC lattice.
2. **Amplitude Scaling (r^1):** Since charge is expressed as wave amplitude A in meters [29], the stability of the soliton requires the amplitude to be geometrically coupled with the radius ($A \propto r$) to maintain the structural integrity.
3. **Resonance Mode Count (r^1):** The number of standing-wave periods N that fit within the soliton scales linearly with its radius ($N \propto r$). A larger soliton supports more resonant modes, each oscillating at the characteristic lattice frequency $f \propto 1/r$.

The product of these factors ($r^3 \times r^1 \times r^1 = r^5$) confirms that mass is a measure of "dynamic information density". However, the divergence between this quintic energy surge and the cubic geometric compensation capacity ($\propto r^3$) explains the precision limits encountered in earlier models (e.g., the 10-16% error for the muon and tau in [37]).

The three contributions combine into two scaling laws that are the formal basis of the Onion Model. The first is the energy–radius identity,

$$\boxed{E \propto r^5} \tag{198}$$

which follows directly from the product of the three factors above and is identical to the mass-to-radius relation in Eq. (197). The second is the volumetric capacity of the three-dimensional lattice,

$$\boxed{V \propto r^3} \tag{199}$$

which expresses the geometric compensation available in real space as the soliton expands. The mismatch between the quintic energy surge, Eq. (198), and the cubic geometric compensation, Eq. (199), is the source of the $r^5/r^3 = r^2$ disparity that forces recursive shell formation in the Onion Model described in 16.3.

15.3 The Geometric Status of Mass: Kilogram as a Derived Unit

The $E \propto r^5$ scaling law acquires its full dimensional significance when combined with the single structural constant that defines the causal structure of the BCC lattice: the maximum propagation speed c.

Step 1 — Metric unification via c. The axiom $c \equiv \lambda_l/t_p$ (Section 9) identifies c as the intrinsic conversion factor between the spatial and temporal steps of the BCC lattice. Adopting $c = 1$ (the natural units of the lattice) unifies the dimensions of space and time:

$$c = 1 \implies [\mathrm{s}] = [\mathrm{m}]. \tag{200}$$

Step 2 — Mass from the r^5 geometric scaling law. The fundamental mass-radius relation of EWT (Section 15.1) states that the energy of a soliton is entirely defined by its geometric extent, scaling as the fifth power of its radius:

$$E \propto r^5. \tag{201}$$

Assuming the proportionality constant is a dimensionless topological feature of the BCC lattice, the geometric dimension of energy is identically $[r^5] = [\mathrm{m}^5]$.

In classical SI units, energy is defined by $E = mc^2$. Equating the geometric and classical dimensions yields the dimensional structure of mass:

$$[m \cdot c^2] = [r^5] \implies [\mathrm{kg}] \cdot [\mathrm{m}^2 \cdot \mathrm{s}^{-2}] = [\mathrm{m}^5]. \tag{202}$$

Solving for the kilogram reveals its true derived nature in the EWT framework:

$$[\mathrm{kg}] = [\mathrm{m}^3 \cdot \mathrm{s}^2]. \tag{203}$$

When evaluated in the natural units of the lattice ($c = 1 \implies [\mathrm{s}] = [\mathrm{m}]$), the dimensional conversion becomes completely spatial:

$$[\mathrm{kg}] = [\mathrm{m}^3 \cdot \mathrm{m}^2] = [\mathrm{m}^5]. \tag{204}$$

Step 3 — Mass hierarchy from r^5. With the mass scale anchored to r_e through the r^5 law, the entire lepton mass hierarchy follows from the topological integer sequence K_{WC}:

$$\frac{m_x}{m_e} = \left(\frac{r_x}{r_e}\right)^5 = \left(\frac{K_x}{K_e}\right)^5, \qquad \text{dimensionless.} \tag{205}$$

Step 4 — Gravitational consistency. The independently derived gravitational identity (Section 11) provides a non-trivial consistency check. The base gravitational scale is defined as $G_{\text{Base}} = c^2 r_e/m_e$. Substituting the dimensional relation $[\text{kg}] = [\text{m}^3 \cdot \text{s}^2]$ from Eq. (204) gives:

$$[G_{\text{Base}}] = \frac{[\text{m}^2 \cdot \text{s}^{-2}] \cdot [\text{m}]}{[\text{m}^3 \cdot \text{s}^2]} = [\text{s}^{-4}]. \tag{206}$$

Adopting $c = 1$ ($[\text{s}] = [\text{m}]$) reduces this to $[G_{\text{Base}}] = [\text{m}^{-4}]$, which is exactly the dimension of the ratio r_e/m_e when mass carries the geometric dimension $[\text{m}^5]$:

$$[r_e/m_e] = \frac{[\text{m}]}{[\text{m}^5]} = [\text{m}^{-4}]. \tag{207}$$

The gravitational constant therefore requires no independent mass scale: its dimension, and ultimately its value, is entirely determined by the geometry of the BCC lattice.

In this framework, the SI kilogram occupies the same logical status as π in Euclidean geometry: it is not a free parameter, but a structural consequence of the underlying geometric medium. The reduction of all three SI mechanical base units to a single geometric primitive (the BCC lattice spacing λ_l) is the dimensional expression of the claim that EWT is a parameter-free geometric theory of matter.

15.4 Geometric Dimension of the Reduced Planck and G Constants

The reduction of mass to a geometric quantity has an immediate consequence for the reduced Planck constant. In SI units $\hbar$ has dimension

$$[\hbar] = [\text{kg}][\text{m}]^2[\text{s}]^{-1}. \tag{208}$$

Using the natural-unit identifications $[\text{s}] = [\text{m}]$ and $[\text{kg}] = [\text{m}]^5$, this becomes

$$[\hbar] = [\text{m}]^5 \cdot [\text{m}]^2.[\text{m}]^{-1} = [\text{m}]^6. \tag{209}$$

In the geometric formulation of EWT, $\hbar$ is therefore not an independent quantum input. It is a derived geometric quantity of dimension length to the sixth power.

The same reduction applies to the gravitational constant. From the defining relation $[G] = [\text{m}]^3/([\text{kg}][\text{s}]^2)$, the natural-unit substitutions give

$$[G] = \frac{[\text{m}]^3}{[\text{m}]^5\,[\text{m}]^2} = [\text{m}]^{-4}. \tag{210}$$

This is consistent with the earlier derivation of the base gravitational scale from the electron soliton.

As a result, the Planck length

$$l_{\text{P}} = \sqrt{\frac{\hbar G}{c^3}} \tag{211}$$

reduces at $c = 1$ to the geometric dimensional identity

$$[l_{\text{P}}] = \sqrt{[\hbar][G]} = \sqrt{[\text{m}]^6\,[\text{m}]^{-4}} = [\text{m}]. \tag{212}$$

Thus l_{P} is dimensionally identical to the primary EMC lattice step λ_l. In the geometric EWT framework, λ_l is not constructed from $\hbar$ and G. The single lattice length λ_l fixes the geometric

units from which $\hbar$ and G inherit their SI dimensions when the system is expressed in laboratory units.

This geometric dimension admits a direct interpretation within the Geometric Ladder of the EWT framework. The sixth degree of freedom is associated with the volumetric bosonic budget $3D+A+f+r$, which is represented by the π^6 rung (Section 2.1). The reduced Planck constant is therefore the geometric measure of the six-dimensional phase-space volume available to a soliton resonance. The dimensional exponent 6 is not accidental: it is the same 6 that appears in the π^6 volumetric coupling operator for neutral weak interactions.

Using the Dimensional Weight Operator formalism, the six factors of π forming the π^6 rung are precisely those that define the six-dimensional phase-space cell whose quantum is $\hbar$.

15.5 The Fundamental Magnetic Deficit ϵ_M

In EWT, the magnetic moment correction is a purely geometric phenomenon resulting from the non-ideal stiffness of the elastic medium. The fundamental magnetic deficit constant, ϵ_M, is defined by the effective geometric stiffness modulus (N_{geom}) and the volumetric factor (π^3):

$$\epsilon_M \equiv \frac{1}{N_{\text{geom}} \cdot \pi^3} \tag{213}$$

The total stiffness deficit (Loss Factor) of the elastic medium is thus expressed as $\epsilon_M \cdot \pi^3 = 1/N_{\text{geom}}$.

15.6 Derivation of the a_e Base Formula

The anomalous magnetic moment of the electron (a_e) is derived as the product of the Ideal Geometric Moment (the Schwinger Term) and the Stiffness Retention Factor:

$$\boxed{a_{\text{Base}}^{\text{Geometric}} = \frac{\alpha_{\text{geom}}}{2\pi} \cdot \left(1 - \epsilon_M \cdot \pi^3\right)} \tag{214}$$

Numerical verification, using the geometric α_{geom} and ϵ_M from the BCC lattice, yields $a_{\text{Base}}^{\text{Geometric}} \approx 1159.916228$ ppm. This result aligns with the experimental value of the electron ($a_e^{\text{exp}} \approx 1159.652182$ ppm) with a relative error of approximately 0.02277%, establishing the electron as the fundamental geometric ground state.

16 The Recursive Lepton Hierarchy: Nodal Shell Resonance Model

Heavier leptons — the Muon (Ψ_μ) and the Tau (Ψ_τ) — emerge through **Recursive Nodal Inclusion** (the "Onion Model"). Each generation is treated as a cumulative wave-packing excitation within the BCC vacuum lattice, where the nodal structure is governed by a characteristic geometric factor $2\pi^2$ (the surface area of a torus, though other topologies yielding the same factor are not excluded).

The numerical values quoted in this section are obtained directly from the geometric Python implementation described in the Appendix.

Methodological Framework: Nodal Metrics and Geometric Determinism

To maintain theoretical rigor, a distinction is established between the discrete lattice and the resonant excitation. While the **Wave Center (Soliton)** (K_{WC}) is the fundamental physical and energy-conserving entity, its internal dynamics involve a level of complexity that makes direct continuous analysis less efficient. In contrast, the **Node** provides a highly effective **topological metric** for the EWT framework. By utilizing the discrete points of the BCC lattice as a reference for **phase space occupancy**, the "Onion Model" derives lepton generations through the precise counting of nodal density (K).

In this approach, nodes do not function as a rigid measurement of spatial distance, but rather as a **quantized index of resonant occupancy**. When the energy surge ($\propto r^5$) exceeds the volumetric capacity ($\propto r^3$) of the base geometry, the system's adaptation is captured by its "latching" onto additional nodal degrees of freedom. By focusing on **nodal counting** as a measure of structural complexity rather than absolute spatial radii, the model replaces the analytical overhead of wave-center dynamics with the deterministic logic of the lattice, enabling sub-percent level predictions without any calibrated parameters.

16.1 Resonance Growth Law and Fibonacci Labeling

The hierarchy is defined by a recursive addition of nested shells. The nodal increment (ΔK) follows a deterministic geometric law based on the factor $2\pi^2$ (which coincides with the surface area of a torus, but may also arise from other resonant configurations) and a decimal scale shift operator. The nodal increment is generated by the discrete operator of Section 16.2, Eq. (215).

In the EWT framework, these shells are parametrized by specific Fibonacci labels (L_n), which serve as classification indices for the magnetic anomaly.

16.2 Discrete Resonance Labels of the BCC Lattice

The hierarchy of lepton generations is defined by the recursive addition of shells, where the nodal count K_n evolves according to the $2\pi^2$ operator. Within the EWT framework, the proposed shell hierarchy is parametrized by discrete Fibonacci labels (L_n), which are used as model-specific shell-closure indices for wave-centre configurations. These labels are dimensionless integers; they are not standard physical quantities, but EWT-internal classification indices:

- **Muon label ($L_\mu = F_5 = 5$):** The first shell ($K = 207$) is labelled by the Fibonacci number $F_5 = 5$. This value follows from the electron core label $F_1 = 1$ by advancing the Fibonacci index by a fixed step of four, as described below. It enters the tau accumulation as an additive shell-coupling term $L_\mu^2 = 25$.
- **Tau label ($L_\tau = F_9 = 34$):** The second shell ($K = 2181$) is labelled by the Fibonacci number $F_9 = 34$. This value follows from $L_\mu = 5$ by the same step of four. It does not enter the numerical computation of a_τ in the three-generation model; its square $L_\tau^2 = 1156$ is reserved as the analogous shell-coupling term for a hypothetical fourth shell.

Discrete operator. The nodal increments are generated by the postulated discrete operator

$$\Delta K_k = \begin{cases} \text{round}(10^0 \cdot \pi^2), & k = 0, \\ \text{round}(10^k \cdot 2\pi^2), & k \geq 1, \end{cases} \tag{215}$$

where k is the shell index. The factor π^2 in the core term is interpreted phenomenologically as a measure of an assumed two-dimensional angular closure, and the additional factor of two in the

Table 10: Nodal Shell Parameters and Fibonacci Labels

Lepton Shell	**Operator**	**Nodal ΔK**	**Total K_n**	**Fibonacci label L_n**
Core (Electron)	$10^0 \cdot \pi^2$	10	10	1
Shell 1 (Muon)	$10^1 \cdot 2\pi^2$	197	207	5
Shell 2 (Tau)	$10^2 \cdot 2\pi^2$	1974	2181	34
Shell 3 (4th gen.)*	$10^3 \cdot 2\pi^2$	19739	21920	233

Here $n = k + 1$ is the cumulative generation index and k is the shell index used in Eq. (215).

* Formally generated candidate. A fourth shell is not observed; it is listed to display the recursive structure of the operator and of the Fibonacci step. Its total nodal count $K_4 = 21920$ and its label $L_4 = F_{13} = 233$ follow from the same rules that generate the three generations represented in the present construction. Whether such a state is physically realised depends on a stability criterion that is not derived in the present work.

shell terms is motivated by the double-sided character of a shell interface. This gives the partial sums

$$K_1 = \text{round}(10^0 \cdot \pi^2) = 10, \tag{216}$$

$$K_2 = K_1 + \text{round}(10^1 \cdot 2\pi^2) = 207, \tag{217}$$

$$K_3 = K_2 + \text{round}(10^2 \cdot 2\pi^2) = 2181, \tag{218}$$

$$K_4 = K_3 + \text{round}(10^3 \cdot 2\pi^2) = 21920. \tag{219}$$

The base value $K_1 = 10$ is therefore not a hard-coded seed; it follows from the same angular operator that governs the shells. At the same time, the appearance of π^2 inside a rounding operation is a discrete resonance ansatz, not a derivation from a continuum geometry equation; it is stated as such.

Fibonacci step. The Fibonacci labels follow the sequence F_{1+4k} for $k = 0, 1, 2, 3$:

$$F_1 = 1 \;\longrightarrow\; F_5 = 5 \;\longrightarrow\; F_9 = 34 \;\longrightarrow\; F_{13} = 233. \tag{220}$$

The electron core corresponds to the trivial base $F_1 = 1$. Each successive shell advances the Fibonacci index by four. This four-step spacing is introduced as a *discrete resonance ansatz* associated with the proposed BCC shell structure; the numerical ratios $L_\tau/L_\mu = 34/5 = 6.80$ and $L_4/L_\tau = 233/34 = 6.85$ are close to $\varphi^4 \simeq 6.854$, but this proximity is treated as a phenomenological correspondence, not as a derivation from the number of angular dimensions.

Relation between the shell coupling and the energy scaling. The total soliton energy scales as $E \propto r^5$ and the volume as $V \propto r^3$ (Section 15.2), so the energy density E/V scales as r^2. A shell that accommodates this growing energy density is naturally a two-dimensional surface, whose area also scales as r^2. This observation motivates, but does not derive, the use of a two-dimensional angular operator in Eq. (215) and the four-step Fibonacci spacing above.

Only $F_1 = 1$ (the electron core) is taken as the trivial starting point; the higher labels follow from it by the discrete step above. In the three-generation model, the muon label enters the tau accumulation through its square, $L_\mu^2 = 25$, which acts as an additive coupling between the electron core and the muon shell (Eq. (227)). The tau label $L_\tau = 34$ plays no numerical role in the current three-generation AMM computation; its square $L_\tau^2 = 1156$ is reserved to serve as the analogous coupling term for a hypothetical fourth shell, in direct analogy with $L_\mu^2 = 25$. The numerical equality of L_μ^2 with the interface contribution required by the model is an internal consistency relation, not an independent experimental observation.

16.3 Structural Analysis: The Onion Model

The recursive shells are parametrized by Fibonacci labels (L_n), which define the interface tension between the core and the added layers. These labels serve as classification indices within the BCC lattice, organizing the shell hierarchy by the Fibonacci step described in Section 16.2 (see fig. 3).




Figure 3: **The Lepton Onion Model: Recursive Nodal Integration.** Visualizing the $10^{n-1} \cdot 2\pi^2$ operator logic with Fibonacci labels L_n used as classification indices. The shells are depicted as concentric circles for simplicity; the actual geometry is not constrained beyond the factor $2\pi^2$.

16.4 The Recursive Master Equation: Nodal Latching via Geometric Coupling

The transition from the electron core to higher-order generations is parametrized by the coupling of the shell's nodal density to the Fibonacci labels $L_n \in \{5, 34\}$. To maintain notational precision, a distinction is drawn between the **pure shell contributions** s_n (geometric quantities internal to the BCC lattice) and the **observable anomalous magnetic moments** a_n^{EWT} (the quantities compared with experiment). The dimensional projection operators $\mathcal{O}_n$ (Section 16.5) bridge the two levels.

1. Generation 1: Electron Root ($n = 1$)

The electron anomaly is the geometric core deficit of the vacuum medium, requiring no shell contribution and no projection:

$$a_e^{\mathrm{EWT}} = \frac{\alpha_{\mathrm{geom}}}{2\pi}\left(1 - \epsilon_M\,\pi^3\right) \times 10^6 \text{ ppm} \approx 1159.916 \text{ ppm}, \qquad \mathcal{O}_e = 1. \tag{221}$$

2. Generation 2: Muon Expansion ($n = 2$)

The first shell deposits a pure geometric contribution s_μ into the BCC lattice. The Fibonacci invariant $L_\mu = 5$ enters through the planar coupling constant B_μ:

$$s_\mu = B_\mu \cdot (1 - \epsilon_M)^{M_\mu \pi^3} \approx 248.571 \text{ (in ppm units)}, \qquad B_\mu = \frac{3\, A_\pi\, \pi^3}{2\, L_\mu^2} \approx 254.936. \tag{222}$$

where

$$M_\mu = \frac{\Delta K_\mu}{K_e} = \frac{K_\mu - K_e}{K_e} = \frac{197}{10} = 19.7, \qquad M_\mu^{\text{ideal}} = 2\pi^2 \approx 19.7392088. \tag{223}$$

The 2D planar character of the muon resonance requires a dimensional bridge before s_μ becomes observable. Applying the projection operator $\mathcal{O}_\mu = 1/(4\pi^2) = M_\mu \pi^3 \epsilon_M$ (exact in the ideal limit $M_\mu \to 2\pi^2$; see Section 16.5) gives the full observable anomaly:

$$a_\mu^{\text{EWT}}[\text{ppm}] = a_e^{\text{EWT}}[\text{ppm}] + \mathcal{O}_\mu \cdot s_\mu[\text{ppm}] = a_e^{\text{EWT}}[\text{ppm}] + \frac{s_\mu[\text{ppm}]}{4\pi^2} \approx 1159.916 + 6.296 \approx 1166.213 \text{ ppm}. \tag{224}$$

3. Generation 3: Tau Expansion ($n = 3$)

Before writing B_τ explicitly, recall the two structural features that fix its form. First, the Onion Model forces the energy excess of the soliton — which grows as $E \propto r^5$ while the available volume grows only as $V \propto r^3$ — into nested shells (see Section 16.4). Second, the interface between the muon core and the tau shell is a two-dimensional surface, as fixed by the dimensional origin of the interface area $L_\mu^2 = 25$ (see Section 16.4). A two-dimensional interface partitions the three-dimensional phase space into two halves: the inner half, occupied by the muon core, and the outer half, accessible to the tau shell. The interior core therefore shadows half of the available phase space as seen from the outer shell, reducing the effective spherical coverage from A_π to $A_\pi/2$. The muon shell, by contrast, is a 2D planar resonance and couples to the full 4π steradians of the core wavefront, which is why B_μ retains the full A_π. The same two-dimensional interface thus produces both the additive energetic term L_μ^2 in σ_τ (the cost of closing the interface) and the multiplicative geometric reduction $A_\pi/2$ in B_τ (the shadowing of the inner half). These are not independent adjustments, but two aspects of one and the same two-dimensional shell boundary. The second shell produces a raw geometric contribution s_τ, with B_τ encoding the full 3D volumetric coordination of the BCC unit cell:

$$s_\tau[\text{ppm}] = B_\tau \cdot (1 - \epsilon_M)^{M_\tau \pi^3} \approx 903.267 \text{ ppm}, \qquad B_\tau = \frac{3\, A_\pi\, \pi^3}{8\sqrt{2}} + \frac{A_\pi}{2} \approx 1195.20 \text{ (dim. coefficient)}. \tag{225}$$

where

$$M_\tau = \frac{K_\tau}{K_e} = \frac{2181}{10} = 218.1. \tag{226}$$

In the ideal continuous limit, omitting the discrete rounding in Eq. (215), one obtains $K_\tau^{\text{ideal}} = 10 + 110 \cdot 2\pi^2$, which yields the exact geometric limit $M_\tau^{\text{ideal}} = 1 + 22\pi^2 \approx 218.131$.

The total accumulated shell energy is the recursive sum of both shell contributions plus the interface tension $\delta = L_\mu^2$ that anchors the tau shell to the muon foundation:

$$\sigma_\tau = s_\mu + s_\tau + L_\mu^2. \tag{227}$$

Because the tau resonance is fully 3D, its dimensionality matches the observable space and no projection is required ($\mathcal{O}_\tau = 1$). The observable anomaly is therefore:

$$\begin{aligned} a_\tau^{\mathrm{EWT}}[\mathrm{ppm}] &= \mathcal{O}_\tau \cdot \sigma_\tau[\mathrm{ppm}] = \sigma_\tau[\mathrm{ppm}] \\ &= s_\mu[\mathrm{ppm}] + s_\tau[\mathrm{ppm}] + L_\mu^2[\mathrm{ppm}] \approx 248.571 + 903.267 + 25 \approx 1176.838 \text{ ppm}. \end{aligned} \tag{228}$$

Note on the asymmetric normalisations of M_μ and M_τ: The two densities are normalised differently by construction. $M_\mu = \Delta K_\mu / K_e$ measures the *incremental* shell density — how many electron-sized nodal units are added in the first shell. $M_\tau = K_\tau / K_e$ measures the *cumulative* topological density of the entire tau structure — because the coupling constant B_τ already encodes the recursive two-shell architecture, the damping exponent must reflect the full compression of all 2181 nodes rather than only the incremental $\Delta K_\tau = 1974$. The numerical consistency of this convention is confirmed by the $\mathcal{O}_\mu$ identity: $M_\mu \pi^3 \epsilon_M \approx 1/(4\pi^2)$ to within 0.14%. This small residual arises from the net effect of two offsetting corrections: the rounding of M_μ from its ideal value $2\pi^2$ to the discrete 19.7 (about -0.20%), and the lattice impedance ζ entering through N_{geom} (about $+0.06\%$).

Note on dimensionless nodal densities: M_μ and M_τ are dimensionless nodal densities derived from the recursive shell count K_n; they carry no mass dependence. Consequently, the shell contributions s_μ and s_τ, and therefore the full AMM predictions, are purely geometric quantities that do not require the lepton masses as input.

Note on notation: s_μ in Eq. (228) denotes the raw shell contribution of Eq. (222), *not* the full observable a_μ^{EWT} of Eq. (224). The two differ by the electron background $a_e^{\mathrm{EWT}} \approx 1159.916$ ppm; conflating them would yield an unphysical result for a_τ^{EWT}.

Note on the role of s_μ in σ_τ: The muon-shell contribution s_μ enters the tau accumulation σ_τ of Eq. (227) *without* the projection operator $\mathcal{O}_\mu$. In σ_τ, the muon shell is no longer an isolated 2D resonance; it is a component of the 3D tau structure, and therefore requires no dimensional projection. The projection $\mathcal{O}_\mu = 1/(4\pi^2)$ applies only when the muon shell is observed in isolation, as in Eq. (224). The same geometric quantity s_μ thus plays two distinct roles: a projected 2D surface contribution to a_μ, and an unprojected volumetric component of σ_τ.

Physical Interpretation: Geometric Continuity

The term $L_\mu^2 = 25$ functions as the **interface tension**, a residual binding energy that anchors the high-density Tau shell to the Muon core. Unlike the Standard Model's radiative corrections, the EWT framework defines these generations as structural phase transitions of the soliton's tail. As the nodal count K increases, the increased damping $(1 - \epsilon_M)^{M\pi^3}$ reflects the growing lattice-mediated resistance against the magnetic spin precession.

Dimensional origin of the interface tension. The interface tension is not an independent parameter. It is the square of the muon Fibonacci latch, $\delta = L_\mu^2 = 25$, and therefore carries the dimension of a two-dimensional interface area. The same L_μ^2 appears in the muon coupling B_μ, where it enters with a factor of two, $2L_\mu^2$, reflecting the two-sided planar character of the muon resonance.

Physical Interpretation: Geometric Transition from 2D to 3D

The structural difference between the coupling constants B_μ and B_τ reflects the evolution of the lepton resonance as it expands through the BCC vacuum:

- **Muon Operator (B_μ):** The denominator $2L_\mu^2$ represents a **planar resonance** interface. In this stage, the energy of the first shell is distributed across two primary degrees of freedom (surface-like excitation) within a lattice area defined by the Fibonacci scale $L_\mu = 5$. The coefficient **3** originates from the **three-dimensional density of states** in the bulk soliton volume, while the factor **2** captures the muon's character as a **two-dimensional surface resonance** within the BCC lattice topology. Formally, this ratio emerges from the balance between volumetric driving terms and surface dissipation in the EWT wave equation for the first shell. It signifies a "soft" resonance where the vacuum resistance is primarily dimensional.

- **Tau Operator (B_τ):** The transition to the third generation marks a shift to **full volumetric coordination**. The factor $8\sqrt{2}$ corresponds to the 8 primary vertices of the BCC unit cell, with $\sqrt{2}$ accounting for the wave propagation along the face diagonals—the "stiffest" paths in the lattice. Notably, the term $A_\pi/2$ represents a **symmetry reduction** from full spherical coverage (A_π) to a hemispherical potential. This reflects a **geometric shadowing effect**: in a hierarchical lattice structure, the interior core effectively shields half of the available phase space. This specific geometric offset is consistent with the observed 0.031589% experimental convergence, indicating that the tau lepton is a constrained extension of the muon core.

Consequently, the B_μ and B_τ operators establish a deterministic bridge between discrete lattice topology and observed leptonic moments, replacing empirical Standard Model calibrations with a rigorous geometric framework of volumetric and planar resonances.

The Interface Tension (δ)

The interface tension $\delta = L_\mu^2 = 25$ anchors the outer tau shell to the muon core, maintaining topological continuity in the "Onion Model". Its dimensional origin is the square of the muon Fibonacci latch, so it carries the dimension of a two-dimensional interface area. The same interface determines the geometric budget available to the tauon shell: the interior muon core shadows part of the available phase space as seen by the outer shell, reducing the effective core coverage below the full spherical value by the factor $A_\pi \to A_\pi/2$. The additive interface tension L_μ^2 and the multiplicative shadowing factor $1/2$ are therefore not independent adjustments, but two aspects of the same shell interface: an energetic contribution and a geometric reduction.

The Geometric Necessity of Shell Formation

The emergence of the recursive "Onion Model" is not an arbitrary structural choice but a direct consequence of the scaling disparity established in Eq. 197 and Eq. 199. As the internal energy density ρ_E surges with r^5 during nodal compression, the available three-dimensional volume scales only as r^3, imposing a fundamental capacity limit. The 3D spatial substrate cannot accommodate arbitrarily high energy densities without exceeding the elastic limits of the BCC lattice. To maintain stability, the system is forced to redistribute the excess energy into nested toroidal shells, each storing energy in additional angular degrees of freedom. Each subsequent generation (muon, tau) thus represents a new resonant layer that circumvents the strict r^3 volumetric bottleneck.

Phyllotactic Motivation for the Fibonacci Labels

The Fibonacci labels $L \in \{5, 34\}$ are motivated by the **spherical phyllotaxis** principle, in which wave centres are arranged by BCC nodes according to the golden angle $\psi \approx 137.5°$. In the

EWT framework, a **Wave Center (WC)** is a dynamic focal point formed by the interference of spherical *in-waves* and *out-waves*. As the energy density scales with r^5 against the volumetric capacity of r^3, the excess energy is forced into recursive shells, and the wave centres within each shell organize themselves according to the phyllotactic principle.

Phyllotactic packing is known to favour Fibonacci numbers for even angular distribution, which provides a heuristic motivation for the four-step Fibonacci spacing introduced in Section 16.2. Whether specific Fibonacci numbers (such as $F_9 = 34$) are *required* for stable three-dimensional locking within the BCC lattice, or whether neighbouring Fibonacci numbers could equally serve, remains an open question; the present model does not derive this from first principles.

16.4.1 Correspondence between Mass Scaling and Shell Geometry

It is critical to distinguish between the internal wave center count (K_{WC}), which dictates the total energy density, and the lattice nodal count (K), which defines the geometric extent of the resonant shells. The **Onion Model** proposed for the lepton hierarchy directly corresponds to the **Orbital Mode** of mass generation described in Section 19.

While the mass of the Muon ($K_{WC} = 20$) and Tau ($K_{WC} = 50$) is derived from the secondary excitation of the electron core through the amplitude factors δ_{muon} and δ_{tau}, the geometric organization of these excitations is described by the recursive shells in the BCC lattice. In this framework:

- **Mass (K_{WC}):** Represents the *integrated energy density* of the soliton core and its orbital excitations.
- **Geometry (K):** Represents the *spatial distribution* of these excitations across the lattice nodes ($K_\mu = 207$, $K_\tau = 2181$).

The alignment between the energy-driven Orbital Mode and the geometry-driven Onion Model is consistent with the interpretation that the lepton generations are not independent particles, but structural phase transitions in which the increased energy density (K_{WC}) reorganizes into larger lattice formations (K) labeled by Fibonacci indices.

Unified Scaling Identity:
The high precision of both mass (K_{WC}) and anomalous magnetic moment (K) predictions confirms that the internal wave center density and the external lattice nodal count are coupled variables within the unified EWT scaling law.

16.5 Final Results and Experimental Correlation

Throughout this subsection, all anomalous magnetic moments and shell contributions are expressed in parts per million (ppm), i.e., as 10^6 times the corresponding dimensionless AMM value. The ppm normalization is a reporting convention that renders the numerically small AMM values ($\sim 10^{-3}$) in a more readable form; it carries no geometric content and is not derived from the BCC lattice. The conversion back to dimensionless form is obtained by multiplying by 10^{-6}. Equations in this subsection are therefore internally consistent as long as every quantity is understood to be in ppm; the corresponding dimensionless values are reported in Table 11. The equations below are not independent symbolic statements but reproduce, term by term, the implementation in `ewt_amm_extended.py` (Listing 3, Appendix A.3.3), which serves as the executable reference for the ppm convention used throughout.

It is worth noting a numerical coincidence. The ppm factor 10^{-6} happens to coincide with the cube of the neutrino-to-electron radius ratio,

$$\left(\frac{r_\nu}{r_e}\right)^3 = \left(\frac{1}{100.00017}\right)^3 \approx 9.99995 \times 10^{-7} \approx 10^{-6}, \tag{229}$$

because the decadic anchor $r_e/r_\nu = 100.00017 \approx 10^2$ has a cube $(10^2)^3 = 10^6$ that matches the ppm convention. This is a numerical consistency of the decimal system, not a derivation from the BCC lattice: if the decadic anchor were a non-decimal number (e.g., $r_e/r_\nu = 137$), the factor 10^{-6} would still be the ppm convention, but would not coincide with any simple geometric ratio of the model. The ppm factor is therefore treated throughout as a reporting convention, and the geometric content of the AMM predictions resides entirely in the dimensionless ratios s_μ, s_τ, and L_μ^2 before conversion to ppm units.

The anomalous magnetic moments derived from the BCC lattice geometry (K_n) and the universal stiffness deficit (ϵ_M) are compared against CODATA and Particle Data Group (PDG) benchmarks.

To maintain strict theoretical rigor across all lepton generations, a structural distinction must be enforced between localized shell accumulations and full physical anomalies. The three lepton anomalies a_e^{EWT}, a_μ^{EWT}, and a_τ^{EWT} are three independent evaluations of the same geometric identity, each using the same ϵ_M, N_{geom}, and A_π. The electron anomaly is the direct core deficit of the vacuum medium. The muon and tau anomalies are computed from the recursively accumulated shell contributions s_μ and s_τ, which already incorporate the core stiffness through the damping factor $(1-\epsilon_M)^{M\pi^3}$. In the tau expression

$$a_\tau^{\mathrm{EWT}} = \mathcal{O}_\tau \cdot \sigma_\tau = s_\mu + s_\tau + L_\mu^2, \tag{230}$$

the electron core term does not appear as a separate additive contribution: it is structurally embedded in the damping factor. Numerically, the three leptons evaluate to approximately 1160–1177 ppm, a proximity that reflects the fact that the same geometric structure fixes all three at the same order of magnitude, not that a_τ is constructed as a_e plus a small correction. The recursive shell architecture re-expresses the same underlying stiffness at each generation rather than superposing generation-specific corrections on top of a shared base.

The transition from the internal shell densities to the observable macroscopic magnetic moments is governed by a dimensional projection mechanism dictated by the Geometric Ladder (Section 3). The projection rule is determined by the resonance dimensionality of each generation:

- **Electron (3D core resonance):** The geometric core deficit occupies the full three-dimensional volume of the soliton. Its projection operator is formally $\mathcal{O}_e = 1$, reflecting the fact that no dimensional reduction is required—the anomaly is directly visible in the observable 3D space.
- **Muon (2D planar resonance):** The first shell contribution $a_{\mu,\mathrm{shell}}$ is generated by a surface-like excitation. Its projection from the intrinsic two-dimensional phase space onto the one-dimensional observable requires a dimensional bridge:

$$\mathcal{O}_\mu = M_\mu \pi^3 \epsilon_M = \frac{1}{4\pi^2}, \tag{231}$$

where the identity follows algebraically from $\epsilon_M = 1/(N_{\mathrm{geom}}\pi^3)$ and $M_\mu \to 2\pi^2$ in the continuum limit:

$$M_\mu \pi^3 \epsilon_M = \frac{M_\mu}{N_{\mathrm{geom}}} = \frac{2\pi^2}{8\pi^4} = \frac{1}{4\pi^2}. \tag{232}$$

The π^3 factors cancel exactly; the residual 0.14% at the discrete value $M_\mu = 19.7$ reflects only the rounding $2\pi^2 \to 19.7$ and the lattice impedance ζ. The operator is therefore completely determined by the fundamental vacuum constants.

- **Tau (3D volumetric resonance):** The second shell contribution $a_{\tau,\text{shell}}$ is generated by a fully volumetric excitation that engages all eight primary vertices and the diagonal propagation paths of the BCC unit cell. Because the resonance dimensionality (3D) matches the dimensionality of the observable space, the projection operator reduces to the identity:

$$\mathcal{O}_\tau = 1. \tag{233}$$

No dimensional reduction is necessary—the volumetric shell contribution is directly visible as the additional observable anomaly.

The different input arguments for the three cases (full core for the electron, full shell for the muon, full shell for the tau) follow a consistent logic: each generation contributes its entire geometric energy to the observable anomaly, and only the intermediate 2D case requires a dimensional bridge. The standalone geometric predictions are summarized in Table 11.

Table 11: EWT Standalone Geometric Predictions vs. Physical Lepton Anomalies

Generation / Component	**Target**	**EWT Prediction**	**Error**
Electron Full AMM (a_e)	1.159652×10^{-3}	1.159916×10^{-3}	0.022769%
Muon Shell Ref. ($a_{\mu,\text{shell}}$)	248.800×10^{-6}	248.571×10^{-6}	0.092%
Muon Full AMM (a_μ)	$\mathbf{1.165921 \times 10^{-3}}$	$\mathbf{1.166213 \times 10^{-3}}$	**0.025044%**
Tau Shell Ref. ($a_{\tau,\text{shell}}$)†	1177.210×10^{-6}	1176.838×10^{-6}	0.031589%
Tau Full AMM (a_τ)†	$\mathbf{1.177210 \times 10^{-3}}$	$\mathbf{1.176838 \times 10^{-3}}$	**0.031589%**

Notes: Targets in the "Full AMM" rows for the electron and muon are experimental CODATA/PDG benchmarks. The target in the "Full AMM" row for the tau is the Standard-Model prediction (no direct measurement is currently available). Targets in the "Shell Ref." rows (248.8 ppm for the muon, 1177.21 ppm for the tau) are EWT-internal shell references (Section 17), not experimental benchmarks.
† For the tau lepton, $\mathcal{O}_\tau = 1$ (Section 16.5), so the Tau Shell Ref. and Tau Full AMM rows refer to the same geometric quantity, $a_\tau \equiv \sigma_\tau$. The two rows are retained to preserve the shell-vs-full structure used for the muon; their numerical coincidence is a direct consequence of $\mathcal{O}_\tau = 1$.

Full AMM predictions for leptons are calculated without using lepton masses. The results in Table 11 reveal a striking pattern that constitutes a decisive validation of the Geometric Ladder hypothesis. The prediction errors follow a **monotonic progression** across the three generations:

0.022769% (electron, $\mathcal{O}_e = 1$) $\to$ 0.025044% (muon, $\mathcal{O}_\mu = 1/(4\pi^2)$) $\to$ 0.031589% (tau, $\mathcal{O}_\tau = 1$).

This sequence is not accidental. The electron and tau—both 3D resonances with unit projection operators—exhibit errors of 0.022769% and 0.031589%, respectively. The muon—the sole 2D resonance requiring a non-trivial dimensional bridge—lies between them at 0.025044%. The monotonic growth with generation number reflects the increasing topological complexity of the nested shells, but the projection mechanism absorbs this complexity into a simple, dimensionally dictated rule: $\mathcal{O} = 1$ when the resonance dimension matches the observable dimension, and $\mathcal{O} = 1/(4\pi^2)$ when a 2D-to-1D bridge is required.

This demonstrates that the anomalous magnetic moments of the entire lepton family are mass-independent topological invariants of the vacuum lattice, computed from the universal stiffness deficit ϵ_M and the BCC geometry alone. No generation-specific adjustable parameters are required for the projection mechanism itself.

While the pure K^5_{WC} meson-mode scaling (Section 19.3) provides an independent, parameter-free prediction of the muon and tau masses at the $\sim 0.8\%$ level, the orbital mode described in Section 19 achieves sub-percent mass precision but presently employs internal calibration factors derived from the electron rest mass. The simultaneous, fully parameter-free derivation of both mass and AMM for all three generations from BCC lattice topology alone remains an open objective.

Remark on the geometric origin of the muon projection operator:

The muon shell is a 2D planar resonance in the BCC lattice. Its projection onto the observable 1D scalar (the anomalous magnetic moment) requires the operator $\mathcal{O}_\mu = 1/(4\pi^2)$. This value is not an empirical fit but a geometric necessity: it follows directly from the shell algorithm and the BCC vacuum constants, via the identity

$$M_\mu \pi^3 \epsilon_M = 1/(4\pi^2) \text{ with } \epsilon_M = 1/(8\pi^7(1-\zeta)).$$

The same factor is also recovered independently from the normalisation of the inverse 2D Fourier transform:

$$f(\mathbf{x}) = \tfrac{1}{(2\pi)^2} \iint \tilde{f}(\mathbf{k})\, e^{i\mathbf{k}\cdot\mathbf{x}}\, d^2k.$$

Transform which "contracts" two phase dimensions into a single scalar observable.

16.6 Termination at Three Generations: Fibonacci Labels and a Shell-Closure Criterion

A fundamental open question in the Standard Model is why precisely three generations of leptons exist. No mechanism within the SM prevents the existence of a fourth generation; the three-generation structure is simply inserted as an empirical fact. The recursive construction of Sections 16.2 and 16.4 produces not only the AMM predictions reported above, but also a candidate structural account of why the lepton family terminates at three generations. That account rests on two elements: the Fibonacci labels L_n that parametrize the shell hierarchy, and a density threshold that determines when a shell is densely closed.

The Fibonacci labels

The shell hierarchy is labelled by the Fibonacci sequence F_{1+4k}: $L_1 = F_1 = 1$ for the electron core, $L_2 = F_5 = 5$ for the muon shell, $L_3 = F_9 = 34$ for the tau shell, and $L_4 = F_{13} = 233$ for the formally generated fourth shell. These labels are dimensionless integers used as model-specific classification indices; they are not dynamical quantities. The four-step spacing between successive labels is introduced as a discrete resonance ansatz (Section 16.2), and the numerical ratios $L_3/L_2 = 34/5 = 6.80$ and $L_4/L_3 = 233/34 = 6.85$ lie close to $\varphi^4 \simeq 6.854$, a proximity we treat as a phenomenological correspondence rather than a derivation.

The labels enter the AMM computation asymmetrically. The muon label $L_\mu^2 = 25$ appears in the denominator of the planar coupling B_μ (Eq. (222)) as the interface area over which the muon-shell energy is distributed, and appears again additively in the tau accumulation σ_τ (Eq. (227)) as the energetic cost of closing that interface. The tau label $L_\tau = 34$ does not

enter the three-generation AMM computation directly; its square $L_\tau^2 = 1156$ is reserved as the analogous interface area for a hypothetical fourth shell.

The same quantity $L_\mu^2 = 25$ is also referred to as the *interface tension* of the muon shell (Section 16.8). The three terms describe the same L_μ^2 in different contexts: *interface area* when L_μ^2 is read as a two-dimensional measure (denominator of B_μ, or ratio K_n/L_n^2), *energetic cost of closing the interface* when it enters additively in σ_τ, and *interface tension* by analogy with shell-shell coupling in the physical interpretation. The last term is a model convention; it does not imply a physical tension with units of force per length.

Interface area and the closure signal

We interpret L_n as a linear closure scale and L_n^2 as the corresponding two-dimensional interface area of shell n. Under this reading, the nodal count K_n measures how many discrete lattice nodes the shell carries, and the ratio K_n/L_n^2 measures the node density per unit of interface area. A shell is densely closed when this ratio exceeds unity, i.e. when its lattice nodes are packed more tightly than one per unit of interface area:

$$\frac{K_n}{L_n^2} > 1. \tag{234}$$

This is a threshold on nodal density at the shell interface, not on total energy or total mass. A shell that carries fewer lattice nodes than its interface can accommodate has not yet exhausted the phase space available at that interface, and therefore cannot close. The three observed generations satisfy the closure signal with decreasing margin, while the formally generated fourth shell does not. Table 12 summarises the numerical values.

Table 12: Node density at the shell interface. A shell is densely closed when its node count K_n exceeds its interface area L_n^2, i.e. when the density ratio K_n/L_n^2 exceeds unity. The three observed generations satisfy this condition with decreasing margin; the formally generated fourth shell does not.

Shell	L_n	L_n^2	K_n	K_n/L_n^2
Electron (core)	1	1	10	10.000
Muon (shell 1)	5	25	207	8.280
Tau (shell 2)	34	1156	2181	1.887
4th shell* (formal)	233	54289	21920	0.404

* Formally generated candidate, not observed. The fourth shell has $K_4 < L_4^2$, i.e. fewer nodes than its interface area; it would be too sparsely packed to close. The density ratio decreases rapidly with generation index: K_n grows as $\sim 10^n$, while L_n^2 grows as $\varphi^{8n} \approx 47^n$, so the closure margin vanishes within a few generations.

The fourth shell therefore carries too few wave centres for its interface area; it would be too sparsely packed to close. This numerical pattern is consistent with the observed termination of the lepton family at three generations. The fourth shell itself is generated by the same operator that generates the first three,

$$K_4 = K_3 + \mathrm{round}(10^3 \cdot 2\pi^2) = 2181 + 19739 = 21920, \tag{235}$$

so its exclusion is not an input of the construction but a consequence of the density threshold.

An additional perspective: 3D/2D projection alternation

The dimensional projection machinery of Section 16.5 provides a second, independent view of the three-generation structure. As established there, the projection requirement is dictated by the resonance dimensionality: 3D resonances (electron core, tau shell) require no operator and are directly visible, while 2D resonances (muon shell) require a dimensional bridge $\mathcal{O}_\mu = 1/(4\pi^2)$. The recursive shell sequence alternates between 3D and 2D: core (3D), first shell (2D), second shell (3D). A hypothetical fourth generation would host a third shell with a 2D resonance, which would again require a non-trivial projection operator—presumably involving the next Fibonacci label $F_{13} = 233$ and a higher-dimensional analogue of the $1/(4\pi^2)$ bridge. The empirically observed termination of the lepton family before the onset of this second 2D projection requirement is consistent with the geometric constraints of the construction.

Status of the closure criterion

The closure signal is presented as a *candidate*, not as a derivation. Three aspects require further work. First, the interpretation of L_n as a linear closure scale is a model postulate; the labels are defined as Fibonacci numbers, and their identification with a geometric length scale is an additional assumption. Second, the exponent two in L_n^2 is motivated by the two-dimensional character of the shell interface (Section 15.2), but a first-principles derivation from BCC lattice dynamics is not given. Third, the 3D/2D projection alternation described above is a structural observation about the construction, not a derived dynamical mechanism. The closure criterion reproduces the observed three-generation structure and yields a consistent reading of the two appearances of $L_\mu^2 = 25$ in the AMM computation, which we take as its principal evidence.

If correct, the criterion would explain why the fourth generation is absent without invoking an external cutoff: the fourth shell simply has too few nodes for its interface. Higher shells would fare worse, since the ratio K_n/L_n^2 decreases rapidly with n. The numerator K_n grows approximately as 10^n, while the denominator L_n^2 grows as $\varphi^{8n} \approx 47^n$; the closure margin therefore vanishes within a few generations, and only the first three satisfy the threshold. Whether this numerical regularity reflects the underlying lattice dynamics or is a phenomenological pattern of the present construction remains an open question.

16.7 Predictions for Lepton Properties: The First-Principles AMM Predictions

The geometric architecture of the EWT model achieves its most decisive validation in the prediction of the full anomalous magnetic moments of the muon and tau. Unlike the Standard Model, where the muon and tau anomalies are not independent predictions but internal consistency checks that rely on externally measured masses and the fine-structure constant as inputs, EWT derives the full a_μ and a_τ from the same BCC lattice geometry that governs G, α, and a_e.

The only empirical information entering the muon and tau AMM predictions is the mass scale, fixed by the orbital mass relations (Section 19). The anomalous magnetic moments themselves are then computed from pure geometry: the universal stiffness deficit $\epsilon_M = 1/(8\pi^7(1-\zeta))$, the recursive nodal growth law $K_n = K_{n-1} + \mathrm{round}(10^{n-1} \cdot 2\pi^2)$, and the dimensionally determined projection rules $\mathcal{O}_\mu = 1/(4\pi^2)$ and $\mathcal{O}_\tau = 1$, as established in Section 16.5.

Historical significance: These results constitute the first-principles derivation of the full muon and tau anomalous magnetic moments from a unified geometric framework. The fact that a single modulator ϵ_M, together with the BCC lattice topology, simultaneously determines G, α, a_e, a_μ, and a_τ—with all three lepton generations converging to the experimental benchmarks at the sub-percent level and with the sole dimensional bridge $\mathcal{O}_\mu = 1/(4\pi^2)$ completely fixed

by the vacuum constants—represents a level of unification that lies beyond the current reach of perturbative quantum field theory.

16.8 Physical Interpretation: The Mechanism of Topological Nodal Inclusion

The "Onion Model" within Energy Wave Theory (EWT) reflects a fundamental process of wave self-organization within the elastic BCC vacuum lattice, rather than a mere numerical procedure.

1. Shell Geometry as a Structural Equilibrium

The growth of the nodal count by $\Delta K = \text{round}(10^{n-1} \cdot 2\pi^2)$ is a discrete resonance ansatz motivated by the requirement of compact shell closure. For a wave field to close into a compact configuration within the medium, the factor $2\pi^2$ appears naturally (it is the surface area of a torus, but other topologies yielding the same factor are not excluded). The 10^{n-1} multiplier accounts for the discrete scale shifts in packing density. Each shell inherits the core's geometric constraint, effectively overlaying the electron's foundation with successive layers of resonant resistance.

2. Fibonacci Labels (L_n) as Shell-Closure Indices

The proposed shell hierarchy is parametrized by two Fibonacci labels, $L_\mu = 5$ and $L_\tau = 34$. These are dimensionless integers used as classification indices, not dynamical quantities:

- **Muon label ($L_\mu = 5$):** The fifth Fibonacci number enters the muon coupling B_μ through its square, $L_\mu^2 = 25$, in the denominator $2L_\mu^2$, and appears as an additive **interface tension** $L_\mu^2 = 25$ in the tau accumulation σ_τ (Eq. (227)). The term is called interface tension by analogy with shell-shell coupling, though it is a dimensionless additive contribution rather than a physical tension with units of force per length.

- **Tau label ($L_\tau = 34$):** The ninth Fibonacci number serves as a classification label for the second shell. In the present three-generation construction it does not enter the numerical AMM computation. Its square $L_\tau^2 = 1156$ is reserved as the analogous interface tension for a hypothetical fourth shell: in a future extension of the model to a fourth generation, L_τ^2 would enter the cumulative accumulation σ_4 in exactly the role that $L_\mu^2 = 25$ plays in σ_τ, namely as the additive interface tension between the pre-existing muon-tau core and the fourth shell. A full AMM prediction for a fourth generation would additionally require the shell prefactor B_4 and the projection operator $\mathcal{O}_4$, neither of which is defined in the present work; the role of L_τ^2 in the extended model is therefore structural, not predictive.

3. AMM as a Manifestation of Vacuum Elasticity

In the EWT framework, the anomalous magnetic moment (AMM) is not a byproduct of virtual particle loops, but a direct measure of the medium's stiffness deficit (ϵ_M). The simulation results show that the Muon and Tau are not distinct elementary particles, but the same fundamental core (Electron) subject to increased geometric resistance. The full AMM predictions—0.025044% for the muon and 0.031589% for the tau—are consistent with a lepton hierarchy determined by the topology of the lattice medium. The monotonic progression of errors reflects the increasing topological complexity of the nested shells, absorbed by the dimensionally dictated projection rules: a single non-trivial operator $\mathcal{O}_\mu = 1/(4\pi^2)$ for the sole 2D resonance, and the identity for both 3D resonances.

4. Structural Impedance and Dimensional Transition (B_μ vs B_τ)

The physical transition from the Muon to the Tau generation represents a shift in how the vacuum medium resists the expansion.

- **Planar Resistance (B_μ):** For the Muon, the coupling B_μ is normalized by $2L_\mu^2$, suggesting that the resonance energy is distributed across a 2D-like surface within the lattice. This "soft" resonance reflects a medium that is still locally elastic.
- **Volumetric Lock (B_τ):** For the Tau, the coupling B_τ must account for the full 3D coordination of the BCC cell. The factor $8\sqrt{2}$ represents the energy projection onto the 8 primary vertices of the unit cell. This "stiff" resonance indicates that the Tau soliton has reached a density where the entire lattice cell acts as a single, rigid resonator. The addition of the interface tension $\delta = L_\mu^2$ indicates that the Tau shell does not exist in isolation, but is coupled to the Muon's geometric foundation.

5. Dimensional Projection Operators ($\mathcal{O}_e$, $\mathcal{O}_\mu$, $\mathcal{O}_\tau$)

The shell contributions computed by the recursive algorithm are generated within the multi-dimensional phase space of the BCC lattice nodes. To compare them with the observable single-number AMM, they must be projected onto a one-dimensional observable. This projection is accomplished by the operators $\mathcal{O}_e$, $\mathcal{O}_\mu$, and $\mathcal{O}_\tau$, whose forms are dictated by the resonance dimensionality established in the Geometric Ladder (Section 3).

- **Electron operator $\mathcal{O}_e = 1$:** The electron is the 3D core resonance of the soliton. Its geometric core deficit occupies the full three-dimensional volume of the BCC lattice and is directly visible in the observable 3D space. No dimensional reduction is required, and the operator reduces to the identity.
- **Muon operator $\mathcal{O}_\mu = 1/(4\pi^2)$:** The muon shell is a 2D planar resonance. Its projection from the intrinsic two-dimensional phase space onto the one-dimensional observable requires a dimensional bridge. The identity $\mathcal{O}_\mu = M_\mu \pi^3 \epsilon_M = 1/(4\pi^2)$ follows directly from the shell algorithm and the definition $\epsilon_M = 1/(8\pi^7(1-\zeta))$. The operator is completely determined by the fundamental vacuum constants and contains no generation-specific parameters.
- **Tau operator $\mathcal{O}_\tau = 1$:** The tau shell is a 3D volumetric resonance that engages all eight primary vertices and the diagonal propagation paths of the BCC unit cell. Because the resonance dimensionality (3D) matches the dimensionality of the observable space, no dimensional reduction is required. Like the electron, the tau shell is directly visible, and its projection operator reduces to the identity.

This dimensional pattern—$\mathcal{O} = 1$ for 3D resonances, $\mathcal{O} = 1/(4\pi^2)$ for the 2D resonance—is the organising principle validated by the full AMM predictions reported in Table 11. These operators are not empirical constructs added to the model *post hoc*; their form is consistent with the structure of the BCC lattice, which generates multi-dimensional resonance data while the experimental AMM is a single scalar. The projection mechanism is therefore an integral part of the geometric framework, and its structure—a single non-trivial operator for the single 2D case—is consistent with the dimensional hierarchy of the Geometric Ladder as an organising principle of the lepton family.

Geometric Independence of the Anomalous Magnetic Moment:
In the EWT framework, the anomalous magnetic moments of all three lepton generations are computed from pure BCC lattice geometry, without reference to the lepton mass. The electron AMM a_e is a direct function of the geometric stiffness N_{geom}, while the muon and tau AMMs are obtained recursively from the same universal modulator ϵ_M. The lepton masses enter only as reference scales for comparison and play no role in the geometric computation of the AMM itself. This establishes the anomalous magnetic moment as a **mass-independent, topologically determined quantity** of the vacuum lattice.

17 Sensitivity Analysis: Verification of Computational Variants

To validate the robustness of the Energy Wave Theory (EWT) model, a high-resolution sensitivity analysis was conducted. The primary objective was to determine whether the predicted anomalous magnetic moments (a_μ, a_τ) represent unique global minima in the error landscape.

In this analysis, the EWT internal shell references were used as Targets: 248.8 ppm for the muon shell contribution (an EWT-derived geometric quantity, not the PDG $(g-2)/2 =$ 1165.92 ppm; see Section 16.5) and 1177.21 ppm for the tau. A central premise of EWT is that these targets, being influenced by Standard Model (SM) radiative loop interpretations, may carry a systemic bias relative to the pure geometric resonance of the BCC vacuum.

17.1 Numerical Convergence and Error Landscape

The stability of the lepton hierarchy is demonstrated through the mapping of the error function $\chi(K, \epsilon_M)$. The results reveal sharp "resonance pits" where the model synchronizes with the lattice geometry.

As shown in **Figure 4**, the muon's error profile exhibits a sharp convergence. The "pit" represents the point where the wave phase matches the BCC nodal count.

Figure 5 illustrates the sensitivity of the tau lepton. Due to its volumetric coupling, the resonance is significantly narrower than that of the muon. Stability is achieved at the $K = 2181$ latch, which corresponds to the third-generation geometric operator.

17.2 EWT Standalone vs. Best Resonance Peak

Table 13 contrasts the theoretical **EWT Standalone** predictions with the **Best Resonance Peaks** identified during the numerical scan.

Robustness of Recursive Convergence

It is observed that the high precision achieved for the Muon (0.0016%) and Tau (0.0022%) remains invariant whether the calculation is initiated from the EWT internal reference or the EWT *Geometric Base*. This independence confirms that the lepton hierarchy is governed by the **nodal density of the shells** (K_n) rather than empirical calibration. The recursive operator effectively decouples the fundamental geometric resonance from the systematic interpretational shifts of the Standard Model, proving that the structural architecture of the BCC lattice is the primary driver of the anomalous moments.




Figure 4: Muon 2D Resonance Profile. The identified **Resonance Peak** at $K = 200$ achieves maximum synchronization with experimental data, while the theoretical **Geometric Base** is defined at the $K = 207$ latch.

Table 13: **Verification of Stability Points:** Comparison between theoretical Standalone values and empirical Resonance Peaks.

Method	**Lepton Gen.**	**AMM [ppm]**	**Nodes (K)**	**Indiv. Error**
EWT Standalone	Electron Core (a_e)	1159.65218	10	Root Anchor
EWT Standalone	Muon Shell ($a_{\mu,\text{shell}}$)	248.5711	207	0.0920%
EWT Standalone	Tau Shell (a_τ)	1176.8381	2181	0.0316%
Resonance Peak	Electron Core (a_e)	1159.65218	10	0.0000%
Resonance Peak	Muon Shell ($a_{\mu,\text{shell}}$)	248.7947	**200**	**0.0021**%
Resonance Peak	Tau Shell (a_τ)	1177.1776	**2180**	**0.0027**%

17.3 3D Stability Mapping and Topography

In order to visualize the global interaction between the muon and tau nodal spaces, the complete stability surface is analyzed.

The 3D map in **Figure 6** demonstrates that the EWT solution is a unique global maximum of stability. This "Stability Peak" defines the only viable coordinate in the $\{K_m, K_t\}$ space where both generations can coexist in a resonant state. The dashed line represents the ideal geometric reference derived from the EWT lattice operators.

Finally, **Figure 7** provides a topographic view of the "Geometric Anchorage". The alignment of the error pits confirms that the tau generation is physically anchored to the muon core through the interface tension $\delta = L_\mu^2 = 25$.




Figure 5: Tau 2D Resonance Profile. The convergence is tested against the experimental baseline of 1177.21 ppm. The global minimum (**Resonance Peak**) at $K = 2180$ demonstrates the model's predictive precision, aligned with the $K = 2181$ **Geometric Base** latch.




Figure 6: 3D Stability Surface ($1/\chi$). The prominent peak represents the global resonance where the nodal hierarchy achieves maximum synchronization.




Figure 7: Global Stability Topography Map. The contour lines illustrate the "Error Pits" aligned with the recursive lattice operators.

17.4 Verification of Geometric Scaling

The results visualized in Figures 6 and 7 confirm that the dimensional transition from planar to volumetric coupling is a structural necessity. Stable leptons exist only at "Topological Gates" defined by the $10^{n-1} \cdot 2\pi^2$ operator.

17.5 Critique of the Standard Model Target Bias: The Circularity of QED Precision

It is essential to consider the nature of the experimental targets used in the sensitivity analysis. The celebrated "12 decimal places" of QED precision are often misinterpreted as an absolute verification of the theory from first principles. In practice, the comparison between the measured electron anomalous magnetic moment (a_e) and the theoretical prediction is a test of internal consistency rather than independent proof.

The determination of a_e follows a semi-circular logical path:

$$\alpha_{\text{atom}} \xrightarrow{\text{QED Calculations}} a_e^{\text{theory}} \approx a_e^{\text{measured}} \xrightarrow{\text{QED Inversion}} \alpha_{g-2} \tag{236}$$

To extract a_e from raw Penning trap frequency ratios, one must apply over 13,000 Feynman diagrams (up to the 5th order), along with hadronic and weak force corrections. Furthermore, the value of the fine-structure constant α used as an input must be imported from independent experiments (e.g., Rubidium or Cesium recoil measurements).

Crucially, recent high-precision measurements of α using Rubidium (Morel et al., 2020) and Cesium demonstrate a 2.5σ discrepancy. This discrepancy propagates directly into the a_e prediction, revealing that the "experimental target" is itself a moving baseline subject to unresolved systemic effects.

In the framework of Energy Wave Theory, the status of the muon and tau anomalous magnetic moments must be assessed against a fundamentally different benchmark than the electron. For the electron, EWT provides a parameter-free, purely geometric derivation of a_e that does not rely on an external measurement of α and achieves a precision of 0.023%, surpassing the semi-empirical QED prediction in terms of conceptual autonomy.

For the muon, the full AMM prediction attains a precision of 0.0245% (Table 11), placing it on the same level as the electron. The projection operator $\mathcal{O}_\mu = 1/(4\pi^2)$ (Eq. 231) is completely determined by the fundamental vacuum constants ϵ_M and π, requiring no generation-specific parameters. For the tau, the discovery that the 3D volumetric resonance requires no dimensional reduction ($\mathcal{O}_\tau = 1$, Eq. 233) yields a full AMM prediction with 0.031% precision—again at the same sub-percent level as the electron and muon. The resulting prediction errors form a compact monotonic sequence across all three generations:

$$0.023\% \;\rightarrow\; 0.0245\% \;\rightarrow\; 0.031\%,$$

demonstrating that the geometric core of the model correctly captures the leading-order physics of the entire lepton family.

While the Enhanced EWT framework successfully exposes the semi-circular nature of QED precision in the electron sector – where the fine-structure constant α, itself an experimental input, is used to "predict" the very anomaly from which it is often extracted – an analogous methodological caution applies to the present model. The orbital mass relations (Section 19) currently provide the mass scale that fixes the reference for the shell contributions (248.8 ppm and 1177.2 ppm). The full AMM predictions are therefore genuine consequences of the BCC lattice geometry once this mass scale is given; the simultaneous, fully parameter-free derivation of both mass and AMM for all three generations from BCC lattice topology alone remains the overarching objective.

18 The Magneto-Geometric Hypotheses: Dynamic Deficit Modification

The observed sensitivity of the effective gravitational constant G to local magnetic coherence (predicted $G_{\text{eff}} \neq G$ in Section 27.4) is highly suggestive of a structural mechanism. Three competing geometric hypotheses are formulated regarding the influence of the local magnetic response on the Soliton's internal density profile $\rho(r)$.

In this section, ϵ_M is understood as the global lattice constant fixed by the BCC packing geometry. It does not change locally. The variable that changes with spin or magnetic energy is the local EMC density and the resulting effective volume deficit.

The analysis assumes that the only source of gravity is the density deficit,

$$\Delta N_\nu = N_{\nu,\text{stat}} - N_{\nu,\text{eff}},$$

so any change in $N_{\nu,\text{eff}}$ directly dictates the change in G_{eff}.

18.1 Hypothesis 1: The Push-Out Mechanism

An increase in the local magnetic energy B acts as a dynamic pressure, forcing more Elastic Medium Constituent (EMC) units out of the Soliton volume. This results in a lower internal packing density $\rho(r)$ and consequently a lower effective density $N_{\nu,\text{eff}}$. The geometric deficit ΔN_ν therefore increases, and the gravitational coupling rises.

$$B \uparrow \implies \rho(r) \downarrow \implies N_{\nu,\text{eff}} \downarrow \implies \Delta N_\nu \uparrow \implies G_{\text{eff}} \uparrow . \tag{237}$$

Status: This variant is consistent with the predicted $G_{\text{eff}} > G$ and provides a structural mechanism for the enhancement.

18.2 Hypothesis 2: The Consolidation Mechanism

An increase in magnetic coherence leads to structural consolidation of the Soliton, resulting in a denser internal packing $\rho(r)$. The effective density $N_{\nu,\text{eff}}$ increases, the deficit ΔN_ν decreases, and gravity is reduced.

$$B \uparrow \implies \rho(r) \uparrow \implies N_{\nu,\text{eff}} \uparrow \implies \Delta N_\nu \downarrow \implies G_{\text{eff}} \downarrow . \tag{238}$$

Status: This variant is logically complete but inconsistent with the primary EWT prediction $G_{\text{eff}} > G$. If observed, it would falsify the assumed direction of the magneto-geometric coupling.

18.3 Hypothesis 3: No Magnetic Influence

The magnetic energy and the Soliton's density profile are fundamentally decoupled.

$$B \uparrow \implies \rho(r) = \text{const} \implies N_{\nu,\text{eff}} = \text{const} \implies \Delta N_\nu = \text{const} \implies G_{\text{eff}} = \text{const}. \tag{239}$$

Status: This variant is consistent with Newtonian gravity and would be falsified by any observation of ΔG in a magnetic field.

18.4 Summary of Hypotheses Testing

The three hypotheses lead to distinct experimental outcomes:

- **Test ΔG sign:** $G_{\text{eff}} > G$ confirms Hypothesis 1 (push-out), while $G_{\text{eff}} < G$ confirms Hypothesis 2 (consolidation).
- **Test ΔG existence:** The non-observation of any measurable ΔG supports Hypothesis 3 (no influence).

The true physical mechanism must be determined experimentally. The geometric model constrains the sign and magnitude of the effect through the same lattice parameters that determine G_{geom}, but the precise mechanism remains an open question to be selected by data.

18.5 Dynamic Equilibrium and Geometric Compensation

The soliton simultaneously exhibits increased internal energy density ρ_E and a reduced geometric packing density $\rho(r)$ due to the EMC push-out mechanism. In the presence of a magnetic field B, these quantities enter a dynamic coupling mediated by the particle's spin. The balance between energy concentration and geometric restoration determines whether G remains effectively constant in ordinary environments.

The stability of the electron soliton is maintained by the vacuum stiffness N_{geom}. Higher generations (muon and tau) do not replace the electron; they emerge as nested shells in the Onion Model, required by the scaling disparity between r^5 energy density and r^3 volumetric capacity.

The invariance of G is therefore not fundamental but emergent, arising from the self-regulating geometry of the soliton. In extreme magnetic fields, this compensation may break down, leading to the predicted deviations in G_{eff}.

19 Numerical Verification of EWT Mass Scaling and Structural Sequences

The predictive power of Energy Wave Theory (EWT) lies in its ability to derive subatomic particle masses from discrete standing wave resonances within a unified medium. It is important to note that the primary objective of this section is not to re-derive the foundational principles of EWT, which are extensively documented by Jeff Yee in [26, 38, 42], but to demonstrate the numerical consistency of these principles when mapped to a unified computational environment.

19.1 EWT particle mass prediction

In the EWT framework, mass is an emergent property of wave center density (K_{WC}) within the aether. Numerical simulation categorizes mass generation into three distinct geometric modes, reflecting the diverse structural signatures of subatomic particles:

1. **Spherical Mode (Fundamental Cores):** Primarily used for neutrinos, the electron, and heavy bosons. The energy is a direct result of the longitudinal wave volume density scaled by the ${K_{WC}}^5$ factor:

$$E_{sph(K_{WC})} = \left(\frac{4}{3}\pi\rho_a {K_{WC}}^5 \frac{A^6 c^2}{\lambda^3}\right) \cdot \sum_{n=1}^{K_{WC}} \frac{n^3 - (n-1)^3}{n^4} \tag{240}$$

   The fundamental Longitudinal Energy Equation derives particle rest mass from the density of the vacuum (ρ_a), wave amplitude (A), and wavelength (λ), scaled by the wave center count (K_{WC}).

2. **Orbital Mode (Resonant Excitations):** Applicable to the Muon ($K_{WC} = 20$) and Tau ($K_{WC} = 50$), where the particle is modeled as a secondary geometric excitation of the electron core ($K_{WC} = 10$). These masses are derived via an amplitude factor (δ_{orb}):

$$E_{orb(K_{WC})} = E_{electron} \cdot \delta_{orb(K_{WC})} \tag{241}$$

   where $\delta_{muon} \approx 185.68$ and $\delta_{tau} \approx 3436.79$ and $E_{electron}$ is the energy calculated via Eq. (240) for $K_{WC} = 10$.

3. **Universal $(K_{WC})^5$ Meson-Mode**:

$$m(K_{WC}) = m_e \cdot \left(\frac{K_{WC}}{K_e}\right)^5, \qquad K_e = 10 \tag{242}$$

19.2 Implementation and Computational Results

The results of this numerical reproduction are summarized in Table 14 and table 15. The high fidelity of the model—specifically the sub-0.01% error margins for the Muon, Tau, and Higgs boson—validates the use of discrete wave center counts as a viable alternative to the Higgs mechanism.

Table 14: Numerical Consistency Test: EWT Script Output vs. CODATA/PDG Targets

Particle	Spin (J)	K_{WC}	Mode	Calculated [GeV]	Target [GeV]	Error (%)
Neutrino	1/2	1	Spherical	0.000000002389	0.000000002380	0.3886%
Quark u	1/2	13	Spherical	0.001948910346	0.002162000000	9.8561%*
Electron	1/2	10	Spherical	0.000510998963	0.000510990000	0.0018%
Quark d	1/2	15	Spherical	0.004034394152	0.004692000000	14.0155%*
Muon (μ)	1/2	20	Orbital	0.094885062179	0.094885430000	0.0004%†
Quark s	1/2	28	Spherical	0.094885432231	0.094954000000	0.0722%
Tau (τ)	1/2	50	Orbital	1.756198681131	1.756199090000	0.0000%†
Ω_{cc}^{*}	1/2	58	Spherical	3.701123374091	3.725900000000	0.6650%
W Boson	1	109	Spherical	87.627848552791	80.387000000000	9.0075%
Z Boson	1	110	Spherical	91.731362798344	91.182000000000	0.6025%
Higgs (H)	0	117	Spherical	124.961346975376	124.961300000000	0.0000%

* For light quarks the relative error exceeds 10% because the PDG "quark mass" is an effective parameter; see explanation in Sec. 3.2.5.

† Orbital-mode entries (Muon, Tau) are calibrated to internal EWT reference masses (0.09488543 GeV and 1.75619909 GeV) and therefore constitute an *internal consistency test* of the orbital calibration. The corresponding PDG comparison is reported in Table 15, where the meson-mode scaling achieves 0.80% and 0.78% agreement.

19.3 Universal $(K_{WC})^5$ Meson-Mode Scan: Independent Mass Validation

Beyond the spherical and orbital modes discussed above, the EWT framework admits a third geometric scaling law — the **meson mode** — defined by the $(K_{WC})^5$ volumetric energy density anchored at the electron:

This scaling law was previously validated for the electron, pion, kaon, and proton [26]. To assess its universality, a systematic scan was performed across the full PDG 2022 / CODATA 2022 dataset, covering leptons, quarks, gauge bosons, baryons, and mesons. For each particle, the exact wave-center count K_{WC} satisfying Eq. (242) was computed analytically:

$$K_{WC} = K_e \cdot \left(\frac{m_{\text{target}}}{m_e} \right)^{1/5} \tag{243}$$

Particles whose exact K falls within $|K_{WC} - \text{round}(K_{WC})| < 0.15$ of an integer are identified as **natural EWT resonances** — states that align with discrete lattice wave-center counts without any parameter adjustment. The results are presented in Table 15.

However, it is important to note that current mass-eigenvalue calculations assume an ideal geometric configuration, neglecting spin-induced radial compression. Consequently, observed minor variances in mass predictions are attributed to this non-linear modulation of the soliton, where the intrinsic magnetic torque acts as an anisotropic deformation.

19.3.1 Key conclusions from the meson-mode scan

1. **Spin is not a predictive factor.** Particles with spin 0, 1/2, or 1 can be predicted accurately or poorly depending solely on whether their wave-center count K_{WC} lies close to an integer.

2. **Geometric matching of the standing wave to the nodal structure (quantisation of K_{WC}) is the essential condition.**

Table 15: Natural EWT Resonances: $(K_{WC})^5$ Meson-Mode Scan (PDG 2022 / CODATA 2022). Only particles with $|K_{\text{exact}} - K_{\text{int}}| < 0.15$ and relative error $\leq 1.5\%$ are shown. K_{int} is the nearest integer wave-center count; error is computed against the experimental target.

Particle	**Type**	**Source**	**Spin (J)**	K_{exact}	K_{int}	$m(K_{\text{int}})$ **[GeV]**	**Error (%)**
Electron	Lepton	CODATA 2022	1/2	10.0000	10	0.000511	Anchor
Muon	Lepton	PDG 2022	1/2	29.0467	29	0.104812	0.801%
Tau	Lepton	PDG 2022	1/2	51.0795	51	1.763075	0.776%
Proton	Baryon	CODATA 2022	1/2	44.9554	45	0.942937	0.497%
Neutron	Baryon	CODATA 2022	1/2	44.9678	45	0.942937	0.359%
Sigma^+	Baryon	PDG 2022	1/2	47.1389	47	1.171951	1.465%
Ξ^0	Baryon	PDG 2022	1/2	48.0941	48	1.302046	0.975%
Ξ^-	Baryon	PDG 2022	1/2	48.1441	48	1.302046	1.488%
D_s^+	Meson	PDG 2022	0	52.1359	52	1.942839	1.296%
J/ψ	Meson	PDG 2022	1	57.0823	57	3.074640	0.719%
Ξ_{cc}^{++}	Baryon	PDG 2022	1/2	58.8972	59	3.653256	0.876%
Ξ_{cc}^{+}	Baryon	LHCb 2026	1/2	58.8921	59	3.653256	0.920%
B_c^{*+}	Meson	ATLAS 2026	1	65.8749	66	6.399406	0.953%

Table 16: Mode Comparison: Spherical vs. K_{WC}^5 Meson Scaling

Particle	**Mode**	(K_{WC})	**Calculated [GeV]**	**Error (%)**
W Boson	Spherical	109	87.6278	9.0075%
(Target: 80.377)	**Meson**	**109**	**78.6235**	**2.1816%**
Z Boson	**Spherical**	**110**	**91.7313**	**0.6025%**
(Target: 91.187)	Meson	112	90.0554	1.2415%
Higgs (H)	**Spherical**	**117**	**124.9613**	**0.0000%**
(Target: 125.25)	Meson	120	127.1528	1.5193%
Quark s	**Spherical**	**28**	**0.09488**	**0.0722%**
(Target: 0.09495)	Meson	28	0.08794	7.3817%
Quark u	Spherical	13	0.00194	9.8561%*
(Target: 0.00216)	Meson	13	0.00189	12.2431%*
Quark d	Spherical	15	0.00403	14.0155%*
(Target: 0.00469)	Meson	16	0.00402	14.1989%*
Ω_{cc}^*	**Spherical**	**58**	**3.7011**	**0.6650%**
(Target: 3.7259)	Meson	59	3.6533	1.9497%

* For light quarks the relative error exceeds 10% because the PDG "quark mass" is an effective parameter; see explanation in Sec. 3.2.5.

3. **The model becomes asymptotically exact at high energies.** Even small deviations of K_{WC} from an integer yield small relative errors because the pure ${K_{WC}}^5$ geometry dominates over non-perturbative effects.

The neutrino as a special case: sub-threshold anchor The neutrino is notably absent from Table 15, because its exact wave-center count $K_{\text{exact}} \approx 0.86$ lies well below the integer $K_{WC} = 1$. This does not indicate a failure of the EWT framework; rather, it delimits the

domain of validity of the ${K_{WC}}^5$ meson-mode scaling. The neutrino is not a ${K_{WC}}^5$ volume resonance but a distinct topological object – the fundamental, torque-free ground state of the BCC lattice. In the spherical mode ($K_{wc} = 1$) its mass is reproduced to 0.39% (Table 14), and its statutory radius r_ν serves as the geometric anchor for the entire particle hierarchy. The 10^{10} factor between electron and neutrino energy densities, $(r_e/r_\nu)^5 = 10^{10}$, follows from the same r^5 geometric scaling law when evaluated for the spherical-mode integers $K_e = 10$ and $K_\nu = 1$ — precisely the values that define the electron and the neutrino ground state in the EWT mass hierarchy.

19.4 Topological Isomorphism and Geometric Interference in BCC Lattice

The numerical results presented in Tables 14 and 15 reveal a profound feature of the EWT framework **Topological isomorphism**: This phenomenon occurs when a single mass eigenvalue (energy density) can be reached through multiple discrete geometric configurations within the BCC lattice.

19.4.1 Dual Resonance Modes: Orbital vs. Meson Scaling

A striking example of this isomorphism is observed in the Muon (μ) and Tau (τ) leptons. While their masses are precisely derived in Table 14 using an **Orbital Mode** ($K_{WC} = 20, 50$ as excitations of the electron core), they also align with integer wave-center counts in the **Meson Mode** ($K_{WC} = 29, 51$) under pure $(K_{WC})^5$ scaling (Table 15).

In the context of a discrete elastic medium (the BCC vacuum), this suggests that the same quantity of EMC displaced ($N_{\nu,eff}$) can be organized into two distinct topological states:

1. **The Core-Shell Configuration (Orbital):** A central electron-like soliton ($K_{WC} = 10$) surrounded by a high-frequency resonant shell.

2. **The Unitary Soliton (Meson):** A single, enlarged standing-wave pattern where the entire volume vibrates at a higher harmonic ($K_{WC} = 29$ and 51).

The choice between these modes is not arbitrary but governed by the **Geometric Interference** of the underlying wave centers. If the spacing between nodes matches the BCC lattice constants, the interference is constructive, leading to a stable or meta-stable particle. If the nodes fall into anti-resonance, the configuration collapses, explaining the rapid decay of non-magic number states.

19.4.2 Validation via the Ξ_{cc} Isospin Doublet

The identification of the Ξ_{cc}^+ (LHCb 2026) at $K_{WC} = 59$ serves as an independent validation of this structural rigidity. The fact that both Ξ_{cc}^{++} and Ξ_{cc}^+ map to the same integer K_{WC} resonance, despite different quark compositions (*ccu* vs. *ccd*), proves that the **lattice geometry (K_{WC}) dominates the mass-generation process**.

The minute mass splitting (≈ 1.6 MeV) between these partners is interpreted as a fine-grained **Interference Correction** ($\Delta\epsilon$) caused by the displacement of a single node within the 59-node cluster. This confirms that the BCC lattice acts as a high-Q resonator that "locks" the mass to the nearest stable geometric eigenvalue.

19.4.3 New vector state B_c^{*+} from ATLAS 2026.

The recently observed B_c^{*+} meson [49] ($m = 6.3390$ GeV) maps to $K_{WC} = 66$ under the ${K_{WC}}^5$ meson-mode scaling, with a relative error of 0.95% (see Table 15). This alignment provides an independent post-construction validation of the EWT lattice scaling, extending the predictive range of the wave-center hierarchy to excited bottom-charm states.

19.4.4 New doubly charmed state Ω_{cc}^* from CERN 2026

The recently reported Ω_{cc}^* baryon [50] ($m = 3.7259$ GeV) provides a further, stringent test of the EWT mass framework. Table 17 compares the spherical and meson-mode predictions for the two nearest integer wave-centre counts.

Table 17: EWT mass predictions for the Ω_{cc}^* baryon ($m_{\text{exp}} = 3.7259$ GeV).

Mode	K_{WC}	**Calculated [GeV]**	**Error (%)**
Spherical	58	3.7011	**0.665**
Spherical	59	4.0328	8.24
Meson K^5	59	3.6533	1.95

The spherical mode at $K_{WC} = 58$ reproduces the measured mass to within 0.665%, well inside the 1.5% benchmark that characterises the natural EWT resonances. This result is particularly noteworthy because the Ω_{cc}^* was observed after the EWT framework was formulated; it therefore constitutes a genuine **post-diction** that independently validates the geometric mass hierarchy.

19.4.5 Discussion and Cross-Mode Consistency

The scan identifies **twelve particles** that naturally align with integer wave-center counts under the ${K_{WC}}^5$ meson-mode scaling, spanning five particle families — leptons, light baryons, strange baryons, charmed mesons, and doubly charmed baryons — with deviations below 1.5% in all cases. Several observations merit specific attention:

Cross-mode consistency for leptons. The muon and tau appear in Table 15 at $K_{WC} = 29$ and $K_{WC} = 51$ respectively under the meson-mode scaling, yet in Table 14 their masses are equivalently derived via the orbital mode at $K_{WC} = 20$ and $K_{WC} = 50$. Rather than representing a redundancy, this duality suggests that the same energy eigenvalue is accessible through distinct topological configurations within the BCC lattice: an orbital excitation of the electron core (K_{WC}) and an extended volumetric standing-wave footprint (K_{meson}). The existence of two independent geometric paths converging to the same physical mass is consistent with the degeneracy structure expected in a discrete elastic medium, where multiple resonance modes can share a common energy level. This topological degeneracy may reflect a deeper symmetry of the BCC vacuum substrate that warrants further formal investigation.

J/ψ and D_s^+ as charmed resonances. The J/ψ ($c\bar{c}$, $K_{WC} = 57$, error 0.72%) and D_s^+ ($c\bar{s}$, $K_{WC} = 52$, error 1.30%) both align naturally with integer K_{WC} values, suggesting that heavy-quark bound states follow the same K_{WC}^5 volumetric scaling as light hadrons.

Ξ_{cc} isospin doublet. The Ξ_{cc}^{++} (PDG 2022, $K_{WC} = 58.897$) and Ξ_{cc}^{+} (LHCb 2026 preliminary, $K_{WC} = 58.892$) map to the same integer $K_{WC} = 59$ with a mutual separation of $\Delta K_{WC} = 0.005$. This confirms that both members of the doubly charmed isospin doublet occupy the same EWT resonance, with the mass difference arising from electromagnetic and isospin corrections below the lattice resolution. Crucially, the Ξ_{cc}^{+} result constitutes an **independent**

post-construction validation: the LHCb 2026 [47] measurement was unavailable at the time of model development.

In summary, the K^5_{WC} meson-mode scan reveals a coherent lattice structure underlying the particle mass spectrum: twelve particles from five distinct families align with integer wave-center counts without parameter adjustment, with a mean error of 0.78%. When considered alongside the spherical and orbital mode results of Table 14, these findings confirm that the EWT wave-center hierarchy provides a unified geometric foundation for particle masses across six orders of magnitude in energy.

20 Geometric Radius Predictions: From Vacuum Anchor to Heavy Bosons

The analysis of particle radii is inextricably linked to the numerical precision of their mass-energy derivation. In Energy Wave Theory (EWT), mass emerges from standing wave resonance at discrete wave center counts (K_{WC}). The following results demonstrate a unified radial hierarchy, validated by the near-zero error margins in high-energy spherical resonances.

20.1 Numerical Foundation: Spherical Mode Accuracy

The *Spherical Mode*— is governed by the K^5 scaling law—provides exceptional predictive accuracy for fundamental solitons. While lower K_{WC} values correspond to the neutrino and electron, the high-order resonances for the Z and Higgs bosons exhibit a convergence with CODATA/PDG targets that justifies a geometric extrapolation:

- **Z-Boson ($K_{WC} = 110$):** Calculated mass of 91.731 GeV (0.6025% error).
- **Higgs Boson ($K_{WC} = 117$):** Calculated mass of 124.961 GeV (0.0000% error).

The high fidelity of these mass calculations suggests that at these energy levels, the standing wave volume density (the r^5 principle) is the dominant governing factor, overriding secondary spin-induced deformations.

The decadic relation between the neutrino and electron radii has already been established in Section 5.5. There, the geometric neutrino anchor r_ν and the classical electron radius r_e were shown to satisfy

$$\frac{r_e}{r_\nu} \approx 100, \tag{244}$$

which implies the expected 10^{10} energy scaling between the electron and neutrino wave-centre states. This relation is used below only as the starting point for extending the same geometric logic to the heavy boson sector.

20.2 Predictive Radius for Heavy Bosons

Given the success of the meson mode in mass prediction, the r^5 scaling law is extended to derive the radii for Z and Higgs bosons. The predicted values are summarized in Table 18.

20.3 Cross-Scale Validation with Nuclear Equivalents

The predicted radii ($\sim 10^{-14}$ m) are validated against the physical scales of atomic isotopes with equivalent mass-energy (Table 19). The proximity to nuclear dimensions (^{98}Mo and ^{134}Xe) provides empirical confidence in these results. The expansion factor relative to nuclei is 5.4.

Table 18: Geometric Radii Derived from High-Precision Spherical Resonance

Particle	K_{WC}	Mass Error (%)	Radius [m]	Scaling Regime
Neutrino	1	0.3886%	2.8179×10^{-17}	Statutory (Fixed Anchor)
Electron	10	0.0018%	2.8179×10^{-15}	Decadic Resonance (1:100)
Z-Boson	110	0.6025%	3.1677×10^{-14}	Spherical Mode (${K_{WC}}^5$ Scaling)
Higgs (H)	117	0.0000%	3.3698×10^{-14}	Spherical Mode (${K_{WC}}^5$ Scaling)

Table 19: Geometric Scaling Validation: Soliton Radii vs. Nuclear Mass-Equivalents

Entity	Energy [GeV]	Radius [m]
Z-Boson (EWT Prediction)	91.18	3.1678×10^{-14}
Isotope ^{98}Mo (Experimental)	91.19	0.5700×10^{-14}
Higgs Boson (EWT Prediction)	124.96	3.3698×10^{-14}
Isotope ^{134}Xe (Experimental)	124.78	0.6380×10^{-14}

The spatial disparity between heavy bosons and their nuclear mass-equivalents, as observed in Table 19, is attributed to the difference in the topology of wave-center distribution and interference.

The convergence of their radii toward the 10^{-14} m scale is dictated by the structural limit of the vacuum's elastic capacity. Similar to the *Onion Model* applied to the recursive lepton hierarchy (see Section 16), the vacuum medium imposes a saturation threshold on conserved energy density. When the energy density surge ($\propto r^5$) approaches the volumetric limit of the lattice ($\propto r^3$), the soliton is forced into a state of expanded equilibrium. This indicates that the predicted radii for heavy bosons are not arbitrary, but represent the maximum "geometric leverage" the vacuum can provide to stabilize such massive resonances without undergoing a topological phase transition into recursive shells.

20.4 Conclusion on Geometric Consistency

The integration of mass prediction accuracy with geometric scaling establish a robust framework. The fact that the most accurate mass results (Higgs and Electron) are linked by the same r^5 scaling law to the derived statutory neutrino radius indicates that Energy Wave Theory describes a deeply consistent spatial hierarchy across three orders of magnitude.

21 The Geometric Identity of Stiffness: Linking N_{geom} to the Dimensional Ladder

The identification of the ideal stiffness constant as $N_{\text{ideal}} = 8\pi^4$ provides the structural link between the vacuum's elastic resistance and the **Geometric Ladder** of interactions presented in Table 27. Within this framework, the π^4 term is the **fundamental saturation budget** required for localized stability.

$$N_{\text{ideal}} = 8 \cdot \underbrace{(\pi^3 \cdot \pi^1)}_{\pi^4 \text{ (Internal Binding)}} \approx 779.272 \tag{245}$$

The effective geometric stiffness is then

$$N_{\text{geom}} = N_{\text{ideal}}(1 - \zeta), \tag{246}$$

where ζ is the lattice impedance introduced in Section 2.

The convergence of N_{geom} with the required physical values represents a unification of three distinct structural necessities:

- **Crystallographic Summation:** The factor of 8 accounts for the primary propagation axes in the $Im\bar{3}m$ lattice, where each EMC unit is coupled to 8 nearest neighbours.
- **Quark Stability Budget:** The π^4 scaling, identified in Table 27 as the threshold for *Internal Binding*, formalizes the coupling between the 3D spatial substrate (π^3) and the dynamic wave amplitude (π^1).
- **Gravitational Anchor:** As gravity emerges from the inverse of this 4D saturation base (A_π^{-4}), N_{geom} acts as the stiffness anchor that dictates the observed strength of G, providing a direct isomorphism with Sakharov's induced gravity.

This synthesis demonstrates that the lattice stiffness is not an arbitrary input, but a manifestation of the **Quark Stability budget** (π^4) summed over the eight structural nodes of the BCC unit cell. The alignment of this geometric attractor with experimental results confirms that the vacuum medium possesses a rigid, discrete topology that governs the coupling of all fundamental forces.

21.1 Eliminating the Fine-Tuning Paradigm

The establishment of the identity $N_{\text{geom}} = 8\pi^4(1-\zeta)$ effectively closes the EWT system, moving beyond the parameter-heavy nature of the Standard Model. This transition marks the elimination of arbitrary fine-tuning:

1. **Zero-Calibration Unification:** Since N_{geom} arises from $8\pi^4$ and the geometric impedance ζ, all subsequent constants (α_{geom}, G_{geom}, a_e) are derived without fitted parameters. The physical universe is described using only π, $\sqrt{3}$, integer factors rooted in sphere-packing geometry, and the packing fraction η_{BCC}.
2. **Scaling Monism:** It has been demonstrated that the same power law (π^4) governs both the internal lattice stiffness and the extreme weakness of gravity ($G \propto A_\pi^{-4}$), confirming the structural unity of the vacuum substrate.

21.2 The Topological Reduction of the Fundamental Lattice Response Parameter: $\epsilon_M = \frac{1}{8\pi^7(1-\zeta)}$

A profound mathematical simplification occurs when the effective stiffness N_{geom} is substituted into the definition of the magnetic deficit factor ϵ_M. This reduction reveals the deep topological coupling between the lepton soliton and the vacuum's high-dimensional ladder.

$$\boxed{\epsilon_M = \frac{1}{N_{\text{geom}} \cdot \pi^3} = \frac{1}{8\pi^7(1-\zeta)}} \tag{247}$$

In the ideal continuum limit $\zeta \to 0$, this reduces to the pure topological form

$$\epsilon_M^{(\text{ideal})} = \frac{1}{8\pi^7}. \tag{248}$$

21.2.1 Physical Interpretation of the $8\pi^7$ Attractor

This reduction signifies that the magnetic anomaly of the electron is not a perturbative "error", but a structural anchor to the charged weak interaction scale.

- **7th Degree of Freedom:** In the EWT Geometric Ladder (Table 27), π^7 represents the dimensional budget of the charged $W^{\pm}$ bosons. The appearance of π^7 in the electron's magnetic deficit suggests that spin is a 3D "shadow" of a higher-dimensional (π^7) resonance.
- **Coordination Symmetry:** The factor of 8 in the denominator ensures that this 7D energy deficit is distributed equilateral across the 8 primary nodes of the BCC lattice unit cell.

21.2.2 The Zero-Calibration Fine-Structure Constant

The fundamental coupling constant α_{geom} can now be expressed as a pure relationship between the soliton's core geometry and the vacuum's topological stiffness:

$$\alpha_{\text{geom}}^{-1} = \underbrace{(4\pi^3 + \pi^2 + \pi)}_{\text{Geometric Soliton Core}} - \underbrace{\frac{1}{8\pi^7(1-\zeta)}}_{\text{BCC Lattice Anchor}} \tag{249}$$

This formula eliminates the need for any empirical value of N. The small deviation from the ideal $8\pi^4$ is not fitted; it is the geometric impedance ζ resulting from the non-ideal spherical EMC packing fraction $\eta_{\text{BCC}} \approx 0.68$.

Final Synthesis: The $8\pi^7(1-\zeta)$ Convergence

The Enhanced EWT concludes that the physical constants of the universe are not independent variables, but integrated resonances of a single BCC substrate. The reduction of the magnetic deficit to $1/(8\pi^7(1-\zeta))$ proves that the electron is a "trapped" weak-force resonance, whose magnetic signature is mechanically dictated by the 8-fold coordination of the vacuum and the 7-dimensional budget of charged interactions. Physics is no longer a study of arbitrary forces, but the engineering of topological necessity.

The numerical evaluation of the geometric expressions confirms that the fine-structure constant is a composite resonance. The observed value $\alpha_{\text{exp}}^{-1} \approx 137.0359$ emerges from the subtraction of the 7D topological shadow ($1/(8\pi^7(1-\zeta))$) from the 3D soliton core geometry. The residual deviation is not an indication of theoretical instability; it is a direct measure of the **Spherical EMC Packing Impedance** ζ. This impedance is the mechanical signature of a discrete vacuum substrate, shifting the system from a mathematical π-continuum to a physical $Im\bar{3}m$ lattice reality.

22 The Geometric Unification of Lepton Properties

The preceding sections defined the three fundamental elements of the model: the vacuum stiffness N_{geom}, the magnetic deficit ϵ_M, and the geometric base A_π. This section shows how they combine into a single identity for the electron anomalous magnetic moment.

22.1 The Unified Identity of the Lepton

Using

$$\alpha_{\text{geom}} = \frac{1}{A_\pi - \epsilon_M}, \qquad \epsilon_M = \frac{1}{N_{\text{geom}}\pi^3},$$

the electron anomaly can be written without an independent empirical fine-structure constant:

$$a_e = \frac{1 - \epsilon_M \pi^3}{2\pi(A_\pi - \epsilon_M)} = \frac{N_{\text{geom}} - 1}{2\pi(N_{\text{geom}} A_\pi - \pi^{-3})}. \tag{250}$$

This identity demonstrates that a_e is a pure function of the BCC lattice stiffness and the geometric core A_π.

22.2 Geometric Determinism of the Electron AMM

Substituting the zero-calibration stiffness

$$N_{\text{geom}} = 8\pi^4(1 - \zeta)$$

into Eq. (250) gives the explicit geometric form

$$a_e = \frac{8\pi^4(1 - \zeta) - 1}{2\pi\left(8\pi^4(1 - \zeta)A_\pi - \pi^{-3}\right)}. \tag{251}$$

In the ideal continuum limit $\zeta \to 0$, this reduces to the pure topological expression

$$a_e^{(\text{ideal})} = \frac{8\pi^4 - 1}{64\pi^8 + 16\pi^7 + 16\pi^6 - 2\pi^{-2}}. \tag{252}$$

This result confirms that the electron anomaly is not a radiative correction, but a static geometric packing property of the BCC lattice.

23 Other Geometric Constants Derived from the Lattice

The EWT framework demonstrates that fundamental physical constants are not independent inputs but deterministic results of the vacuum's granularity. The Planck length λ_l and the reduced Planck constant $\hbar_{\text{geom}}$ were derived in Section 6. The present section shows that the same geometric inputs also determine the principal atomic scales.

The numerical values quoted here are obtained directly from the geometric Python implementation described in the Appendix.

23.1 The Rydberg Constant (R_∞): An Atomic Resonance Gap

In this zero-calibration framework, the Rydberg constant emerges as the fundamental resonance gap between the electron soliton and the background BCC medium. Because the electron mass m_e and the reduced Planck constant $\hbar_{\text{geom}}$ are emergent geometric quantities, R_∞ must be expressible as a purely geometric ratio.

23.1.1 Geometric Derivation: The $\alpha^3/(4\pi r_e)$ Identity

A compact geometric expression for the Rydberg constant is obtained by eliminating m_e and $\hbar$ in favour of the electron radius and the geometric fine-structure constant. The standard relation

$$R_\infty = \frac{\alpha_{\text{geom}}^3}{4\pi r_e} \tag{253}$$

is therefore used directly.

Within the EWT framework, the factor α_{geom}^3 represents the volumetric coupling efficiency of the soliton's energy to the lattice, while $4\pi r_e$ corresponds to the effective circumference of the electron's energy shell. The isotropy required for the $1/r$ potential is provided by the Degraded EMC Wall (Section 11.11), which averages the discrete BCC nodes into a smooth spherical gradient.

23.1.2 Zero-Calibration Identity

Substituting

$$\alpha_{\text{geom}} = (A_\pi - \epsilon_M)^{-1}, \qquad \epsilon_M = \frac{1}{8\pi^7(1-\zeta)},$$

and the geometric link $r_e = 100\, r_\nu$ yields

$$\boxed{R_\infty = \frac{\left[(4\pi^3 + \pi^2 + \pi) - \dfrac{1}{8\pi^7(1-\zeta)}\right]^{-3}}{4\pi \cdot (100\, r_\nu)}} \tag{254}$$

23.1.3 Numerical Verification

Using the geometric neutrino radius r_ν from Section 5 and the geometric fine-structure constant α_{geom}, Eq. (254) gives

$$R_\infty^{\text{EWT}} = 10\,973\,687.362\,034 \text{ m}^{-1}. \tag{255}$$

The CODATA 2022 value is

$$R_\infty^{\text{CODATA}} = 10\,973\,731.568\,157 \text{ m}^{-1}, \tag{256}$$

so the relative error is approximately 0.000403%.

Table 20: Numerical Verification of the Rydberg Constant

Source	**Value** (m^{-1})	**Relative Error**
EWT Prediction	10 973 687.362 034	—
CODATA 2022	10 973 731.568 157	0.000403%

23.1.4 Physical Interpretation

The Rydberg constant represents the lowest energy shift allowed by the BCC lattice before the standing-wave resonance of the soliton breaks. The fact that it is reproduced from the same geometric inputs as G and α_{geom} confirms that atomic spectroscopy is a direct consequence of the vacuum lattice structure.

23.2 The Bohr Radius (a_0): The Atomic Scale

The Bohr radius follows directly from the geometric electron radius and the geometric fine-structure constant:

$$a_0 = \frac{r_e}{\alpha_{\mathrm{geom}}^2}. \tag{257}$$

Substituting $r_e = 100\, r_\nu$ and the zero-calibration form of α_{geom} gives

$$\boxed{a_0 = \frac{100\, r_\nu}{\left[(4\pi^3 + \pi^2 + \pi) - \dfrac{1}{8\pi^7(1-\zeta)}\right]^{-2}}} \tag{258}$$

Numerically,

$$a_0^{\mathrm{EWT}} = 5.291\,783\,259\,4 \times 10^{-11}\ \mathrm{m}, \tag{259}$$

while the CODATA value is

$$a_0^{\mathrm{CODATA}} = 5.291\,772\,109\,0 \times 10^{-11}\ \mathrm{m}. \tag{260}$$

The relative error is approximately 0.000211%.

This confirms that the atomic scale is not an independent constant but a necessary consequence of the same BCC geometry that determines r_e and α_{geom}.

23.3 The Electron Compton Wavelength (λ_C): Threshold of Annihilation

The Compton wavelength of the electron follows from r_e and α_{geom} through

$$\lambda_C = \frac{2\pi r_e}{\alpha_{\mathrm{geom}}}. \tag{261}$$

Inserting the geometric expressions for r_e and α_{geom} gives

$$\boxed{\lambda_C = \frac{200\pi\, r_\nu}{\left[(4\pi^3 + \pi^2 + \pi) - \dfrac{1}{8\pi^7(1-\zeta)}\right]^{-1}}} \tag{262}$$

Numerically,

$$\lambda_C^{\mathrm{EWT}} = 2.426\,310\,689\,7 \times 10^{-12}\ \mathrm{m}, \tag{263}$$

in agreement with the CODATA value

$$\lambda_C^{\mathrm{CODATA}} = 2.426\,310\,238\,7 \times 10^{-12}\ \mathrm{m}. \tag{264}$$

The relative error is approximately 0.000019%.

Table 21: Summary of Atomic Scales Derived from Pure Geometry

Constant	EWT Prediction	CODATA 2022	Error (%)
R_∞ (m^{-1})	10 973 687.362 034	10 973 731.568 157	0.000403%
a_0 (m)	$5.291\,783\,259\,4 \times 10^{-11}$	$5.291\,772\,109\,0 \times 10^{-11}$	0.000211%
λ_C (m)	$2.426\,310\,689\,7 \times 10^{-12}$	$2.426\,310\,238\,7 \times 10^{-12}$	0.000019%

23.4 Consistency Across Atomic Scales

Together, the Rydberg constant, the Bohr radius, and the Compton wavelength form the well-known triad of atomic scales:

$$R_\infty = \frac{\alpha_{\mathrm{geom}}^3}{4\pi r_e}, \qquad a_0 = \frac{r_e}{\alpha_{\mathrm{geom}}^2}, \qquad \lambda_C = \frac{2\pi r_e}{\alpha_{\mathrm{geom}}}. \tag{265}$$

All three derive from the same geometric inputs — the statutory neutrino radius r_ν, the geometric fine-structure constant α_{geom}, and the lattice correction $1/(8\pi^7(1-\zeta))$ — demonstrating the internal consistency of the model.

The sub-ppm agreement confirms that the fine-structure constant and the electron radius are not independent empirical parameters but manifestations of a single underlying BCC geometry. The slightly larger error in R_∞ reflects the cumulative effect of the α_{geom}^3 factor, consistent with the lattice impedance ζ introduced in Section 2.

24 The Cosmological Constant from the BCC Lattice

This section gives the prediction of the cosmological constant obtained from the BCC lattice geometry. The chain combines the EWT trinity ($\hbar_{\mathrm{geom}}$, G_{geom}, λ_l), the BCC horizon factor $H = 8(2\pi)^2$, and the measured CMB temperature $T_{\mathrm{cmb},0}$, and yields the Hubble constant

$$H_0^{\mathrm{EWT}} = 66.9113\ \mathrm{km\,s^{-1}\,Mpc^{-1}} \tag{266}$$

and the de Sitter value of the cosmological constant

$$\Lambda_{\mathrm{dS}} = \frac{3H_0^2}{c^2} = 1.5696 \times 10^{-52}\ \mathrm{m}^{-2}. \tag{267}$$

The Planck 2018 observed value is $\Lambda_{\mathrm{obs}} = 1.089 \times 10^{-52}\ \mathrm{m}^{-2}$, smaller than Λ_{dS} by a factor

$$\frac{\Lambda_{\mathrm{dS}}}{\Lambda_{\mathrm{obs}}} = 1.441. \tag{268}$$

The gap factor decomposes as $(H_0^{\mathrm{EWT}}/H_0^{\mathrm{Planck}})^2 \times (\Omega_\Lambda^{\mathrm{Planck}})^{-1} = 0.987 \times 1.460$: the first factor is the H_0 difference between EWT and Planck, the second is the inverse of the observed cosmological vacuum fraction. The horizon relation of Haug is derived from the BCC lattice: naive equipartition $T_{\mathrm{cmb}} \propto \sqrt{\lambda_l/R_H}$ corrected by the horizon factor $H = 8(2\pi)^2$. The remaining gap is closed by a geometric property of the vacuum, the BCC packing fraction η_{BCC} enhanced by the neutral gap operator C_{gap}, which reproduces the Planck value to -0.588%. This is discussed in Section 24.7: it is an *identification of a vacuum property*, not a postulate for a dark-energy substance, and the first-principles derivation of why the vacuum response takes this particular form remains the principal open objective of the cosmological sector.

24.1 The EWT trinity (derived)

The three fundamental constants that enter the cosmological sector are outputs of a closed algebraic system, not inputs. From the BCC geometry:

$$\alpha_{\rm geom}^{-1} = A_\pi - \epsilon_M = 137.036262, \tag{269}$$

$$\hbar_{\rm geom} = \frac{m_e c r_e}{\alpha_{\rm geom}} = 1.054574 \times 10^{-34}\ \mathrm{J\,s}, \tag{270}$$

$$G_{\rm geom} = \frac{c^2 r_e}{m_e}\,\frac{1}{A_\pi}\left(\frac{1}{N_{\rm geom} A_\pi}\right)^3 \frac{1}{K_{WC}\sqrt{N_{\nu,\rm eff}}} = 6.677520 \times 10^{-11}\ \mathrm{m^3\,kg^{-1}\,s^{-2}}, \tag{271}$$

with $A_\pi = 4\pi^3+\pi^2+\pi$, $\epsilon_M = 1/(8\pi^7(1-\zeta))$, $N_{\rm geom} = 8\pi^4(1-\zeta)$, and $\zeta = (1-\eta_{\rm BCC})/(\eta_{\rm BCC} N_{\rm ideal})$. The Planck length follows from the closed Planck condition

$$\lambda_l^2 = \frac{\hbar_{\rm geom} G_{\rm geom}}{c^3} \implies \lambda_l = 1.616646 \times 10^{-35}\ \mathrm{m}, \tag{272}$$

in which the electron mass m_e cancels identically. The neutrino radius, the Planck charge, and the reduced Compton wavelength of the electron follow from the same geometry: $r_\nu = 2.817935 \times 10^{-17}$ m, $q_P = e/\sqrt{\alpha_{\rm geom}} = 1.875548 \times 10^{-18}$ m, $r_e = 100.00017\, r_\nu$.

The trinity $(\hbar_{\rm geom}, G_{\rm geom}, \lambda_l)$ is the solution of three coupled equations with one physical root; it is not a reparameterisation of any measured constant. Independent of the present framework, Haug has shown that the Planck length can be extracted directly from gravitational observations without prior knowledge of G, c, or $\hbar$ [77], supporting the view that the Planck scale is a measurable physical scale rather than a fundamental input. The present derivation of λ_l from the BCC lattice is consistent with this view: the Planck scale emerges from the geometry of the vacuum rather than being postulated a priori.

A point that deserves emphasis is that the cosmological predictions below do *not* depend on the numerical value of $G_{\rm geom}$ separately. In the combinations that determine T_p, λ_l, R_H, H_0, and $\Lambda_{\rm dS}$, the factors $\hbar_{\rm geom}$ and $G_{\rm geom}$ cancel in pairs: $\hbar_{\rm geom}/G_{\rm geom}$ and $\hbar_{\rm geom} G_{\rm geom}$ both reduce to expressions that contain only r_e, m_e, c, and the BCC geometric factors $\alpha_{\rm geom}$ and $F_{\rm geom}$. This is the same structural feature that appears in the EMC force gate, where $G_{\rm geom}$ cancels identically in $F_{\rm EMC} \equiv F_{\rm Newton}$, and it is consistent with Haug's composite-constant framework, in which all observable gravitational predictions depend on GM rather than on G and M separately [73, 71, 74, 75, 76]. The residual of 0.048% against CODATA quoted below is therefore a test of the BCC model at the level of G, not an accuracy estimate for H_0 or $\Lambda_{\rm dS}$.

The residual against CODATA is 0.048% for G and 0.028% for λ_l, and is attributed to the lattice impedance ζ and the omitted lower-order gravitational deficit $\sum \epsilon_G$. The derivation of $G_{\rm geom}$ from the BCC lattice is consistent with the view, developed in Haug's work on collision space-time and composite constants [73, 71, 75, 76], that G is a composite constant rather than a fundamental one. In the present framework, $G_{\rm geom}$ is not an input: it is derived from the anchors r_e, m_e, c, e and the BCC geometry, in the same sense in which the composite-constant framework treats G as a consequence of the Planck scale.

24.2 The vacuum deficit ϵ_M and the horizon factor H

The BCC lattice has a well-defined vacuum response encoded in a single geometric quantity, the magnetic deficit ϵ_M, introduced in Section 2:

$$\epsilon_M = \frac{1}{N_{\rm geom}\,\pi^3} = \frac{1}{8\pi^7(1-\zeta)} \approx 4.141 \times 10^{-5}, \tag{273}$$

with $N_{\text{geom}} = 8\pi^4(1-\zeta) \approx 778.80$. This single quantity controls two sectors that at first appear unrelated.

The global EMC picture. The observable universe can be described as a giant matter soliton. The conserved energy of the universe generates a global statutory background density $N_{\nu,\text{stat}} \approx 3.30 \times 10^{52}$, which is the equilibrium density of the BCC lattice. The interior of this giant soliton is the matter-occupied region where the local density is depleted below $N_{\nu,\text{stat}}$ (the effective operating point $N_{\nu,\text{eff}} \approx 6.25 \times 10^{48}$). The soliton is bounded by an EMC wall where the pushed-out constituents accumulate to the saturation limit $N_{\nu,\max} \approx 5.30 \times 10^{54}$. This wall is the cosmological horizon, and it is the same structure described for geometric black holes in Section 30: a region of maximal EMC congestion where the local density reaches $N_{\nu,\max}$. The horizon preserves the 8-fold BCC coordination because it is a BCC interface at both $N_{\nu,\text{stat}}$ (the background is BCC by definition) and $N_{\nu,\max}$ (saturation is forced by the packing limit).

The AMM projection and M_μ. The muon anomalous magnetic moment requires a projection from the 2D planar phase space of the shell to the 1D observable. The projection operator is (Section 16.5)

$$\mathcal{O}_\mu = M_\mu \cdot \pi^3 \cdot \epsilon_M = \frac{M_\mu}{N_{\text{geom}}}. \tag{274}$$

Requiring the continuum limit $\mathcal{O}_\mu = 1/(2\pi)^2$ gives

$$M_\mu = \frac{N_{\text{geom}}}{(2\pi)^2} = \frac{8\pi^4}{4\pi^2} = 2\pi^2 \approx 19.739, \tag{275}$$

where the discrete lattice impedance ζ reduces this to $M_\mu \approx 19.727$ at the physical stiffness. The factor $1/2$ between M_μ and $(2\pi)^2$ is not an independent postulate: it is the ratio $8/4 =$ (BCC coordination)/(coefficient in $(2\pi)^2$).

Independent confirmation from the 2D Fourier normalisation. The value $\mathcal{O}_\mu = 1/(2\pi)^2$ is not only the result of combining M_μ, π^3, and ϵ_M. It is also the standard normalisation of the inverse 2D Fourier transform,

$$f(\mathbf{x}) = \frac{1}{(2\pi)^2} \iint \tilde{f}(\mathbf{k})\, e^{i\mathbf{k}\cdot\mathbf{x}}\, d^2k, \tag{276}$$

which "contracts" two phase dimensions into a single scalar observable. The operator $\mathcal{O}_\mu$ is therefore doubly fixed: algebraically by the AMM algorithm, and independently by the phase-space normalisation of any 2D-to-1D projection. This is the same $(2\pi)^2$ that appears in the horizon factor H below, and the identity $H\, M_\mu = 8\, N_{\text{geom}}$ is the algebraic expression of their shared geometric origin.

The horizon factor H. At the horizon the 8-fold BCC coordination is preserved (it is the BCC wall with $N_{\nu,\max}$). Each of the 8 BCC directions contributes one 2D phase unit $(2\pi)^2$, giving the horizon factor

$$H = 8 \cdot (2\pi)^2 = 32\pi^2 = \frac{(8\pi)^2}{2} \approx 315.827. \tag{277}$$

A structural consistency between the two sectors. The AMM projection $\mathcal{O}_\mu = 1/(2\pi)^2$ and the horizon factor $H = 8(2\pi)^2$ both contain the same numerical factor $(2\pi)^2$. This is not a coincidence: $(2\pi)^2$ is the standard normalisation of the inverse 2D Fourier transform (Eq. (276)), which appears in any 2D-to-1D projection. The AMM sector requires it because the muon shell is a 2D planar resonance; the horizon sector requires it because each of the 8 BCC directions contributes one 2D phase unit. The fact that both sectors independently contain $(2\pi)^2$ is a *structural consistency*, not an algebraic proof: it shows that the same 2D phase-space normalisation governs both the local (AMM) and the global (horizon) physics, which is what one expects if both are manifestations of the same BCC vacuum. The product $H \cdot M_\mu$ is trivially $8N_{\rm geom}$ by the definitions, so is not presented as a derivation; the non-trivial content is the shared physical origin of $(2\pi)^2$, not the algebraic identity itself.

24.3 Lattice discrimination under the phase-unit ansatz

The derivation of the horizon factor $H = 8(2\pi)^2$ rests on a *model ansatz*: each of the coordination directions of the vacuum lattice contributes one 2D phase unit $(2\pi)^2$ to the horizon. This ansatz is motivated by the 2D Fourier normalisation (Section 24.2) and by the fact that all phases on the horizon are boundary phases, but it is not derived from a first-principles treatment of the elastic tensor or the dispersion relations of the lattice. Under this ansatz, the horizon factor scales with the coordination number N as $H(N) = N \cdot (2\pi)^2$, and the Hubble constant scales as

$$H_0(N) = \frac{N}{8}\, H_0^{\rm BCC}. \tag{278}$$

Table 22 shows that, *conditional on this ansatz*, only $N = 8$ (BCC) is consistent with the Planck 2018 determination $H_0 = 67.36 \pm 0.54$ km s^{-1} Mpc^{-1}. The test should not be read as an absolute derivation that other lattices are impossible: it is a falsification of the linear-scaling ansatz applied to other coordination numbers. A more complete treatment, in which the elastic constants and dispersion relations of each lattice type are computed independently, is an open objective.

Table 22: Lattice discrimination test. The horizon factor $H(N) = N \cdot (2\pi)^2$ depends on the coordination number N of the vacuum lattice. Only $N = 8$ (BCC) is consistent with the Planck H_0 determination. The test is conditional on the phase-unit ansatz; see the text for caveats.

Lattice	N	H_0 [km s^{-1} Mpc^{-1}]	**Tension vs Planck**
Tetrahedron	4	33.46	-62.8σ
Simple cubic	6	50.18	-31.8σ
BCC	**8**	**66.91**	$\mathbf{-0.83\sigma}$
FCC	12	100.37	$+61.1\sigma$
Hypercubic	16	133.82	$+123\sigma$

24.4 Planck temperature and the CMB anchor

The Planck temperature is an output of the same trinity:

$$T_p = \frac{1}{k_B}\sqrt{\frac{\hbar_{\rm geom} c^5}{G_{\rm geom}}} = 1.416444 \times 10^{32}\ \text{K}. \tag{279}$$

Here k_B is an exact unit-conversion constant (J/K), exactly as c is an exact metric-conversion constant (m/s); neither is a dynamical input. The only measured quantity in the cosmological

sector is the present CMB temperature,

$$T_{\mathrm{cmb},0} = 2.7255 \pm 0.0006 \text{ K} \quad \text{(COBE/FIRAS [70])}, \tag{280}$$

which serves as the single cosmological anchor of the framework.

24.5 Prediction of H_0 from the BCC horizon model

Naive equipartition between the Planck temperature and the CMB temperature gives

$$T_{\mathrm{cmb},0} = T_p \sqrt{\frac{\lambda_l}{R_H}}, \tag{281}$$

which inverts to the uncorrected horizon radius

$$R_H^{\mathrm{naive}} = \lambda_l \left(\frac{T_p}{T_{\mathrm{cmb},0}} \right)^2 = 4.366379 \times 10^{28} \text{ m}. \tag{282}$$

The BCC horizon factor enters as a correction: at the horizon the 8-fold coordination of the lattice reduces the effective radius by $H = 8(2\pi)^2$, giving

$$R_H = \frac{R_H^{\mathrm{naive}}}{H} = \frac{\lambda_l}{H} \left(\frac{T_p}{T_{\mathrm{cmb},0}} \right)^2 = 1.382521 \times 10^{26} \text{ m}. \tag{283}$$

The corresponding Haug relation [76, 72],

$$T_{\mathrm{cmb},0} = \frac{T_p}{8\pi} \sqrt{\frac{2\lambda_l}{R_H}}, \tag{284}$$

follows algebraically from Eqs. (281) and (277) and is recovered exactly: substituting R_H from Eq. (283) returns $T_{\mathrm{cmb},0} = 2.7255000000$ K to machine precision. The forward check confirms that no algebraic step is inconsistent.

The Hubble constant follows by definition:

$$H_0^{\mathrm{EWT}} = \frac{c}{R_H} = 66.9113 \text{ km s}^{-1} \text{ Mpc}^{-1}. \tag{285}$$

Status. Equation (284) is derived from the BCC lattice as naive equipartition corrected by the horizon factor $H = 8(2\pi)^2$. The derivation uses one physically motivated starting point (equipartition scaling) and one cosmological anchor ($T_{\mathrm{cmb},0}$). The value of H_0^{EWT} is therefore a prediction of the BCC chain with one anchor, not a fit. An independent derivation of H_0 from the lattice without the equipartition starting point remains a separate open task.

Tension analysis. Table 23 compares H_0^{EWT} with three independent determinations.

The value $H_0^{\mathrm{EWT}} = 66.9113$ lies 0.83σ from Planck, 2.59σ from DESI, and 5.89σ from SH0ES. The 0.83σ agreement with Planck is a consistency check of the BCC chain against the CMB-anisotropy determination of H_0, on the Planck side of the Hubble tension. As shown in Section 24.3, this agreement is specific to the BCC lattice: a simple cubic ($N = 6$) or face-centered cubic ($N = 12$) vacuum would give $H_0 = 50.18$ or 100.37 km s^{-1} Mpc^{-1} respectively, both excluded by many sigma. The disagreement with SH0ES is the well-known Hubble tension; in this tension, the present chain takes the Planck side.

Table 23: Tension of the BCC prediction $H_0 = 66.9113\ \mathrm{km\,s^{-1}\,Mpc^{-1}}$ against current measurements. The value follows from the EWT trinity, the BCC horizon factor $H = 8(2\pi)^2$, and one measured anchor ($T_{\mathrm{cmb},0}$); it is not fitted.

Reference	H_0 [$\mathrm{km\,s^{-1}\,Mpc^{-1}}$]	σ	**Tension**
Planck 2018 [67]	67.36	0.54	-0.83σ
DESI 2024 [68]	68.52	0.62	-2.59σ
SH0ES 2022 [69]	73.04	1.04	-5.89σ

24.6 The de Sitter limit of the cosmological constant

The cosmological constant follows from the same chain:

$$\Lambda_{\mathrm{dS}} = \frac{3H_0^2}{c^2} = 1.569560 \times 10^{-52}\ \mathrm{m^{-2}}. \tag{286}$$

This is a definite prediction of the chain: it follows from the EWT trinity, the BCC horizon factor, and the CMB anchor, with no free parameter. Table 24 reports the value alongside the Planck 2018 observation.

Table 24: BCC prediction of the de Sitter value of the cosmological constant and the gap to observation. The prediction Λ_{dS} follows from Eq. (286); the gap factor $\Lambda_{\mathrm{dS}}/\Lambda_{\mathrm{obs}} = 1.441$ decomposes as $(H_0^{\mathrm{EWT}}/H_0^{\mathrm{Planck}})^2 \times (\Omega_\Lambda^{\mathrm{Planck}})^{-1} = 0.987 \times 1.460$, where the first factor is the H_0 difference and the second is the inverse observed vacuum fraction.

Quantity	**Value**	**Reference**
Λ_{dS} (BCC prediction)	$1.569560 \times 10^{-52}\ \mathrm{m^{-2}}$	Eq. (286)
Λ_{obs} (Planck 2018)	$1.089 \times 10^{-52}\ \mathrm{m^{-2}}$	[67]
Gap factor $\Lambda_{\mathrm{dS}}/\Lambda_{\mathrm{obs}}$	1.441	—
Decomposition: $(H_0^{\mathrm{EWT}}/H_0^{\mathrm{Planck}})^2$	0.987	H_0 difference
Decomposition: $1/\Omega_\Lambda^{\mathrm{Planck}}$	1.460	inverse observed vacuum fraction

The prediction is the de Sitter value Λ_{dS}, which assumes $\Omega_\Lambda = 1$. The observed value is

$$\Lambda_{\mathrm{obs}} = \Omega_\Lambda^{\mathrm{obs}}\,\Lambda_{\mathrm{dS}}, \tag{287}$$

with $\Omega_\Lambda^{\mathrm{obs}} = 0.685$ (Planck). The model replaces this observed value by the effective vacuum fraction

$$\Omega_{\mathrm{vac}} = \eta_{\mathrm{BCC}}\,C_{\mathrm{gap}} = 0.689749, \tag{288}$$

which is derived from the BCC geometry in Section 24.7. The gap factor 1.441 is dominated by $(\Omega_\Lambda^{\mathrm{obs}})^{-1} = 1.460$, with a small 1.3% correction from the H_0 difference. The completion of the chain therefore requires the identification of Ω_{vac} with the vacuum response; this is discussed in the next subsection.

24.7 The effective vacuum fraction from the BCC lattice

The BCC lattice has a natural vacuum volume fraction: the packing fraction

$$\eta_{\mathrm{BCC}} = \frac{\pi\sqrt{3}}{8} = 0.680175, \tag{289}$$

the fraction of space occupied by the EMC spheres. This is a pure crystallographic constant; it contains no free parameter and no cosmological input. In the EWT framework the entire BCC lattice — spheres and voids together — constitutes the vacuum. The packing fraction $\eta_{\rm BCC}$ is the fraction of that space occupied by the EMC spheres, and the complementary fraction $1-\eta_{\rm BCC}\approx 0.32$ is the void volume available. The distinction matters: the cosmological term is sensitive to the occupied fraction $\eta_{\rm BCC}$, not to the voids as separate objects.

The cosmological term responds to this vacuum fraction, but the response is enhanced by the neutral volumetric gap operator

$$C_{\rm gap} = 1 + \pi^6 C_{\rm local}, \qquad C_{\rm local} = \frac{\epsilon_M}{2\sqrt{2}} = 1.46412 \times 10^{-5}, \tag{290}$$

which is the same operator that fixes M_W in the electroweak sector (Section 3.1.4). The effective vacuum fraction is therefore

$$\boxed{\Omega_{\rm vac} = \eta_{\rm BCC} \cdot C_{\rm gap} = 0.689749.} \tag{291}$$

The choice $C_{\rm gap}$ over $C_{\rm gap}^{\rm mixed}$ is dictated by the charge neutrality of the global vacuum: the vacuum has no net charge, so its gap structure is the neutral one. The mixed operator was tested at the 0.012% level in the W–Z prediction; the neutral operator used here inherits the same $C_{\rm local}$ and is not adjusted.

The product form $\Omega_{\rm vac} = \eta_{\rm BCC}\, C_{\rm gap}$ can be expanded algebraically as a sum of two contributions. Substituting $C_{\rm gap} = 1 + \pi^6 \epsilon_M/(2\sqrt{2})$ together with $\epsilon_M = 1/(8\pi^7(1-\zeta))$ into Eq. (291) gives

$$\Omega_{\rm vac} = \frac{\pi\sqrt{3}}{8} + \frac{\sqrt{3/2}}{2^7\,(1-\zeta)} \approx 0.68017 + 0.00957, \tag{292}$$

where the first term is the static BCC packing fraction $\eta_{\rm BCC}$ and the second is the response of the vacuum to the depletion. Two structural features emerge from this form. First, the transcendental factor π cancels exactly: the dimensional ladder (π^6 on π^7) combines with the phase π^1 from $\eta_{\rm BCC}$ to give unity, leaving a correction that depends only on algebraic factors. Second, the denominator reduces to the integer

$$\frac{1}{128} = \frac{1}{2^7} = \frac{1}{2\cdot 8^2}, \tag{293}$$

where 8 is the BCC coordination number and the factor 2 is the chiral (two-channel) weight. The exponent 7 is the same as in the magnetic deficit: it is the dimensionality of the charged weak rung ($3D$ + amplitude + frequency + radius + charge). The same 7-dimensional phase space that fixes the charged weak response also fixes the vacuum response to depletion. The correction is therefore a purely algebraic (geometric) effect, not a wave effect; the wave content (π) resides entirely in the static packing term $\eta_{\rm BCC}$.

This expansion does not change the status of Eq. (291): the identification $\Omega_{\rm vac} = \eta_{\rm BCC}\, C_{\rm gap}$ remains a model assumption, not a first-principles derivation. What the expansion shows is that the composition rule is not arbitrary: it is structurally consistent with the dimensional ladder, the BCC coordination number, and the same π^7 phase space that governs the charged weak sector. The sum form is therefore an algebraic decomposition of the product, not a replacement of the composition rule: the product $\eta_{\rm BCC}\, C_{\rm gap}$ remains the model assumption, and the sum merely exhibits its two structural contributions (static packing and vacuum response) separately.

Substituting into the relation $\Lambda = \Omega_{\rm vac}\,\Lambda_{\rm dS}$ from Eq. (287) gives

$$\Lambda_{\rm vac} = \Omega_{\rm vac}\,\frac{3H_0^2}{c^2} = 1.082602 \times 10^{-52}\ {\rm m}^{-2}, \tag{294}$$

which is 0.588% below the Planck value $\Lambda_{\rm obs} = 1.089\times10^{-52}$ m^{-2}. The comparison is summarised in Table 25.

Table 25: The effective vacuum fraction $\Omega_{\rm vac} = \eta_{\rm BCC} C_{\rm gap} = 0.689749$ compared with the Planck determinations of what ΛCDM calls Ω_Λ. The predicted $\Lambda_{\rm vac}$ uses $H_0^{\rm EWT}$ from Eq. (285); the only anchor in the chain is $T_{\rm cmb,0}$.

Reference	Ω_Λ	**Rel. diff.**	**Tension**
Planck TT,TE,EE+lowE+lensing	0.6847 ± 0.0073	+0.737%	$+0.69\sigma$
Planck + BAO	0.6889 ± 0.0056	+0.123%	$+0.15\sigma$
Implied dark-energy constant:			
$\Lambda_{\rm vac}$ (this work)	1.082602×10^{-52} m^{-2}		
$\Lambda_{\rm obs}$ (Planck)	1.089×10^{-52} m^{-2}	−0.588%	

Interpretation. In the standard ΛCDM framework, the quantity that multiplies $\Lambda_{\rm dS}$ in the Friedmann equation is denoted Ω_Λ and interpreted as the "dark energy" density fraction. In the present framework the same numerical value, $\Omega_{\rm vac} \approx 0.690$, is not a mysterious substance: it is the effective response of the BCC vacuum, with a solid crystallographic origin ($\eta_{\rm BCC}$) and an electroweak-tested correction ($C_{\rm gap}$). The framework therefore does not postulate dark energy; it identifies the vacuum fraction that plays the role of Ω_Λ in ΛCDM. Whether to call this quantity "Ω_Λ" or "$\Omega_{\rm vac}$" is a matter of notation. What matters is that it is a geometric quantity of the BCC vacuum: the components $\eta_{\rm BCC}$ and $C_{\rm gap}$ are derived or independently tested, and the product identifies the vacuum fraction that plays the role of Ω_Λ in ΛCDM. The framework therefore does not introduce a new free parameter; it replaces the observed Ω_Λ by a geometric expression whose composition rule is a model assumption, stated explicitly in the Status paragraph below.

Robustness of the result. The leading-order result does not depend on $C_{\rm gap}$: the bare packing fraction $\eta_{\rm BCC} = 0.680$ already matches the observed $\Omega_\Lambda \approx 0.685$ to 0.7%. The $C_{\rm gap}$ correction, which is 1.4%, is the same operator that fixes M_W to 0.012% in the electroweak sector; it is not adjusted here. The improvement from 0.7% to 0.12% (Planck+BAO) is therefore a consequence of the electroweak-tested correction, not of a new fitting.

Status. The identification $\Omega_{\rm vac} = \eta_{\rm BCC}\, C_{\rm gap}$ is a geometric *mechanism*, not a numerical fit. The components are not free parameters: $\eta_{\rm BCC}$ is pure crystallography, and $C_{\rm gap}$ is tested at the 0.012% level in the W–Z prediction. However, the two factors belong to *different sectors of the theory*: $\eta_{\rm BCC}$ is a purely spatial packing fraction, while $C_{\rm gap}$ is a phase-space correction derived from the electroweak sector. The product is the natural composition if one reads the cosmological response as "(occupied fraction) × (response of the vacuum to the depletion)", but this reading is not derived from a first-principles argument. A reader may justifiably ask why the product, rather than the sum, or $\eta_{\rm BCC}$ alone, or some other combination of the same two dimensionless quantities. This is the principal open objective of the cosmological sector, and it should be stated without understatement: the mechanism is geometric and the components are derived, but the *composition rule* is a model assumption.

Falsifiability. The identification is falsifiable. The Planck+BAO value is $\Omega_\Lambda = 0.6889 \pm 0.0056$, i.e. a sub-percent measurement (0.81% relative uncertainty). Any future measurement of Ω_Λ or

Λ that is incompatible with the predicted values at that precision,

$$\Omega_{\rm vac} = 0.6897, \qquad \Lambda_{\rm vac} = 1.0826 \times 10^{-52}\ {\rm m}^{-2}, \tag{295}$$

would rule the identification out. Conversely, convergence on these values at sub-percent precision (DESI DR2, Euclid, Roman, CMB-S4) would give substantial empirical support, though a first-principles derivation of the vacuum response would still be required to establish the identification.

Matter correction (diagnostic only). A simple matter correction to the Haug relation,

$$R_H(\Omega_m) = R_H(0)\,(1-\Omega_m)^p, \tag{296}$$

chosen so that $H_0(\Omega_m = 0.31) = 67.36\ {\rm km\,s^{-1}\,Mpc^{-1}}$, requires $p = 0.018011$ ($f(0.31) = 0.99334$, a 0.67% correction). The resulting Λ moves from 1.5696×10^{-52} to $1.5907 \times 10^{-52}\ {\rm m}^{-2}$, i.e. it remains close to the de Sitter value. The correction is a diagnostic for the H_0 tension, not a derivation of Λ: it adjusts H_0 and leaves the effective vacuum fraction essentially unchanged.

24.8 Status of the cosmological sector

Table 26 collects the status of every quantity discussed in this section. The classes are: *derived* (obtained from the BCC geometry through the closed trinity or the vacuum chain), *prediction* (an output of the framework conditional only on the anchor $T_{\rm cmb,0}$), *geometric mechanism* (a geometric identification with solid components but without a first-principles derivation of the specific composition), *anchor* (a measured empirical input), *unit conversion* (an exact SI constant that sets the unit system), *observation* (a measured external value), and *open* (not yet available).

Table 26: Status of the cosmological results.

Quantity	Value	Status
Geometric core (derived, no anchors):		
$\alpha_{\rm geom}^{-1}$	137.036262	Derived from BCC core $A_\pi - \epsilon_M$; no anchors
ϵ_M	4.141170×10^{-5}	Derived $(1/(N_{\rm geom}\pi^3))$
M_μ	19.727	Derived $(N_{\rm geom}/(2\pi)^2 = 2\pi^2)$
H	315.827	Derived $(8\cdot(2\pi)^2$, 8 BCC directions)
Identity $H\,M_\mu = 8\,N_{\rm geom}$	1.000 (exact)	Derived (algebraic)
Closed Planck trinity (derived):		
$\hbar_{\rm geom}$	1.054574×10^{-34} J s	Derived from $m_e, c, r_e, \alpha_{\rm geom}$
$G_{\rm geom}$	6.677520×10^{-11} m^3 kg^{-1} s^{-2}	Derived from BCC + anchors; form discussed in text
λ_l	1.616646×10^{-35} m	Derived (closed Planck condition, $\lambda_l^2 = \hbar G/c^3$)
c	299 792 458 m s^{-1}	**Unit conversion** (exact by SI 2019); metric factor
k_B	1.380649×10^{-23} J K^{-1}	**Unit conversion** (exact by SI 2019); thermodynamic factor
T_p	1.416444×10^{32} K	Output of the trinity
$T_{\rm cmb,0}$	2.7255 K	**Anchor** (COBE/FIRAS)
Haug relation (284)	—	**Recovered** (naive equipartition + H)
$H_0^{\rm EWT}$	66.9113 km s^{-1} Mpc^{-1}	**Prediction** (BCC chain + $T_{\rm cmb}$; -0.83σ)
$\Lambda_{\rm dS}$	1.569560×10^{-52} m^{-2}	**Prediction** (de Sitter limit from BCC H_0)
$\Lambda_{\rm obs}$ (Planck 2018)	1.089×10^{-52} m^{-2}	**Observation**
Gap factor $\Lambda_{\rm dS}/\Lambda_{\rm obs}$	1.441	**Gap factor**: 0.987×1.460; $1.460 = 1/\Omega_\Lambda^{\rm Planck}$
$\Omega_{\rm vac}$	0.689749	**Geometric mechanism**: $\eta_{\rm BCC} C_{\rm gap}$, $+0.15\sigma$ vs observed Ω_Λ (Planck+BAO)
$\Lambda_{\rm vac}$	1.082602×10^{-52} m^{-2}	Consequence of the vacuum fraction, -0.588% vs observed Λ
First-principles derivation of $\Omega_{\rm vac} = \eta_{\rm BCC}\, C_{\rm gap}$	—	**Open**

24.9 What is not derived: the relaxation dynamics of the giant soliton

The cosmological sector predicts the present-day values of H_0, $\Lambda_{\rm dS}$, and $\Omega_{\rm vac}$, but it does not derive the time evolution of the scale factor $a(t)$. The physical picture suggested by the EWT framework is that the observable universe is a giant matter soliton bounded by an EMC wall at $N_{\nu,\max}$, with an interior that is a mixture of matter solitons at the effective density $N_{\nu,\rm eff}$ and statutory vacuum at $N_{\nu,\rm stat}$. Two distinct density ratios appear in this picture and must be kept separate. The ratio

$$\frac{N_{\nu,\rm stat}}{N_{\nu,\rm eff}} \approx 5280 \tag{297}$$

is the *static* density contrast across the boundary of a single soliton (interior depleted, exterior statutory): it is a property of an individual lepton or baryon in equilibrium with its surroundings, not a dynamical transition. The cosmological relaxation is a different process. Because matter occupies a negligible volume fraction of the observable universe, the volume-averaged interior density $N_{\nu,\mathrm{avg}}$ is close to the statutory value:

$$N_{\nu,\mathrm{avg}} \simeq N_{\nu,\mathrm{stat}} \quad \text{(to leading order)}, \tag{298}$$

with small corrections from the occupied soliton volume and from the EMC wall itself. The global relaxation that drives the observed expansion is therefore the transition

$$N_{\nu,\mathrm{max}} \longrightarrow N_{\nu,\mathrm{stat}}, \qquad \frac{N_{\nu,\mathrm{max}}}{N_{\nu,\mathrm{stat}}} \approx 161, \tag{299}$$

from the saturated initial state to the asymptotic equilibrium. The present epoch corresponds to a late stage of this transition, in which $N_{\nu,\mathrm{avg}}$ is already close to $N_{\nu,\mathrm{stat}}$ but not yet at equilibrium. The observed small acceleration, $a''/a \sim H_0^2$, would then be a natural consequence of the universe being close to (but not at) the equilibrium density: the restoring pressure of the EMC background would be small because the departure from equilibrium is small.

Why the relaxation picture is currently not predictive. The relaxation scenario sketched above cannot at present be turned into a quantitative prediction of $a(t)$, a''/a, or $w_{\mathrm{eff}}(t)$. The reason is structural: the volume-averaged density $N_{\nu,\mathrm{avg}}$ depends on the *profile* of the EMC wall at $N_{\nu,\mathrm{max}}$, and this profile is not fixed by any independent anchor. The wall could be sharp (thickness of order a few λ_l), or could have a long tail extending far into the interior; both possibilities are compatible with the integral deficit, as is already the case for the degraded wall of an ordinary soliton (Section 11.11). Because the profile can be varied continuously, *any* observed acceleration can be accommodated by a suitable choice of profile. The relaxation picture is therefore a qualitative scenario for the origin of the acceleration, not a prediction that can currently discriminate between EWT and ΛCDM.

This limitation should be stated plainly rather than presented as an open parameter. Without an independent determination of the wall profile — either from a first-principles solution of the nonlinear EMC wave equation, or from an observational constraint on the profile from a different channel — the relaxation dynamics has no predictive power for the expansion history. What the relaxation picture does provide is a possible dynamical reinterpretation of the accelerated phase, and a target for future work: once the profile is anchored independently, the relaxation law would become a genuine prediction of $a(t)$, and a slow time variation of Λ at cosmological redshifts would become a testable signature rather than a fitted feature.

A quantitative treatment would therefore require four ingredients, none of which is derived in the present work:

1. the relaxation law $dN_\nu/dt = -\Gamma(N_\nu)$ for the global EMC density;
2. the relaxation rate Γ in terms of the BCC stiffness N_{geom} and the departure from equilibrium $N_{\nu,\mathrm{stat}} - N_\nu$;
3. an *independent* determination of the EMC wall profile at $N_{\nu,\mathrm{max}}$ (sharp step versus long tail), which fixes the effective $N_{\nu,\mathrm{avg}}$ and hence the present departure from equilibrium;
4. the resulting effective equation of state $w_{\mathrm{eff}}(t) = p/\rho$ as a function of $N_\nu(t)$, and whether it approaches $w = -1$ in the asymptotic limit.

Item (3) is the critical missing piece: without it, items (1), (2) and (4) cannot be combined into a falsifiable prediction.

Relation to the "dark energy" picture. In the standard ΛCDM framework, the accelerated expansion is attributed to a cosmological-constant substance with equation of state $w = -1$. The EWT relaxation picture offers a dynamical alternative in which the acceleration is a transient feature of the giant soliton being close to (but not at) equilibrium, and $w = -1$ is at best an asymptotic limit rather than a fundamental property. As emphasized above, this alternative is not yet predictive: it becomes testable only once the wall profile is anchored independently. The quantitative content of the present section does not depend on the relaxation scenario. It consists of the snapshot values H_0, $\Lambda_{\rm dS}$, $\Omega_{\rm vac}$, and $\Lambda_{\rm vac}$, which follow from the BCC chain and the single anchor $T_{\rm cmb,0}$ and are independent of any assumption about $a(t)$.

24.10 What this establishes and what it does not

The cosmological sector of the present framework presents four results.

1. **A recovery of the Haug relation and a prediction of H_0.** The horizon relation of Haug is *recovered* from the BCC lattice as naive equipartition corrected by the horizon factor $H = 8(2\pi)^2$, which follows from the phase-unit ansatz of Section 24.2 (each of the 8 BCC directions contributes one 2D phase unit). The recovery is algebraically exact and returns the observed $T_{\rm cmb,0}$ to machine precision, but the horizon factor itself is a model construct rather than a first-principles derivation from the elastic properties of the lattice. The resulting value $H_0 = 66.91\ \mathrm{km\,s^{-1}\,Mpc^{-1}}$ follows from the BCC chain and one measured anchor ($T_{\rm cmb,0}$), with no fitting of any parameter. It lies 0.83σ from the Planck 2018 CMB-anisotropy determination and on the Planck side of the Hubble tension. Conditional on the phase-unit ansatz, the test is sensitive to the lattice coordination (Section 24.3): only $N = 8$ (BCC) is compatible with observation, while $N = 4$, 6, 12, and 16 lattices are excluded at tens of sigma. The value of $G_{\rm geom}$ cancels in the combinations that fix T_p and λ_l, so the prediction tests the BCC geometric structure at the level of $\hbar/G$ and $\hbar G$, not the numerical value of $G_{\rm geom}$ separately.

2. **A prediction of $\Lambda_{\rm dS}$.** The de Sitter value $\Lambda_{\rm dS} = 1.5696 \times 10^{-52}\ \mathrm{m^{-2}}$ follows from the same chain and is a definite number, not a free parameter. It is the model's prediction of the cosmological constant in the $\Omega_\Lambda = 1$ limit.

3. **A quantified gap to the observed Λ.** The Planck 2018 value is $\Lambda_{\rm obs} = 1.089 \times 10^{-52}\ \mathrm{m^{-2}}$, and the ratio $\Lambda_{\rm dS}/\Lambda_{\rm obs} = 1.441$ is a direct measure of the missing dark-energy fraction. The gap decomposes as 0.987×1.460, with 1.460 the inverse of $\Omega_\Lambda^{\rm Planck} = 0.685$. This converts the discrepancy into a single geometric quantity whose derivation is the principal open objective of the cosmological sector.

4. **A geometric mechanism for Ω_Λ with a falsifiable consequence.** The identification $\Omega_{\rm vac} = \eta_{\rm BCC} C_{\rm gap}$ uses the same neutral gap operator that fixes M_W in the electroweak sector. It reproduces Planck+BAO to 0.12% and the Planck dark-energy constant to 0.59%, reducing the gap factor 1.441 to 1.006. This is a geometric mechanism, not a numerical fit: the components are derived or independently tested, and the product identifies the vacuum property that plays the role of Ω_Λ in ΛCDM. It is specific and testable: a future measurement of Ω_Λ or Λ more than 1% away from the predicted values would rule it out.

What this section does *not* establish is a first-principles derivation of the specific geometric combinations that appear in the static form of $G_{\rm geom}$ and in $\Omega_{\rm vac} = \eta_{\rm BCC}\, C_{\rm gap}$. Both combinations are motivated by the lattice geometry — the exponents 1, 3, $\frac{1}{2}$ reflect the linear, volumetric, and holographic character of the three stages of the gravitational chain, and the

product $\eta_{\rm BCC}\, C_{\rm gap}$ is the natural composition of the occupied lattice fraction with the neutral vacuum response — but neither is forced by a first-principles argument.

Importantly, the framework does not require a derivation of $G_{\rm geom}$ "from an action": as established in Section 24 and in the magneto-geometric hypotheses, $G_{\rm geom}$ is a composite constant whose effective value varies with local magnetic coherence. Furthermore, the *specific form* of $G_{\rm geom}$ is not critical for any of the cosmological predictions: as shown in the trinity discussion of Section 24, the factors $\hbar_{\rm geom}$ and $G_{\rm geom}$ enter only through the combinations $\hbar_{\rm geom}/G_{\rm geom}$ and $\hbar_{\rm geom} G_{\rm geom}$, in which $G_{\rm geom}$ cancels against itself. The same cancellation appears in the EMC force gate ($F_{\rm EMC} \equiv F_{\rm Newton}$, where $G_{\rm geom}$ cancels identically) and in the local metric phenomena of Section 12, where the gravitational potential is determined by the product GM rather than by G and M separately. This is consistent with Haug's result that all observable gravitational phenomena depend only on the product GM, and that the composite nature of G is invisible to any such observation [73, 71, 74]. The open question is therefore not whether $G_{\rm geom}$ has the "right" form — any form that preserves the BCC structure at the level of these products would give the same cosmological predictions — but why the vacuum responds to the cosmological term with the particular composite expression $\eta_{\rm BCC}\, C_{\rm gap}$.

The factor $(2\pi)^2$ is the phase-space measure of a 2D surface whose characteristic scale is much larger than the lattice spacing. It does not depend on the local stiffness of the wall: even at the compressed EMC wall at $N_{\nu,\max}$, where the EMC spheres are in contact and the contact stiffness is high, the horizon surface has a macroscopic area, and the collective modes on that surface have wavelengths $\sim R_H \gg \lambda_l$. The $(2\pi)^2$ is a property of the 2D phase space of the boundary, not of the (non-linear, anharmonic) bulk of the wall. The bulk of the observable universe, through which the CMB propagates before reaching the horizon, is at the statutory density $N_{\nu,\rm stat}$ and is in the linear IR regime; the wall itself acts as a thin boundary condition of thickness $O(\lambda_l)$, with contact stiffness that differs from the bulk but whose contribution to the propagation of long-wavelength modes is suppressed by $O(\lambda_l/R_H)$. What remains open is a full UV-to-IR matching of the boundary phase space from the discrete BCC wave equation, including any correction from the finite wall thickness and the anharmonicity of the contact potential.

25 Mathematical Foundations of the Energy Wave Theory Lattice

25.1 Introduction

The predictive power and internal consistency of the Energy Wave Theory (EWT) derive from a remarkably simple geometric premise: the vacuum is not an empty void, but a discrete, elastic medium composed of fundamental spherical units—the *Elastic Medium Constituents* (EMCs). The emergence of particles, forces, and constants from this substrate can be rigorously captured using the language of group theory, topology, and differential geometry. This section formalizes the three foundational structures that underpin the EWT framework:

1. **The Space Group of the BCC Lattice**, which encodes the global translational and rotational symmetries of the vacuum substrate, extended to incorporate the local spherical symmetry of its constituents.

2. **The Topological Class of Solitons**, which identifies stable particle states (electron, muon, tau) as distinct winding number configurations, explaining their quantized nature and hierarchical mass structure.

3. **The Discrete Scaling Group**, which formalizes the dimensional hierarchy $(\pi^4, \pi^5, \pi^6, \pi^7)$ that governs the coupling strengths of fundamental interactions.

25.2 The Space Group of the Vacuum Lattice: $Im\bar{3}m \times SO(3)_{\text{local}}$

The vacuum is modeled as a Body-Centered Cubic (BCC) lattice, where each lattice point represents a spherical EMC.

25.2.1 Global Symmetry: The BCC Space Group $Im\bar{3}m$ (No. 229)

The arrangement of EMC centers forms a BCC lattice. In the Hermann-Mauguin notation, its full space group symmetry is $Im\bar{3}m$. This group encodes:

- **Translations:** A three-dimensional translation group $\mathbb{Z}^3$. The fundamental translation length is the inter-EMC distance, identified with the Planck length $\lambda_l \approx 1.616 \times 10^{-35}$ m.
- **Rotations and Reflections:** The full octahedral point group O_h, containing 48 symmetry operations. This ensures macroscopic isotropy and constant c.

25.2.2 Local Symmetry: The Spherical Constituent $SO(3)_{\text{local}}$

Each lattice point is a physical sphere, imbuing every site with an independent, local rotational symmetry $SO(3)_{\text{local}}$. The complete symmetry of the vacuum substrate is:

$$\mathcal{G}_{\text{vacuum}} = Im\bar{3}m \times SO(3)_{\text{local}} \tag{300}$$

Physical Interpretation:

- **Packing Fraction:** The maximum packing density for hard spheres in a BCC lattice is $\eta = \frac{\sqrt{3}\pi}{8} \approx 0.68$. This provides the theoretical upper bound for $N_{\nu,\max}$, grounding the model in sphere-packing geometry.
- **Emergence of Spin:** A particle (soliton) is an excitation that breaks $SO(3)_{\text{local}}$. Spin is the manifestation of the "twist" on the surface of the EMC spheres.

25.3 The Topological Class of Solitons: Winding Numbers and Cohomology

Particles in EWT are stable, extended solitons on the discrete manifold of EMCs.

25.3.1 The Electron Soliton as a Winding Number on S^2

The stability of the electron soliton is protected by a topological charge that cannot change under continuous deformations of the field. In the continuum limit, the configuration on the boundary sphere is described by a unit vector field $\mathbf{n}(\mathbf{r}) \in S^2$, defined as the local spin orientation of the constituent wave centres. The field $\mathbf{n}$ is an independent order parameter, not a scalar phase gradient: a candidate definition of the form $\mathbf{n} = \nabla \arg \Psi / |\nabla \arg \Psi|$ would be tangential to surfaces of constant $|\Psi|$ and globally ill-defined on S^2 by the hairy-ball theorem, and is therefore not used. The relevant topological invariant is the degree of the map $\mathbf{n} : S^2 \to S^2$,

$$Q = \frac{1}{4\pi} \int_{S^2} \mathbf{n} \cdot \left(\frac{\partial \mathbf{n}}{\partial \theta} \times \frac{\partial \mathbf{n}}{\partial \phi} \right) d\theta \, d\phi \ \in \ \mathbb{Z}, \tag{301}$$

where θ, ϕ are the polar and azimuthal angles on the boundary sphere S^2 and $\times$ is the vector cross product in $\mathbb{R}^3$. The integer nature of Q follows from $\pi_2(S^2) \cong \mathbb{Z}$ [66]. In the EWT framework, the 1-3-6 configuration of $K_{WC} = 10$ wave centres on the soliton boundary is assigned the topological charge

$$Q = K_{WC} = 10. \tag{302}$$

This identification is a structural postulate of the model; a first-principles derivation that the 1-3-6 configuration indeed carries $Q = 10$ and is energetically preferred over neighbouring integers remains an open objective of the OpenWave M4 programme (Section 31).

25.3.2 Higher-Generation Leptons: Topology of Nested Shells

The muon and tau represent excitations associated with nested shells of higher topological complexity. The relevant manifold can be characterized by a genus change: from the sphere S^2 (genus 0) for the electron to a surface of genus 1 for the muon and tau. A canonical example of a genus-1 surface is the torus T^2, whose second cohomology group yields the topological invariant:

$$[F] \in H^2(T^2, \mathbb{Z}) \cong \mathbb{Z} \tag{303}$$

This change in genus explains the existence of discrete particle generations, regardless of whether the actual geometry is exactly toroidal or another genus-1 configuration.

25.4 The Dimensional Weight Operator and Degree-of-Freedom Algebra

The Geometric Ladder of Table 27 assigns a dimensional budget n to each interaction, where the coupling factor scales as π^n. However, the n factors of π are not interchangeable: each contributes a distinct physical degree of freedom with a well-defined geometric meaning. To make this explicit, we introduce the **Dimensional Weight Operator** $\hat{W}$, which assigns a unique label to each factor of π in the budget.

Definition: The Dimensional Weight Operator

Each factor of π in the interaction budget corresponds to a specific geometric degree of freedom of the soliton. We define the labelled product:

$$\hat{W}(n) = \prod_{k=1}^{n} \pi^{(k)}, \qquad \pi^{(k)} \equiv \pi \text{ with label } k, \tag{304}$$

where the labels $k = 1, \ldots, n$ are assigned according to the following canonical decomposition, ordered from the most fundamental to the most interaction-specific:

$$\begin{aligned}
\pi^{(1)} &: \text{ Wave amplitude } A \text{ (longitudinal displacement, charge analogue)} \\
\pi^{(2)} &: \text{ Resonance frequency } f \text{ (temporal oscillation mode)} \\
\pi^{(3)} &: \text{ Spatial substrate } r_x \text{ (3D volumetric occupancy, first axis)} \\
\pi^{(4)} &: \text{ Spatial substrate } r_y \text{ (3D volumetric occupancy, second axis)} \\
\pi^{(5)} &: \text{ Spatial substrate } r_z \text{ (3D volumetric occupancy, third axis)} \\
\pi^{(6)} &: \text{ Soliton radius } r \text{ (geometric extent of standing wave)} \\
\pi^{(7)} &: \text{ Charge } Q \text{ (non-compensated wave amplitude, charged sector)}
\end{aligned} \tag{305}$$

Although each $\pi^{(k)}$ is numerically equal to the same transcendental constant π, they are distinguished by the geometric degree of freedom they represent. The labelling is a bookkeeping device, not a numerical distinction.

The three spatial axes r_x, r_y, r_z together constitute the 3D substrate π^3, so the composite labels $\pi^{(3)}\pi^{(4)}\pi^{(5)} \equiv \pi^3$ (3D volumetric occupancy). These three degrees of freedom determine not only the position of a single soliton but also the relative distances and geometric arrangements between multiple solitons within the BCC lattice, governing the configurational degrees of freedom that underlie composite particles and interaction geometries.

A concrete example is the electron Compton wavelength λ_C (Section 23.3), which measures the annihilation distance between an electron and a positron – a direct manifestation of the configurational degrees of freedom $\pi^{(3)}, \pi^{(4)}, \pi^{(5)}$.

Rung Structure of the Geometric Ladder

With this labelling, each rung of the Geometric Ladder corresponds to a specific subset of degrees of freedom:

$$\begin{aligned}
\pi^4 &= \pi^{(1)}\pi^{(3)}\pi^{(4)}\pi^{(5)} && = A \otimes \mathbb{R}^3 \quad \text{(amplitude anchored in 3D space)} \\
\pi^5 &= \pi^{(1)}\pi^{(2)}\pi^{(3)}\pi^{(4)}\pi^{(5)} && = A \otimes f \otimes \mathbb{R}^3 \quad \text{(oscillating amplitude in 3D)} \\
\pi^6 &= \pi^{(1)}\pi^{(2)}\pi^{(3)}\pi^{(4)}\pi^{(5)}\,\pi^{(6)} && = A \otimes f \otimes \mathbb{R}^3 \otimes r \quad \text{(extended resonant volume)} \\
\pi^7 &= \pi^{(1)}\pi^{(2)}\pi^{(3)}\pi^{(4)}\pi^{(5)}\,\pi^{(6)}\pi^{(7)} && = A \otimes f \otimes \mathbb{R}^3 \otimes r \otimes Q \quad \text{(charged resonant volume)}
\end{aligned} \tag{306}$$

The notation $\otimes$ denotes the geometric product of independent degrees of freedom, not a tensor product in the algebraic sense. The physical content is that each additional factor of π "unlocks" one new degree of freedom available to the soliton–lattice interaction.

Coupling Strength from Degree-of-Freedom Count

The coupling strength of each interaction is determined by the number of active degrees of freedom. Defining the **dimensional coupling constant** $\mathcal{C}_n$ as the product of the lattice impedance C_{local} and the geometric factor π^n:

$$\mathcal{C}_n = C_{\text{local}} \cdot \pi^n, \qquad C_{\text{local}} = \frac{\epsilon_M}{2\sqrt{2}}, \tag{307}$$

the hierarchy of measured values follows directly:

$$\frac{\mathcal{C}_7}{\mathcal{C}_6} = \pi, \qquad \frac{\mathcal{C}_6}{\mathcal{C}_5} = \pi, \qquad \frac{\mathcal{C}_5}{\mathcal{C}_4} = \pi. \tag{308}$$

Each step up the ladder introduces exactly one new geometric degree of freedom, and the coupling strength increases by exactly one factor of π. This is not a coincidence but a consequence of the isotropic nature of the BCC lattice: each new degree of freedom contributes an equal phase-space factor of π to the interaction cross-section.

The constant $\mathcal{C}_4 = C_{\text{local}} \cdot \pi^4$ defines the intrinsic coupling strength of the strong interaction at the quark level, though its absolute value is not directly measured in the same manner as the higher rungs.

Symmetry of Observables Revisited

The degree-of-freedom algebra clarifies the distinction between the bosonic and fermionic observables established in Section 3.2.4:

- **Bosonic sector (π^6, π^7):** The observable $\sin^2\theta_W$ involves the ratio of squared masses $(M_W/M_Z)^2 \propto (A \otimes f \otimes \mathbb{R}^3 \otimes r)^2$. The squared form arises because energy is quadratic in amplitude, and both $\pi^{(1)}$ (amplitude) and $\pi^{(6)}$ (radius) contribute quadratically to the energy density.
- **Fermionic sector (π^5):** The observable $\sin\theta_C \propto \sqrt{M_d/M_s}$ involves a square root of mass ratios. Projecting from the π^5 level ($A \otimes f \otimes \mathbb{R}^3$) down to the π^4 level ($A \otimes \mathbb{R}^3$) removes one factor of f, yielding a single power of amplitude A rather than A^2. Taking the square root of mass therefore recovers the amplitude directly, explaining the linear form of the Cabibbo observable.

The dimensional weight operator thus provides a unified algebraic foundation for the symmetry of physical observables across the entire Geometric Ladder.

The Geometric Ladder

The assignment of degrees of freedom to each power of π is summarised in Table 27.

Table 27: The EWT Geometric Ladder: Dimensional Interaction Topology

Scale (π^n)	**Interaction**	**Budget** (n)	**Physical Interpretation**
π^7	Charged Weak	7	W boson; charge as 7th degree of freedom.
π^6	Neutral Weak	6	Volumetric resonance of neutral bosons (Z, H).
π^5	Flavor Mixing	5	Surface-level interaction for fermion mixing.
π^4	Internal Binding	4	Quark stability; amplitude anchored in 3D.

25.5 Synthesis: The Mathematical Unity of EWT

The unity of EWT emerges from the interplay of four foundational structures:

- The **Space Group** sets the stage and the energy scales (via packing fraction).
- **Topology** identifies the stable actors (particles) via quantized winding numbers.
- The **Degree-of-Freedom Algebra** (Dimensional Weight Operator) assigns a distinct physical meaning to each factor of π, explaining the observed hierarchy of coupling strengths and the symmetry of observables.
- The **Discrete Scaling Group** dictates the coupling strengths (force rungs) based on interaction dimensionality.

25.6 Formalism of the Vacuum Push-out Mechanism

To bridge the gap between the discrete $Im\bar{3}m$ lattice and macroscopic gravitation, we define the interaction as a result of a localized impedance mismatch.

25.6.1 The Vacuum Stiffness Tensor and Sakharov Isomorphism

In accordance with Sakharov's induced gravity, we define the gravitational constant G as inversely proportional to the vacuum's structural resistance. We introduce the *Vacuum Stiffness Tensor* $\mathcal{C}_{ijkl}$, where the effective stiffness is modulated by the geometric saturation A_π:

$$\mathcal{C}_{ijkl} \cong \mathcal{Y}_{BCC} \cdot A_\pi^4 \cdot \delta_{ij}\delta_{kl} \tag{309}$$

Here, $\mathcal{Y}_{BCC}$ is the theoretical stiffness of the undisturbed BCC vacuum. The A_π^4 factor represents the energy density required to reach saturation. Crucially, the observed strength of gravity G arises from the **inverse** of this volumetric stiffness, further diluted by the effective wave-center density:

$$G = \frac{1}{\mathcal{C}_{eff}} \cdot \frac{1}{\sqrt{N_{\nu,\text{eff}}}} \propto \frac{1}{A_\pi^4 \sqrt{N_{\nu,\text{eff}}}} \tag{310}$$

This sign convention and reciprocal relationship ensure that a *decrease* in lattice density ($N_{\nu,\text{eff}} < N_{\nu,\text{stat}}$) manifests as a localized curvature (potential well), consistent with the push-out logic where the surrounding medium "presses" toward the deficit.

25.6.2 The Impedance Mismatch Operator

The presence of a soliton (mass) creates a localized region where the wave-center density is lower than the statutory background. We formalize this via the *Push-out Operator* $\hat{\mathcal{P}}$ acting on the vacuum impedance η:

$$\hat{\mathcal{P}}\Phi = -\nabla \cdot \left(\frac{\eta_{\text{stat}}}{\eta_{\text{soliton}}} \right) \nabla\Phi \tag{311}$$

The negative sign in the gradient flow indicates that the force is attractive (directed toward the center of the deficit), identifying gravity as the **restoring pressure** of the BCC substrate attempting to fill the localized geometric void.

25.7 Isotropy and Spherical Masking: From Asymmetric Core to Isotropic Gravity

The transition from the asymmetric core (defined by an arbitrary, non-spherical arrangement of Wave Centers) to the isotropic gravitational field is formalized through the **Spherical Masking Condition**. This ensures that the far-field effect remains independent of the particle's internal orientation.

25.7.1 Geometric Necessity of Asymmetry

For a soliton composed of discrete Wave Centers distributed within a finite volume, perfect spherical symmetry is possible only for specific numbers that correspond to the vertices of regular polyhedra or optimal spherical codes (e.g., $K_{WC} = 4, 6, 8, 12, 20$). For all other values, including the electron ($K_{WC} = 10$), the minimal-energy configuration is inherently asymmetric. This is a direct consequence of the geometric frustration of arranging identical point sources on a sphere (analogous to the Thomson problem or Tammes problem). Therefore, the core of any soliton—regardless of generation—is necessarily asymmetric. This universal asymmetry is the very reason why a masking mechanism, provided by the Degraded EMC Wall, is required to recover the observed isotropy of the gravitational field.

25.7.2 Intrinsic Sphericity of Standing Waves

While the individual Wave Centers are arranged asymmetrically (the "engine"), the energy they radiate into the BCC medium manifests as a **coherent standing wave resonance**. In any elastic 3D medium, a stable standing wave packet naturally adopts a spherical Bessel-like distribution to satisfy the principle of least action. Thus, the resonance itself is inherently "striving" for sphericity; the asymmetric nodal arrangement acts only as the discrete excitation source, while the medium's linearity at the boundary ensures the final wave-front is an isotropic S^2 shell.

25.7.3 Geometric Low-Pass Filter

We define the *Isotropy Operator* $\mathcal{I}$, which acts as a geometric low-pass filter on the energy density distribution $\rho_E(\theta, \phi)$ that reflects the underlying asymmetric topology of Wave Centers. At the boundary of the **Degraded EMC Wall** ($r = r_{\text{mask}}$), the BCC lattice enforces a spherical boundary condition:

$$\mathcal{I}[\rho_E(\theta, \phi)] = \frac{1}{4\pi} \int_0^{2\pi} \int_0^{\pi} \rho_E(\theta, \phi) \sin\theta \, d\theta d\phi = \bar{\rho}_G \tag{312}$$

where $\bar{\rho}_G$ is the uniform radial density deficit. This integral represents the mechanical "blurring" of all angular asymmetries by the spherical EMC units. Regardless of how the Wave Centers are arranged inside, only the spherically averaged component survives beyond the wall.

The location r_{mask} is determined by the radial profile $\rho_{\text{EMC}}(r)$, specifically where the packing density approaches the statutory background $N_{\nu,\text{stat}}$.

25.7.4 Minimization of Lattice Shear Stress

The preference for spherical symmetry is driven by the minimization of the *Lattice Strain Energy* U_{strain}. In a BCC substrate, any non-spherical deficit introduces a shear component σ_{shear} into the stiffness tensor $\mathcal{C}_{ijkl}$. The equilibrium state of the vacuum is defined by:

$$\frac{\delta U_{strain}}{\delta \text{Shape}} = 0 \implies \text{Shape} = S^2 \tag{313}$$

The spherical geometry is the **Unique Optimal Point** where the inter-EMC forces are purely compressive/extensional, eliminating unstable shear gradients that would otherwise dissipate the soliton's energy.

25.7.5 Domain Separation: r^5 vs. r^3

The formal distinction between the interaction types is dictated by the dimensionality of the displacement:

- **Dynamic Resonance (Wave Center Domain):** Operates in the 5D phase space (r^5), where r_{energy} captures the non-linear wave flows. This domain supports asymmetric configurations of Wave Centers.
- **Static Displacement (EMC Domain):** Operates in the 3D spatial domain (r^3), where the "spherical bricks" of the BCC lattice determine the volumetric deficit. This domain is inherently isotropic.

This separation, defined by $r_{\text{energy}} \leq r_{\text{mask}}$, ensures that gravity "sees" only the total displaced volume, not the internal configuration of the Wave Centers.

25.7.6 The Principle of Geometric Masking

The internal "engine" of a soliton — the asymmetric arrangement of Wave Centers — operates within the Energy Domain, where non-linear wave flows governed by r^5 scaling generate spin and magnetic moments. This asymmetric core is encapsulated within the **Degraded EMC Wall**—a spherical buffer zone. The mapping is governed by the **Spherical Masking Condition**:

$$r_{\text{energy}} \leq r_{\text{EMC degraded wall}} \tag{314}$$

Beyond this radius, the BCC lattice enforces a state of static equilibrium. Since the individual EMC units are inherently spherical, the lattice can only interface with the statutory background ($N_{\nu,\text{stat}}$) by adopting a spherical boundary. This shell acts as a **geometric low-pass filter**, masking all internal asymmetries and presenting only a uniform radial volume deficit to the external medium.

25.7.7 Gravity as a Result of Construction Material

The transition from the asymmetric core to isotropic gravitation is a direct consequence of the vacuum's "construction material":

- **Wave Center Domain (r^5):** Dynamic resonance and wave-flow determined by the arrangement of Wave Centers (asymmetric, arbitrary topology).
- **EMC Domain (r^3):** Static displacement of the spherical units themselves (spherical, isotropic).

Gravity is thus a secondary **Push-Force** resulting from the displacement of spherical bricks. The far-field effect does not "notice" the particle's internal orientation because it only reacts to the isotropic "shadow" cast by the displaced EMCs.

25.7.8 Universal Applicability: From Leptons to Hadrons

This mechanical necessity applies universally. While a lepton's core may exhibit an asymmetric arrangement of Wave Centers and a hadron's core may be a complex multi-centered system, both are dictated into a spherical gravitational manifestation by the structural constraints of the vacuum units. The **Effective Volume Deficit** ($N_{\nu,\text{eff}}$) is the integrated result of this masking:

$$N_{\nu,\text{eff}} = \int_0^{r_\nu} 4\pi r^2 \rho_{\text{EMC}}(r) dr \tag{315}$$

where $\rho_{\text{EMC}}(r)$ represents the finalized spherical density gradient of the EMC packing. The far-field gravitation is not a signal emitted by the core, but a static structural deformation of the BCC substrate forced into symmetry by its own constituent geometry.

25.8 The Uniqueness of the BCC Topology: Why $N_{geometric} = 8\pi^4$ Excludes Other Lattices

The identification of the stiffness constant as $N_{geometric} = 8\pi^4$ serves as a selection rule that excludes other lattice geometries (such as FCC or SC) from being viable candidates for the vacuum substrate. This uniqueness is rooted in the interplay between the coordination number and the topological requirements of soliton stability.

25.8.1 The Coordination Number and Stiffness Distribution

In the $Im\bar{3}m$ (BCC) symmetry, each EMC unit possesses a coordination number of 8. The identity $N_{geometric} = 8\pi^4$ implies that the fundamental saturation budget π^4 is distributed precisely along the 8 primary axes of the unit cell.

- **FCC and SC Incompatibility:** In an Face-Centered Cubic (FCC) lattice (coordination 12) or Simple Cubic (SC) lattice (coordination 6), the resulting stiffness constants ($12\pi^4$ or $6\pi^4$) would lead to coupling strengths and a gravitational constant G that deviate from observed reality by orders of magnitude.
- **Structural Optimality:** Only the BCC coordination of 8 aligns with the internal energy density requirements of the π^4 Quark Stability budget.

25.8.2 Packing Fraction and the ζ Correction

The small residual correction ζ (where $N_{\text{geom}} = 8\pi^4(1-\zeta)$) is a direct consequence of the BCC packing fraction ($\eta_{\text{BCC}} \approx 0.68$). Unlike a hypothetical ideal continuum, the discrete BCC lattice is "near-optimal" but not perfect. The ζ factor represents the geometric impedance of a lattice that is 68% saturated, providing a physical basis for the minute deviations in the fine-structure constant and anomalous moments.

The magnitude of ζ is consistent with the per-node void impedance, normalized to the occupied fraction:

$$\zeta \sim \frac{1-\eta}{\eta \cdot N_{\text{ideal}}} = \frac{1-\frac{\sqrt{3}\pi}{8}}{\frac{\sqrt{3}\pi}{8} \cdot 8\pi^4} = \frac{8-\sqrt{3}\pi}{\sqrt{3}\pi \cdot 8\pi^4} \approx 6.0 \times 10^{-4}, \tag{316}$$

within 3.4% of the observed $\zeta \approx 5.8 \times 10^{-4}$. The normalization by η reflects that geometric impedance acts on the *occupied* sublattice, not the total volume.

25.8.3 Topological Consistency: The $2\pi^2$ Factor and BCC Axes

The consistency of the lepton generation numbers (K_μ, K_τ) further confirms the BCC choice. The topological invariant for higher generations involves the characteristic geometric factor $2\pi^2$ (which equals the surface area of a torus, but may also arise from other genus-1 configurations) and the 10^{n-1} scaling:

- **Resonance Matching:** The 8 primary directions of the BCC lattice provide the necessary degrees of freedom to support the $2\pi^2$ flux without introducing destructive interference in the medium.
- **Operator Universality:** The Isotropy Operator $\mathcal{I}$ achieves perfect spherical masking only in the BCC structure, where the equilateral distribution of the 8 nodes allows for the complete cancellation of shear stresses (σ_{shear}), a feat impossible in less symmetric or over-constrained lattices (like FCC).

This convergence identifies the BCC lattice as the **Unique Geometric Solution** capable of supporting the observed particle spectrum and interaction strengths. The vacuum is not merely a medium; it is a mathematically optimized 8-fold resonator.

25.9 Mechanical Resolution of Field Paradoxes

Through the formalism of *Spherical Masking*, a purely mechanical solution is provided for the long-standing paradox of field isotropy: how an asymmetric, rotating soliton (with non-spherical internal topology) manifests a perfectly isotropic gravitational field.

Two complementary mechanisms operate in concert:

- **Intrinsic Sphericity of the Standing Wave:** In any elastic 3D medium, a stable standing wave packet naturally adopts a spherical Bessel-like distribution to satisfy the principle of least action. This inherent tendency toward spherical symmetry operates independently of the discrete arrangement of Wave Centers, ensuring that the radiated field is already nearly isotropic before any lattice effects come into play.
- **Geometric Low-Pass Filtering (Lattice Enforcement):** The BCC lattice, composed of spherical EMC units, reinforces this natural sphericity by imposing a strict spherical boundary condition at the Degraded EMC Wall, where the dynamic wave regime transitions into static displacement. This eliminates any residual angular asymmetries that might survive from the near field. Moreover, any departure from spherical symmetry would introduce shear stresses (σ_{shear}) into the stiffness tensor $\mathcal{C}_{ijkl}$. The spherical geometry is the **Unique Optimal Operating Point** where the inter-EMC forces are purely compressive/extensional, minimizing internal shear stress and avoiding unstable energy dissipation.
- **Gravity as Restoring Pressure:** The derivation of gravity as a "Push-out" force is finalized. The constant G is revealed not as an independent coupling, but as a manifestation of the geometric density deficit, explaining its weakness as a holographic projection of 4D saturation into 3D space.

The separation between the asymmetric core and the isotropic gravitational field is formalized by the **Spherical Masking Condition**:

$$r_{\text{energy}} \leq r_{\text{EMC degraded wall}} \tag{317}$$

This condition states that the asymmetric energy core (r^5 domain) is contained within the radius of the Degraded EMC Wall (r^3 domain), enforcing gravitational isotropy through the structural constraints of the spherical vacuum units.

26 Nonlinear Stabilisation of the Electron Soliton

The geometric identities derived in Sections 11 and 2 determine the coupling constants of the vacuum, but they do not by themselves constitute a dynamical stability proof. A stable soliton requires a nonlinear term $\mathcal{F}$ in the wave equation that counteracts dispersion. The form of $\mathcal{F}$ is constrained by the BCC lattice geometry, the topological class of the electron, and the native 1-3-6 wave-centre arrangement inherited from the EWT standing-wave model.

26.1 General Soliton Wave Equation

The scalar amplitude $\Psi(\mathbf{r}, t)$ of the standing wave satisfies

$$\Big(\frac{\partial^2}{\partial t^2} - c^2\nabla^2\Big)\Psi(\mathbf{r}, t) + \mathcal{F}(\Psi) = 0. \tag{318}$$

The first two terms describe free wave propagation in the elastic BCC medium; the third term encodes the nonlinear self-interaction that makes the soliton possible.

Units convention. Throughout this section, the wave equation is written in the *natural units of the BCC lattice*. The lattice spacing λ_l, the lattice time step $t_p = \lambda_l/c$, and the lattice wave speed $c \equiv \lambda_l/t_p$ are all set to unity, and the wave amplitude Ψ is measured in units of the Planck charge amplitude q_P (which in the EWT framework has dimension of length; Section 5.1). With $c = 1$, the operator $(\partial_t^2 - c^2\nabla^2)$ reduces to $(\partial_t^2 - \nabla^2)$: the factor c disappears, because c has been absorbed into the unit system (Section 9, Eq. (90)). Every quantity in Eq. (318) — the field Ψ, the cubic coupling $\gamma = 1/\epsilon_M \approx 2.414 \times 10^4$, and the operator $(\partial_t^2 - \nabla^2)$ — is therefore a pure number, and the equation is *numerically homogeneous*. No dimensional discussion is required at the level of the natural-unit equation; the dimensional structure of the physical equation is analysed separately below (see Eq. (319)).

To translate Eq. (318) to SI units, one restores the lattice scales, $\Psi = q_P\varphi$, $t = t_p\tau$, $x = \lambda_l\xi$, and rescales the nonlinear coefficient accordingly:

$$\Big(\frac{\partial^2}{\partial t^2} - c^2\nabla^2\Big)\Psi_{\rm SI} - \gamma_{\rm SI}\Psi_{\rm SI}^3 = 0, \qquad \gamma_{\rm SI} = \frac{\gamma}{q_P^2\, t_p^2}. \tag{319}$$

Here $\Psi_{\rm SI}$ is the physical displacement amplitude of dimension $[L]$, and the SI coefficient $\gamma_{\rm SI}$ has dimension $[L^{-2}T^{-2}]$, consistent with the dimensional structure of the linear operator $(\partial_t^2 - c^2\nabla^2)$ acting on a length amplitude: both sides of Eq. (319) then carry dimension $[L/T^2]$. Numerically, with $q_P \approx 1.876 \times 10^{-18}$ m and $t_p = \lambda_l/c \approx 5.39 \times 10^{-44}$ s, the SI coefficient is $\gamma_{\rm SI} \approx 2.36 \times 10^{126}$ m^{-2} s^{-2}. The large numerical value is simply a consequence of the tiny lattice scales: the physically meaningful quantity is the dimensionless product $\gamma_{\rm SI}\Psi_0^2 t_p^2 = \gamma \approx 2.41 \times 10^4$, where $\Psi_0 = q_P$ is the characteristic amplitude.

All equations in Sections 26–31 are written in natural lattice units, unless explicitly stated otherwise.

26.2 Coupling Coefficient from BCC Geometry

The magnetic deficit ϵ_M is the fundamental lattice response parameter derived in Section 2.2:

$$\epsilon_M = \frac{1}{N_{\rm geom}\,\pi^3} = \frac{1}{8\pi^7(1-\zeta)}. \tag{320}$$

In the ideal continuum limit $\zeta \to 0$, this reduces to $\epsilon_M = 1/(8\pi^7)$. The approximate equality reflects the 0.058% lattice impedance δ between the calibrated stiffness $N_{\rm final}$ and the ideal geometric limit $8\pi^4$, as discussed in Section 21.1. Physically, ϵ_M measures the fractional loss of stiffness caused by the soliton's spin-induced magnetic torque. To eliminate any systemic ambiguity with the standard wave vector k, the natural nonlinear coupling strength is designated by the parameter γ, defined as the inverse of this deficit:

$$\gamma \equiv \frac{1}{\epsilon_M} = N_{\rm geom}\,\pi^3 \approx 2.414 \times 10^4. \tag{321}$$

This coefficient is not a free parameter; it follows from the BCC coordination number (8) and the dimensional budget of the charged weak scale (π^7).

For numerical implementation, two values of γ are available:

- $\gamma_{\rm geo} = 8\pi^7 \approx 2.4162 \times 10^4$ — the pure geometric limit corresponding to ideal BCC packing ($N_{\rm geo} = 8\pi^4$);

- $\gamma_{\text{final}} = N_{\text{final}}\pi^3 \approx 2.4148 \times 10^4$ — the value calibrated to the measured fine-structure constant, differing from γ_{geo} by the lattice impedance $\delta \approx 0.058\%$ (see Section 21.1).

The choice between them is not arbitrary: the 0.058% difference encodes the non-ideal spherical packing fraction of the physical BCC lattice, and it may prove decisive for discriminating $K = 10$ from neighbouring configurations in a numerical simulation. The OpenWave implementation should initially use γ_{final} (anchored to the measured α) and, if necessary, explore the geometric limit as a consistency check.

26.3 Proposed Forms of the Nonlinear Term

Three complementary variants are proposed, ordered by increasing structural fidelity to the EWT 1-3-6 architecture.

Variant A — Pure cubic nonlinearity

The simplest form compatible with NLS-type soliton equations is the cubic (self-focusing) nonlinearity:

$$\boxed{\mathcal{F}(\Psi) = -\gamma\,\Psi^3 = -\frac{1}{\epsilon_M}\,\Psi^3 = -N_{\text{final}}\,\pi^3\,\Psi^3}. \tag{322}$$

A term proportional to $-\Psi^3$ (for $\gamma > 0$) is the lowest-order odd nonlinearity that produces self-focusing; it corresponds to the inverted quartic potential $V(\Psi) = -\frac{1}{4}\gamma\Psi^4$. As is well known from the theory of nonlinear Schrödinger equations, the pure cubic nonlinearity is subject to Derrick-type scaling instability in three spatial dimensions; the additional structure of Variants B and C is introduced to circumvent this instability.

Variant B — Nonlinearity modulated by EMC packing density

In the EWT framework the nonlinear stiffness should be active only where the granular (EMC) density is depleted relative to the statutory vacuum background ρ_0. Let $\rho(r)$ be the local EMC density inside the soliton; the deficit fraction is $(1 - \rho(r)/\rho_0)$. The nonlinear term then becomes

$$\boxed{\mathcal{F}(\Psi, \rho) = -\gamma\left(1 - \frac{\rho(r)}{\rho_0}\right)\Psi^3}. \tag{323}$$

Equation (323) directly couples the wave equation to the push-out mechanism (Section 11.10): the self-focusing nonlinearity is maximal in the soliton core where $\rho(r) \ll \rho_0$, and vanishes in the surrounding statutory vacuum.

Variant C — Explicit 1-3-6 source-driven dynamics

In the native EWT model the wave centres (WCs) are not an abstract continuum but ten discrete, mobile sources arranged in the 1-3-6 configuration. The vector wave equation of M4 then takes the driven form

$$\boxed{\left(\frac{\partial^2}{\partial t^2} - c^2\nabla^2\right)\boldsymbol{\Psi}(\mathbf{r},t) - \gamma\,\|\boldsymbol{\Psi}\|^2\,\boldsymbol{\Psi} = \mathbf{J}_{\text{1-3-6}}(\mathbf{r},t)}, \tag{324}$$

where the source operator $\mathbf{J}_{\text{1-3-6}}$ is the sum of ten vectorial centres partitioned into three topological classes:

$$\mathbf{J}_{\text{1-3-6}}(\mathbf{r},t) = \mathbf{j}_{\text{core}}(\mathbf{r} - \mathbf{r}_0) + \sum_{i=1}^{3}\mathbf{j}_{\text{inner}}(\mathbf{r} - \mathbf{r}_i(t)) + \sum_{j=1}^{6}\mathbf{j}_{\text{outer}}(\mathbf{r} - \mathbf{r}_j(t)). \tag{325}$$

Each class carries a distinct phase offset, and the kinematic rule inherited from the Yee standing-wave model applies: every centre migrates toward the local minimum of the field amplitude $\|\mathbf{\Psi}\|$. The phase asymmetry between the inner (3) and outer (6) groups forces a synchronised rotation of the vertices, generating the 720° spin characteristic of the electron.

26.4 Structural Emergence of the 1-3-6 Geometry

Why the 1-3-6 arrangement, and not another partition of ten? The answer follows from the interplay of the nonlinearity and the BCC lattice topology:

1. The **core** (1) anchors the soliton at a single BCC lattice node, providing the central phase reference.

2. The **inner shell** (3) forms an equilateral triangle. Three is the minimal number of vertices that can define a plane; the plane's normal coincides with the spin axis. Any other number would either fail to define a unique axis (1 or 2) or, it is proposed, introduce internal phase frustration (4 or more on a single shell at this radius) — a physical claim whose verification requires the same numerical proof as the rest of the stability argument.

3. The **outer shell** (6) completes the octahedral coordination of the BCC unit cell. Six wave centres placed at the face-centre-equivalent positions are proposed to saturate the remaining propagation axes, thereby guaranteeing isotropic far-field emission after spherical masking by the Degraded EMC Wall (Section 25.7) — a claim that, like the others in this section, requires numerical verification.

It is proposed that a configuration such as 2-3-5 or 2-2-6 would either misalign the spin axis or leave some BCC axes unmatched, producing a net multipole moment that the nonlinearity cannot cancel — a prediction to be tested in the OpenWave M4 simulation. The 1-3-6 triad is therefore the unique self-consistent candidate that simultaneously satisfies: (a) the topological winding number $Q = 10$, (b) the cubic nonlinearity with coefficient γ, and (c) the octahedral symmetry of the $Im\bar{3}m$ vacuum lattice. A rigorous proof that this configuration indeed minimises the energy functional constructed from $\gamma\Psi^3$ and the $Im\bar{3}m$ projection remains a task for the OpenWave M4 simulation.

26.5 Topological Selection Rule and the Electron Ground State

The nonlinear wave equation alone does not select a specific number of wave centres. The selection of $K_{\mathrm{WC}} = 10$ is enforced by the topological class of the soliton (Section 19.4). On the boundary sphere S^2 that encloses the soliton, the configuration is described by a unit vector field $\mathbf{n} : S^2 \to S^2$ (the local spin axis of the wave centres, or equivalently the local wavefront normal), and the topological invariant is the integer degree of this map (Eq. 301):

$$Q = \frac{1}{4\pi} \int_{S^2} \mathbf{n} \cdot \left(\frac{\partial \mathbf{n}}{\partial \theta} \times \frac{\partial \mathbf{n}}{\partial \phi} \right) d\theta \, d\phi \; \in \; \mathbb{Z}. \tag{326}$$

For the electron, the 1-3-6 configuration is assigned $Q = K_{\mathrm{WC}} = 10$ (Eq. 302). Because Q is a topological invariant, it cannot change under continuous deformations of the wave field; a transition to $Q = 9$ or $Q = 11$ would require a discontinuous rupture of the elastic medium and is therefore energetically forbidden. The soliton is protected against perturbations that would alter its internal multiplicity.

This topological winding number is not an empirical adjustment; it emerges from the discrete projection of the BCC lattice's $Im\bar{3}m$ space group symmetry onto the continuous spherical boundary S^2. While the 8 primary propagation axes define the innermost volumetric kernel, the configuration of $Q = 10$ wave centres is a natural candidate for the lowest-energy, stable homotopy class capable of establishing a fully isotropic boundary condition on S^2 (a problem analogous to the global minimum of the spherical Thomson packing problem for discrete phase nodes). A rigorous proof that $Q = 10$ yields a deeper potential well than $Q = 8, 9, 11$ requires an explicit evaluation of the energy functional built from the nonlinear term $\gamma\Psi^3$ and the $Im\bar{3}m$ projection, which remains a task for the OpenWave M4 simulation.

Because Q is a topological invariant, it cannot change under continuous deformations of the wave field; a transition to $Q = 9$ or $Q = 8$ would require a discontinuous rupture of the elastic medium, requiring infinite energy. The soliton is therefore permanently protected against perturbations that would alter its internal multiplicity.

Equation (318) together with any of the three nonlinear variants and the topological condition $Q = 10$ constitutes a well-posed, falsifiable model of the electron core. The stability of the 1-3-6 configuration within this model is at present a postulate, not a proof; its numerical verification is the immediate objective of the ongoing OpenWave M4 implementation described in Section 31.

1. The **nonlinearity** (γ) creates a deep, narrow potential well that resists dispersion.
2. The **topological winding number** is proposed to fix the ground state at exactly ten wave centres, pending numerical confirmation that $Q = 10$ is energetically preferred over neighbouring integers $Q = 8, 9, 11$ within the $Im\bar{3}m$ projection.
3. The **BCC stiffness constant** N_{final} (or its geometric limit $8\pi^4$) fixes the numerical value of the coupling without adjustable parameters.

26.6 Connection to the Dimensional Weight Operator

The Dimensional Weight Operator $\hat{W}(n)$ (Section 25.4) assigns a distinct degree of freedom to each factor of π in the coupling budget. The nonlinear coefficient (321) can be written as

$$\gamma = \frac{1}{\epsilon_M} = N_{\text{final}}\,\pi^3 \;\approx\; 8\pi^7 = 8 \;\cdot\; \pi^{(1)}\pi^{(2)}\pi^{(3)}\pi^{(4)}\pi^{(5)}\pi^{(6)}\pi^{(7)}, \tag{327}$$

where the seven factors correspond to the seven degrees of freedom of the charged weak scale (Table 27). The approximate equality with $8\pi^7$ reflects the geometric limit γ_{geo}, while the exact factorisation in terms of labelled $\pi^{(k)}$ degrees of freedom is a formal bookkeeping device that holds irrespective of the numerical value of the prefactor. Thus the stabilisation of the electron is not an isolated electromagnetic phenomenon but a direct mechanical consequence of the full 7-dimensional phase space that the BCC lattice makes available to a charged soliton. The factor 8 reflects the eight primary propagation axes of the $Im\bar{3}m$ unit cell, ensuring that the nonlinear restoring force acts isotropically.

26.7 Geometric Estimate of the Degraded EMC Wall Radius

The nonlinear stabilisation described above depends on the extent of the EMC deficit region, which is bounded by the Degraded EMC Wall (Section 25.7). A natural length scale for the wall radius r_{wall} is provided by the electron Compton wavelength λ_C, whose geometric derivation is given in Section 23.3 (see Eq. 261). Numerically $\lambda_C \approx 2.426 \times 10^{-12}$ m. Since the classical

electron radius is $r_e = \lambda_C \cdot \alpha/2\pi$, the reduced Compton wavelength $\lambda_C/2\pi$ is approximately $r_e/\alpha \approx 137\, r_e$. A plausible scale is

$$\boxed{r_{\text{wall}} \sim \frac{\lambda_C}{2\pi} = \frac{r_e}{\alpha}}. \tag{328}$$

Because α is itself derived from the BCC lattice geometry (Eq. 18), the relation $r_{\text{wall}} \sim r_e/\alpha$ does not introduce a new parameter; it is algebraically equivalent to the standard QED definition of the classical electron radius. The significance of this estimate lies not in its novelty but in its consistency: the same geometric framework that yields α and λ_C from r_ν and the $8\pi^7$ correction also predicts a natural length scale for the region in which the nonlinearity is active, providing a unified description of particle stability, forces, and annihilation. The considerable width of the wall — roughly $137\, r_e$ — is not an arbitrary feature but a geometric consequence of the ratio $1/\alpha \approx 137$. Physically, this extended transition zone provides a broad buffer within which the nonlinearity can adiabatically relax the EMC density from its deep core deficit to the statutory background, suppressing the sharp gradients that would otherwise destabilise the standing wave.

Role of a Possible EMC Density Peak

The preceding estimate fixes the radial scale of the wall, but leaves open the question of its internal density profile. As noted in Section 11.11, two scenarios are compatible with the integral deficit $N_{\nu,\text{eff}}$: a monotonic approach to the statutory background, or a local density peak ($\rho_{\text{wall}} > \rho_0$) caused by the accumulation of EMCs expelled by the push-out mechanism. If such a peak exists, it would actively assist the nonlinear stabilisation in two ways. First, in Variant B the factor $(1 - \rho/\rho_0)$ would change sign within the peak, turning the self-focusing nonlinearity into a local defocusing barrier that prevents the standing wave from leaking outward. Second, even for the unmodulated Variants A and C, a density peak implies a locally higher elastic modulus, which acts as a partial reflector, increasing the Q-factor of the soliton resonator. Whether the peak is indispensable or merely auxiliary can only be decided once the full radial EMC profile is computed from the nonlinear wave equation, which remains a task for the OpenWave M4 simulation.

Internal Wave-Centre Dynamics and Spin

In the foundational Energy Wave Theory (Yee, 2019), an additional hypothesis was put forward that could further clarify the electron's stability: the electron's ten wave centres are not perfectly static but undergo continuous micro-motion around their nodal positions. According to this picture, a wave centre displaced from its node experiences a restoring force toward the local amplitude minimum; this perpetual shifting of centres is the origin of the 720° spin and, at the same time, a dynamic stabilisation mechanism. The tetrahedral 1-3-6 geometry ensures that only a fraction of the centres are off-node at any instant, while the remainder anchor the structure. Configurations such as $K_{WC} = 9$ or $K_{WC} = 11$ lack this balance and break apart. This hypothesis has not yet been proven mathematically and should be tested in the OpenWave M4 model once the nonlinear term and the Degraded EMC Wall are in place: comparing simulations with and without WC motion could reveal whether the internal dynamics is an essential ingredient or a secondary effect. In the language of the Dimensional Weight Operator (Section 25.4), this internal dynamics may be tentatively associated with the degree of freedom $\pi^{(6)}$ (soliton radius), which would thereby acquire a dynamic interpretation beyond its static geometric extent — a reinterpretation that remains speculative, is not used in deriving any numerical result in this paper, and applies to this one degree of freedom only.

26.8 Relation Between the Energetic and Topological Arguments

The present section advances two complementary conditions that together single out $K_{WC} = 10$: (i) an **energetic** condition based on the r^5/r^3 scaling disparity, which creates a deep potential well for intermediate wave-centre counts (Section 15.1); and (ii) a **topological** condition requiring an integer winding number Q on S^2 (Section 19.4). These are not independent post-hoc justifications of the same number; they are mutually reinforcing constraints. The r^5/r^3 argument explains why K_{WC} cannot be too small (shallow well) or too large (volumetric overload), while the topological argument explains why, within the allowed range, only discrete integer values are permitted. The specific value $K_{WC} = 10$ emerges as the unique integer that simultaneously satisfies both constraints under the octahedral symmetry of the $Im\bar{3}m$ BCC lattice. A rigorous proof that no other integer passes both filters remains a task for the OpenWave M4 simulation.

26.9 Summary

- The general soliton wave equation is $(\partial_t^2 - c^2\nabla^2)\Psi + \mathcal{F}(\Psi) = 0$.
- The coupling coefficient is fixed by BCC geometry: $\gamma = 1/\epsilon_M = N_{\text{final}}\pi^3 \approx 2.41 \times 10^4$. Two variants are available: γ_{final} (anchored to α) and $\gamma_{\text{geo}} = 8\pi^7$ (ideal BCC limit); the 0.058% difference may be critical for K-selectivity.
- Three complementary forms for $\mathcal{F}$ are proposed:

Variant A: $-\gamma\Psi^3$ (pure cubic self-focusing, minimal implementation);

Variant B: $-\gamma(1-\rho/\rho_0)\Psi^3$ (density-modulated self-focusing, links stability to the EMC push-out deficit);

Variant C: $-\gamma\|\mathbf{\Psi}\|^2\mathbf{\Psi}$ with an explicit 1-3-6 source operator $\mathbf{J}_{\text{1-3-6}}(\mathbf{r}, t)$ (directly encodes the discrete wave-centre architecture of the electron).

- The Degraded EMC Wall is proposed to provide the necessary boundary condition for the nonlinearity: its radius, estimated as $r_{\text{wall}} \sim r_e/\alpha$, is consistent with the Compton wavelength and may ensure a broad adiabatic transition zone ($\sim 137\, r_e$) in which the density-modulated term can act without sharp gradients. A possible density peak within the wall would further assist stabilisation by creating a local defocusing barrier.
- The winding number $Q = 10$ on S^2 is proposed as the topological selection rule that would isolate the electron ground state under $Im\bar{3}m$ constraints; rigorous proof that $Q = 10$ is energetically preferred over other integers remains a task for the OpenWave M4 simulation.
- The 1-3-6 partition is the unique self-consistent candidate that simultaneously satisfies the topological winding, the cubic nonlinearity, and the octahedral symmetry of the BCC lattice; its stability is a postulate pending numerical verification.
- The energetic (r^5/r^3) and topological arguments are complementary: the former restricts the range of allowed K_{WC}, the latter restricts K_{WC} to integer values; together they single out $K_{WC} = 10$ as the only candidate satisfying both constraints.
- The coupling $\gamma \approx 8\pi^7$ (in its geometric limit) directly expresses the 7-dimensional charged-weak budget of the BCC lattice.
- The model is well-posed and falsifiable; the stability of the 1-3-6 configuration is a postulate whose numerical verification is the immediate objective of the OpenWave M4 simulation.

27 Predictive Power

The EWT model yields several precise, quantitative predictions that follow directly from the geometry of the Soliton and the BCC lattice. The results summarized in Table 28 demonstrate the internal consistency of the zero-calibration framework.

Table 28: Principal Geometric Predictions of the EWT Framework

Physical Quantity	**Geometric Prediction**	**Target**	**Relative Error**
G_{geom}	$6.6775199755 \times 10^{-11}$	6.674305×10^{-11}	0.048169%
$\alpha_{\text{geom}}^{-1}$	137.036262364	137.035999084	0.000192%
a_e^{EWT}	1159.916228 ppm	1159.652182 ppm	0.022769%
a_μ^{EWT}	1166.212608 ppm	1165.920610 ppm	0.025044%
a_τ^{EWT}	1176.838130 ppm	1177.210000 ppm	0.031589%

Note: Values for a_e, a_μ, and a_τ are full anomalous magnetic moments. All predictions are obtained from the geometric Python implementation described in the Appendix. Error for a_τ is compared with SM prediction.

27.1 Prediction of the Fundamental Value of G

The model predicts the value of G as a direct function of microscopic geometric parameters through the operator $\mathcal{U}$ (Section 11.5). For the zero-calibration inputs used in this work, the geometric identity gives

$$G_{\text{geom}} \approx 6.6775199755 \times 10^{-11}\ \text{m}^3\,\text{kg}^{-1}\,\text{s}^{-2}. \tag{329}$$

Precise experimental measurements of G (e.g., torsion balance, atom interferometry) directly test this prediction.

27.2 Predictions for Lepton Properties: The First-Principles AMM Predictions

The full anomalous magnetic moments of the entire lepton family are computed from the same BCC lattice geometry that governs G and α_{geom}. The results form a compact monotonic sequence:

$$\boxed{a_e^{\text{EWT}} = 1.159916 \times 10^{-3}} \qquad \boxed{a_\mu^{\text{EWT}} = 1.166213 \times 10^{-3}} \qquad \boxed{a_\tau^{\text{EWT}} = 1.176838 \times 10^{-3}} \tag{330}$$

with relative errors 0.022769%, 0.025044%, and 0.031589%.

The projection rules $\mathcal{O}_e = \mathcal{O}_\tau = 1$ and $\mathcal{O}_\mu = 1/(4\pi^2)$ are fixed by the dimensional hierarchy and contain no generation-specific parameters. A fourth generation would require a second 2D projection and is geometrically forbidden by the saturation limit of the BCC lattice.

27.3 Cosmological Predictions: Recovered Relation and Predicted Constants

The cosmological sector (Section 24) extends the same geometric framework to the large-scale vacuum. The results fall into four categories, summarised in Table 29.

Table 29: Cosmological results by status. The first column gives the symbol, the second the value, the third the status (in the sense used throughout this work), and the fourth the source or key assumption.

Quantity	Value	Status	Source / assumption
Haug relation	$T_{\mathrm{cmb},0} = \dfrac{T_p}{8\pi}\sqrt{\dfrac{2\lambda_l}{R_H}}$	**Recovered**	From naive equipartition corrected by $H = 8(2\pi)^2$; phase-unit ansatz is a model assumption
H_0^{EWT}	$66.9113\ \mathrm{km\,s^{-1}\,Mpc^{-1}}$	**Predicted**	BCC chain + one anchor $T_{\mathrm{cmb},0}$; -0.83σ vs Planck 2018
Λ_{dS}	$1.5696 \times 10^{-52}\ \mathrm{m^{-2}}$	**Predicted**	$3H_0^2/c^2$; $1.441\times$ the observed Λ
Ω_{vac}	0.6897	**Mechanism**	$\eta_{\mathrm{BCC}} C_{\mathrm{gap}}$; plays the role of Ω_Λ; $+0.15\sigma$ vs Planck+BAO; composition rule is a model assumption
Λ_{vac}	$1.0826 \times 10^{-52}\ \mathrm{m^{-2}}$	**Consequence**	$\Omega_{\mathrm{vac}}\Lambda_{\mathrm{dS}}$; -0.59% vs Planck; inherits the assumption on Ω_{vac}

The two *predictions* are directly testable: they depend on the BCC geometry and the single anchor $T_{\mathrm{cmb},0}$, with no calibrated parameter. The *recovered* Haug relation is a reformulation of the equipartition scaling; the horizon factor $H = 8(2\pi)^2$ is a model construct, not a first-principles derivation. The *mechanism* $\Omega_{\mathrm{vac}} = \eta_{\mathrm{BCC}} C_{\mathrm{gap}}$ and its *consequence* Λ_{vac} are conditional on the composition rule, which is stated as a model assumption in Section 24.7.

As with the lepton AMM predictions, none of these results depends on the numerical value of G_{geom} separately: the factors $\hbar_{\mathrm{geom}}$ and G_{geom} cancel in the combinations that fix T_p and λ_l.

27.4 Magnetic Sensitivity $G(B)$ and Hypotheses Testing

The model predicts a dependence of the effective gravitational coupling on local magnetic coherence, as described in Section 18. In the zero-calibration formulation, the push-out operator is expressed through the geometric stiffness:

$$\mathbf{T}_{\mathrm{II}} = \left(\frac{1}{A_\pi \cdot N_{\mathrm{geom}}} \right)^3 . \tag{331}$$

The observable shift is expected to be small in ordinary environments:

$$\left(\frac{\Delta G}{G} \right)_{\mathrm{obs}} = \xi(B) \cdot \mathbf{T}_{\mathrm{II}}, \tag{332}$$

where $\xi(B)$ is the coupling efficiency of the local magnetic field to the soliton's geometry. The sign of the effect distinguishes the hypotheses of Section 18.4.

27.5 AMM as a Strong Test of Geometric Universality

Because AMM measurements are exceptionally precise, they provide an ideal testing ground for the geometric mechanism. A single universal modulator ϵ_M, together with the recursive nodal structure, replaces the perturbative loop corrections of the Standard Model.

27.6 Correlations with Neutrino Flux

Since G emerges from the effective density $N_{\nu,\text{eff}}$, the model allows minute correlations between measured G and local neutrino flux densities. High neutrino flux may momentarily alter the local EMC background, leading to sub-ppm modulations of G.

27.7 Spectral and Geometric Modal Tests

The modal nature of the π^3 factor implies that internal geometric excitations may leave traces in the frequency spectrum of small fluctuations of G. The model predicts characteristic nodal frequencies of the soliton's standing modes.

27.8 Tests using Gravitational Wave Observations

If G is emergent from the BCC lattice, gravitational waves may exhibit small dispersion effects:

$$v_{\text{GW}} \simeq c\,(1 - \delta_{\text{dispersion}})\,, \tag{333}$$

where $\delta_{\text{dispersion}}$ depends on the local nodal density. Arrival-time comparisons between GW and electromagnetic signals constrain this effect.

27.9 Test on Bose-Einstein Condensates

A BEC state synchronises the local magnetic coherence across a macroscopic volume. The model predicts a shift

$$G_{\text{eff}}^{(\text{BEC})} = G\,(1 + \delta_{\text{BEC}})\,, \qquad \delta_{\text{BEC}} \propto \mathbf{T}_{\text{II}}. \tag{334}$$

Consistent with the push-out mechanism, the primary prediction is $G_{\text{eff}} > G$. Atom interferometry on ultra-cold clouds before and after BEC formation can test this directly.

27.10 The Higgs Soliton vs. Fundamental Scalar Field

The EWT model identifies the Higgs as a resonant state ($K_{WC} = 117$) rather than a fundamental scalar field. This leads to specific predictions for Higgs-vector mixing:

$$\sin^2\theta_{ZH} \approx 0.4686, \qquad \sin^2\theta_{WH} \approx 0.5906. \tag{335}$$

28 Falsifiability and Model Constraints

The following outcomes represent critical tests that can either **falsify the core tenets of the EWT model** or place stringent constraints on specific parameters and hypotheses. These tests are focused on the **existence or non-existence** of predicted effects, independent of the sign of the dynamic coupling (Hypotheses 1 and 2).

- **Falsification of Fundamental Constants (G):** Precise measurements of G consistently rejecting G_{model} (Eq. 27.1) beyond the level of uncertainty while maintaining the assumed values of microscopic parameters.

- **Falsification of Dynamic Coupling ($G(B)$):** The observation of a null effect, i.e., the absence of any observed $G(B)$ dependence above the detection threshold, assuming the technical feasibility of such a test. This outcome would support **Hypothesis 3 (No Influence)** and refute the core idea of magneto-geometric coupling.

- **Falsification of Correlations:** The absence of statistical correlations between fluctuations in G and external factors (neutrino flux, magnetic fields) within the range predicted by the model (ref. Section 27.6).

- **Falsification of Full Lepton AMM Predictions (a_e, a_μ, a_τ):** Future high-precision measurements of the electron, and muon, anomalous magnetic moments that deviate significantly from the full AMM predictions $a_e^{\text{EWT}} = 1.159916 \times 10^{-3}$, $a_\mu^{\text{EWT}} = 1.166213 \times 10^{-3}$, and $a_\tau^{\text{EWT}} = 1.176838 \times 10^{-3}$ (Table 11) would falsify the model in its present formulation. For the tau, the current reference value is the Standard Model prediction $a_\tau^{\text{SM}} = 1.177210 \times 10^{-3}$, not a direct measurement. Therefore the tau comparison is a consistency test against the SM prediction, not a direct experimental falsification. The projection rules underlying these predictions are completely determined by the dimensional hierarchy of the Geometric Ladder: $\mathcal{O}_e = 1$ (3D core), $\mathcal{O}_\mu = 1/(4\pi^2)$ (2D planar shell), and $\mathcal{O}_\tau = 1$ (3D volumetric shell). A significant deviation of any of the three measured anomalies from the predicted values would therefore directly challenge the geometric core of the model. The compact monotonic sequence of prediction errors—$0.022769\% \to 0.025044\% \to 0.031589\%$—provides a stringent, correlated test: any future measurement that breaks this monotonicity would also indicate a structural tension with the BCC lattice framework.

28.1 Practical Considerations and Detection Limits

In practice, the expected magnitude of the relative change $\Delta G/G$ is governed by the robustness of the dynamic geometric equilibrium between energy scaling (r^5) and volumetric displacement (r^3). Due to this compensatory mechanism, the predicted deviations in near-equilibrium (low-field) environments are expected to be extremely subtle, likely requiring a detection sensitivity in the range of 10^{-9} **to** 10^{-12}.

While these requirements are stringent, they align precisely with the current state-of-the-art in quantum metrology and atom interferometry. Modern platforms, such as MAGIS-100 or advanced gravity gradiometers, are specifically designed to operate within these detection limits to probe dark matter and gravitational waves.

The model further posits that the detection threshold is fundamentally tied to the **magnetic flux density** or **quantum coherence level** of the source. In extreme high-field regimes or coherent macroscopic states (such as BEC), the compensatory mechanism is hypothesized to reach its non-linear limit, potentially shifting the magnitude of $\Delta G/G$ towards the 10^{-7} range. Consequently, even a null result within current experimental sensitivities would provide critical boundary values for the volumetric "stiffness" of the EMC mediu governed by $\mathbf{T}_{\text{II}}$ and the saturation point of the geometric equilibrium. This ensures that the framework remains fully falsifiable, providing a clear mapping for the transition from the "low-field artefact" of invariant G to the dynamic G_{eff} regime predicted by EWT.

28.2 Two Readings of the Higgs Sector: What a Future Measurement Can Settle

The Higgs-sector tests proposed in Section 3.4 rest on a specific question: which of two readings of the Higgs sector is correct. The two readings make different commitments about the nature of the Standard Model, and stating those commitments explicitly clarifies what the proposed measurements can and cannot settle.

The three Standard-Model theses

The interpretation of the electroweak sector commonly presented as the Standard Model rests on three theses about the Higgs:

1. **The Higgs field is fundamental.** A scalar $SU(2)$ doublet Φ with hypercharge $1/2$ transforms under the gauge group as a matter field and carries no substructure at any energy scale.
2. **The vacuum expectation value is a fundamental constant.** The value $v = (\sqrt{2}\,G_F)^{-1/2} \approx 246$ GeV is a property of the vacuum of the underlying theory, fixed by the parameters of the Lagrangian and the renormalisation condition.
3. **The Higgs mechanism is the fundamental source of mass.** The masses of the W and Z bosons and of the charged leptons arise from the coupling of those fields to Φ. No deeper mechanism contributes to the values of those masses at any order.

The electroweak Lagrangian, understood as a quantum field theory, is one formal realisation of these theses. The theses are not logically equivalent to the formalism. Each can be denied while the formalism continues to describe the same low-energy physics, at least as an effective field theory.

What Energy Wave Theory claims instead

The model presented in this work replaces the three theses as follows:

1. The Higgs boson is a composite soliton of the BCC lattice with wave-centre count $K_{WC} = 117$. The field Φ is the low-energy projection of the collective degrees of freedom of that soliton, not a fundamental object.
2. The value $v \approx 246$ GeV is emergent from the same lattice geometry that determines α, the lepton anomalous magnetic moments, and the gravitational coupling G. It is not a free parameter of the vacuum.
3. The masses of the W and Z bosons and of the charged leptons are derived from the geometric parameters of the soliton states (the wave-centre count). The Higgs mechanism is not required to generate these masses.

What the proposed measurements can settle

The two readings coincide at the level of the tree-level HZZ and HWW couplings, under the restricted hypothesis of Eq. (43). They differ in the presence or absence of internal structure at a characteristic length scale, which manifests as a momentum-dependent deformation of the HZZ vertex at energies comparable to the inverse of that scale. The explicit form of the deformation is not derived in this work; it awaits the soliton form factor calculation.

The observables proposed in Section 3.4 test precisely this difference. Pillar I is a null test on the absolute ZH production rate: a statistically significant deviation from Eq. (44), after the specified corrections, falsifies the restricted hypothesis. Pillar II is a search for a momentum-dependent or tensorial deformation of the HZZ vertex in the Legendre moments of the Z production angle. Two outcomes are possible:

- **The Standard-Model theses hold.** A null result on the angular moments, at a precision sufficient to exclude any deformation of the magnitude that a composite soliton of the known internal size would produce, confirms the three theses at the tested scale. It leaves the Energy Wave Theory reading constrained but not excluded, since the EWT angular prediction is not currently quantified.

- **The Standard-Model theses fail.** A statistically significant deformation of the angular moments, consistent with a soliton-derived form factor, falsifies the three theses at the tested scale. It does not falsify the Standard Model as an effective field theory: the electroweak sector continues to describe physics below the compositeness scale, with the soliton replacing the scalar field at higher energies. What is falsified is the interpretation of the electroweak Lagrangian as a fundamental description of nature.

The distinction between falsifying the Lagrangian and falsifying its interpretation is central to the logic of this subsection. The Standard Model has been reinterpreted many times in response to new observations, from the discovery of the muon to the discovery of neutrino masses. The three theses above are the current interpretation of the Higgs sector. If they are wrong, the correct response is a reinterpretation, not a discarding of the framework. This is what the Energy Wave Theory proposes, and the measurements proposed in Section 3.4 are the way to decide between the two readings.

29 Robustness Analysis: Geometric Stability and Sensitivity

To demonstrate that the Enhanced EWT model is not a result of numerical coincidence, a series of sensitivity tests were performed. This analysis examines how small perturbations in the fundamental geometric inputs affect the physical outputs. By isolating each primary variable, the structural rigidity of the vacuum lattice is verified against experimental CODATA/PDG benchmarks.

29.1 Robustness Analysis of the Neutrino Soliton Radius and Geometric Identity

The structural integrity of the EWT framework is evaluated through the stability of the fundamental identity

$$\Delta N_\nu \sim \frac{1}{\epsilon_G}. \tag{336}$$

where ϵ_G is the gravitational deficit associated with the reference soliton. This equivalence suggests that the statutory constituent count, derived from the ratio of the neutrino radius to the Planck scale, must inherently align with the geometric deficit required to scale gravity from its classical base G_{Base} to the observed value.

To verify the absence of numerical fine-tuning, a sensitivity analysis was conducted by applying relative perturbations of $\pm 20\%$ to the primary geometric inputs: the statutory neutrino

radius r_ν and the fundamental EMC spacing λ_l. The resulting compliance curves are shown in Fig. 8. They demonstrate that even under significant variations in the input scales, the compliance ratios remain strictly within the $\pm 30\%$ tolerance boundaries.




Figure 8: **Structural Robustness Analysis of the Statutory Density $N_{\nu,\mathbf{stat}}$.** The plot tracks the stability of the statutory background population $N_{\nu,\text{stat}}$ against relative fluctuations in the neutrino radius r_ν and the EMC spacing λ_l. Both curves remain within the established $\pm 30\%$ compliance boundaries (0.7 to 1.3 ratio). This stability ensures that the geometric vacuum parameters do not require fine-tuning.

This performance confirms that the EWT model is not a product of precise numerical coincidence, but is instead rooted in a robust geometric power-law relationship.

29.2 Gravitational Surface Transition and $N_{\nu,\mathbf{eff}}$

The robustness test focuses on the relationship between the geometric volume deficit and the emergent gravitational constant G. As detailed in Section 11.10, gravity is interpreted as a surface effect resulting from a three-dimensional packing deficit within the EMC medium.

The results illustrated in Fig. 9 confirm the following physical constraints:

- **Transition trajectory:** The evolution of the curve follows the $G \propto N_\nu^{-1/2}$ scaling law. This confirms the gravitational interaction as a surface manifestation of the internal volumetric dilution.
- **Deterministic convergence:** The intersection with the G_{CODATA} threshold justifies the transition to the effective volume deficit

$$N_{\nu,\text{eff}} \approx 6.2472 \times 10^{48}, \tag{337}$$




Figure 9: Analysis of the G-constant surface transition. The green curve illustrates the evolution of G as a function of the effective volume deficit N_ν. The red intersection point marks the deterministic alignment with G_{CODATA} at the calculated $N_{\nu,\text{eff}}$.

obtained from the geometric dilution factor X_{eff}.

- **Model rigidity:** The steep slope of the transition curve indicates that the system is highly sensitive to the internal geometry. A deviation of even 1% in the value of N_ν results in a significant departure from G, proving that the EWT unification is not a result of arbitrary curve-fitting.

29.3 Sensitivity of α^{-1} to the Geometric Modulator N_{geom}

This test evaluates the stability of the fine-structure constant derivation by examining the influence of the dimensionless geometric stiffness N_{geom}. In the EWT framework, $\alpha^{-1}_{\text{geom}}$ is determined by the fixed vacuum geometry A_π modulated by the magnetic deficit $\epsilon_M = 1/(N_{\text{geom}}\pi^3)$.

The analysis of the results presented in Fig. 10 confirms:

- **Geometric necessity:** Precise alignment with $\alpha^{-1}_{\text{CODATA}}$ occurs only when the complete stability factor is applied. The "golden point" intersection at

$$N_{\text{geom}} \approx 778.8025 \tag{338}$$

proves that the fine-structure constant emerges from a rigid geometric base corrected by a finite magnetic deficit.




Figure 10: Sensitivity analysis of the inverse fine-structure constant. The plot illustrates the convergence of the model toward the CODATA value as a function of the geometric stiffness coefficient N_{geom}.

- **Force hierarchy indication:** While the modulator ϵ_M is shared by both G and α^{-1}, the gravitational constant operates on a significantly different scale through the cubic factor $(\epsilon_M/A_\pi)^3$ and the surface projection $N_{\nu,\text{eff}}^{-1/2}$.

29.4 Phase Space Analysis: EWT Trajectory vs Standard Model Interpretation

The phase-space mapping between the inverse fine-structure constant α^{-1} and the electron anomalous magnetic moment a_e reveals a fundamental geometric trajectory. By varying the vacuum stiffness parameter N, the model demonstrates that α^{-1} and a_e are not independent constants, but emergent properties of the same underlying geometric modulator.

The experimental CODATA point lies close to this EWT trajectory, as shown in Fig. 11, with a residual offset of approximately 0.022769%. In the EWT framework, this offset is not interpreted as model failure, but as a consequence of the structural difference between the geometric vacuum picture and the point-like renormalisation programme of the Standard Model.

29.5 Parametric Unification: The Stability Path of G and α

A parametric analysis of the relationship between G and α^{-1} was performed by varying the geometric modulator N. The resulting trajectory, derived from the geometric operator $\mathcal{U}$, shows that α^{-1} remains tightly constrained by the core geometry A_π, whereas G is significantly more




Figure 11: Phase space correlation between the inverse fine-structure constant and the electron anomalous magnetic moment. The curve represents the EWT geometric trajectory, showing that α^{-1} and a_e emerge from the same vacuum stiffness.

sensitive to changes in the stiffness parameter.

The resulting trajectory is shown in Fig. 12. This sensitivity difference follows directly from the algebraic structure:

$$\alpha_{\text{geom}}^{-1} \sim A_\pi - \frac{1}{N\pi^3}, \qquad G_{\text{geom}} \sim \frac{1}{N^3\sqrt{N_{\nu,\text{eff}}}}. \tag{339}$$

Thus, small changes in N affect G much more strongly than α. This explains why the electromagnetic sector appears stable while the gravitational sector is more sensitive to microscopic geometric parameters.

29.6 EWT Unification: Convergence of G and a_e on the Vacuum Stiffness Isentrope

A fundamental consistency check is the interdependence between G and a_e at a single geometric stiffness value

$$N_{\text{geom}} \approx 778.8025. \tag{340}$$

At this point, G_{geom} and a_e^{EWT} are both determined from the same BCC lattice parameters, as illustrated in Fig. 13. This is not a fit: the value of N_{geom} is fixed by the packing geometry and the impedance ζ.




Figure 12: The EWT Unification Path. The magenta line represents the geometric correlation between gravity and electromagnetism as a function of the vacuum stiffness parameter N.

29.7 Robustness Analysis: Geometric Stability and Vacuum Elasticity

To evaluate the reliability of the derived gravitational constant, a sensitivity analysis of the coupling between the soliton volume and the vacuum stiffness was performed. The structural stability is governed by the relationship

$$N \propto N_{\nu,\text{eff}}^{-1/6}, \tag{341}$$

which follows from the volumetric scaling $N_\nu \propto r^3$ and the stiffness-radius relation $N \propto r^{-1/2}$.

The resulting stability relationship is illustrated in Fig. 14. This damping factor protects the geometric vacuum parameters from large fluctuations of the EMC density. In the current model, the relevant stiffness is N_{geom}, not an arbitrary fitted value.

29.8 Structural Robustness and Resonance Stability of the Lepton Hierarchy

The same geometric stiffness N_{geom} anchors the recursive lepton hierarchy. The sensitivity gradients of a_μ and a_τ are shown in Fig. 15. They differ because the muon is a 2D surface resonance, while the tau is a 3D volumetric resonance. This is consistent with the dimensional hierarchy established in Section 3.




Figure 13: EWT Unification: Direct convergence of the gravitational constant G and the electron anomaly a_e at the geometric stiffness N_{geom}.

29.9 Electroweak and CKM Coincidence: Geometric Robustness of $\sin^2\theta_W$ and $\sin\theta_C$

The electroweak and Cabibbo mixing sectors converge at the same geometric anchor N_{geom}, as shown in Fig. 16. The figure uses the EWT-derived spherical-mode quark masses (Variant A of Section 3.4); the corresponding Cabibbo trajectory is therefore slightly offset from the Weinberg one. The π^6 bosonic coupling is steeper than the π^5 fermionic coupling, reflecting the difference between volumetric and surface resonances.

The Cabibbo sector remains sensitive to the quality of the quark mass inputs. When PDG 2024 quark masses are used, the flavor deficit $\epsilon_{\text{flavor}} = 1/(8\pi^5(1-\zeta))$ reproduces $\sin\theta_C$ to within 0.002%. The remaining discrepancy with EWT-derived quark masses originates from the mass prediction sector, not from the mixing geometry itself.

29.10 Lock-In Test: Independent Determination of N_{geom}

The robustness analysis presented so far demonstrates that the model is stable under perturbations of its inputs. A complementary and more stringent test concerns whether the geometric stiffness N_{geom} is *over-determined* by independent physical sectors, or merely fitted to a single observable. The results of such a test are reported below.




Figure 14: Gravity Stability Equilibrium: The $r^{-1/2}$ trajectory of nodal stiffness N relative to the cubic scaling of soliton volume N_ν.

29.10.1 Method

For each of seven physical sectors — α^{-1}, G, a_e, a_μ, a_τ, $\sin^2\theta_W$, and $\sin\theta_C$ — the following quantities are computed:

1. the value of the sector at $N_{\text{geom}} = 778.8025$;
2. the *drift* of that value when N is changed by ± 10;
3. the logarithmic derivative $\beta = d\ln O / d\ln N$ at N_{geom}, obtained by central difference in $\log N$ with step 10^{-6};
4. the *resolution*, defined as the relative error of the sector at N_{geom} divided by $|\beta|$, which measures how narrowly the sector can locate N;
5. the *optimal* value N_{opt} for each sector, found by log-space scan followed by bisection.

The procedure is implemented in the OpenWave script `m4_11_lock_in_n_scan.py` [54]. The sectors are classified into three groups according to their resolution:

- **PRIMARY** — resolution $< 1\%$. These sectors can independently locate N at the sub-percent level.
- **WEAK** — resolution between 1% and 100%. These sectors are consistent with N_{geom} but cannot discriminate it from neighbouring values.




Figure 15: Real sensitivity of the lepton AMMs to a perturbation of the universal modulator ϵ_M. The muon (2D planar shell, projected through $\mathcal{O}_\mu = 1/(4\pi^2)$) is almost insensitive to ϵ_M; the tau (3D volumetric shell, $\mathcal{O}_\tau = 1$) shows the full sensitivity. The baseline point at N_{geom} is marked in red.

- **NO CONSTRAINT** — resolution $> 100\%$. These sectors provide no information about N.

29.10.2 Results

The resolution analysis is summarised in Table 30. The grouping of sectors follows directly from the resolution column.

29.10.3 PRIMARY sectors: G, a_τ, and $\sin\theta_C$

Three sectors resolve N at the sub-percent level. The two strongest are the gravitational constant G (0.004%) and the tau anomalous magnetic moment a_τ (0.14%). Their separately optimal values are

$$N_{\text{opt}}(G) = 778.83, \qquad N_{\text{opt}}(a_\tau) = 779.92, \tag{342}$$

which agree to within

$$\Delta = 0.14\% \quad \text{of } N_{\text{geom}}. \tag{343}$$

Crucially, the two sectors are constructed from completely different chains. G arises from the gravitational self-consistent derivation involving λ_l, $N_{\nu,\text{eff}}$, and the surface projection. a_τ arises from the recursive lepton shell damping governed by ϵ_M and the Fibonacci latching. Neither chain uses the other's inputs. The agreement between them is therefore not tautological.




Figure 16: Shift-Normalized Robustness Analysis: The intersection of the $\sin^2\theta_W$ and $\sin\theta_C$ trajectories at the geometric lock-in point N_{geom}.

Table 30: Resolution of N_{geom} by independent physical sectors. The three groups are separated by horizontal rules.

Sector	$\beta = d\ln O/d\ln N$	**Resolution**	N_{opt}
PRIMARY (resolves N, $< 1\%$)			
G	-12.000002	4.0×10^{-5}	778.83
a_τ	$+0.220297$	1.4×10^{-3}	779.92
$\sin\theta_C$	-0.008921	6.2×10^{-3}	774.03
WEAK (consistent, 1%–100%)			
$\sin^2\theta_W$	-0.045754	1.5×10^{-1}	910.04
a_e	$+0.001285$	1.8×10^{-1}	661.63
a_μ	$+0.001415$	1.8×10^{-1}	661.69
NO CONSTRAINT ($> 100\%$)			
α^{-1}	$+3.0 \times 10^{-7}$	6.4×10^{0}	105.85

The exponent $\beta_G = -12$ (rather than the naive -3) is a structural consequence of the self-consistent derivation of λ_l: the lattice spacing scales as $\lambda_l \propto N^{-6}$, and G inherits an additional N^{-9} dependence through $N_{\nu,\text{eff}}$, compounding with the explicit N^{-3} factor. This makes gravity by far the most sensitive probe of the lattice stiffness.

The Cabibbo angle $\sin\theta_C$ formally lands in the PRIMARY group with a resolution of 0.62%. This result is, however, conditioned on the central PDG values for the light quark masses. The

experimental uncertainty on m_d and m_s spans approximately 14%, roughly fifteen times larger than the geometric correction being tested. At the low corner of that range no positive N reproduces $\sin\theta_C$. The Cabibbo leg is therefore treated as *conditionally* primary, not as a fully independent constraint.

29.10.4 WEAK sectors: a_e, a_μ, and $\sin^2\theta_W$

Three sectors are consistent with $N_{\rm geom}$ but cannot discriminate it from neighbouring values. Their resolutions of approximately 15–18% reflect different suppression mechanisms.

For the electron anomalous magnetic moment a_e, the sensitivity to N is suppressed because the prediction is dominated by the nearly N-independent Schwinger term. The electron AMM therefore confirms the overall framework but does not constrain the lattice stiffness.

For the muon anomalous magnetic moment a_μ, the projection operator $\mathcal{O}_\mu = 1/(4\pi^2)$ suppresses the sensitivity of the muon shell. The muon leg is consistent with $N_{\rm geom}$ but, like the electron, cannot discriminate between neighbouring values.

For the Weinberg angle $\sin^2\theta_W$, the logarithmic derivative is small ($\beta \approx -0.046$), giving a resolution of 14.6%. The comparison between the model value $\sin^2\theta_W = 0.23276$ and the PDG reference uses the MS-bar scheme, while the on-shell value derived from the PDG masses is 0.22305. The geometric correction $C_{\rm gap}$ occupies the same slot as a known scheme-conversion factor of approximately 1.06%. Consequently, the Weinberg leg cannot yet carry independent weight.

29.10.5 NO CONSTRAINT: α^{-1}

The inverse fine-structure constant provides no constraint on N. Its logarithmic derivative is $\beta \approx 3 \times 10^{-7}$, giving a resolution of 635%. This is expected and consistent with the model structure: α^{-1} is dominated by the geometric core A_π, and the ϵ_M correction is tiny. α therefore confirms the model but does not constrain its central parameter. Its inclusion in the lock-in analysis serves as a consistency check: the absence of a constraint is itself a prediction of the model's structure.

29.10.6 Interpretation

The lock-in test demonstrates that $N_{\rm geom}$ is not a fitted parameter in the usual sense. It is a *node* at which two independent physical sectors — gravity and the tau anomalous magnetic moment — converge to within 0.14%, with completely different construction chains and different sensitivities to the lattice stiffness.

The PRIMARY sectors are those whose prediction chains contain N at high powers ($\beta_G = -12$, $\beta_{a_\tau} = 0.22$) and whose targets are known with sufficient precision. The WEAK sectors are those whose sensitivity is suppressed by projection operators or by dominant N-independent terms. The NO CONSTRAINT sector is dominated by the geometric core A_π and is essentially insensitive to N.

This hierarchy is a falsifiable structural feature of the model. If future measurements of a_τ (currently only bounded experimentally) or of G (currently known to 2.2×10^{-5}) were to shift either sector's optimal N beyond the common interval, the model would be challenged. If, on the other hand, the PRIMARY sectors were to remain in agreement while a WEAK sector moved into the PRIMARY group, the model would be strengthened by the addition of a third independent constraint.

30 Geometric Phase Transitions: From Neutrino to Black Hole with EMC Wall

In the EWT framework, the vacuum lattice is a physical medium with a finite redistribution capacity. The formation of a **Geometric Black Hole (GBH)** represents a phase transition where the vacuum is pushed to its absolute elastic and geometric limits.

30.1 Hierarchy of Vacuum States and the G-Operating Point

The hierarchy of states is governed by the depth of the density deficit relative to the statutory equilibrium:

- **Effective Point ($N_{\nu,\mathbf{eff}} \approx 6.25 \times 10^{48}$):** The standard operating point of attractive gravity (approx. 99.98% deficit relative to background).
- **Statutory Limit ($N_{\nu,\mathbf{stat}} \approx 3.30 \times 10^{52}$):** The saturation ceiling where gravity inverts into repulsion (**EMC Wall**).
- **Max Packing ($N_{\nu,\mathbf{max}} \approx 5.30 \times 10^{54}$):** The asymptotic structural limit of the BCC lattice at the Planck scale.

The GBH Core: Finite Vacuum Exhaustion and Pure Wave Centers

Unlike General Relativity, EWT defines the interior of a GBH as a domain of **maximal individual energy concentration** (ρ_E) and **maximal medium deficit** (ρ). This state is realized by high-energy neutrino solitons acting as pure Wave Centers (WC).

Empirical Validation (Supernovae): The observation that **99% of supernova energy is released as neutrinos** constitutes empirical confirmation that the collapse of matter under extreme gravity leads to the conversion of the dominant mass into the **minimal geometric state** of the Neutrino (r_ν) at the point d_{crit}. This fact justifies the adoption of the Neutrino as the fundamental WC for the GBH model.

The EMC Wall and the Push-Out Mechanism

The EMC Wall is a physical density barrier in a vacuum created when the medium (BCC network) cannot keep up with the redistribution of displaced components (EMC). In ordinary solitons (such as leptons) this barrier exists in a **degraded** form – the *Degraded EMC Wall* discussed in Sec. 11.11 – where the local EMC density exceeds the statutory background $N_{\nu,\mathrm{stat}}$ only moderately, and the wall acts as a passive geometric low-pass filter. In a Geometric Black Hole (GBH), however, the push-out mechanism is driven to its extreme: the core exhausts the vacuum so severely that the accumulated EMC density at the wall **significantly exceeds** the statutory background $N_{\nu,\mathrm{stat}}$, turning the wall into an insurmountable barrier. The EMC wall then defines the geometric event horizon of the black hole. The relationship between the black hole core's energy and the wall's density is defined by the displacement flux:

$$\Delta\rho_{E(\mathrm{core})} \xrightarrow{\text{push-out}} \Delta\rho_{(\mathrm{wall})} \geq N_{\nu,\mathrm{stat}} \tag{344}$$

Energy Dissipation and Degradation

Energy exchange between the Geometric Black Hole and the external environment remains possible; however, the emitted energy must adopt a **degraded form**. The EMC Wall acts as a

mandatory transducer, where the extreme resistance of the saturated lattice forces all outgoing flux into states compatible with the medium's localized stiffness.

Astrophysical Jets and the Density Wall

The maximal vacuum exhaustion within the GBH core necessitates a **massive reciprocal accumulation** of EMCs at the event horizon's boundary—the **EMC Density Wall**. This wall possesses extreme nodal stiffness, anchoring ultra-strong magnetic fields. Astrophysical jets act as **structural discharge mechanisms**, where the immense potential of the compressed lattice is released along the axes of least resistance (the poles).

The EMC Barrier and Orbital Stability

Beyond the **Statutory Limit** ($N_{\nu,\text{stat}}$), the density gradient undergoes a sign reversal ($\nabla\rho > 0$). This creates a localized **Repulsion Zone** (anti-gravity). The event horizon thus functions as a buffer; matter is either stabilized by this "gravitational conjunction" or **actively ejected** into jets upon encountering the super-statutory pressure of the wall.

The EMC Wall: Selective Frequency Response

The extreme nodal stiffness of the **EMC Wall** (approaching $N_{\nu,\text{max}}$) establishes a **high-pass filter**. Because the lattice bonds in the wall are pushed to their elastic limit, only waves with **extremely high frequencies** (Gamma and X-rays) have the energy density required to overcome the medium's inertia.

The EMC Wall: Decomposition of Matter

As leptonic matter approaches the boundary, the transition on the wall's congestion creates an insurmountable **impedance mismatch**. The vacuum lattice becomes too rigid to maintain the wave structure of leptons, leading to the **structural breakdown** of the particle soliton.

Hypothesis: Gravitational Waves as Background Density Shifts

Within the EWT framework, gravitational waves can be interpreted as propagating pulses that momentarily increase the **statutory background density** ($N_{\nu,\text{stat}}$) of the BCC lattice. Crucially, the soliton's internal push-out mechanism is blocking EMCs; the matter does not absorb additional EMCs. Instead, the passing wave momentarily elevates the surrounding medium's density toward the EMC Wall threshold.As the background density $N_{\nu,\text{stat}}$ surges, the relative gradient between the soliton's core (the vacuum exhaustion zone) and the statutory environment is altered. This transient shift in the background reference point modulates the $\mathbf{\Omega}_{\text{Push}}$ operator, manifesting as a temporary change in the "gravitational shadow". Once the peak of the background EMC pulse passes, the lattice undergoes relaxation, restoring the statutory equilibrium. This interpretation suggests that gravitational waves are not a change in matter itself, but a moving "impedance spike" in the vacuum background that matter inhabits.

In this scenario, the matter "does not notice" the wave because its internal structural identity is preserved relative to the shifting background. The wave is only detectable via phase-sensitive measurements (interferometry), where the transient change in lattice impedance affects the propagation velocity of massless wave packets (photons) without disrupting the integrity of the leptonic solitons.

Hypothesis: Stellar Stability and Local EMC Saturation

In the context of stellar evolution, the EWT framework suggests that the equilibrium of a star is not solely a result of thermodynamic radiation pressure, but rather a consequence of the **structural impedance of the vacuum**. As stellar mass accumulates, the central region approaches a state of extreme vacuum exhaustion, while the surrounding layers may experience a density surge toward the **EMC Wall threshold** ($N_{\nu,\mathrm{tmp}} \to N_{\nu,\mathrm{stat}}$).This creates a "mechanical cushion" of high nodal density that effectively supports the matter against gravitational collapse. In this scenario, the stellar plasma "floats" on a layer of saturated lattice, where the push-out mechanism of the core solitons meets the resistance of a localized EMC Wall. This hypothesis could provide a more robust mechanical explanation for the stability of stars and the anomalous heating of the solar corona, interpreted here as a region of non-linear lattice relaxation and high-frequency nodal friction.

Hypothesis: Variable Vacuum Impedance in Stellar Media

It has been hypothesized that stars possess an EMC precursor wall. This local increase in $N_{\nu,\mathrm{tmp}}$ would manifest itself as a measurable change in vacuum impedance, potentially altering the local speed of light. Unlike general relativity (GR), which attributes such delays to metric curvature, EWT theory proposes a mechanical slowdown caused by increased BCC lattice stiffness near the saturation point. It remains an open question how far from the stellar core such a precursor could exist and what magnitude of deflection it assumes.

Hypothesis: Residual EMC Walls and the Geometry of Weak Interactions

It is hypothesized that every matter soliton is bounded by a **residual or "degraded" EMC wall**, defined by a local density gradient $N_{\nu,\mathrm{stat}} > x > N_{\nu,\mathrm{eff}}$. While the soliton core represents a zone of vacuum exhaustion, its boundary serves as a high-impedance interface where the push-out mechanism meets the statutory background of the BCC lattice. This boundary layer can provides a physical site for weak interactions.

Hypothesis: The EMC Wall as a Dimensional Transformer ($\pi^6 \to \pi^7$)

It is hypothesized that the residual EMC wall acts as a dimensional interface. Within this gradient, the interaction scales from the **volumetric bosonic coupling** (π^6) to the **full charge density resonance** (π^7). In this view, the electric charge is not an intrinsic "point-property" but the highest dimensional manifestation of the soliton's engagement with the BCC lattice.

30.2 The Erasure of "Dark" Placeholders

The success of the structural EWT framework renders several foundational "mysteries" of the Standard Model obsolete. By grounding physics in the properties of the BCC lattice, we eliminate the following "dark" placeholders:

- **Dark Matter as a Geometric Push-Force:** Rather than invoking undiscovered WIMPs, EWT identifies galactic rotation anomalies as a consequence of the **massive EMC volume deficit**. The near-total geometric gap at the G-operating point creates a pervasive external pressure gradient that "pushes" matter toward regions of lower nodal density.
- **Dark Energy as Vacuum Static Pressure:** Accelerated expansion is reinterpreted as the **restorative elastic response** of the BCC lattice: the statutory static pressure of the

EMC background acting upon the structural voids created by matter solitons, the vacuum's tendency to return to its statutory equilibrium. The quantitative realization of this picture — the effective vacuum fraction and the corresponding cosmological constant — is given in Section 24.7, where the composition rule is stated explicitly as a model assumption.

- **Virtual Particles and Loop Complexity:** The thousands of Feynman diagrams are replaced by the **Nodal Stiffness** N. Vacuum fluctuations are revealed to be the deterministic, high-frequency oscillations of the lattice nodes.
- **The Hierarchy Problem:** The discrepancy between gravity and electromagnetism is a simple ratio: gravity is a statistical effect of **missing nodes** (EMC deficit), while electromagnetism is a dynamic effect of **node workload** (energetic load ρ_E).

30.3 Resolution of Historically "Unsolvable" Paradoxes

The mechanical properties of the BCC lattice, combined with the threshold-based dynamics of the **EMC Wall**, provide deterministic solutions to several long-standing paradoxes that have remained unresolved within the probabilistic framework of the Standard Model.

The Horizon Problem: Nodal Saturation and Thermalization

The large-scale uniformity of the universe is a consequence of the **Initial Saturation State**. In the earliest epoch, the vacuum medium existed at the **Max Packing limit** ($N_{\nu,\max} \approx 10^{54}$), where the BCC lattice is structurally incompressible.

In this state of maximal density, nodal stiffness is absolute, and the velocity of structural equilibrium is effectively infinite. This "Super-Statutory" phase allowed the entire medium to act as a single, coupled resonator, achieving perfect thermal uniformity. The subsequent relaxation of the lattice toward the **Statutory Limit** ($N_{\nu,\text{stat}}$) and eventually the current **Effective Operating Point** ($N_{\nu,\text{eff}}$) preserved this initial homogeneity, rendering the speculative "Inflation" field unnecessary.

The Information Paradox: Nodal Memory

The "loss" of information in a black hole is prevented by the physical nature of the event horizon. The **EMC Wall** acts as a high-density storage medium. The decomposition of matter at the boundary is not an erasure, but a **transduction**: the soliton's wave frequency is encoded into the **Nodal Memory** of the saturated BCC lattice ($N_{\nu,\max}$). Information is preserved as a specific vibrational state of the wall's nodes.

The GZK Limit: Lattice-Induced Drag

The Greisen-Zatsepin-Kuzmin (GZK) limit, which restricts the energy of cosmic rays, is revealed to be a **structural speed limit**. As a particle soliton approaches extreme energies, it creates a "Micro-EMC Wall" (a local compression wave) directly ahead of its path. This results in a non-linear increase in vacuum resistance, effectively capping the maximum energy a soliton can maintain while moving through the BCC substrate.

Matter-Antimatter Asymmetry: Asymmetric Phase-Lattice Transition

In the EWT framework, following the geometric interpretation proposed by Yee and Gardi [26], antimatter is defined as a soliton state characterized by a 180° (π) phase-shift. The observable

dominance of matter is reinterpreted as a result of **Non-linear Phase Conversion** occurring at the **trans-statutory boundary** ($N > N_{\nu,\text{stat}}$) of EMC Walls.

In this model, the transition through an EMC Wall is not phase-symmetrical. While the wall acts as a resonant processor that can theoretically shift phases in both directions, the current G-operating point of the global BCC lattice creates a bias that favors "locking" solitons into the matter-phase. Consequently, antimatter is a phase that less frequently exits the high-EMC density regions in its original form, being resonantly re-tuned to the dominant background. This suggests that the universe's matter-bias is a dynamic equilibrium maintained by the lattice's tuning, where high-EMC density regions act as filters that preferentially stabilize the primary soliton phase.

30.4 Density Duality: From Local Refraction to Cosmological Redshift

The Energy Density Profile $\rho_{\mathbf{E}}(\mathbf{r})$ as the Origin of Local Refractive Index and EM Wave Slowing

While the **geometric packing density function** $\rho(\mathbf{r})$ describes the non-uniform distribution of the elastic medium and is fundamental to the emergence of the gravitational constant G, a profound and distinct consequence of the Soliton's internal structure is the role of its **localized energy concentration** in the propagation of electromagnetic (EM) waves.

The hypothesis is presented that the local **energy density** $\rho_{\mathbf{E}}(\mathbf{r})$ (which defines the Soliton's mass $E = mc^2$) acts as a local, variable refractive index for EM waves. The speed of light c is modified locally to $v(r)$ based on the energy density profile:

$$v(r) \;=\; \frac{c}{n(r)}, \qquad n(r) \approx 1 + C_{\text{E}} \cdot \rho_{\text{E}}(r), \tag{345}$$

where C_{E} is a constant of proportionality related to the electromagnetic field coupling with the Soliton's high energy density.

Compatibility with Density Duality (Clear Separation of Roles). This change establishes clear and distinct physical roles:

- **Refraction:** The **high energy density ($\rho_{\mathbf{E}}$)** ensures $n(r) > 1$ and correctly predicts the **slowing of EM waves** inside the Soliton/Matter.
- **Gravity:** The **low geometric packing density ($\rho(\mathbf{r})$)** remains solely responsible for the geometric deficit ($N_{\nu,\text{effective}}$), which drives the **push force gravity**.

This framework suggests that the fundamental mechanism for the **slowing of EM waves in matter** (refraction) is the interaction with the combined, non-uniform $\rho_{\text{E}}(\mathbf{r})$ profiles of the atomic constituents. The geometric parameter that governs gravitational weakness ($\mathbf{N}_{\nu,\text{effective}}$, which is derived from the **integral of** $\rho(\mathbf{r})$) is thus linked to a core electromagnetic phenomenon (refraction) controlled by a **separate density component ($\rho_{\mathbf{E}}$)**.

Refractive index and Nodal Delay. The slowing of EM waves inside the Soliton is not a function of the number of nodes (geometric density $\rho(r)$), but the "workload" of each node (energy density ρ_E). While the massive deficit of EMC components (the gap between statutory and effective N_ν) drives the gravitational push-force, the nodes are subject to extreme oscillatory stress. This increases the **nodal response time**, effectively lowering the propagation velocity $v(r) = c/n(r)$ as a function of $\rho_E(r)$.

Cosmological Implications and the Geometric Origin of Redshift

The geometric derivation of **G** and the structural formalization of the Soliton ($\rho(\mathbf{r})$) carry profound consequences for cosmology. The EWT model fundamentally shifts the interpretation of observed phenomena away from dynamic spacetime expansion toward **wave propagation dynamics within the Elastic Medium Constituent (EMC)**.

The concept of gravity as a **push force** resulting from the volume deficit ($\mathbf{N}_\nu$) suggests an alternative framework for the accelerating expansion of the universe. This expansion, typically attributed to Dark Energy, can be re-interpreted as the **external pressure of the EMC background** acting on regions of matter deficit.

Furthermore, the dual role of the Soliton's energy density ($\rho_{\mathrm{E}}(\mathbf{r})$) in governing both mass and local EM wave slowing (refraction) opens the possibility that **cosmological redshift** ($\mathbf{z}$) may be partially or entirely a consequence of **"tired light" effects**—minimal energy loss of EM waves due to interaction with the bulk properties or temporal evolution of the EMC background over vast distances.

Synthesis of Geometric and Temporal Properties: While ϵ_M defines the fundamental geometric permeability (stiffness) of the vacuum in solitons, the **Nodal Delay** represents the temporal manifestation of this stiffness when under energetic load (ρ_E). This unifies the EMC deficit (ρ) with the dynamic slowing of light (refraction) and the cumulative energy loss over cosmological distances (redshift). This perspective effectively replaces the notion of "empty space" with a dynamic, elastic medium whose response time is intrinsically coupled to its energy density.

Although this model focuses on leptonic structure, the unification of geometric parameters suggests that the Nodal Delay mechanism may have a cumulative component at the cosmological scale. This allows for the hypothesis of a non-Doppler origin for a portion of the redshift, representing a promising direction for future research into the nature of the EMC background.

31 Outlook and Computational Tools

The formal establishment of the Magneto-Geometric Hypotheses (Section 18) provides clear, testable predictions for the effective gravitational constant G (Section 27.4). Beyond direct experimental verification, the computational modelling of the Soliton geometry and its dynamic coupling to the Elastic Medium Constituent (EMC) is being pursued on the open-source platform **OpenWave** (`https://github.com/openwave-labs/openwave`).

Progress in the Scalar M3 Model

In the scalar Wolff-LaFreniere model (M3), a spherical-phyllotaxis (golden-angle, $\sim 137.5°$) configuration of wave centres was implemented and tested as a proof-of-concept. The results provided the first in-platform demonstration that a spherical, minimally interfering geometry can create K-selectivity: only $K_{WC} = 10$ formed a stable standing wave, while $K_{WC} = 9$ and $K_{WC} = 11$ decayed systematically. This validated the underlying principle that phyllotactic packing suppresses destructive interference, a principle that will be essential for the high-multiplicity recursive shells of the muon and tau. For the electron core itself, the native EWT topology (1-3-6 tetrahedron) remains the physically required configuration, as it alone guarantees the correct multipole structure for far-field forces. The golden-angle module, together with the logged simulation data, is available in the `m3_wolff_lafreniere` directory of the OpenWave repository.

Diagnostics in the Vector M4 Model and Implementation Plan

The same golden-angle arrangement does *not* stabilise the electron core in the vector model M4 (neither with nor without an initial spin), indicating that the missing ingredient in M4 is not geometric but nonlinear. The immediate goal is to stabilise the native 1-3-6 electron topology by introducing the nonlinear self-trapping term derived in Section 26. Once the electron core is stable with its proper geometry, the golden-angle phyllotaxis will be applied to the high-multiplicity recursive shells of the muon and tau, where it is expected to be indispensable. Consequently, the following implementation roadmap has been adopted:

1. **Nonlinear term $\mathcal{F}(\Psi)$:** A cubic self-trapping term $\gamma\,\Psi^3$ will be added to the M4 wave kernel, with the coupling coefficient $\gamma = 1/\epsilon_M = N_{\text{final}}\pi^3 \approx 2.41 \times 10^4$ derived from the BCC geometry.
2. **Degraded EMC Wall:** A radial density gradient will be introduced to suppress shear instabilities and to provide a natural boundary where the phyllotaxis can later be introduced for the outer shells.
3. **Validation:** Once the nonlinearity is in place, the native EWT topology (1-3-6 tetrahedron) will be tested for K-selectivity.
4. **Shell application:** After the core is stabilised, the golden-angle arrangement will be deployed for the muon and tau shells.

Connection to the Nonlinear Stability Equation

A sketch of the required nonlinear wave equation was proposed in part by the OpenWave team during the collaborative development of the M3 and M4 engines. Building on that foundation, a formal nonlinear wave equation that guarantees the electron's stability is derived in Section 26, where the geometric constants ϵ_M and $\gamma = 1/\epsilon_M$ determine the self-trapping term. The full implementation of this equation in the OpenWave M4 kernel is the immediate next step.

This computational framework serves as a **theoretical complement to physical experiments**, allowing rapid iteration and falsification of the Soliton's micro-geometric constraints within the EWT Model.

32 Conclusion: The Geometric Paradigm, Correlated Lepton Anomalies, and Emergent Gravity

The formalization within the Energy Wave Theory establishes the **Minimal Wave Center** — the Neutrino Soliton — as a fundamental geometric structure. The central finding of this work is the derivation of the gravitational constant G as a geometric identity, together with the prediction of the lepton anomalous magnetic moments from the same underlying BCC lattice parameters. This necessitates a critical re-evaluation of the Soliton's geometric parameters.

This achievement realizes a principle that is increasingly recognized as a necessity for resolving the deepest tensions in modern physics. As highlighted by the CosmoVerse Network [48], addressing the observational crises in cosmology may ultimately require abandoning the notion of rigid, immutable fundamental constants in favour of dynamic, relational parameters. The Enhanced EWT model does not merely accommodate this principle; it provides its precise geometric mechanism. The observed "constants" — the gravitational coupling G, the electromagnetic coupling α, and the electron anomalous magnetic moment a_e — are not independent

inputs but are scale-dependent projections of a single invariant: the structural impedance of the BCC vacuum lattice.

For the muon and tau generations, the model presently derives the anomalous magnetic moments as genuine predictions of the BCC lattice geometry, while the reference scale for comparison is obtained from the orbital mass relations. This constitutes a high-precision internal consistency test between the geometric and mass-generation sectors; the simultaneous, fully independent derivation of both mass and AMM for all generations remains an open objective.

Constants are not constant; only the relations between them are.
This relation, dictated by the fixed topology of the vacuum substrate, is the true fundamental law from which all measurable quantities emerge.

32.1 Geometric Unification in a Nutshell: The ϵ_M Flowchart

The Enhanced EWT Model operates on the principle of **Structural Monism**: all fundamental coupling constants are derived from the same vacuum stiffness deficit. Rather than treating gravitation, electromagnetism, and lepton anomalies as independent phenomena, they are expressed as direct geometric functions of the primary modulator ϵ_M.

$$\epsilon_M \implies \begin{cases} \mathbf{G} = \dfrac{\mathbf{G}_{\text{Base}}}{A_\pi} \cdot \left(\dfrac{\epsilon_M \cdot \pi^3}{A_\pi} \right)^3 \cdot \dfrac{1}{K_{WC} \cdot \sqrt{N_{\nu,\text{eff}}}} & \text{(Gravitational Scale)} \\ \alpha_{\text{geom}}^{-1} = A_\pi - \epsilon_M & \text{(Electromagnetic Scaling)} \\ a_e = \dfrac{\alpha_{\text{geom}}}{2\pi} \cdot \left(1 - \epsilon_M \cdot \pi^3\right) & \text{(Nodal Magnetic Deficit)} \\ C_{\text{local}} = \dfrac{\epsilon_M}{2\sqrt{2}} & \text{(Geometric Seed for Mixing Hierarchy)} \\ R_l = \text{recursive}(\epsilon_M, L_{\text{Fib}}) & \text{(Leptonic Shell Resonance)} \end{cases} \tag{346}$$

In this zero-calibration formulation,

$$\epsilon_M = \frac{1}{N_{\text{geom}} \pi^3} = \frac{1}{8\pi^7(1-\zeta)}, \tag{347}$$

where $N_{\text{geom}} = 8\pi^4(1-\zeta)$ is the effective geometric stiffness and ζ is the BCC packing impedance.

32.2 The Unitary Geometric Path

As illustrated in the schematic, ϵ_M acts as the **Universal Gear Ratio** of the vacuum lattice. A single modulator simultaneously determines:

- the gravitational coupling G,
- the electromagnetic coupling α_{geom},
- the damping of the magnetic precession across generations, yielding a_e, a_μ, a_τ.

This flowchart confirms that the Enhanced EWT Model is not a collection of independent equations, but a **Unitary Geometric Circuit**. Any change in the value of ϵ_M would simultaneously shift all fundamental constants derived from it.

32.3 The Geometry of the Constituent: From Cubic Grids to Spherical Units

While the initial formalization of the EWT lattice utilizes the Body-Centered Cubic framework for its nodal efficiency, the physical reality of the **Elastic Medium Constituent** is fundamentally spherical. This transition from a "cubic cell" to a "spherical grain" is a natural consequence of the model's convergence toward a minimum energy state and isotropic wave propagation.

Packing Fraction and the Vacuum Void

In a BCC lattice composed of hard spherical constituents, the maximum packing factor is

$$\eta_{\text{BCC}} = \frac{\sqrt{3}\pi}{8} \approx 0.680. \tag{348}$$

This inherent geometric property implies that:

- The vacuum is not a continuous "solid block" but a structured, granular medium with a defined interstitial capacity.
- The volume deficit N_ν identified in the gravitational derivation represents the displacement of these spherical units.
- The spherical nature of the EMC ensures isotropic nodal stiffness.

The Transition of Density Levels

The granular nature of the EMC directly dictates the three levels of vacuum density identified in the EWT hierarchy:

- **Maximum packing:** $N_{\nu,\text{max}} \approx 5.30 \times 10^{54}$.
- **Statutory background:** $N_{\nu,\text{stat}} \approx 3.30 \times 10^{52}$.
- **Effective gravitational density:** $N_{\nu,\text{eff}} \approx 6.25 \times 10^{48}$.

32.4 Pillar 1: The N_ν Deficit and Soliton Stability

A cornerstone of the EWT model is the derivation of the gravitational constant G from the microscopic geometry of the neutrino soliton and the electron soliton. In the zero-calibration formulation, the transition from the statutory background to the effective density is governed by the geometric dilution factor X_{eff}:

$$N_{\nu,\text{eff}} = \frac{N_{\nu,\text{stat}}}{X_{\text{eff}}}, \tag{349}$$

where X_{eff} is fixed by A_π, $K_{WC} = 10$, and the ideal BCC projection factor $L_p^{\text{geom}} = 2/\sqrt{3}$. The resulting density hierarchy is:

$$\rho_{\text{eff}} < \rho_{\text{stat}} < \rho_{\text{max}}. \tag{350}$$

Macroscopic gravitational attraction is the response of the external, higher-density vacuum medium ($N_{\nu,\text{stat}}$) attempting to restore equilibrium by pressing toward the lower-density soliton core.

32.5 Pillar 2: The Dynamic Scaling Operator Ω_{Push}

The dynamic push-out operator,

$$\Omega_{\text{Push}} \equiv G_{\text{Base}} \cdot A_\pi^{-1} \cdot (N_{\text{geom}} A_\pi)^{-3}, \tag{351}$$

is the central mechanism in the G equation.

The final gravitational identity is:

$$G \equiv \underbrace{G_{\text{Base}} \cdot \frac{1}{A_\pi} \cdot \left(\frac{1}{N_{\text{geom}} A_\pi}\right)^3}_{\Omega_{\text{Push}}} \cdot \underbrace{\frac{1}{K_{WC}\sqrt{N_{\nu,\text{eff}}}}}_{T_{\text{Surface}}}. \tag{352}$$

The factor $\epsilon_M = (N_{\text{geom}}\pi^3)^{-1}$ is the master key of this operator. It dictates both the gravitational scale and the lepton anomalous magnetic moments.

32.6 Pillar 3: The Recursive Paradigm — Hierarchical Nodal Integration

The transition from the electron core to the muon and tau generations is viewed as a hierarchical integration of nested nodal shells, parametrized by the Fibonacci labels

$$L_\mu = F_5 = 5, \qquad L_\tau = F_9 = 34. \tag{353}$$

The full AMM predictions are:

$$a_e^{\text{EWT}} = 1.159916 \times 10^{-3}, \qquad a_\mu^{\text{EWT}} = 1.166213 \times 10^{-3}, \qquad a_\tau^{\text{EWT}} = 1.176838 \times 10^{-3}. \tag{354}$$

The recursive architecture provides a unified geometric framework that offers an alternative to the perturbative loop approach of the Standard Model, deriving all three lepton anomalies from a single common modulator ϵ_M rather than from independent loop corrections.

32.7 The Soliton Radius and Neutrino Interaction Cross-Section

A common concern regarding geometric particle models is the apparent tension between a comparatively large geometric radius and the extremely small neutrino interaction cross-sections. This model resolves the issue by distinguishing between:

- the **geometric soliton radius** r_{geom},
- the **interaction cross-section** σ_ν.

The weakness of the neutrino interaction reflects the fact that only a minute fraction of the soliton's internal energy is available at its boundary for weak interactions. This dilution is quantified by the effective volume deficit:

$$\sigma_\nu \propto \frac{1}{N_{\nu,\text{eff}}}, \tag{355}$$

so that even a soliton with a substantial geometric radius can possess an exceedingly small interaction cross-section.

32.8 Paradigm Shift: The Planck Length as a Physical Boundary

In EWT, the Planck length is assigned a definitive physical and geometric role: it is the approximate magnitude of the base quantum distance λ_l, which defines the physical separation of the Elastic Medium Constituents.

In this framework, λ_l is not an external input; it is the self-consistent geometric Planck length derived in Section 11.6. All physics operates consistently at and above this scale, allowing the direct geometric calculation of macroscopic constants like G and α.

32.9 The ϵ_M Factor: Universal Geometric Modulator

The core tenet of EWT is that mass, charge, and spin are manifestations of the Soliton's wave geometry. The unified role of the magnetic deficit factor is governed by a fundamental geometric duality: while the Soliton's internal energy storage follows $E \propto r^5$, the volume of the displaced elastic medium follows $V \propto r^3$.

The factor ϵ_M acts as the universal modulator that reconciles these two scaling laws. It performs three distinct functions:

1. **Gravity:** It scales the volumetric deficit to the observed gravitational constant G.
2. **Electromagnetism:** It bridges the gap between α_{geom} and the soliton core.
3. **Spin/AMM:** It defines the recursive nodal integration of lepton anomalies.

The necessity of using the same factor ϵ_M across all these domains is the definitive proof of geometric unification.

32.10 Geometric Self-Consistency of the Planck–Gravity–Metric Triangle

A central result of the present work is the recognition that G_{geom}, $\hbar_{\text{geom}}$, λ_l, and c do not form a hierarchy of independent constants. They are linked by the closed geometric condition

$$\lambda_l = \sqrt{\frac{\hbar_{\text{geom}}\, G_{\text{geom}}}{c^3}}, \tag{356}$$

where

$$\hbar_{\text{geom}} = \frac{m_e c r_e}{\alpha_{\text{geom}}}, \qquad G_{\text{geom}} = \frac{G_{\text{Base}}}{A_\pi}\left(\frac{1}{N_{\text{geom}} A_\pi}\right)^3 \frac{1}{K_{WC}\sqrt{N_{\nu,\text{eff}}}}. \tag{357}$$

The effective density $N_{\nu,\text{eff}}$ itself depends on λ_l through the statutory background,

$$N_{\nu,\text{eff}} = \frac{1}{X_{\text{eff}}}\left(\frac{r_\nu}{2e\lambda_l}\right)^3. \tag{358}$$

Thus Eq. (356) is not a definition imported from outside the model. It is an internal algebraic constraint that simultaneously fixes the gravitational scale, the quantum scale, and the fundamental lattice spacing from the same BCC geometry.

In natural units ($c = 1$), this constraint reduces to the purely geometric statement

$$\lambda_l^2 = \hbar_{\text{geom}}\, G_{\text{geom}}, \tag{359}$$

showing that the square of the fundamental EMC length is the product of the geometric quantum of action and the geometric gravitational coupling.

This closure is a direct consequence of the BCC lattice structure. It is not an import from the Standard Model: the Planck scale, the reduced Planck constant, and the gravitational constant are reconstructed here as derived quantities, not assumed inputs.

32.11 Final Synthesis

The Enhanced EWT model demonstrates that the fundamental constants are integrated resonances of a single BCC substrate. The geometric vacuum stiffness N_{geom}, the magnetic deficit ϵ_M, and the geometric core A_π together determine the fine-structure constant, the gravitational constant, the lepton anomalous magnetic moments, and the principal atomic scales.

The model is not claimed to be absolutely parameter-free. It currently uses four experimental anchors:

$$r_e, \quad m_e, \quad c, \quad e. \tag{360}$$

These anchors are fixed by measurement and by the metric convention. They are not adjustable calibration parameters: if any one of them were changed independently while keeping the BCC geometry fixed, the entire set of geometric predictions would break. In addition to these four measured anchors, the model contains one discrete postulate: the muon Fibonacci label $L_\mu = F_5 = 5$, from which all higher shell labels follow by the four-step rule of Section 16.2. Apart from this discrete postulate, the model contains no free calibration constants; it is zero-free-parameter in the operational sense.

Of these anchors, c is a metric conversion factor rather than a dynamical parameter. The electron mass m_e may also be derivable in the future through the $E \propto r^5$ scaling law and the $K_{WC} = 10$ stability condition. The true irreducible anchors would then reduce to r_e and e.

The cosmological sector extends this accounting by one further empirical anchor, the present CMB temperature $T_{\mathrm{cmb},0}$ (COBE/FIRAS [70]), and by one further exact unit-conversion constant, the Boltzmann constant k_B, which, like c, carries no dynamical information. With these two additions the same BCC geometry recovers the thermodynamic horizon relation of Haug and predicts the Hubble constant and the de Sitter value of the cosmological constant. The horizon factor $H = 8(2\pi)^2$ that closes the recovered relation is a model construct, not a first-principles derivation. The observed cosmological constant is obtained by combining the de Sitter value with a geometric mechanism for the effective vacuum fraction, $\Omega_{\mathrm{vac}} = \eta_{\mathrm{BCC}} C_{\mathrm{gap}}$; the composition rule of this mechanism is a model assumption, stated explicitly in Section 24.7. No continuously adjustable parameter is introduced in the cosmological sector: the only new inputs are $T_{\mathrm{cmb},0}$ and the exact constant k_B, and the only new assumption is the composition rule for Ω_{vac}.

A Appendix

A.1 List of Model Parameters and Physical Constants

The following table lists the fundamental physical constants and key dimensionless geometric factors used in the Geometric Soliton Model. The numerical consistency of the model relies on the exact values shown below, combining CODATA 2022 constants with the derived EWT geometric parameters.

Table 31: **EWT Parameters and CODATA 2022 Values Used in the Geometric Derivations**

Parameter	Symbol	Numerical Value	Source / Justification
I. Fundamental Constants (CODATA 2022)			
Speed of Light	c	299792458	Defined constant (m/s).
Electron Mass	m_e	$9.1093837015 \times 10^{-31}$	CODATA 2022 reference (kg).
Classical Radius	r_e	$2.8179403262 \times 10^{-15}$	CODATA 2022 reference (m).
II. Hierarchical Volume Deficit Parameters			
Absolute Maximum	$N_{\nu,\max}$	5.3004×10^{54}	Theoretical EMC packing limit.
Statutory Background	$N_{\nu,\text{stat}}$	3.2986×10^{52}	Eulerian vacuum capacity.
Effective Deficit	$N_{\nu,\text{eff}}$	6.2472×10^{48}	Integrated Soliton density.
Geometric Bridge Operator	X_{eff}	5275.7683	Transforms $N_{\nu,\text{stat}}$ into $N_{\nu,\text{eff}}$.
III. Geometric Operators			
Geometric Base	A_π	137.0363037759	$4\pi^3 + \pi^2 + \pi$. Static foundation.
Effective Stiffness	N_{geom}	778.8025179	$8\pi^4(1-\zeta)$.
Lattice Impedance	ζ	0.0006033964	From BCC packing fraction.
Magnetic Deficit	ϵ_M	$4.141169769 \times 10^{-5}$	$1/(N_{\text{geom}}\pi^3)$.
Lattice Projection	L_p^{geom}	1.1547005384	$2/\sqrt{3}$, ideal BCC projection.
Unified Coupling	C_{unif}	1.1020116168	Using L_p^{geom}.

A.2 Inventory of Inputs, Postulates, and Parameter-Free Forms

The following table classifies every element of the model according to its logical status: whether it is a measured input (anchor), a test anchor used only for consistency checks, a discrete postulate, a geometrically fixed integer, a parameter-free functional form, or a rounding of a derived quantity. The purpose is to provide the reader with an explicit, section-by-section accounting of where each element enters and what role it plays.

Element	Section	Status	Role in the model
I. Primary inputs and exact conversions			
I.A Primary empirical anchors (measured, CODATA 2022 / observational)			
r_e	11	Anchor	Classical electron radius, fixes G_{Base}
m_e	11	Anchor	Electron mass, fixes G_{Base} and the $E \propto r^5$ scale
e	7	Anchor	Elementary charge amplitude $e = 1.602176634 \times 10^{-19}$ m (SI numerical value, dimension length); fixes $q_P = e/\sqrt{\alpha_{\mathrm{geom}}}$ and r_ν. **Notation warning:** the symbol e also denotes Euler's number ($e = 2.71828\ldots$) in $S_{\mathrm{tot}} = \alpha/(8+\pi)+e+(1-g_v)(\sqrt{2}-1)$ and in $r_\nu = 2q_P e^2/g_v$. The two uses are distinguished by context: elementary charge appears in the anchor and in q_P; Euler's number appears in the additive decay term of S_{tot} and in the neutrino-radius prefactor. In the executable implementation the two are separate constants (`E_CHARGE_CODATA` vs `EULER`).
$T_{\mathrm{cmb},0}$	24	Anchor (cosmological)	Present CMB temperature 2.7255 ± 0.0006 K (COBE/FIRAS [70]); single cosmological anchor of the framework
I.B Exact SI conversions (2019 revision, no uncertainty)			
c	9	Unit conversion (metric)	Exact by SI 2019 ($299\,792\,458\ \mathrm{m\,s^{-1}}$); conversion between spatial and temporal lattice steps; not a dynamical input
k_B	24	Unit conversion (thermodynamic)	Exact by SI 2019 ($1.380649\times10^{-23}\ \mathrm{J\,K^{-1}}$); converts Planck energy to Planck temperature T_p; not a dynamical input
II. Test anchors (consistency checks, not predictions)			
M_Z, $\sin^2\theta_W$	3.1.4	Test anchor	Evaluate $C_{\mathrm{gap}}^{\mathrm{mixed}}$; M_W prediction is compared with ATLAS 2024, CMS 2024, and PDG 2024
M_d, M_s	3	Test anchor	Evaluate $\epsilon_{\mathrm{flavor}}$; $\sin\theta_C$ tests the operator form
a_e, a_μ	16.5	Test anchor	PDG/CODATA values used as external test of the AMM predictions

continued on next page

Element	Section	Status	Role in the model
a_τ^{SM}	16.5	Test anchor (SM)	Standard-Model prediction, not measurement; no direct experimental anchor available
H_0 (Planck/DESI/SH0ES)	24	Test anchor	External determinations: Planck 2018 67.36 ± 0.54, DESI 2024 68.52 ± 0.62, SH0ES 2022 73.04±1.04 in $\mathrm{km\,s^{-1}\,Mpc^{-1}}$; compared with the chain prediction H_0^{EWT}
Ω_Λ, Λ_{obs} (Planck 2018)	24	Test anchor	Planck determinations $\Omega_\Lambda = 0.6889 \pm 0.0056$ (TT+BAO) and $\Lambda_{\mathrm{obs}} = 1.089 \times 10^{-52}$ $\mathrm{m^{-2}}$; compared with Ω_{vac} and Λ_{vac}
III. Discrete postulates and geometrically fixed integers			
$L_\mu = F_5 = 5$	16.2	Discrete postulate	Muon Fibonacci latch; $L_\mu^2 = 25$ is the interface area
$K_{WC} = 10$	26	Geometric integer	1-3-6 tetrahedral configuration of the BCC lattice; stability proof open in M4
$Q = 10$	25.3.1	Topological invariant (structural assignment)	Degree of the map $n : S^2 \to S^2$, $Q = (1/4\pi)\int_{S^2} n \cdot (\partial_\theta n \times \partial_\phi n)\, d\theta\, d\phi \in \mathbb{Z}$ (Eq. 301); assigned structurally to the 1-3-6 wave-centre configuration with $K_{WC} = 10$ (Eq. 302); first-principles derivation that $Q = 10$ is energetically preferred over neighbouring integers remains open (OpenWave M4)
IV. Model functional forms and chain outputs			
IV.A Parameter-free functional forms			
ζ	2	Functional form	Lattice impedance $(1-\eta)/(\eta N_{\mathrm{ideal}})$ from BCC packing fraction
C_{unif}	8.1	Functional form	Unified coupling $1/K_{WC} + 1 + \alpha/(\pi L_p)$
X_{eff}	8.1	Functional form	Dilution factor $A_\pi \cdot 3K_{WC}\sqrt{2}/C_{\mathrm{unif}}$
S_{tot}	5.2	Geometric ansatz	$\alpha/(8+\pi) + e + (1-g_v)(\sqrt{2}-1)$; validated by g_v agreement and $r_e/r_\nu = 100$
g_v	5.3	Derived fixed point	Unique root of the S_{tot} quadratic equation; $g_v \approx 0.983594$; determines $r_\nu = q_P S_{\mathrm{tot}}$; no independent parameter
r_ν	5.4	Derived quantity	Neutrino radius $r_\nu = q_P S_{\mathrm{tot}} \approx 2.8179 \times 10^{-17}$ m; derived from q_P (class I anchor e and α_{geom}) and S_{tot} (class IV, depends on g_v); enters $N_{\nu,\mathrm{stat}}$, $N_{\nu,\mathrm{eff}}$, X_{eff}, and the Planck-length derivation (Section 11.6); no independent parameter
C_{gap}	3	Functional form	$1+\pi^6 C_{\mathrm{local}}$; applies to pure neutral vertices (Z–Z, Z–H)
$C_{\mathrm{gap}}^{\mathrm{mixed}}$	3.1.4	Methodologically motivated ansatz	$1 + O_{\mathrm{mixed}} C_{\mathrm{local}}$, where $O_{\mathrm{mixed}} = \pi^6\pi^7/(\pi^6+\pi^7)$ is the parallel-impedance combination of the neutral and charged weak rungs; unique simple form with correct single-rung limits; derivation from BCC dynamics is open

continued on next page

Element	Section	Status	Role in the model
ϵ_{flavor}	3	Functional form	$1/(8\pi^5(1-\zeta))$; flavor deficit at the π^5 rung, analogous to $\epsilon_M = 1/(8\pi^7(1-\zeta))$ at the charged weak rung
B_μ, B_τ	16.4	Functional form	Shell prefactors: $3A_\pi\pi^3/(2L_\mu^2)$ and $3A_\pi\pi^3/(8\sqrt{2}) + A_\pi/2$
ΔK_k	16.2	Functional form	Shell-rounding operator round$(10^k \cdot 2\pi^2)$; $K_n = 10, 207, 2181$ follow deterministically
γ	26	Functional form	Nonlinear coefficient $1/\epsilon_M$ in the $\gamma\Psi^3$ term
n_γ	12.3	Postulate	Null-probe refractive index $(1 - r_s/r)^{-1}$; full derivation open
q_P	5.1	Derived quantity	Planck charge $q_P = e/\sqrt{\alpha_{\text{geom}}} \approx 1.8755 \times 10^{-18}$ m. Parameter-free combination of the anchor e (class I) and the geometric α_{geom} defined in Section 2. Inherits the same relative deviation as α_{geom} vs. α_{CODATA} ($\sim$ 1 ppm in q_P). Not an independent prediction; sets the amplitude scale for r_ν.
IV.B Cosmological chain			
Haug relation	24.5	Recovered	$T_{\text{cmb},0} = (T_p/8\pi)\sqrt{2\lambda_l/R_H}$; recovered from BCC horizon model (naive equipartition corrected by $H = 8(2\pi)^2$). Phase-unit ansatz is a model assumption
T_p	24	Derived	1.416444×10^{32} K; output of the EWT trinity $(1/k_B)\sqrt{\hbar_{\text{geom}}c^5/G_{\text{geom}}}$
R_H	24.5	Derived	1.382521×10^{26} m; $\lambda_l(T_p/T_{\text{cmb},0})^2/H$; conditional on the phase-unit ansatz
H_0^{EWT}	24.5	Predicted	66.9113 km s^{-1} Mpc^{-1}; from BCC chain + one anchor $T_{\text{cmb},0}$; -0.83σ vs Planck
Λ_{dS}	24.6	Predicted	1.5696×10^{-52} m^{-2}; de Sitter value $3H_0^2/c^2$; 1.441$\times$ Planck (structural)
Ω_{vac}	24.7	Geometric mechanism	$\eta_{\text{BCC}}C_{\text{gap}} = 0.689749$; plays the role of Ω_Λ in ΛCDM; $+0.15\sigma$ vs Planck+BAO. Composition rule is a model assumption; first-principles derivation open
Λ_{vac}	24.7	Derived (consequence)	$\Omega_{\text{vac}}\Lambda_{\text{dS}} = 1.0826 \times 10^{-52}$ m^{-2}; -0.588% vs Planck
V. Roundings of derived quantities (not postulates)			
$r_e = 100\, r_\nu$	5.5	Rounding	Exact ratio 100.0001735 derived from g_v; 100 is a numerical convenience
F_{1+4k}	16.6	Phenomenological label	Classifies the three-generation closure; does not enter AMM numerics
VI. Legacy baseline-EWT constants and auxiliary calibration factors (diagnostic consistency with the baseline EWT model only; not used in any prediction)			

continued on next page

Element	Section	Status	Role in the model
ρ_a	19	Legacy wave constant	Aether density $\rho_a \approx 3.86 \times 10^{22}$ kg/m^3 from Yee's original EWT; enters only `mass_spherical()` as a dimensional prefactor, for diagnostic comparison with the baseline EWT mass formula
A_l	19	Legacy wave constant	Longitudinal wave amplitude $A_l \approx 9.22 \times 10^{-19}$ m from Yee's original EWT; enters only `mass_spherical()`
λ_l (baseline)	19	Legacy wave constant	Longitudinal wavelength $\lambda_l \approx 2.85 \times 10^{-17}$ m from Yee's original EWT; enters only `mass_spherical()`. Note: this is a *different* quantity from the geometric Planck length λ_l of Section 11.6, which is derived from the BCC geometry
δ_μ	19	Orbital multiplier	Empirical amplitude ratio 185.68543 for muon ($K = 20$); enters only `mass_orbital()`; not derived from the BCC lattice
δ_τ	19	Orbital multiplier	Empirical amplitude ratio 3436.795 for tau ($K = 50$); enters only `mass_orbital()`; not derived from the BCC lattice
M_K^{ref}	17	Internal reference	Fixed reference values 248.8 ppm (muon shell) and 1177.21 ppm (tau shell) used in the diagnostic resonance scanner only; not inputs to any AMM prediction

No continuously adjustable parameter appears in the core predictive content of this table. The model's predictive content is fixed once the primary anchors of class I are given, together with the discrete structures of classes II–IV and the roundings of class V. Classes II–V either test the geometry externally, fix discrete structures, or specify parameter-free functional forms.

Class VI lists the *legacy baseline-EWT constants and auxiliary calibration factors*. These appear only in the auxiliary spherical and orbital mass modes and in the diagnostic resonance scanner. They serve two purposes: (i) to allow direct numerical comparison with the baseline EWT mass formula of Yee (whose results are reproduced by `mass_spherical()` and `mass_orbital()`), and (ii) to provide reference benchmarks for the diagnostic resonance scanner of Section 17. None of the class VI entries enters any core prediction of the model. Removing the auxiliary mass modes and the diagnostic scanner would leave the core predictive content unchanged.

The distinction between the classes is therefore tripartite: *primary anchors* (class I: dimensional conversion), *structural choices* (classes II–IV: tested externally, fixed discretely, or given as parameter-free forms), and *auxiliary diagnostic quantities* (class VI: used only to cross-check against the baseline EWT formulation, not used in predictions).

A.3 Computational Scripts (Python)

All numerical results in this work are produced by a set of Python scripts that implement the zero-calibration geometric model described in the main text. The scripts are self-contained and use only standard scientific libraries. The source code and console outputs are listed below. Persistent identifiers for each script will be added in the final published version.

A.3.1 Emergence Engine

`ewt_emergence_engine.py` is the core geometric engine. It derives the fine-structure constant, magnetic deficit, neutrino radius, Planck length, gravitational constant, lepton anomalous magnetic moments, and atomic scales from the BCC lattice geometry. The script also performs the rigidity tests described in Section 29.

```
# From manuscript version: 5.2.4 or leter
# ============================================================================
# SECTION 1: GLOBAL INPUTS & CONSTANTS
# ----------------------------------------------------------------------------
# This section contains:
#   - mathematical constants
#   - experimental reference values (CODATA 2022 / PDG 2022)
#   - pure BCC lattice geometry (independent of the EWT model)
#
# Every quantity is explicitly labelled:
#   INPUT         : experimental value used as an anchor or metric input
#   MATHEMATICAL : pure mathematical constant
#   BCC_GEOMETRY : crystallographic fact about the BCC lattice
#
# The model itself will not define any additional hidden numerical parameters
# in this section. All later parameters must be either DERIVED, POSTULATED,
# or explicitly marked as CALIBRATION (to be eliminated in future versions).
# ============================================================================

import math

# ----------------------------------------------------------------------------
# 1.1 MATHEMATICAL CONSTANTS
# ----------------------------------------------------------------------------

PI      = math.pi                    # circle ratio
EULER   = math.e                     # Euler's number
SQRT2   = math.sqrt(2.0)             # diagonal of unit square
SQRT3   = math.sqrt(3.0)             # diagonal of unit cube face

# ----------------------------------------------------------------------------
# 1.2 CODATA 2022 / PDG 2022 EXPERIMENTAL INPUTS
# ----------------------------------------------------------------------------

# Metric and electromagnetic scales
C0                = 299792458.0              # speed of light [m/s] (exact SI)
M_E               = 9.1093837015e-31         # electron mass [kg]
R_E               = 2.8179403262e-15         # classical electron radius [m]
ALPHA_INV_CODATA  = 137.035999084             # inverse fine-structure constant
A_E_CODATA        = 0.0011596521816           # electron AMM (g-2)/2

# Gravitational scale
G_CODATA          = 6.674305e-11             # gravitational constant [m^3 kg^-1 s^-2]

# Electroweak and flavour scales
M_Z_CODATA        = 91.1876                  # Z boson mass [GeV]
M_W_PDG           = 80.377                   # W boson mass, PDG 2022 average [GeV]
M_W_CDFII         = 80.4335                  # W boson mass, CDF II 2022 [GeV]
M_W_ATLAS_2024    = 80.3600                  # ATLAS 2024 average [GeV]
M_W_CMS_2024      = 80.3602                  # CMS 2024 average [GeV]
M_W_PDG_2024      = 80.3692                  # PDG 2024 average [GeV]
M_H_CODATA        = 125.25                   # Higgs mass [GeV]
SIN2_THETA_W      = 0.23122                  # Weinberg angle sin^2(theta_W)
SIN_THETA_C_PDG   = 0.2243                   # Cabibbo angle sin(theta_C)
# PPM_CONVENTION = 10^-6 is the parts-per-million reporting convention.
# It happens to coincide numerically with (r_nu/r_e)^3 because the decadic
# anchor r_e/r_nu ~ 100 = 10^2 has a cube (10^2)^3 = 10^6. This is a
# numerical coincidence of the decimal system, not a derivation from the
# BCC lattice: if the decadic anchor were non-decimal, the ppm convention
# would be unchanged but would not coincide with any simple geometric
# ratio of the model.
PPM_CONVENTION = 1.0e-6
R_INF_CODATA      = 10973731.568157          # Rydberg constant [m^-1]
A0_CODATA         = 5.29177210903e-11        # Bohr radius [m]
LAMBDA_C_CODATA   = 2.42631023867e-12        # electron Compton wavelength [m]
```

```
4818
4819 # Quark masses (MS-bar, PDG 2022) used in flavour mixing test
4820 M_D_PDG            = 0.00470                    # d-quark mass [GeV], PDG 2024
4821 M_S_PDG            = 0.09350                    # s-quark mass [GeV], PDG 2024
4822
4823 A_MU_EXP           = 1.1659206100e-3        # muon AMM from Fermilab/Brookhaven
4824 A_TAU_EXP          = 1.177210e-3            # tau AMM: Standard Model prediction, not a direct
4825       measurement
4826
4827 # Reference value of the Planck charge (interpreted as a geometric amplitude).
4828 # The DERIVED value is q_P_geom = e / sqrt(alpha_geom); see derive_planck_charge_from_e().
4829 # Q_P_REFERENCE is retained only as a cross-check anchor for the main() report.
4830 Q_P_REFERENCE      = 1.87554603778e-18          # Planck charge reference value [m]
4831 # Fundamental EMC spacing (Planck length) [m]
4832 LAMBDA_L           = 1.6162e-35
4833 # Elementary charge (CODATA 2022) -- empirical anchor of the amplitude scale
4834 E_CHARGE_CODATA    = 1.602176634e-19
4835
4836 # ------------------------------------------------------------------------------
4837 # 1.3 PURE BCC LATTICE GEOMETRY
4838 # ------------------------------------------------------------------------------
4839 # These are purely mathematical / crystallographic facts about the BCC lattice.
4840 # They are independent of the Enhanced EWT model and contain no physics inputs.
4841 # ------------------------------------------------------------------------------
4842
4843 BCC_NEIGHBOURS           = 8                       # nearest neighbours in BCC
4844 BCC_PACKING_FRACTION     = (SQRT3 * PI) / 8.0      # ~0.6801748
4845 BCC_IDEAL_PROJECTION_LP  = 2.0 / SQRT3             # inverse nearest-neighbour distance
4846
4847 # ------------------------------------------------------------------------------
4848 # 1.4 STATUS LABELS
4849 # ------------------------------------------------------------------------------
4850 # These labels will be used in all reporting functions to make the origin
4851 # of each quantity transparent.
4852
4853 STATUS = {
4854     "INPUT":        "Experimental␣input␣(CODATA/PDG)",
4855     "MATHEMATICAL": "Pure␣mathematical␣constant",
4856     "BCC_GEOMETRY": "Pure␣BCC␣lattice␣geometry",
4857     "POSTULATE":    "Geometric␣postulate␣of␣the␣model",
4858     "DERIVED":      "Derived␣from␣model␣postulates/constraints",
4859     "CALIBRATION":  "Calibrated␣parameter␣(to␣be␣eliminated)",
4860 }
4861
4862 # ==============================================================================
4863 # SECTION 2: GEOMETRIC POSTULATES & ALPHA CORE EMERGENCE
4864 # ------------------------------------------------------------------------------
4865 # The alpha core emerges from three geometric postulates:
4866 #   P1: charge = amplitude (x)
4867 #   P2: emission surface = sphere + cone
4868 #   P3: natural length ratios r = x, l = pi*x
4869 #
4870 # These postulates lead directly to the geometric ratio:
4871 #   alpha_core^{-1} = S_total / x^2 = 4*pi^3 + pi^2 + pi
4872 # ==============================================================================
4873
4874 def compute_emission_surface(x: float) -> dict:
4875     """
4876     Compute the total emission surface and the resulting alpha core
4877     from the geometric postulates.
4878
4879     Parameters
4880     ----------
4881     x : float
4882         Wave amplitude (charge). The final ratio is independent of x.
4883
4884     Returns
4885     -------
4886     dict with:
4887         r          : base radius of the cone (equals x)
4888         l          : propagation distance (equals pi*x)
4889         S_sphere   : surface area of the spherical part
4890         S_cone     : total surface area of the conical part
```

```
        S_total   : total emission surface
        A_pi      : alpha core inverse = S_total / x^2
    """
    r = x
    l = PI * x

    S_sphere = 4.0 * PI * l**2
    S_cone = PI * r * l + PI * r**2

    S_total = S_sphere + S_cone
    A_pi = S_total / x**2

    return {
        "r": r,
        "l": l,
        "S_sphere": S_sphere,
        "S_cone": S_cone,
        "S_total": S_total,
        "A_pi": A_pi,
    }

def compute_alpha_core() -> float:
    """
    Compute the purely geometric core of the fine-structure constant.
    It is obtained from the emission surface ratio for a unit amplitude.

    Returns
    -------
    float
        A_pi = 4*pi^3 + pi^2 + pi
    """
    return compute_emission_surface(1.0)["A_pi"]

# ============================================================================
# SECTION 2B: GEOMETRIC LADDER EMERGENCE
# ----------------------------------------------------------------------------
# The geometric ladder is not an input. It is derived from the active
# degrees of freedom of a soliton embedded in the BCC vacuum lattice.
#
# The central modulator eps_M is treated as an INPUT to this module.
# In the current zero-calibration version it is obtained from pure BCC geometry:
#   - eps_M derived from packing fraction eta and lattice impedance zeta
#   - historical calibrated variant (N_final) is no longer used
#
# Later sections will compute eps_M and pass it here.
# ============================================================================

def build_geometric_ladder(eps_M: float, source_label: str = "",
                           zeta: float = 0.0) -> dict:
    """
    Build the geometric ladder from the magnetic deficit eps_M.

    Parameters
    ----------
    eps_M : float
        Magnetic deficit parameter.
    source_label : str
        Optional label describing the source of eps_M.
    zeta : float
        Lattice impedance (default 0.0 for ideal BCC packing).

    Returns
    -------
    dict with:
        eps_M       : magnetic deficit
        zeta        : lattice impedance
        source      : source label
        ...
    """
    C_local = eps_M / (2.0 * SQRT2)

    pi6 = PI**6
    pi7 = PI**7
```

```
    O_mixed = (pi6 * pi7) / (pi6 + pi7)

    # Flavor deficit (analogous to eps_M, at the pi^5 rung)
    eps_flavor = 1.0 / (8.0 * pi7 * (1.0 - zeta) / PI**2)  # = 1/(8*pi^5*(1-zeta))
    # Simplified: eps_flavor = 1 / (8 * PI**5 * (1 - zeta))

    return {
        "eps_M": eps_M,
        "zeta": zeta,
        "eps_flavor": eps_flavor,
        "source": source_label,
        "C_local": C_local,
        "C_gap": 1.0 + pi6 * C_local,
        "C_gap_mixed": 1.0 + O_mixed * C_local,
        "O_mixed": O_mixed,
        "ladder": {n: PI**n for n in range(3, 8)},
    }


def report_ladder(lad: dict) -> None:
    """Print the contents of a ladder dictionary in a readable form."""
    print(f"  Source       : {lad.get('source', 'unknown')}")
    print(f"  eps_M        : {lad['eps_M']:.15e}")
    print(f"  zeta         : {lad.get('zeta', 0.0):.10e}")
    print(f"  C_local      : {lad['C_local']:.15e}")
    print(f"  C_gap        : {lad['C_gap']:.15f}   (pure pi^6, Z-Z and Z-H)")
    print(f"  C_gap_mixed  : {lad['C_gap_mixed']:.15f}   (mixed W-Z)")
    print(f"  O_mixed      : {lad['O_mixed']:.6f}   (harmonic mean pi^6*pi^7/(pi^6+pi^7))")
    print(f"  eps_flavor   : {lad['eps_flavor']:.15f}")
    print("  Ladder:")
    for n in sorted(lad["ladder"]):
        print(f"    pi^{n} = {lad['ladder'][n]:.15f}")

# ============================================================================
# SECTION 3: BCC-DERIVED LATTICE IMPEDANCE
# ----------------------------------------------------------------------------
# Functions that derive eps_M from pure BCC geometry using the packing
# fraction and lattice impedance.
# ============================================================================

def lattice_impedance(N: float, eta: float = BCC_PACKING_FRACTION) -> float:
    """
    Compute the lattice impedance zeta for a given geometric stiffness N.

    Formula:
        zeta = (1 - eta) / (eta * N)

    Parameters
    ----------
    N : float
        Geometric stiffness modulus (e.g. 8*pi^4, 8*pi^7).
    eta : float
        Packing fraction of the BCC lattice.

    Returns
    -------
    float
        Lattice impedance.
    """
    return (1.0 - eta) / (eta * N)


def derive_eps_M_from_BCC(N_input: float = 8.0 * PI**4) -> dict:
    """
    Derive eps_M from BCC geometry using the lattice impedance.

    Parameters
    ----------
    N_input : float
        Ideal geometric stiffness to use as base.
        Default is 8*pi^4 (BCC saturation stiffness).

    Returns
```

```
    -------
    dict with:
        eta       : BCC packing fraction
        N_ideal   : input ideal stiffness
        zeta      : lattice impedance
        N_geom    : effective stiffness after impedance correction
        eps_M     : magnetic deficit derived from BCC geometry
    """
    eta = BCC_PACKING_FRACTION
    zeta = lattice_impedance(N_input, eta)
    N_geom = N_input * (1.0 - zeta)
    eps_M = 1.0 / (N_geom * PI**3)

    return {
        "eta": eta,
        "N_ideal": N_input,
        "zeta": zeta,
        "N_geom": N_geom,
        "eps_M": eps_M,
    }

# ============================================================================
# SECTION 4: TEST FUNCTIONS (EXPERIMENTAL VALIDATION)
# ----------------------------------------------------------------------------
# These functions compare model predictions with experimental data.
# They do NOT influence the model itself.
# ============================================================================

def weinberg_sector(C_gap: float, M_Z: float, sin2_W: float) -> dict:
    """
    Compute the W boson mass from the Weinberg angle using a given C_gap operator.

    Since v5.2.0, the main M_W prediction uses the mixed W-Z operator
    (see weinberg_sector_mixed). This function remains for:
      - pure pi^6 calculations (neutral vertices Z-Z, Z-H),
      - backward compatibility with older scripts.
    """
    M_W_pred = M_Z * math.sqrt((1.0 - sin2_W) * C_gap)

    rel_err_PDG = abs(M_W_pred - M_W_PDG) / M_W_PDG * 100.0
    rel_err_CDFII = abs(M_W_pred - M_W_CDFII) / M_W_CDFII * 100.0
    rel_err_PDG_2024 = abs(M_W_pred - M_W_PDG_2024) / M_W_PDG_2024 * 100.0

    return {
        "M_W_pred": M_W_pred,
        "rel_err_PDG": rel_err_PDG,
        "rel_err_CDFII": rel_err_CDFII,
        "rel_err_PDG_2024": rel_err_PDG_2024,
    }


def weinberg_sector_mixed(lad: dict, M_Z: float, sin2_W: float) -> dict:
    """
    Compute W mass via the mixed W-Z operator (harmonic mean pi^6 * pi^7).

    The W-Z vertex is a mixed process: one charged (W) and one neutral
    (Z) soliton. The harmonic combination
        O_mixed = pi^6 * pi^7 / (pi^6 + pi^7)
    is the unique simple parameter-free form that reduces to the correct
    single-rung limit when one channel decouples. It is a methodologically
    motivated ansatz, not a first-principles derivation.

    Parameters
    ----------
    lad : dict
        Ladder dictionary from build_geometric_ladder(); must contain
        "C_gap_mixed" and "O_mixed".
    M_Z : float
        Z-boson mass [GeV].
    sin2_W : float
        MS-bar Weinberg angle.

    Returns
```

```
    -------
    dict with:
        M_W_pred          : predicted W mass [GeV]
        C_gap_mixed       : the mixed operator value used
        O_mixed           : harmonic mean of pi^6 and pi^7
        rel_err_PDG       : relative error vs PDG 2022 [%]
        rel_err_CDFII     : relative error vs CDF II 2022 [%]
        rel_err_PDG_2024  : relative error vs PDG 2024 [%]
        rel_err_ATLAS     : relative error vs ATLAS 2024 [%]
        rel_err_CMS       : relative error vs CMS 2024 [%]
    """
    C_gap_mixed = lad["C_gap_mixed"]
    M_W_pred = M_Z * math.sqrt((1.0 - sin2_W) * C_gap_mixed)

    return {
        "M_W_pred": M_W_pred,
        "C_gap_mixed": C_gap_mixed,
        "O_mixed": lad["O_mixed"],
        "rel_err_PDG": abs(M_W_pred - M_W_PDG) / M_W_PDG * 100.0,
        "rel_err_CDFII": abs(M_W_pred - M_W_CDFII) / M_W_CDFII * 100.0,
        "rel_err_PDG_2024": abs(M_W_pred - M_W_PDG_2024) / M_W_PDG_2024 * 100.0,
        "rel_err_ATLAS": abs(M_W_pred - M_W_ATLAS_2024) / M_W_ATLAS_2024 * 100.0,
        "rel_err_CMS": abs(M_W_pred - M_W_CMS_2024) / M_W_CMS_2024 * 100.0,
    }


def cabibbo_sector(
    M_d: float, M_s: float,
    eps_flavor: float = None,
    target: float = SIN_THETA_C_PDG,
) -> dict:
    """
    Compute the Cabibbo angle from the bare Gatto-Sartori-Tonin
    relation with the EWT flavor deficit correction:

        sin(theta_C) = sqrt(M_d/M_s) * (1 + eps_flavor)

    where eps_flavor = 1/(8*pi^5*(1-zeta)) by default, or
    1/(8*pi^5) in the ideal limit (zeta = 0).

    Parameters
    ----------
    M_d, M_s : float
        Down and strange quark masses [GeV].
    eps_flavor : float, optional
        Flavor deficit. If None, uses ideal value 1/(8*pi^5).
    target : float
        Reference sin(theta_C), default PDG value.

    Returns
    -------
    dict with keys:
        sin_C_pred : predicted sin(theta_C)
        rel_err    : relative error vs target [%]
        eps_flavor : the flavor deficit value used
    """
    if eps_flavor is None:
        eps_flavor = 1.0 / (8.0 * PI**5)  # ideal limit
    sin_C_pred = math.sqrt(M_d / M_s) * (1.0 + eps_flavor)
    rel_err = abs(sin_C_pred - target) / target * 100.0
    return {
        "sin_C_pred": sin_C_pred,
        "rel_err": rel_err,
        "eps_flavor": eps_flavor,
    }


def run_sector_tests(lad: dict) -> None:
    """
    Run experimental validation tests on a given ladder dictionary.

    Since v5.2.0, the W-mass prediction uses the mixed W-Z operator
    (harmonic mean of pi^6 and pi^7). The pure pi^6 result is shown
```

```
    for comparison and corresponds to a hypothetical neutral W boson.

    Parameters
    ----------
    lad : dict
        Ladder dictionary produced by build_geometric_ladder().
    """
    # Mixed operator (main prediction, v5.2.0)
    result_W_mixed = weinberg_sector_mixed(lad, M_Z_CODATA, SIN2_THETA_W)
    print(f"  M_W prediction (mixed)    : {result_W_mixed['M_W_pred']:.6f} GeV")
    print(f"  rel. err PDG 2024 (mixed) : {result_W_mixed['rel_err_PDG_2024']:.4f} %")
    print(f"  rel. err ATLAS 2024       : {result_W_mixed['rel_err_ATLAS']:.4f} %")
    print(f"  rel. err CMS 2024         : {result_W_mixed['rel_err_CMS']:.4f} %")
    print(f"  rel. err CDF II 2022      : {result_W_mixed['rel_err_CDFII']:.4f} %   (outlier)"
        )

    # Pure pi^6 (for comparison / neutral vertices)
    result_W_pure = weinberg_sector(lad["C_gap"], M_Z_CODATA, SIN2_THETA_W)
    print(f"  M_W (pure pi^6)           : {result_W_pure['M_W_pred']:.6f} GeV")
    print(f"  rel. err PDG 2024 (pure)  : {result_W_pure['rel_err_PDG_2024']:.4f} %")

    result_C = cabibbo_sector(
        M_D_PDG, M_S_PDG,
        eps_flavor=lad.get("eps_flavor"),
    )
    print(f"  sin theta_C               : {result_C['sin_C_pred']:.10f}")
    print(f"  rel. err PDG              : {result_C['rel_err']:.6f} %")
    print(f"  eps_flavor                : {result_C['eps_flavor']:.6e}")
    print(f"  zeta                      : {lad.get('zeta', 0.0):.6e}")


# ============================================================================
# SECTION 5: ALPHA CORE EMERGENCE
# ----------------------------------------------------------------------------
# The alpha core is a purely geometric object emerging from three postulates:
#   P1: charge = amplitude
#   P2: cone--sphere emission geometry
#   P3: natural length ratios (r = x, l = pi*x)
#
# This section contains only two pure functions:
#   compute_alpha_core()  -> A_pi
#   compute_alpha_geometric(eps_M) -> full geometric inverse alpha
# ============================================================================

def compute_alpha_core() -> float:
    """
    Compute the purely geometric core of the fine-structure constant
    from the emission surface ratio.

    The derivation uses three geometric postulates:
      P1: charge = amplitude (x)
      P2: emission surface = sphere + cone
      P3: natural length ratios: r = x, l = pi*x

    Returns
    -------
    float
        A_pi = S_total / x^2 = 4*pi^3 + pi^2 + pi
    """
    x = 1.0
    r = x
    l = PI * x

    S_sphere = 4.0 * PI * l**2
    S_cone   = PI * r * l + PI * r**2

    S_total = S_sphere + S_cone
    A_pi = S_total / x**2

    return A_pi


def compute_alpha_geometric(eps_M: float) -> float:
```

```
    """
    Compute the full geometric inverse fine-structure constant.

    Formula:
        alpha^{-1} = A_pi - eps_M

    Parameters
    ----------
    eps_M : float
        Magnetic deficit parameter.

    Returns
    -------
    float
        Geometric inverse fine-structure constant.
    """
    A_pi = compute_alpha_core()
    return A_pi - eps_M

def test_alpha_sector(alpha_inv_pred: float) -> dict:
    """
    Compare the predicted inverse fine-structure constant with CODATA 2022.

    Parameters
    ----------
    alpha_inv_pred : float
        Predicted inverse fine-structure constant.

    Returns
    -------
    dict with:
        alpha_inv_pred : predicted value
        abs_err        : absolute error vs CODATA
        rel_err        : relative error in percent
    """
    abs_err = abs(alpha_inv_pred - ALPHA_INV_CODATA)
    rel_err = abs_err / ALPHA_INV_CODATA * 100.0

    return {
        "alpha_inv_pred": alpha_inv_pred,
        "abs_err": abs_err,
        "rel_err": rel_err,
    }

# ============================================================================
# SECTION 6: NEUTRINO & ELECTRON ANCHORS
# ----------------------------------------------------------------------------
# This section derives hbar, Planck charge, uncorrected wavelength, and
# the neutrino radius from geometric alpha and electron anchors.
# ============================================================================

def derive_planck_charge_from_e(alpha_geom: float, e_charge: float) -> float:
    """
    Derive Planck charge (fundamental geometric amplitude) from
    geometric alpha and the elementary charge amplitude e.

    Formula:
        q_P = e / sqrt(alpha_geom)

    Note: This is NOT a tautology because alpha_geom is derived
    from pure BCC geometry (A_pi - eps_M), not from e or q_P.
    """
    return e_charge / math.sqrt(alpha_geom)


def derive_hbar(alpha_geom: float, r_e: float, m_e: float, c0: float) -> float:
    """
    Derive the reduced Planck constant from geometric alpha and electron data.

    Formula:
        hbar = m_e * c * r_e / alpha_geom

    Parameters
```

```
    ----------
    alpha_geom : float
        Geometric fine-structure constant (dimensionless).
    r_e : float
        Classical electron radius [m].
    m_e : float
        Electron mass [kg].
    c0 : float
        Speed of light [m/s].

    Returns
    -------
    float
        Reduced Planck constant [kg m^2 / s].
    """
    return (m_e * c0 * r_e) / alpha_geom


def derive_lambda_uncorr(q_P: float) -> float:
    """
    Compute the uncorrected neutrino wavelength from Planck charge.

    Formula:
        lambda_uncorr = 2 * q_P * e^2

    where e is Euler's number.

    Parameters
    ----------
    q_P : float
        Planck charge (in EWT interpreted as length).

    Returns
    -------
    float
        Uncorrected wavelength [m].
    """
    return 2.0 * q_P * EULER**2


def derive_neutrino_radius(alpha_geom: float, q_P: float) -> dict:
    """
    Derive the neutrino radius r_nu from the geometric fixed point of g_v.

    The derivation uses the self-consistent equation:
        S_tot = alpha_geom_inv/(8+pi) + e + (1-g_v)*(sqrt2-1)
        S_tot = 2 e^2 / g_v
    which leads to a quadratic equation for g_v.

    Parameters
    ----------
    alpha_geom : float
        Geometric fine-structure constant.
    q_P : float
        Planck charge / geometric amplitude [m].

    Returns
    -------
    dict with:
        alpha_inv_geom : geometric inverse alpha
        S_proj         : static lattice projection
        S_exp          : dynamic wave expansion (e)
        delta_imp      : lattice impedance term
        gv_pred        : geometric fixed point for g_v
        S_tot          : total scaling factor
        r_nu           : neutrino radius [m]
    """
    alpha_inv_geom = 1.0 / alpha_geom

    S_proj = alpha_inv_geom / (BCC_NEIGHBOURS + PI)
    S_exp = EULER

    a_coef = SQRT2 - 1.0
```

```
    b_coef = -(S_proj + EULER + SQRT2 - 1.0)
    c_coef = 2.0 * EULER**2

    discriminant = b_coef**2 - 4.0 * a_coef * c_coef
    if discriminant < 0:
        raise ValueError("Negative discriminant in g_v fixed point equation")

    sqrt_disc = math.sqrt(discriminant)
    root1 = (-b_coef + sqrt_disc) / (2.0 * a_coef)
    root2 = (-b_coef - sqrt_disc) / (2.0 * a_coef)

    if 0.0 < root1 < 1.0:
        gv_pred = root1
    elif 0.0 < root2 < 1.0:
        gv_pred = root2
    else:
        raise ValueError("No physical root for g_v in (0,1)")

    delta_imp = (1.0 - gv_pred) * (SQRT2 - 1.0)
    S_tot = S_proj + S_exp + delta_imp
    r_nu = q_P * S_tot

    return {
        "alpha_inv_geom": alpha_inv_geom,
        "S_proj": S_proj,
        "S_exp": S_exp,
        "delta_imp": delta_imp,
        "gv_pred": gv_pred,
        "S_tot": S_tot,
        "r_nu": r_nu,
    }


def test_decadic_resonance(r_e: float, r_nu: float) -> dict:
    """
    Test the 1:100 radial resonance between electron and neutrino.

    Parameters
    ----------
    r_e : float
        Classical electron radius [m].
    r_nu : float
        Neutrino radius [m].

    Returns
    -------
    dict with:
        ratio      : r_e / r_nu
        expected   : 100.0
        rel_err    : relative error in percent
        K_implied  : ratio^5
    """
    ratio = r_e / r_nu
    rel_err = abs(ratio - 100.0) / 100.0 * 100.0
    K_implied = ratio**5
    return {
        "ratio": ratio,
        "expected": 100.0,
        "rel_err": rel_err,
        "K_implied": K_implied,
    }

def compute_C_unif(alpha_geom: float, L_p_geom: float, K_WC: int) -> float:
    """
    Compute the unified coupling operator C_Unif.

    Formula:
        C_Unif = 1/K_WC + 1 + alpha_geom / (pi * L_p_geom)

    Parameters
    ----------
    alpha_geom : float
        Geometric fine-structure constant.
```

```
    L_p_geom : float
        Ideal BCC projection factor.
    K_WC : int
        Number of wave centres in the electron.

    Returns
    -------
    float
        Unified coupling operator.
    """
    return (1.0 / K_WC) + 1.0 + (alpha_geom / (PI * L_p_geom))


def compute_X_eff(alpha_geom: float, L_p_geom: float, K_WC: int) -> float:
    """
    Compute the geometric dilution factor X_eff.

    Formula:
        X_eff = A_pi * 3 * K_WC * sqrt(2) / C_Unif

    Parameters
    ----------
    alpha_geom : float
        Geometric fine-structure constant.
    L_p_geom : float
        Ideal BCC projection factor.
    K_WC : int
        Number of wave centres in the electron.

    Returns
    -------
    float
        Dilution factor X_eff.
    """
    A_pi = compute_alpha_core()
    C_Unif = compute_C_unif(alpha_geom, L_p_geom, K_WC)
    return (A_pi * 3.0 * K_WC * SQRT2) / C_Unif

def derive_lambda_l_geometric(
    alpha_geom: float,
    r_e: float,
    r_nu: float,
    N_geom: float,
    L_p_geom: float,
    K_WC: int,
) -> float:
    """
    Derive the fundamental EMC length lambda_l from pure geometry.

    The formula is the analytic solution of the self-consistent equation
    obtained by combining the geometric G with the Planck length definition.
    It does NOT use hbar or G as input.

    Formula:
        lambda_l = [ r_e^2 * sqrt(X_eff) /
                     ( alpha_geom * A_pi^4 * N_geom^3 * K_WC *
                       (r_nu/(2*e))^(3/2) ) ]^2

    Parameters
    ----------
    alpha_geom : float
        Geometric fine-structure constant.
    r_e : float
        Classical electron radius [m].
    r_nu : float
        Neutrino radius [m].
    N_geom : float
        Effective BCC stiffness.
    L_p_geom : float
        Ideal BCC projection factor.
    K_WC : int
        Number of wave centres in the electron.

```

```
    Returns
    -------
    float
        Derived lambda_l [m].
    """
    A_pi = compute_alpha_core()
    X_eff = compute_X_eff(alpha_geom, L_p_geom, K_WC)

    r_nu_factor = (r_nu / (2.0 * EULER)) ** (3.0 / 2.0)

    bracket = (
        r_e**2 * math.sqrt(X_eff)
        / (alpha_geom * A_pi**4 * N_geom**3 * K_WC * r_nu_factor)
    )

    return bracket**2

# ============================================================================
# SECTION 7: GRAVITY SECTOR EMERGENCE
# ----------------------------------------------------------------------------
# Gravity is derived from the electron soliton and the BCC lattice geometry.
# No calibrated N_final and no calibrated L_p are used.
#
#   N_geom       : derived from BCC packing impedance
#   L_p_geom     : ideal BCC projection factor 2/sqrt(3)
#   alpha_geom   : derived from A_pi and eps_M
#
# The effective volume deficit is computed via the unified coupling:
#   C_Unif = 1/K_WC + 1 + alpha_geom/(pi * L_p_geom)
# ============================================================================

def gravity_sector(
    alpha_geom: float,
    r_nu: float,
    N_geom: float,
    L_p_geom: float,
    K_WC: int,
    lambda_l: float,
    r_e: float,
    m_e: float,
    c0: float,
) -> dict:
    """
    Derive the gravitational constant G from pure geometry and electron anchors.

    Parameters
    ----------
    alpha_geom : float
        Geometric fine-structure constant.
    r_nu : float
        Neutrino radius [m].
    N_geom : float
        Effective BCC stiffness derived from packing impedance.
    L_p_geom : float
        Ideal BCC projection factor (2/sqrt(3)).
    K_WC : int
        Number of wave centres in the electron (10).
    lambda_l : float
        Fundamental EMC spacing derived from geometry [m].
    r_e : float
        Classical electron radius [m].
    m_e : float
        Electron mass [kg].
    c0 : float
        Speed of light [m/s].

    Returns
    -------
    dict with:
        G_Base        : base soliton scaling
        N_nu_statutory: statutory background density
        C_Unif        : unified coupling operator
        X_eff         : dilution factor
```

```
        N_nu_eff      : effective volume deficit
        G_EWT         : predicted gravitational constant
        rel_err       : relative error vs CODATA
    """
    G_Base = (c0**2 * r_e) / m_e
    A_pi = compute_alpha_core()
    C_Unif = compute_C_unif(alpha_geom, L_p_geom, K_WC)
    X_eff = compute_X_eff(alpha_geom, L_p_geom, K_WC)
    N_nu_statutory = (r_nu / (2.0 * lambda_l * EULER)) ** 3
    N_nu_eff = N_nu_statutory / X_eff

    G_EWT = (
        (G_Base / A_pi)
        * (1.0 / (N_geom * A_pi)) ** 3
        * (1.0 / (K_WC * math.sqrt(N_nu_eff)))
    )

    rel_err = abs(G_EWT - G_CODATA) / G_CODATA * 100.0

    return {
        "G_Base": G_Base,
        "N_nu_statutory": N_nu_statutory,
        "C_Unif": C_Unif,
        "X_eff": X_eff,
        "N_nu_eff": N_nu_eff,
        "G_EWT": G_EWT,
        "rel_err": rel_err,
    }


def report_gravity(res: dict) -> None:
    """Print gravity sector results."""
    print(f"  G_Base        : {res['G_Base']:.15e}")
    print(f"  N_nu_statutory: {res['N_nu_statutory']:.15e}")
    print(f"  C_Unif        : {res['C_Unif']:.15f}")
    print(f"  X_eff         : {res['X_eff']:.10f}")
    print(f"  N_nu_eff      : {res['N_nu_eff']:.15e}")
    print(f"  G_EWT         : {res['G_EWT']:.15e}")
    print(f"  G_CODATA      : {G_CODATA:.15e}")
    print(f"  rel_error     : {res['rel_err']:.6f} %")

# ============================================================================
# MAIN
# ============================================================================

def rigidity_test(
    alpha_geom: float,
    r_e: float,
    r_nu: float,
    N_geom: float,
    L_p_geom: float,
    K_WC: int,
    lambda_l_geom: float,
    m_e: float,
    c0: float,
) -> None:
    """
    Test the geometric rigidity of the model by perturbing key parameters.

    Shows how relative changes in N, L_p, or lambda_l propagate into
    alpha, G, and the derived lambda_l.
    """
    print("\n" + "=" * 78)
    print("RIGIDITY TEST")
    print("=" * 78)

    print("\n[1] Vary N_geom")
    print("-" * 78)
    print(f"{'N_geom':>12} {'alpha_inv':>14} {'lambda_l [m]':>16} {'G_EWT [m^3/kg/s^2]':>22} {'G_err %':>10}")
    for factor in [0.995, 0.998, 1.0, 1.002, 1.005]:
        N_test = N_geom * factor
        eps_M_test = 1.0 / (N_test * PI**3)
```

```
        alpha_inv_test = compute_alpha_geometric(eps_M_test)
        alpha_test = 1.0 / alpha_inv_test

        lambda_test = derive_lambda_l_geometric(
            alpha_geom=alpha_test,
            r_e=r_e,
            r_nu=r_nu,
            N_geom=N_test,
            L_p_geom=L_p_geom,
            K_WC=K_WC,
        )

        res = gravity_sector(
            alpha_geom=alpha_test,
            r_nu=r_nu,
            N_geom=N_test,
            L_p_geom=L_p_geom,
            K_WC=K_WC,
            lambda_l=lambda_test,
            r_e=r_e,
            m_e=m_e,
            c0=c0,
        )

        print(
            f"{N_test:12.5f} {alpha_inv_test:14.9f} {lambda_test:16.6e} "
            f"{res['G_EWT']:22.15e} {res['rel_err']:10.6f}"
        )

    print("\n[2] Vary L_p_geom")
    print("-" * 78)
    print(f"{'L_p':>10} {'alpha_inv':>14} {'lambda_l [m]':>16} {'G_EWT [m^3/kg/s^2]':>22} {'G_err %':>10}")
    for factor in [0.995, 0.998, 1.0, 1.002, 1.005]:
        Lp_test = L_p_geom * factor

        lambda_test = derive_lambda_l_geometric(
            alpha_geom=alpha_geom,
            r_e=r_e,
            r_nu=r_nu,
            N_geom=N_geom,
            L_p_geom=Lp_test,
            K_WC=K_WC,
        )

        res = gravity_sector(
            alpha_geom=alpha_geom,
            r_nu=r_nu,
            N_geom=N_geom,
            L_p_geom=Lp_test,
            K_WC=K_WC,
            lambda_l=lambda_test,
            r_e=r_e,
            m_e=m_e,
            c0=c0,
        )

        print(
            f"{Lp_test:10.6f} {1.0/alpha_geom:14.9f} {lambda_test:16.6e} "
            f"{res['G_EWT']:22.15e} {res['rel_err']:10.6f}"
        )

    print("\n[3] Vary lambda_l directly")
    print("-" * 78)
    print(f"{'lambda_l [m]':>16} {'G_EWT [m^3/kg/s^2]':>22} {'G_err %':>10}")
    for factor in [0.99, 1.0, 1.01]:
        lambda_test = lambda_l_geom * factor
        res = gravity_sector(
            alpha_geom=alpha_geom,
            r_nu=r_nu,
            N_geom=N_geom,
            L_p_geom=L_p_geom,
            K_WC=K_WC,
```

```
            lambda_l=lambda_test,
            r_e=r_e,
            m_e=m_e,
            c0=c0,
        )
        print(
            f"{lambda_test:16.6e} {res['G_EWT']:22.15e} {res['rel_err']:10.6f}"
        )

    print("=" * 78)

def input_rigidity_test(base_alpha_geom, base_r_nu, base_N_geom, base_Lp):
    print("\n[4] Vary fundamental inputs r_e and m_e")
    print("-" * 78)
    print(f"{'factor':>8} {'r_e [m]':>16} {'m_e [kg]':>16} {'lambda_l [m]':>18} {'G_err
        %':>10}")

    for factor in [0.998, 0.999, 1.0, 1.001, 1.002]:
        r_e_test = R_E * factor
        m_e_test = M_E * factor

        lambda_test = derive_lambda_l_geometric(
            alpha_geom=base_alpha_geom,
            r_e=r_e_test,
            r_nu=base_r_nu,
            N_geom=base_N_geom,
            L_p_geom=base_Lp,
            K_WC=10,
        )

        res = gravity_sector(
            alpha_geom=base_alpha_geom,
            r_nu=base_r_nu,
            N_geom=base_N_geom,
            L_p_geom=base_Lp,
            K_WC=10,
            lambda_l=lambda_test,
            r_e=r_e_test,
            m_e=m_e_test,
            c0=C0,
        )

        print(
            f"{factor:8.3f} {r_e_test:16.6e} {m_e_test:16.6e} "
            f"{lambda_test:18.6e} {res['rel_err']:10.6f}"
        )

# ============================================================================
# SECTION 8: LEPTON ANOMALOUS MAGNETIC MOMENTS (AMM)
# ============================================================================

def get_AMMi_K(n: int) -> int:
    """
    Recursive nodal count for the lepton generations.

    K_1 = 10
    K_n = K_{n-1} + round(10^(n-1) * 2*pi^2)

    Parameters
    ----------
    n : int
        Generation number (1 = electron, 2 = muon, 3 = tau).

    Returns
    -------
    int
        Total nodal count K_n.
    """
    if n == 1:
        return 10
    else:
        delta_K = round(10 ** (n - 1) * (2.0 * PI**2))
        return get_AMMi_K(n - 1) + delta_K
```

```python


def compute_lepton_amms(
    alpha_geom: float,
    eps_M: float,
    L_mu_dim: int = 5,
    L_tau_dim: int = 34,  # Reserved for the 4th-generation extension
                          # (see Section sec:three_generations in the manuscript);
                          # not used in the three-generation AMM computation.
    K_WC_e: int = 10,
) -> dict:
    """
    Compute the full anomalous magnetic moments for the lepton family.
    """
    A_pi = compute_alpha_core()

    a_e_ppm = (alpha_geom / (2.0 * PI)) * (1.0 - eps_M * (PI**3)) / PPM_CONVENTION

    K_e = K_WC_e
    K_mu_total = get_AMMi_K(2)
    K_mu_delta = K_mu_total - K_e
    M_mu_shell = K_mu_delta / K_e

    B_mu_scale = (3.0 * A_pi * PI**3) / (2.0 * L_mu_dim**2)
    a_mu_shell_ppm = B_mu_scale * (1.0 - eps_M) ** (M_mu_shell * PI**3)

    O_mu = 1.0 / (4.0 * PI**2)
    a_mu_shell_correction = a_mu_shell_ppm * O_mu
    a_mu_ppm = a_e_ppm + a_mu_shell_correction

    K_tau_total = get_AMMi_K(3)
    M_tau_rel = K_tau_total / K_e

    B_tau_base = ((3.0 * A_pi * PI**3) / (8.0 * SQRT2)) + (A_pi / 2.0)
    a_tau_shell_raw_ppm = B_tau_base * (1.0 - eps_M) ** (M_tau_rel * PI**3)

    a_tau_shell_total_ppm = a_mu_shell_ppm + a_tau_shell_raw_ppm + L_mu_dim**2

    O_tau = 1.0
    a_tau_ppm = a_e_ppm + O_tau * (a_tau_shell_total_ppm - a_e_ppm)

    return {
        "a_e_ppm": a_e_ppm,
        "a_mu_ppm": a_mu_ppm,
        "a_tau_ppm": a_tau_ppm,
        "a_mu_shell_ppm": a_mu_shell_ppm,
        "a_tau_shell_ppm": a_tau_shell_total_ppm,
    }


def report_lepton_amms(res: dict) -> None:
    """Print lepton AMM results and compare with experimental / SM references."""
    print("--- Lepton Anomalous Magnetic Moments ---")
    print(f"  Electron a_e   : {res['a_e_ppm']:.6f} ppm  (CODATA: {A_E_CODATA/PPM_CONVENTION
        :.6f} ppm)")
    print(f"  Muon shell     : {res['a_mu_shell_ppm']:.6f} ppm")
    print(f"  Muon full a_mu : {res['a_mu_ppm']:.6f} ppm  (Exp: {A_MU_EXP/PPM_CONVENTION:.6f}
        ppm)")
    print(f"  Tau shell      : {res['a_tau_shell_ppm']:.6f} ppm")
    print(f"  Tau full a_tau : {res['a_tau_ppm']:.6f} ppm  (SM prediction: {A_TAU_EXP/
        PPM_CONVENTION:.6f} ppm)")

def amm_rigidity_test(
    base_alpha_geom: float,
    base_eps_M: float,
    L_mu_dim: int = 5,
    L_tau_dim: int = 34,
) -> None:
    """
    Test the rigidity of the lepton AMM sector against changes in eps_M.
    """
    print("\n" + "=" * 78)
    print("AMM RIGIDITY TEST (vary eps_M)")
```

```
    print("=" * 78)
    print(f"{'factor':>8} {'eps_M':>18} {'alpha_inv':>14} {'a_e [ppm]':>14} {'a_mu [ppm]':>14} {'a_tau [ppm]':>14}")
    print("-" * 78)

    for factor in [0.998, 0.999, 1.0, 1.001, 1.002]:
        eps_test = base_eps_M * factor
        alpha_inv_test = compute_alpha_geometric(eps_test)
        alpha_test = 1.0 / alpha_inv_test

        res = compute_lepton_amms(
            alpha_geom=alpha_test,
            eps_M=eps_test,
            L_mu_dim=L_mu_dim,
            L_tau_dim=L_tau_dim,
        )

        print(
            f"{factor:8.3f} {eps_test:18.6e} {alpha_inv_test:14.9f} "
            f"{res['a_e_ppm']:14.6f} {res['a_mu_ppm']:14.6f} {res['a_tau_ppm']:14.6f}"
        )

    print("-" * 78)
    print("Reference experimental:")
    print(f"  a_e   : {A_E_CODATA/PPM_CONVENTION:.6f} ppm")
    print(f"  a_mu  : {A_MU_EXP/PPM_CONVENTION:.6f} ppm")
    print(f"  a_tau : {A_TAU_EXP/PPM_CONVENTION:.6f} ppm")
    print("=" * 78)

# ============================================================================
# SECTION 9: ATOMIC SCALES FROM PURE GEOMETRY
# ============================================================================

def compute_atomic_scales(alpha_geom: float, r_e_geom: float) -> dict:
    """
    Compute atomic scales from geometric alpha and geometric electron radius.
    """
    R_inf = (alpha_geom**3) / (4.0 * PI * r_e_geom)
    a0 = r_e_geom / (alpha_geom**2)
    lambda_C = (2.0 * PI * r_e_geom) / alpha_geom

    return {
        "R_inf": R_inf,
        "a0": a0,
        "lambda_C": lambda_C,
    }


def report_atomic_scales(res: dict) -> None:
    """Print atomic scale results and compare with CODATA 2022."""
    print("--- Atomic Scales from Pure Geometry ---")
    print(f"  Rydberg R_inf    : {res['R_inf']:.8f} m^-1")
    print(f"  CODATA           : {R_INF_CODATA:.8f} m^-1")
    print(f"  rel_error        : {abs(res['R_inf'] - R_INF_CODATA)/R_INF_CODATA*100:.6f} %")

    print(f"  Bohr radius a0   : {res['a0']:.15e} m")
    print(f"  CODATA           : {A0_CODATA:.15e} m")
    print(f"  rel_error        : {abs(res['a0'] - A0_CODATA)/A0_CODATA*100:.6f} %")

    print(f"  Compton lambda_C : {res['lambda_C']:.15e} m")
    print(f"  CODATA           : {LAMBDA_C_CODATA:.15e} m")
    print(f"  rel_error        : {abs(res['lambda_C'] - LAMBDA_C_CODATA)/LAMBDA_C_CODATA*100:.6f} %")

def main():
    print("ENHANCED EWT - EMERGENCE ENGINE (PROTOTYPE)")

    A_pi = compute_alpha_core()
    print(f"\nAlpha core A_pi = 4*pi^3 + pi^2 + pi = {A_pi:.15f}")

    bcc_derived = derive_eps_M_from_BCC(8.0 * PI**4)
    eps_M_bcc = bcc_derived["eps_M"]
```

```
    print("\nDerived eps_M from BCC packing:")
    print(f"  eta       = {bcc_derived['eta']:.10f}")
    print(f"  N_ideal   = {bcc_derived['N_ideal']:.10f}")
    print(f"  zeta      = {bcc_derived['zeta']:.10f}")
    print(f"  N_geom    = {bcc_derived['N_geom']:.10f}")
    print(f"  eps_M     = {bcc_derived['eps_M']:.15e}")

    alpha_inv_bcc = compute_alpha_geometric(eps_M_bcc)
    alpha_geom = 1.0 / alpha_inv_bcc

    res_alpha_bcc = test_alpha_sector(alpha_inv_bcc)

    print("\nAlpha inverse fine-structure constant test:")
    print(f"  Pure BCC geometry  : pred = {res_alpha_bcc['alpha_inv_pred']:.12f} | rel_err = {res_alpha_bcc['rel_err']:.6f} %")
    print(f"  CODATA 2022        : {ALPHA_INV_CODATA:.12f}")

    hbar_geom = derive_hbar(alpha_geom, R_E, M_E, C0)
    q_P_geom = derive_planck_charge_from_e(alpha_geom, E_CHARGE_CODATA)

    print(f"Derived q_P          : {q_P_geom:.15e}")

    ratio_qP_e_derived = q_P_geom / E_CHARGE_CODATA
    ratio_qP_e_expected = 1.0 / math.sqrt(alpha_geom)

    print(f"\nGeometric ratio q_P / e:")
    print(f"  derived   q_P / e = {ratio_qP_e_derived:.10f}")
    print(f"  expected  1/sqrt(alpha_geom) = {ratio_qP_e_expected:.10f}")
    print(f"  difference                   = {abs(ratio_qP_e_derived - ratio_qP_e_expected):.6e}")

    lambda_unc = derive_lambda_uncorr(q_P_geom)

    print(f"\nDerived hbar         : {hbar_geom:.15e}")
    print(f"Using q_P input      : {q_P_geom:.15e}")
    print(f"lambda_uncorr        : {lambda_unc:.15e}")

    nu_result = derive_neutrino_radius(alpha_geom, q_P_geom)
    print(f"\nNeutrino radius r_nu : {nu_result['r_nu']:.15e} m")
    print(f"g_v fixed point      : {nu_result['gv_pred']:.15f}")

    res_nu = test_decadic_resonance(R_E, nu_result['r_nu'])
    print(f"\nElectron/neutrino ratio : {res_nu['ratio']:.12f}")
    print(f"Expected                : {res_nu['expected']}")
    print(f"Relative error          : {res_nu['rel_err']:.6f} %")
    print(f"K_implied (r^5)         : {res_nu['K_implied']:.12e}")

    lambda_l_geom = derive_lambda_l_geometric(
        alpha_geom=alpha_geom,
        r_e=R_E,
        r_nu=nu_result["r_nu"],
        N_geom=bcc_derived["N_geom"],
        L_p_geom=BCC_IDEAL_PROJECTION_LP,
        K_WC=10,
    )

    print(f"\nDerived lambda_l (geometry): {lambda_l_geom:.15e} m")
    print(f"CODATA lambda_l            : {LAMBDA_L:.15e} m")
    print(f"Relative error             : {abs(lambda_l_geom - LAMBDA_L)/LAMBDA_L*100:.6f} %")

    print("\n--- Gravity Sector (Geometric L_p, derived lambda_l) ---")
    gravity_res = gravity_sector(
        alpha_geom=alpha_geom,
        r_nu=nu_result["r_nu"],
        N_geom=bcc_derived["N_geom"],
        L_p_geom=BCC_IDEAL_PROJECTION_LP,
        K_WC=10,
        lambda_l=lambda_l_geom,
        r_e=R_E,
        m_e=M_E,
        c0=C0,
    )
    report_gravity(gravity_res)
```

```
    input_rigidity_test(alpha_geom, nu_result["r_nu"], bcc_derived["N_geom"],
        BCC_IDEAL_PROJECTION_LP)

    rigidity_test(
        alpha_geom=alpha_geom,
        r_e=R_E,
        r_nu=nu_result["r_nu"],
        N_geom=bcc_derived["N_geom"],
        L_p_geom=BCC_IDEAL_PROJECTION_LP,
        K_WC=10,
        lambda_l_geom=lambda_l_geom,
        m_e=M_E,
        c0=C0,
    )

    ladder_bcc = build_geometric_ladder(eps_M_bcc, "BCC packing derived", zeta=bcc_derived["
        zeta"])

    print(f"\n--- Geometric Ladder (Pure BCC) ---")
    report_ladder(ladder_bcc)
    run_sector_tests(ladder_bcc)

    amm_results = compute_lepton_amms(alpha_geom, eps_M_bcc)
    report_lepton_amms(amm_results)
    amm_rigidity_test(alpha_geom, eps_M_bcc)

    r_e_geom = 100.0 * nu_result["r_nu"]
    atomic_res = compute_atomic_scales(alpha_geom, r_e_geom)
    report_atomic_scales(atomic_res)

    print("\n" + "=" * 78)
    print("END OF TEST")
    print("=" * 78)

if __name__ == "__main__":
    main()
```

Listing 1: Core geometric emergence engine.

A.3.2 Emergence Engine Output

The console output of `ewt_emergence_engine.py` is reproduced below.

```
ENHANCED EWT - EMERGENCE ENGINE (PROTOTYPE)

Alpha core A_pi = 4*pi^3 + pi^2 + pi = 137.036303775878423

Derived eps_M from BCC packing:
  eta       = 0.6801747616
  N_ideal   = 779.2727282720
  zeta      = 0.0006033964
  N_geom    = 778.8025178842
  eps_M     = 4.141169769303932e-05

Alpha inverse fine-structure constant test:
  Pure BCC geometry  : pred = 137.036262364181 | rel_err = 0.000192 %
  CODATA 2022        : 137.035999084000
Derived q_P          : 1.875547839472885e-18

Geometric ratio q_P / e:
  derived  q_P / e = 11.7062488597
  expected 1/sqrt(alpha_geom) = 11.7062488597
  difference               = 0.000000e+00

Derived hbar         : 1.054573843737794e-34
Using q_P input      : 1.875547839472885e-18
lambda_uncorr        : 2.771705640418664e-17

Neutrino radius r_nu : 2.817935436016830e-17 m
g_v fixed point      : 0.983594444710376
```

```
Electron/neutrino ratio : 100.000173537800
Expected                : 100.0
Relative error          : 0.000174 %
K_implied (r^5)         : 1.000008676920e+10

Derived lambda_l (geometry): 1.616646406608278e-35 m
CODATA lambda_l            : 1.616200000000000e-35 m
Relative error             : 0.027621 %

--- Gravity Sector (Geometric L_p, derived lambda_l) ---
  G_Base         : 2.780252259136399e+32
  N_nu_statutory: 3.295904039818979e+52
  C_Unif         : 1.102011616800948
  X_eff          : 5275.7683235656
  N_nu_eff       : 6.247249381851994e+48
  G_EWT          : 6.677519975508460e-11
  G_CODATA       : 6.674305000000000e-11
  rel_error      : 0.048169 %

[4] Vary fundamental inputs r_e and m_e
--------------------------------------------------------------------------------
  factor           r_e [m]          m_e [kg]        lambda_l [m]     G_err %
   0.998      2.812304e-15      9.091165e-31        1.603752e-35    1.146422
   0.999      2.815122e-15      9.100274e-31        1.610190e-35    0.550621
   1.000      2.817940e-15      9.109384e-31        1.616646e-35    0.048169
   1.001      2.820758e-15      9.118493e-31        1.623123e-35    0.649961
   1.002      2.823576e-15      9.127602e-31        1.629618e-35    1.254766

================================================================================
RIGIDITY TEST
================================================================================

[1] Vary N_geom
--------------------------------------------------------------------------------
      N_geom       alpha_inv      lambda_l [m]      G_EWT [m^3/kg/s^2]     G_err %
   774.90851   137.036262156      1.666006e-35   7.091501922019398e-11    6.250792
   777.24491   137.036262281      1.636183e-35   6.839883420425040e-11    2.480834
   778.80252   137.036262364      1.616646e-35   6.677519975508460e-11    0.048169
   780.36012   137.036262447      1.597382e-35   6.519323594110790e-11    2.322061
   782.69653   137.036262570      1.568985e-35   6.289591749060191e-11    5.764095

[2] Vary L_p_geom
--------------------------------------------------------------------------------
       L_p       alpha_inv      lambda_l [m]      G_EWT [m^3/kg/s^2]     G_err %
  1.148927   137.036262364      1.616632e-35   6.677397472932486e-11    0.046334
  1.152391   137.036262364      1.616640e-35   6.677471121371631e-11    0.047437
  1.154701   137.036262364      1.616646e-35   6.677519975508460e-11    0.048169
  1.157010   137.036262364      1.616652e-35   6.677568635151728e-11    0.048899
  1.160474   137.036262364      1.616661e-35   6.677641262474325e-11    0.049987

[3] Vary lambda_l directly
--------------------------------------------------------------------------------
    lambda_l [m]      G_EWT [m^3/kg/s^2]     G_err %
    1.600480e-35   6.577608001792828e-11    1.448795
    1.616646e-35   6.677519975508460e-11    0.048169
    1.632813e-35   6.777932766352434e-11    1.552638
================================================================================

--- Geometric Ladder (Pure BCC) ---
  Source        : BCC packing derived
  eps_M         : 4.141169769303932e-05
  zeta          : 6.0339643713e-04
  C_local       : 1.464124612959771e-05
  C_gap         : 1.014075935809472  (pure pi^6, Z-Z and Z-H)
  C_gap_mixed   : 1.010677258781862  (mixed W-Z)
  O_mixed       : 729.258882  (harmonic mean pi^6*pi^7/(pi^6+pi^7))
  eps_flavor    : 0.000408717073808
  Ladder:
    pi^3 = 31.006276680299816
    pi^4 = 97.409091034002415
    pi^5 = 306.019684785281413
    pi^6 = 961.389193575304262
    pi^7 = 3020.293227776791355
```

```
  M_W prediction (mixed)    : 80.379087 GeV
  rel. err PDG 2024 (mixed) : 0.0123 %
  rel. err ATLAS 2024       : 0.0238 %
  rel. err CMS 2024         : 0.0235 %
  rel. err CDF II 2022      : 0.0676 %  (outlier)
  M_W (pure pi^6)           : 80.514122 GeV
  rel. err PDG 2024 (pure)  : 0.1803 %
  sin theta_C               : 0.2242955157
  rel. err PDG              : 0.001999 %
  eps_flavor                : 4.087171e-04
  zeta                      : 6.033964e-04
--- Lepton Anomalous Magnetic Moments ---
  Electron a_e   : 1159.916228 ppm  (CODATA: 1159.652182 ppm)
  Muon shell     : 248.571133 ppm
  Muon full a_mu : 1166.212608 ppm  (Exp: 1165.920610 ppm)
  Tau shell      : 1176.838130 ppm
  Tau full a_tau : 1176.838130 ppm  (SM prediction: 1177.210000 ppm)

================================================================================
AMM RIGIDITY TEST (vary eps_M)
================================================================================
  factor              eps_M      alpha_inv      a_e [ppm]     a_mu [ppm]    a_tau [ppm]
--------------------------------------------------------------------------------
   0.998       4.132887e-05  137.036262447    1159.919210    1166.215909    1177.356781
   0.999       4.137029e-05  137.036262406    1159.917719    1166.214259    1177.097420
   1.000       4.141170e-05  137.036262364    1159.916228    1166.212608    1176.838130
   1.001       4.145311e-05  137.036262323    1159.914737    1166.210958    1176.578912
   1.002       4.149452e-05  137.036262281    1159.913246    1166.209308    1176.319764
--------------------------------------------------------------------------------
Reference experimental:
  a_e   : 1159.652182 ppm
  a_mu  : 1165.920610 ppm
  a_tau : 1177.210000 ppm
================================================================================
--- Atomic Scales from Pure Geometry ---
  Rydberg R_inf   : 10973687.36203397 m^-1
    CODATA        : 10973731.56815700 m^-1
    rel_error     : 0.000403 %
  Bohr radius a0  : 5.291783259434870e-11 m
    CODATA        : 5.291772109030000e-11 m
    rel_error     : 0.000211 %
  Compton lambda_C: 2.426310689655160e-12 m
    CODATA        : 2.426310238670000e-12 m
    rel_error     : 0.000019 %

================================================================================
END OF TEST
================================================================================
```

Listing 2: Console output from `ewt_emergence_engine.py`.

A.3.3 Extended Lepton AMM

`ewt_amm_extended.py` computes the full anomalous magnetic moments of the electron, muon, and tau using the geometric fine-structure constant and the magnetic deficit. It also reports the internal shell references and the corresponding consistency errors.

```python
#!/usr/bin/env python3
"""
From manuscript version: 5.2.4 or leter

Enhanced EWT -- Extended Lepton AMM Module
==========================================
Provides the extended lepton anomalous magnetic moments with internal
shell references and consistency tests between the geometric AMM
sector and the orbital mass sector.

This module complements ewt_emergence_engine.py, which contains the
basic full AMM predictions and rigidity tests.
"""
```

```python

import math

try:
    from ewt_emergence_engine import (
        PI, SQRT2, SQRT3, EULER,
        A_MU_EXP,
        A_TAU_EXP,
        A_E_CODATA,
        compute_alpha_core,
        derive_eps_M_from_BCC,
        get_AMMi_K,
    )
except ImportError:
    raise ImportError("This module requires ewt_emergence_engine.py in the same directory.")


def compute_lepton_amms_extended(alpha_geom: float, eps_M: float) -> dict:
    """
    Compute extended lepton AMMs with internal shell references.

    Parameters
    ----------
    alpha_geom : float
        Geometric fine-structure constant.
    eps_M : float
        Magnetic deficit from BCC geometry.

    Returns
    -------
    dict with:
        a_e_ppm
        a_mu_shell_ppm
        a_mu_full_ppm
        a_tau_shell_total_ppm
        a_tau_full_ppm
        ref_a_mu_shell_ppm
        ref_a_tau_shell_ppm
        err_a_mu_shell_pct
        err_a_tau_shell_pct
    """
    A_pi = compute_alpha_core()
    L_mu_dim = 5
    L_tau_dim = 34  # Reserved for the 4th-generation extension; see note in
                    # ewt_emergence_engine.compute_lepton_amms.
    K_e = 10

    # Electron
    a_e_ppm = (alpha_geom / (2.0 * PI)) * (1.0 - eps_M * (PI**3)) * 1e6

    # Muon shell
    K_mu_total = get_AMMi_K(2)
    K_mu_delta = K_mu_total - K_e
    M_mu_shell = K_mu_delta / K_e

    B_mu_scale = (3.0 * A_pi * PI**3) / (2.0 * L_mu_dim**2)
    a_mu_shell_ppm = B_mu_scale * (1.0 - eps_M) ** (M_mu_shell * PI**3)

    # Projection operator for muon (2D -> 1D)
    O_mu = 1.0 / (4.0 * PI**2)

    a_mu_full_ppm = a_e_ppm + a_mu_shell_ppm * O_mu

    # Tau shell
    K_tau_total = get_AMMi_K(3)
    M_tau_rel = K_tau_total / K_e

    B_tau_base = ((3.0 * A_pi * PI**3) / (8.0 * SQRT2)) + (A_pi / 2.0)
    a_tau_shell_raw_ppm = B_tau_base * (1.0 - eps_M) ** (M_tau_rel * PI**3)

    # Recursive accumulation with interface tension L_mu^2
    a_tau_shell_total_ppm = a_mu_shell_ppm + a_tau_shell_raw_ppm + L_mu_dim**2

```

```
    # Projection operator for tau (3D -> 3D, unity).
    # With O_tau = 1, this expression reduces algebraically to
    # a_tau_shell_total_ppm. It is written explicitly to show that no
    # dimensional projection is applied for the tau, in contrast to the
    # muon's O_mu = 1/(4*pi^2).
    a_tau_full_ppm = a_e_ppm + (a_tau_shell_total_ppm - a_e_ppm)

    # Internal references (from orbital mass relations)
    ref_a_mu_shell_ppm = 248.8
    ref_a_tau_shell_ppm = 1177.21

    err_a_mu_shell_pct = abs(a_mu_shell_ppm - ref_a_mu_shell_ppm) / ref_a_mu_shell_ppm *
        100.0
    err_a_tau_shell_pct = abs(a_tau_shell_total_ppm - ref_a_tau_shell_ppm) /
        ref_a_tau_shell_ppm * 100.0

    return {
        "a_e_ppm": a_e_ppm,
        "a_mu_shell_ppm": a_mu_shell_ppm,
        "a_mu_full_ppm": a_mu_full_ppm,
        "a_tau_shell_total_ppm": a_tau_shell_total_ppm,
        "a_tau_full_ppm": a_tau_full_ppm,
        "ref_a_mu_shell_ppm": ref_a_mu_shell_ppm,
        "ref_a_tau_shell_ppm": ref_a_tau_shell_ppm,
        "err_a_mu_shell_pct": err_a_mu_shell_pct,
        "err_a_tau_shell_pct": err_a_tau_shell_pct,
    }


def report_extended_amms(res: dict) -> None:
    """Print the extended AMM report in Scilab style."""
    print("\n---␣EXTENDED␣LEPTON␣ANOMALOUS␣MAGNETIC␣MOMENTS␣---")
    print(f"␣␣Electron␣full␣a_e␣␣␣␣␣␣␣:␣{res['a_e_ppm']:.6f}␣ppm")
    print(f"␣␣␣␣CODATA␣reference␣␣␣␣␣␣:␣{A_E_CODATA*1e6:.6f}␣ppm")
    print(f"␣␣␣␣Full␣rel.␣error␣␣␣␣␣␣:␣{abs(res['a_e_ppm']␣-␣A_E_CODATA*1e6)/(A_E_CODATA*1e6
        )*100:.6f}␣%")
    print()
    print(f"␣␣Muon␣shell␣(internal)␣␣␣:␣{res['a_mu_shell_ppm']:.6f}␣ppm")
    print(f"␣␣␣␣EWT␣shell␣reference␣␣␣:␣{res['ref_a_mu_shell_ppm']:.6f}␣ppm")
    print(f"␣␣␣␣Internal␣rel.␣error␣␣␣:␣{res['err_a_mu_shell_pct']:.6f}␣%")
    print(f"␣␣Muon␣full␣a_mu␣␣␣␣␣␣␣␣␣:␣{res['a_mu_full_ppm']:.6f}␣ppm")
    print(f"␣␣␣␣Experimental␣target␣␣␣:␣{A_MU_EXP*1e6:.6f}␣ppm")
    print(f"␣␣␣␣Full␣rel.␣error␣␣␣␣␣␣:␣{abs(res['a_mu_full_ppm']␣-␣A_MU_EXP*1e6)/(A_MU_EXP*1
        e6)*100:.6f}␣%")
    print()
    print(f"␣␣Tau␣shell␣total␣␣␣␣␣␣␣␣:␣{res['a_tau_shell_total_ppm']:.6f}␣ppm")
    print(f"␣␣␣␣EWT␣shell␣reference␣␣␣:␣{res['ref_a_tau_shell_ppm']:.6f}␣ppm")
    print(f"␣␣␣␣Internal␣rel.␣error␣␣␣:␣{res['err_a_tau_shell_pct']:.6f}␣%")
    print(f"␣␣Tau␣full␣a_tau␣␣␣␣␣␣␣␣␣:␣{res['a_tau_full_ppm']:.6f}␣ppm")
    print(f"␣␣␣␣SM␣prediction␣␣␣␣␣␣␣␣:␣{A_TAU_EXP*1e6:.6f}␣ppm␣␣(no␣direct␣measurement)")
    print(f"␣␣␣␣Full␣rel.␣error␣␣␣␣␣␣:␣{abs(res['a_tau_full_ppm']␣-␣A_TAU_EXP*1e6)/(
        A_TAU_EXP*1e6)*100:.6f}␣%")


def main():
    print("EWT␣EXTENDED␣LEPTON␣AMM␣MODULE")

    # Use pure BCC geometry from the main engine
    bcc = derive_eps_M_from_BCC(8.0 * PI**4)
    eps_M = bcc["eps_M"]
    alpha_geom = 1.0 / (compute_alpha_core() - eps_M)

    res = compute_lepton_amms_extended(alpha_geom, eps_M)
    report_extended_amms(res)


if __name__ == "__main__":
    main()
```

Listing 3: Extended lepton anomalous magnetic moment computation.

A.3.4 Extended Lepton AMM Output

The console output of `ewt_amm_extended.py` is reproduced below.

```
EWT EXTENDED LEPTON AMM MODULE

--- EXTENDED LEPTON ANOMALOUS MAGNETIC MOMENTS ---
  Electron full a_e        : 1159.916228 ppm
    CODATA reference       : 1159.652182 ppm
    Full rel. error        : 0.022769 %

  Muon shell (internal)    : 248.571133 ppm
    EWT shell reference    : 248.800000 ppm
    Internal rel. error    : 0.091988 %
  Muon full a_mu           : 1166.212608 ppm
    Experimental target    : 1165.920610 ppm
    Full rel. error        : 0.025044 %

  Tau shell total          : 1176.838130 ppm
    EWT shell reference    : 1177.210000 ppm
    Internal rel. error    : 0.031589 %
  Tau full a_tau           : 1176.838130 ppm
    SM prediction          : 1177.210000 ppm  (no direct measurement)
    Full rel. error        : 0.031589 %
```

Listing 4: Console output from `ewt_amm_extended.py`.

A.3.5 AMM Resonance Scanner

`ewt_amm_resonance_scanner.py` scans the (K_μ, K_τ) nodal-count space and identifies the resonance minima. It also generates the PDF figures used in the sensitivity analysis.

```
#!/usr/bin/env python3
"""
From manuscript version: 5.2.4 or leter

Enhanced EWT -- AMM Resonance Scanner (Onion Model)
====================================================
Python port of the Scilab AMM_find.sc script, adapted to the
zero-calibration geometric eps_M from BCC packing impedance.

Scans the (K_muon, K_tau) space and identifies resonance points
with minimal internal error against the EWT shell references.

Generates:
  Fig1_Muon_Profile.pdf
  Fig2_Tau_Profile.pdf
  Fig3_3D_Peak.pdf
  Fig4_Topography.pdf
"""

import math
import os
import numpy as np
import matplotlib
matplotlib.use("pdf")  # non-interactive backend
import matplotlib.pyplot as plt

try:
    from ewt_emergence_engine import (
        PI, SQRT2, SQRT3, EULER,
        compute_alpha_core,
        compute_alpha_geometric,
        derive_eps_M_from_BCC,
    )
except ImportError:
    raise ImportError("This module requires ewt_emergence_engine.py in the same directory.")


# ============================================================================
```

```
# 1. Core EWT calculation using geometric eps_M
# ============================================================================

def calculate_EWT_from_geometry(Kn_m_in: int, Kn_t_in: int):
    """
    Compute the predicted AMMs and their internal errors for given
    nodal counts.

    Returns
    -------
    tuple (am_ppm, at_ppm, e_m, e_t)
    """

    # Use pure BCC geometry
    bcc = derive_eps_M_from_BCC(8.0 * PI**4)
    eps_M = bcc["eps_M"]

    A_pi = compute_alpha_core()
    alpha_geom = 1.0 / compute_alpha_geometric(eps_M)

    Kn_e = 10

    # Experimental / internal shell references
    target_amu = 248.8 / 1e6
    target_atau = 1177.21 / 1e6

    # Invariants from Onion Model
    L_mu = 5
    L_tau = 34

    # Shell 1: Muon Resonance
    M_mu_shell = (Kn_m_in - Kn_e) / Kn_e
    B_mu_scale = (3.0 * A_pi * PI**3) / (2.0 * L_mu**2)
    amu_shell = B_mu_scale * (1.0 - eps_M) ** (M_mu_shell * PI**3)

    # The shell comparison uses the pure muon-shell contribution (in ppm).
    # The electron AMM (ae_pred = (alpha/2pi)(1 - eps_M*pi^3), dimensionless
    # ~0.00116) is NOT added here: it is negligible at the ppm scale
    # (~5e-6 relative) and is already accounted for in the full muon AMM of
    # ewt_amm_extended.py. Including it would contaminate the shell reference
    # and break consistency with Table final_results_confirmed (a_mu_shell_ppm).
    am_ppm = amu_shell

    e_m = abs(am_ppm / 1e6 - target_amu) / target_amu * 100.0

    # Shell 2: Tau Resonance (Recursive addition)
    M_tau_rel = Kn_t_in / Kn_e
    B_tau_base = ((3.0 * A_pi * PI**3) / (8.0 * SQRT2)) + (A_pi / 2.0)
    at_shell = B_tau_base * (1.0 - eps_M) ** (M_tau_rel * PI**3)

    at_ppm = am_ppm + at_shell + L_mu**2  # interface tension L_mu^2 = 25

    e_t = abs(at_ppm / 1e6 - target_atau) / target_atau * 100.0

    return am_ppm, at_ppm, e_m, e_t


# ============================================================================
# 2. Scan and output
# ============================================================================

def main():
    print("=== EWT RECURSIVE HIERARCHY ANALYSIS ===")
    print("Reference: Onion Model")
    print()

    # Scan ranges
    Km_scan = list(range(195, 216))      # 195..215
    Kt_scan = list(range(2170, 2191))    # 2170..2190

    # Meshgrid equivalent using nested lists
    Z_err = [[0.0 for _ in Km_scan] for _ in Kt_scan]
    am_arr = [[0.0 for _ in Km_scan] for _ in Kt_scan]
```

```
    at_arr = [[0.0 for _ in Km_scan] for _ in Kt_scan]

    special_points = [(207, 2181), (200, 2181), (207, 2180), (200, 2180)]

    for i, Kt in enumerate(Kt_scan):
        for j, Km in enumerate(Km_scan):
            am, at, em, et = calculate_EWT_from_geometry(Km, Kt)
            Z_err[i][j] = (em + et) / 2.0
            am_arr[i][j] = am
            at_arr[i][j] = at

            if (Km, Kt) in special_points:
                print(
                    f"POINT: Km={Km}, Kt={Kt} | "
                    f"Err_mu: {em:.6f}% | Err_tau: {et:.6f}% | "
                    f"Mean: {(em+et)/2:.6f}% | amu: {am:.4f} | atau: {at:.4f}"
                )

    # Z_plot = 1 / (Z_err + 0.0001) transposed for plotting
    Z_plot = [[1.0 / (Z_err[i][j] + 0.0001) for i in range(len(Kt_scan))]
              for j in range(len(Km_scan))]

    # Ensure output directory
    out_dir = os.path.dirname(os.path.abspath(__file__))

    # ------------------------------------------------------------------------
    # FIG 1: Muon Profile
    # ------------------------------------------------------------------------
    m_idx = Kt_scan.index(2181)
    muon_err = [Z_err[m_idx][j] for j in range(len(Km_scan))]

    plt.figure()
    plt.plot(Km_scan, muon_err, "r-o")
    plt.grid(True)
    plt.xlabel("Nodes Km")
    plt.ylabel("Total Error %")
    plt.title("Muon 2D Resonance Profile")
    plt.savefig(os.path.join(out_dir, "Fig1_Muon_Profile.pdf"))
    plt.close()
    print("EXPORTED: Fig1_Muon_Profile.pdf")

    # ------------------------------------------------------------------------
    # FIG 2: Tau Profile
    # ------------------------------------------------------------------------
    t_idx = Km_scan.index(207)
    tau_err = [Z_err[i][t_idx] for i in range(len(Kt_scan))]

    plt.figure()
    plt.plot(Kt_scan, tau_err, "b-s")
    plt.grid(True)
    plt.xlabel("Nodes Kt")
    plt.ylabel("Total Error %")
    plt.title("Tau 2D Resonance Profile")
    plt.savefig(os.path.join(out_dir, "Fig2_Tau_Profile.pdf"))
    plt.close()
    print("EXPORTED: Fig2_Tau_Profile.pdf")

    # ------------------------------------------------------------------------
    # FIG 3: 3D Stability Surface
    # ------------------------------------------------------------------------
    from mpl_toolkits.mplot3d import Axes3D  # noqa: F401

    fig = plt.figure()
    ax = fig.add_subplot(111, projection="3d")

    # Create meshgrid for plotting
    X, Y = np.meshgrid(Km_scan, Kt_scan)  # X: Km (shape len(Kt) x len(Km)), Y: Kt

    # Z_err has shape (len(Kt), len(Km))
    # We want Z shape also (len(Kt), len(Km))
    Z_for_3d = np.array(Z_err)

    # For 1/error, compute directly from Z_err
```

```
    Z_inv = 1.0 / (Z_for_3d + 0.0001)

    ax.plot_surface(X, Y, Z_inv, cmap="viridis", alpha=0.8)
    ax.set_xlabel("Km")
    ax.set_ylabel("Kt")
    ax.set_zlabel("1 / error")
    ax.set_title("Recursive Stability Surface (Onion Model)")
    ax.view_init(elev=35, azim=45)
    plt.savefig(os.path.join(out_dir, "Fig3_3D_Peak.pdf"))
    plt.close()
    print("EXPORTED: Fig3_3D_Peak.pdf")

    # ------------------------------------------------------------------------
    # FIG 4: Topographic Contour Map (3D version)
    # ------------------------------------------------------------------------
    fig = plt.figure()
    ax = fig.add_subplot(111, projection="3d")

    # Create meshgrid for plotting (same as FIG 3)
    X, Y = np.meshgrid(Km_scan, Kt_scan)

    # Z_inv was computed earlier as 1 / (Z_err + 0.0001)
    Z_for_contour = np.array(Z_inv)  # shape (len(Kt), len(Km))

    # Draw the surface
    ax.plot_surface(
        X, Y, Z_for_contour,
        cmap="viridis",
        alpha=0.6,
        edgecolor="none",
        antialiased=True
    )

    # Draw contour lines on the surface
    ax.contour3D(
        X, Y, Z_for_contour,
        15,
        cmap="Reds",
        linewidths=1.5
    )

    ax.set_xlabel("Km")
    ax.set_ylabel("Kt")
    ax.set_zlabel("1 / error")
    ax.set_title("Topographic Lattice Latches (Km vs Kt)")

    # Adjust viewing angle for better depth perception
    ax.view_init(elev=35, azim=45)

    plt.savefig(os.path.join(out_dir, "Fig4_Topography_3D.pdf"))
    plt.close()
    print("EXPORTED: Fig4_Topography_3D.pdf")

    print("\n--- ALL DATA SYNCHRONIZED WITH LATEX DOCUMENT ---")


if __name__ == "__main__":
    main()
```

Listing 5: AMM resonance scanner and figure generator.

A.3.6 AMM Resonance Scanner Output

The console output of `ewt_amm_resonance_scanner.py` is reproduced below.

```
=== EWT RECURSIVE HIERARCHY ANALYSIS ===
Reference: Onion Model

POINT: Km=200, Kt=2180 | Err_mu: 0.002147% | Err_tau: 0.002748% | Mean: 0.002448% | amu:
    248.7947 | atau: 1177.1776
```

```
POINT: Km=207, Kt=2180 | Err_mu: 0.091988% | Err_tau: 0.021736% | Mean: 0.056862% | amu:
     248.5711 | atau: 1176.9541
POINT: Km=200, Kt=2181 | Err_mu: 0.002147% | Err_tau: 0.012601% | Mean: 0.007374% | amu:
     248.7947 | atau: 1177.0617
POINT: Km=207, Kt=2181 | Err_mu: 0.091988% | Err_tau: 0.031589% | Mean: 0.061789% | amu:
     248.5711 | atau: 1176.8381
EXPORTED: Fig1_Muon_Profile.pdf
EXPORTED: Fig2_Tau_Profile.pdf
EXPORTED: Fig3_3D_Peak.pdf
EXPORTED: Fig4_Topography_3D.pdf

--- ALL DATA SYNCHRONIZED WITH LATEX DOCUMENT ---
```

Listing 6: Console output from `ewt_amm_resonance_scanner.py`.

A.3.7 Electroweak and Boson Module

`ewt_electroweak_bosons.py` computes the mixed W–Z vertex operator (parallel-impedance combination of π^6 and π^7, Eq. 23), the resulting W-boson mass prediction, the pure-π^6 neutral attractor for comparison, the Higgs-vector mixing angles (ZH via pure π^6, WH via mixed), the Cabibbo angle, heavy boson radii, and the benchmark comparison described in Table 3.

```python
#!/usr/bin/env python3
# -*- coding: utf-8 -*-
"""
From manuscript version: 5.2.4 or leter

Enhanced EWT -- Electroweak and Heavy Boson Module
==================================================
Extends the main emergence engine with:
  - Mixed W-Z vertex operator (harmonic pi^6 - pi^7 combination)
  - Weinberg sector (W mass via mixed operator, default)
  - Pure pi^6 W mass (for comparison / neutral vertices)
  - Higgs-vector boson mixing angles (ZH pure, WH mixed)
  - Cabibbo angle (surface mixing)
  - heavy boson geometric radii (Z, H)
  - comparison with experimental W-mass benchmarks (PDG 2024, ATLAS, CMS)

This module supersedes the standalone ewt_mixed_interaction_operator.py,
which is now integrated here.

All functions that return a computed quantity return a dict
(matching the convention of ewt_emergence_engine.py).

This module should be used together with ewt_emergence_engine.py
and ewt_particle_masses.py.
"""

import math

# --- Import shared constants and functions ---
try:
    from ewt_emergence_engine import (
        PI, SQRT2, SQRT3, EULER,
        M_Z_CODATA, M_H_CODATA,
        BCC_PACKING_FRACTION,
        compute_alpha_core,
        derive_eps_M_from_BCC,
        build_geometric_ladder,
        lattice_impedance,
    )
    from ewt_particle_masses import mass_spherical
except ImportError:
    raise ImportError(
        "This module requires ewt_emergence_engine.py and ewt_particle_masses.py."
    )


# ============================================================
```

```
# Physical constants (CODATA / PDG) and experimental benchmarks
# ============================================================

SIN2_THETA_W_MSBAR = 0.23122   # MS-bar at M_Z (CODATA)
# Flavor deficit is computed at runtime from the lattice impedance zeta:
# eps_flavor = 1 / (8 * pi^5 * (1 - zeta))
# analogous to eps_M = 1 / (8 * pi^7 * (1 - zeta)) for charged weak.
# The ideal value (zeta = 0) is 1/(8*pi^5) ~ 4.085e-4; with zeta ~ 6e-4,
# the correction shifts eps_flavor by ~0.06%.

# Standard Model reference values
M_W_SM_TREE = M_Z_CODATA * math.sqrt(1.0 - SIN2_THETA_W_MSBAR)
M_W_SM_FULL = 80.360           # radiative-corrected SM reference

# Experimental W-mass benchmarks: (mass [GeV], 1-sigma uncertainty [GeV])
EXP_BENCHMARKS = {
    "CDF␣II␣2022":      (80.4335, 0.0094),
    "PDG␣2022␣average": (80.377,  0.012),
    "ATLAS␣2024":       (80.360,  0.016),
    "CMS␣2024":         (80.3602, 0.0099),
    "PDG␣2024␣average": (80.3692, 0.0133),
}
REFERENCE_BENCHMARK = "PDG␣2024␣average"


# ============================================================
# Mixed interaction operator (harmonic mean of pi^6 and pi^7)
# ============================================================

def mixed_operator(pi6: float, pi7: float) -> float:
    """
    Harmonic combination of the neutral (pi^6) and charged (pi^7) rungs.

    O_mixed = pi^6 * pi^7 / (pi^6 + pi^7)

    Limiting behavior:
      - pi^7 -> infinity : O_mixed -> pi^6  (neutral channel dominant)
      - pi^6 -> infinity : O_mixed -> pi^7  (charged channel dominant)

    This is the unique simple parameter-free combination that reduces
    to the correct single-rung limit in either decoupling case.
    Arithmetic and geometric means both diverge in these limits.
    """
    return (pi6 * pi7) / (pi6 + pi7)


def c_gap_mixed(c_local: float, pi6: float, pi7: float) -> dict:
    """
    Return the mixed C_gap operator and the underlying O_mixed value.

    Returns
    -------
    dict with keys:
        C_gap   : 1 + O_mixed * C_local
        O_mixed : harmonic mean of pi^6 and pi^7
    """
    o_mixed = mixed_operator(pi6, pi7)
    return {"C_gap": 1.0 + o_mixed * c_local, "O_mixed": o_mixed}


# ============================================================
# Weinberg sector (W mass)
# ============================================================

def weinberg_sector(C_gap: float, M_Z: float, sin2_W: float) -> dict:
    """
    Compute W boson mass for a given C_gap operator.

        M_W = M_Z * sqrt( (1 - sin^2(theta_W)) * C_gap )

    Parameters
    ----------
    C_gap : float
```

```
        Either the pure pi^6 operator or the mixed operator; the caller
        chooses which one based on the physics (neutral vertex vs mixed
        W-Z vertex).
    M_Z : float
        Z-boson mass [GeV].
    sin2_W : float
        MS-bar Weinberg angle.

    Returns
    -------
    dict with keys:
        M_W_pred : predicted W mass [GeV]
        C_gap    : the operator value used
        sin2_W   : the Weinberg angle input
    """
    M_W_pred = M_Z * math.sqrt((1.0 - sin2_W) * C_gap)
    return {"M_W_pred": M_W_pred, "C_gap": C_gap, "sin2_W": sin2_W}


def weinberg_sector_mixed(
    c_local: float, pi6: float, pi7: float,
    M_Z: float, sin2_W: float,
) -> dict:
    """
    Convenience wrapper: compute W mass via the mixed W-Z operator.

    Returns
    -------
    dict with keys:
        M_W_pred : predicted W mass [GeV]
        C_gap    : the mixed operator value used
        O_mixed  : the harmonic mean of pi^6 and pi^7
        sin2_W   : the Weinberg angle input
    """
    cg = c_gap_mixed(c_local, pi6, pi7)
    result = weinberg_sector(cg["C_gap"], M_Z, sin2_W)
    result["O_mixed"] = cg["O_mixed"]
    return result


# ============================================================
# Higgs-vector boson mixing angles
# ============================================================

def compute_higgs_mixing_angles(
    C_gap_ZH: float,
    C_gap_WH: float,
    M_Z_EWT: float,
    M_H_EWT: float,
    M_W_EWT: float,
) -> dict:
    """
    Compute Higgs-vector mixing angles.

    Z-H uses the pure pi^6 operator (both solitons neutral);
    W-H uses the mixed operator (one charged, one neutral soliton).

    Returns
    -------
    dict with keys:
        sin2_ZH : sin^2(theta_ZH) using C_gap_ZH (pure pi^6)
        sin2_WH : sin^2(theta_WH) using C_gap_WH (mixed)
    """
    sin2_ZH = 1.0 - ((M_Z_EWT / M_H_EWT) ** 2) * (1.0 / C_gap_ZH)
    sin2_WH = 1.0 - ((M_W_EWT / M_H_EWT) ** 2) * (1.0 / C_gap_WH)
    return {"sin2_ZH": sin2_ZH, "sin2_WH": sin2_WH}


# ============================================================
# Cabibbo sector (surface mixing)
# ============================================================

def cabibbo_sector(
```

```
    M_d_EWT: float,
    M_s_EWT: float,
    M_d_PDG: float,
    M_s_PDG: float,
    eps_flavor: float = None,
    target: float = 0.22430,
) -> dict:
    """
    Compute the Cabibbo angle with the flavor deficit correction.

    Formula:
        sin(theta_C) = sqrt(M_d/M_s) * (1 + eps_flavor)

    where eps_flavor = 1/(8*pi^5*(1-zeta)) is the flavor deficit at
    the pi^5 rung, analogous to eps_M = 1/(8*pi^7*(1-zeta)) at the
    charged weak rung. If eps_flavor is None, the ideal limit
    (zeta = 0) value 1/(8*pi^5) is used.

    Parameters
    ----------
    M_d_EWT, M_s_EWT : float
        Spherical-mode quark masses [GeV] (Variant A).
    M_d_PDG, M_s_PDG : float
        PDG light-quark masses [GeV] (Variant B/C).
    eps_flavor : float, optional
        Flavor deficit. Default None -> ideal 1/(8*pi^5).
    target : float
        Reference sin(theta_C), default PDG 0.22430.

    Returns
    -------
    dict with keys:
        eps_flavor   : the flavor deficit value
        correction   : (1 + eps_flavor)
        sinC_A       : prediction with EWT masses
        sinC_B       : prediction with PDG masses
        err_A_pct    : relative error of Variant A [%]
        err_B_pct    : relative error of Variant B [%]
        target       : reference value
    """
    if eps_flavor is None:
        eps_flavor = 1.0 / (8.0 * math.pi**5)  # ideal limit (zeta = 0)
    correction = 1.0 + eps_flavor
    sinC_A = math.sqrt(M_d_EWT / M_s_EWT) * correction
    sinC_B = math.sqrt(M_d_PDG / M_s_PDG) * correction
    return {
        "eps_flavor": eps_flavor,
        "correction": correction,
        "sinC_A": sinC_A,
        "sinC_B": sinC_B,
        "err_A_pct": abs(sinC_A - target) / target * 100.0,
        "err_B_pct": abs(sinC_B - target) / target * 100.0,
        "target": target,
        "M_d_EWT": M_d_EWT,
        "M_s_EWT": M_s_EWT,
        "M_d_PDG": M_d_PDG,
        "M_s_PDG": M_s_PDG,
    }


# ============================================================
# Heavy boson radii
# ============================================================

def compute_boson_radii(r_e_geom: float) -> dict:
    """
    Predict radii of heavy neutral bosons from spherical mode masses
    using r = r_e * (m / m_e)^(1/5).

    Returns
    -------
    dict with keys:
        r_Z, r_H : predicted radii [m]
```

```
        m_Z, m_H : spherical-mode masses [GeV]
    """
    m_e_GeV = mass_spherical(10)
    m_Z = mass_spherical(110)
    m_H = mass_spherical(117)

    r_Z = r_e_geom * (m_Z / m_e_GeV) ** (1.0 / 5.0)
    r_H = r_e_geom * (m_H / m_e_GeV) ** (1.0 / 5.0)

    return {"r_Z": r_Z, "r_H": r_H, "m_Z": m_Z, "m_H": m_H}


# ============================================================
# Weinberg angle consistency test
# ============================================================

def test_weinberg_angle(
    M_Z: float, M_W: float, C_gap: float,
    label: str, target_sin2_W: float = SIN2_THETA_W_MSBAR,
) -> dict:
    """
    Compute sin^2(theta_W) from given boson masses and C_gap.

        sin^2(theta_W) = 1 - (M_W/M_Z)^2 * (1/C_gap)

    Returns
    -------
    dict with keys:
        sin2_W_pred  : reconstructed Weinberg angle
        target       : reference value
        rel_err_pct  : relative error [%]
        label        : input label
    """
    sin2_W_pred = 1.0 - ((M_W / M_Z) ** 2) * (1.0 / C_gap)
    rel_err_pct = abs(sin2_W_pred - target_sin2_W) / target_sin2_W * 100.0
    return {
        "sin2_W_pred": sin2_W_pred,
        "target": target_sin2_W,
        "rel_err_pct": rel_err_pct,
        "label": label,
    }


# ============================================================
# Benchmark comparison
# ============================================================

def compare_w_mass_benchmarks(
    M_W_EWT: float,
    benchmarks: dict = None,
    reference_name: str = REFERENCE_BENCHMARK,
) -> dict:
    """
    Compare an EWT W-mass prediction with experimental benchmarks.

    Returns
    -------
    dict with keys:
        M_W_EWT    : the EWT prediction [GeV]
        reference  : name of the reference benchmark
        deviations : {benchmark_name: {M_W, sigma, delta_mev, sigma_dev}}
    """
    if benchmarks is None:
        benchmarks = EXP_BENCHMARKS
    ref_M, _ = benchmarks[reference_name]
    deviations = {}
    for name, (M, sigma) in benchmarks.items():
        deviations[name] = {
            "M_W": M,
            "sigma": sigma,
            "delta_mev": (M_W_EWT - M) * 1000.0,
            "delta_vs_ref_mev": (M - ref_M) * 1000.0,
            "sigma_dev": (M_W_EWT - M) / sigma,
```

```
        }
    return {
        "M_W_EWT": M_W_EWT,
        "reference": reference_name,
        "deviations": deviations,
    }


# ============================================================
# Sector runners (encapsulate geometry + computation + print)
# ============================================================

def run_geometry_summary(geom: dict) -> None:
    print("--- GEOMETRY ---")
    print(f"  C_local              = {geom['C_local']:.6e}")
    print(f"  pi^6                 = {geom['pi6']:.6f}")
    print(f"  pi^7                 = {geom['pi7']:.6f}")
    print(f"  C_gap (pure pi^6)    = {geom['C_gap_pure']:.10f}")
    print(f"  C_gap (mixed)        = {geom['C_gap_mixed']:.10f}")
    print(f"  O_mixed              = {geom['O_mixed']:.4f}")
    print()


def run_mixed_w_sector(geom: dict) -> dict:
    result = weinberg_sector_mixed(
        geom["C_local"], geom["pi6"], geom["pi7"],
        M_Z_CODATA, SIN2_THETA_W_MSBAR,
    )
    print("--- MIXED W-Z VERTEX (main prediction) ---")
    print(f"  O_mixed        = {result['O_mixed']:.4f}")
    print(f"  C_gap          = {result['C_gap']:.10f}")
    print(f"  M_W_EWT        = {result['M_W_pred']:.4f} GeV")
    print()
    return result


def run_pure_w_sector(geom: dict) -> dict:
    result = weinberg_sector(geom["C_gap_pure"], M_Z_CODATA, SIN2_THETA_W_MSBAR)
    print("--- PURE pi^6 W MASS (for comparison) ---")
    print(f"  C_gap (pure pi^6)  = {result['C_gap']:.10f}")
    print(f"  M_W (pure pi^6)    = {result['M_W_pred']:.4f} GeV")
    print()
    return result


def run_benchmark_comparison(M_W_EWT: float) -> dict:
    result = compare_w_mass_benchmarks(M_W_EWT)
    print("--- BENCHMARK COMPARISON (EWT mixed vs experiments) ---")
    print(f"  Reference: {result['reference']}")
    print(f"  {'Benchmark':<20} {'M_W [GeV]':<22} {'EWT - bench [MeV]':>16} {'sigma':>8}")
    print("  " + "-" * 66)
    for name, d in result["deviations"].items():
        print(f"  {name:<20} {d['M_W']:.4f} +/- {d['sigma']:.4f}    "
              f"{d['delta_mev']:>+10.1f} {d['sigma_dev']:>+7.1f} sigma")
    print()
    return result


def run_higgs_sector(geom: dict, M_W_EWT: float) -> dict:
    M_Z_EWT = mass_spherical(110)
    M_H_EWT = mass_spherical(117)
    mixing = compute_higgs_mixing_angles(
        C_gap_ZH=geom["C_gap_pure"],
        C_gap_WH=geom["C_gap_mixed"],
        M_Z_EWT=M_Z_EWT,
        M_H_EWT=M_H_EWT,
        M_W_EWT=M_W_EWT,
    )
    print("--- HIGGS-VECTOR MIXING ---")
    print(f"  Z (K=110)    = {M_Z_EWT:.10f} GeV  (spherical mode)")
    print(f"  H (K=117)    = {M_H_EWT:.10f} GeV  (spherical mode)")
    print(f"  sin^2(theta_ZH) [pure pi^6] = {mixing['sin2_ZH']:.10f}")
    print(f"  sin^2(theta_WH) [mixed]     = {mixing['sin2_WH']:.10f}")
```

```
    print()
    mixing["M_Z_EWT"] = M_Z_EWT
    mixing["M_H_EWT"] = M_H_EWT
    return mixing


def run_weinberg_tests(geom: dict, M_W_EWT: float, M_Z_EWT: float) -> dict:
    print("--- WEINBERG ANGLE CONSISTENCY TESTS ---")
    results = {}

    r1 = test_weinberg_angle(M_Z_EWT, M_W_EWT, geom["C_gap_mixed"],
                             "EWT spherical masses + mixed")
    print(f"  [{r1['label']}]")
    print(f"    M_Z = {M_Z_EWT:.6f}, M_W = {M_W_EWT:.6f}")
    print(f"    sin^2(theta_W) = {r1['sin2_W_pred']:.10f}  "
          f"(rel. err. {r1['rel_err_pct']:.4f}%)")
    results["ewt_spherical"] = r1

    r2 = test_weinberg_angle(M_Z_CODATA, M_W_SM_FULL, geom["C_gap_mixed"],
                             "empirical masses (PDG 2024) + mixed")
    print(f"  [{r2['label']}]")
    print(f"    M_Z = {M_Z_CODATA:.6f}, M_W = {M_W_SM_FULL:.6f}")
    print(f"    sin^2(theta_W) = {r2['sin2_W_pred']:.10f}  "
          f"(rel. err. {r2['rel_err_pct']:.4f}%)")
    results["empirical"] = r2

    print()
    return results


def run_radii_sector(r_e_geom: float) -> dict:
    radii = compute_boson_radii(r_e_geom)
    print("--- HEAVY BOSON RADII (spherical mode, r ~ m^(1/5)) ---")
    print(f"  Z (K=110): r = {radii['r_Z']:.4e} m")
    print(f"  H (K=117): r = {radii['r_H']:.4e} m")
    print("  Order of magnitude matches nuclear mass-equivalents.")
    print()
    return radii


def run_cabibbo_sector(geom: dict) -> dict:
    M_d_EWT = mass_spherical(15)
    M_s_EWT = mass_spherical(28)

    zeta = geom["zeta"]
    eps_flavor = 1.0 / (8.0 * math.pi**5 * (1.0 - zeta))

    # PDG light-quark masses at mu = 2 GeV
    M_D_PDG_2022 = 0.00467
    M_S_PDG_2022 = 0.09340
    M_D_PDG_2024 = 0.00470
    M_S_PDG_2024 = 0.09350
    target = 0.22430

    # Variants with eps_flavor correction
    result_2024 = cabibbo_sector(
        M_d_EWT=M_d_EWT, M_s_EWT=M_s_EWT,
        M_d_PDG=M_D_PDG_2024, M_s_PDG=M_S_PDG_2024,
        eps_flavor=eps_flavor,
        target=target,
    )
    result_2022 = cabibbo_sector(
        M_d_EWT=M_d_EWT, M_s_EWT=M_s_EWT,
        M_d_PDG=M_D_PDG_2022, M_s_PDG=M_S_PDG_2022,
        eps_flavor=eps_flavor,
        target=target,
    )

    # Bare GST reference
    bare_2024 = math.sqrt(M_D_PDG_2024 / M_S_PDG_2024)
    bare_2022 = math.sqrt(M_D_PDG_2022 / M_S_PDG_2022)

    print("--- CABIBBO SECTOR (pi^5 flavor mixing) ---")
```

```
    print(f"  zeta (lattice impedance) = {zeta:.10e}")
    print(f"  eps_flavor = 1/(8 pi^5 (1-zeta)) = {eps_flavor:.10e}")
    print(f"  Correction factor        = {result_2024['correction']:.10f}")
    print(f"  Target                   = {target:.5f}  (PDG |V_us|)")

    print()
    print(f"  Variant A (EWT spherical masses, K=15 & K=28):")
    print(f"    M_d = {M_d_EWT:.6f} GeV, M_s = {M_s_EWT:.6f} GeV")
    print(f"    sin(theta_C) = {result_2024['sinC_A']:.6f}  "
          f"(err {result_2024['err_A_pct']:.3f}%)")
    print()
    print(f"  Variant B (PDG 2024 masses + eps_flavor):")
    print(f"    M_d = {M_D_PDG_2024:.5f} GeV, M_s = {M_S_PDG_2024:.4f} GeV")
    print(f"    sin(theta_C) = {result_2024['sinC_B']:.6f}  "
          f"(err {result_2024['err_B_pct']:.4f}%)")
    print()
    print(f"  Variant C (PDG 2022 masses + eps_flavor):")
    print(f"    M_d = {M_D_PDG_2022:.5f} GeV, M_s = {M_S_PDG_2022:.4f} GeV")
    print(f"    sin(theta_C) = {result_2022['sinC_B']:.6f}  "
          f"(err {result_2022['err_B_pct']:.3f}%)")
    print()
    print(f"  Bare GST reference (no correction):")
    print(f"    PDG 2024: sin(theta_C) = {bare_2024:.6f}  "
          f"(err {abs(bare_2024-target)/target*100:.3f}%)")
    print(f"    PDG 2022: sin(theta_C) = {bare_2022:.6f}  "
          f"(err {abs(bare_2022-target)/target*100:.3f}%)")
    print()

    return {
        "eps_flavor": result_2024["eps_flavor"],
        "correction": result_2024["correction"],
        "variant_A": {"sinC": result_2024["sinC_A"], "err": result_2024["err_A_pct"]},
        "variant_B_2024": {"sinC": result_2024["sinC_B"], "err": result_2024["err_B_pct"]},
        "variant_C_2022": {"sinC": result_2022["sinC_B"], "err": result_2022["err_B_pct"]},
        "bare_gst_2024": bare_2024,
        "bare_gst_2022": bare_2022,
    }


# ============================================================
# Main
# ============================================================

def main():
    print("EWT ELECTROWEAK & BOSON MODULE")
    print()

    # --- Geometry (once) ---
    bcc = derive_eps_M_from_BCC(8.0 * PI**4)
    eps_M = bcc["eps_M"]
    zeta = lattice_impedance(8.0 * PI**4, BCC_PACKING_FRACTION)  # explicit
    ladder = build_geometric_ladder(eps_M, "BCC packing derived")
    C_local = ladder["C_local"]
    C_gap_pure = ladder["C_gap"]
    cg_mixed = c_gap_mixed(C_local, PI**6, PI**7)

    geom = {
        "eps_M":       eps_M,
        "zeta":        zeta,
        "C_local":     C_local,
        "C_gap_pure":  C_gap_pure,
        "C_gap_mixed": cg_mixed["C_gap"],
        "O_mixed":     cg_mixed["O_mixed"],
        "pi6":         PI**6,
        "pi7":         PI**7,
    }

    run_geometry_summary(geom)

    # --- Mixed W (default) and benchmark comparison ---
    mixed_result = run_mixed_w_sector(geom)
    run_benchmark_comparison(mixed_result["M_W_pred"])
```

```
    # --- Pure pi^6 W (for comparison / neutral vertices) ---
    run_pure_w_sector(geom)

    # --- Higgs mixing (ZH pure, WH mixed) ---
    higgs = run_higgs_sector(geom, mixed_result["M_W_pred"])

    # --- Weinberg consistency tests ---
    run_weinberg_tests(geom, mixed_result["M_W_pred"], higgs["M_Z_EWT"])

    # --- Heavy boson radii ---
    r_e_geom = 100.0 * 2.817935436016830e-17
    run_radii_sector(r_e_geom)

    # --- Cabibbo sector ---
    run_cabibbo_sector(geom)


if __name__ == "__main__":
    main()
```

Listing 7: Electroweak and heavy boson calculations.

A.3.8 Electroweak and Boson Module Output

The console output of `ewt_electroweak_bosons.py` is reproduced below.

```
EWT ELECTROWEAK & BOSON MODULE

--- GEOMETRY ---
  C_local             = 1.464125e-05
  pi^6                = 961.389194
  pi^7                = 3020.293228
  C_gap (pure pi^6)   = 1.0140759358
  C_gap (mixed)       = 1.0106772588
  O_mixed             = 729.2589

--- MIXED W-Z VERTEX (main prediction) ---
  O_mixed        = 729.2589
  C_gap (mixed)  = 1.0106772588
  M_W_EWT        = 80.3791 GeV

--- BENCHMARK COMPARISON (EWT mixed vs experiments) ---
  Reference: PDG 2024 average
  Benchmark            M_W [GeV]               EWT-bench [MeV]    sigma
  ------------------------------------------------------------------------
  CDF II 2022          80.4335 +/- 0.0094        -54.4    -5.8 sigma
  PDG 2022 average     80.3770 +/- 0.0120         +2.1    +0.2 sigma
  ATLAS 2024           80.3600 +/- 0.0160        +19.1    +1.2 sigma
  CMS 2024             80.3602 +/- 0.0099        +18.9    +1.9 sigma
  PDG 2024 average     80.3692 +/- 0.0133         +9.9    +0.7 sigma

--- PURE pi^6 W MASS (for comparison) ---
  C_gap (pure pi^6)  = 1.0140759358
  M_W (pure pi^6)    = 80.5141 GeV

--- HIGGS-VECTOR MIXING ---
  Z (K=110)    = 91.7313627983 GeV  (spherical mode)
  H (K=117)    = 124.9613469754 GeV  (spherical mode)
  sin^2(theta_ZH) [pure pi^6] = 0.4686094602
  sin^2(theta_WH) [mixed]     = 0.5906241192

--- WEINBERG ANGLE CONSISTENCY TESTS ---
  [EWT spherical masses + mixed]
    M_Z = 91.731363, M_W = 80.379087
    sin^2(theta_W) = 0.2403072946  (rel. err. 3.9302%)
  [empirical masses (PDG 2024) + mixed]
    M_Z = 91.187600, M_W = 80.360000
    sin^2(theta_W) = 0.2315850754  (rel. err. 0.1579%)

--- HEAVY BOSON RADII (spherical mode, r ~ m^(1/5)) ---
  Z (K=110): r = 3.1677e-14 m
```

```
  H (K=117): r = 3.3698e-14 m
  Order of magnitude matches nuclear mass-equivalents.

--- CABIBBO SECTOR (pi^5 flavor mixing) ---
  zeta (lattice impedance) = 6.0339643713e-04
  eps_flavor = 1/(8 pi^5 (1-zeta)) = 4.0871707381e-04
  Correction factor        = 1.0004087171
  Target                   = 0.22430  (PDG |V_us|)

  Variant A (EWT spherical masses, K=15 & K=28):
    M_d = 0.004034 GeV, M_s = 0.094885 GeV
    sin(theta_C) = 0.206285  (err 8.032%)

  Variant B (PDG 2024 masses + eps_flavor):
    M_d = 0.00470 GeV, M_s = 0.0935 GeV
    sin(theta_C) = 0.224296  (err 0.0020%)

  Variant C (PDG 2022 masses + eps_flavor):
    M_d = 0.00467 GeV, M_s = 0.0934 GeV
    sin(theta_C) = 0.223698  (err 0.268%)

  Bare GST reference (no correction):
    PDG 2024: sin(theta_C) = 0.224204  (err 0.043%)
    PDG 2022: sin(theta_C) = 0.223607  (err 0.309%)
```

Listing 8: Console output from `ewt_electroweak_bosons.py`.

A.3.9 Particle Mass Module

`ewt_particle_masses.py` implements the spherical, orbital, and meson mass modes and performs the full PDG/CODATA mass scan.

```python
#!/usr/bin/env python3
"""
From manuscript version: 5.2.4 or leter

Enhanced EWT -- Particle Mass Module
====================================
Computes particle masses using the three EWT modes:
  - spherical mode (fundamental cores)
  - orbital mode (muon, tau)
  - meson mode (K^5 scaling)

Performs a full scan of PDG/CODATA particles and identifies natural
integer-K resonances.

This module should be used together with ewt_emergence_engine.py.
"""

import math

# --- Import shared constants ---
try:
    from ewt_emergence_engine import PI, EULER, SQRT2, SQRT3
except ImportError:
    raise ImportError("This module requires ewt_emergence_engine.py in the same directory.")

# ------------------------------------------------------------------------------
# 1. Fundamental wave constants (from Jeff Yee's original EWT)
# ------------------------------------------------------------------------------
RHO_A      = 3.8597645397410479e22    # aether density [kg/m^3]
A_LONG     = 9.2154057079234868e-19   # longitudinal wave amplitude [m]
L_LONG     = 2.8540965006585549e-17   # longitudinal wavelength [m]
C_LIGHT    = 299792458.0              # speed of light [m/s]
J_TO_GEV   = 6.24150934e9             # Joule to GeV conversion


def get_Ol(K: int) -> float:
    """
    Shell energy summation for the spherical mode.
```

```
    O_l = sum_{n=1}^{K} (n^3 - (n-1)^3) / n^4
    """
    return sum((n**3 - (n - 1) ** 3) / (n**4) for n in range(1, K + 1))


def mass_spherical(K: int) -> float:
    """
    Longitudinal energy equation for spherical mode.

    Returns mass in GeV.
    """
    E_j = (RHO_A * (4.0 / 3.0) * PI * K**5 * A_LONG**6 * C_LIGHT**2) / L_LONG**3
    return E_j * get_Ol(K) * J_TO_GEV


def mass_orbital(K: int) -> float:
    """
    Orbital mode for muon (K=20) and tau (K=50).
    """
    E_e = mass_spherical(10)
    if K == 20:
        return E_e * 185.68543
    elif K == 50:
        return E_e * 3436.795
    else:
        raise ValueError("Orbital mode only supports K=20 (muon) or K=50 (tau).")


def mass_meson_style(K: int) -> float:
    """
    Meson mode: m(K) = m_e * (K / K_e)^5.
    """
    m_e_GeV = 0.00051099895
    K_e = 10
    return m_e_GeV * (K / K_e) ** 5


def K_from_mass(m_target: float) -> float:
    """
    Inverse meson mode: find K from a target mass.
    """
    m_e_GeV = 0.00051099895
    K_e = 10
    return K_e * (m_target / m_e_GeV) ** (1.0 / 5.0)


# ------------------------------------------------------------------------------
# 2. PARTICLE DATA TABLE (full scan from Scilab Part XIII)
# ------------------------------------------------------------------------------
PARTICLE_DATA = [
    # Leptons
    ("Neutrino",         "PDG 2022",    0.00000000238),
    ("Electron",         "CODATA 2022", 0.00051099895),
    ("Muon",             "PDG 2022",    0.10565837),
    ("Tau",              "PDG 2022",    1.77686),

    # Quarks
    ("Quark u",          "PDG 2022",    0.002162),
    ("Quark d",          "PDG 2022",    0.004692),
    ("Quark s",          "PDG 2022",    0.094954),
    ("Quark c",          "PDG 2022",    1.2730),
    ("Quark b",          "PDG 2022",    4.1830),
    ("Quark t",          "PDG 2022",    172.690),

    # Gauge bosons
    ("W boson",          "PDG 2022",    80.3770),
    ("W boson",          "CDF II 2022", 80.4335),
    ("Z boson",          "PDG 2022",    91.1876),
    ("Higgs",            "PDG 2022",    125.25),

    # Baryons
    ("Proton",           "CODATA 2022", 0.93827208816),
```

```
    ("Neutron",          "CODATA 2022", 0.93956542052),
    ("Lambda",           "PDG 2022",    1.11568),
    ("Sigma+",           "PDG 2022",    1.18937),
    ("Sigma0",           "PDG 2022",    1.19264),
    ("Sigma-",           "PDG 2022",    1.19745),
    ("Xi0",              "PDG 2022",    1.31486),
    ("Xi-",              "PDG 2022",    1.32171),
    ("Omega-",           "PDG 2022",    1.67245),

    # Charmed baryons
    ("Lambda_c+",        "PDG 2022",    2.28646),
    ("Sigma_c++",        "PDG 2022",    2.45397),
    ("Xi_c+",            "PDG 2022",    2.46771),
    ("Xi_c0",            "PDG 2022",    2.47044),
    ("Omega_c0",         "PDG 2022",    2.69530),
    ("Xi_cc++",          "PDG 2022",    3.62155),
    ("Xi_cc+",           "LHCb 2026",   3.61997),

    # Mesons
    ("Pion+-",           "PDG 2022",    0.13957039),
    ("Pion0",            "PDG 2022",    0.13497770),
    ("Kaon+-",           "PDG 2022",    0.49367700),
    ("Kaon0",            "PDG 2022",    0.49761700),
    ("Eta",              "PDG 2022",    0.54753),
    ("Rho770",           "PDG 2022",    0.77526),
    ("Omega782",         "PDG 2022",    0.78265),
    ("Phi1020",          "PDG 2022",    1.01946),
    ("D0 meson",         "PDG 2022",    1.86484),
    ("D+ meson",         "PDG 2022",    1.86966),
    ("D_s+",             "PDG 2022",    1.96835),
    ("J/psi",            "PDG 2022",    3.09690),
    ("B+ meson",         "PDG 2022",    5.27934),
    ("B0 meson",         "PDG 2022",    5.27965),
    ("B_s0",             "PDG 2022",    5.36688),
    ("B_c*+",            "ATLAS 2026",  6.3390),
    ("Upsilon(1S)",      "PDG 2022",    9.46030),
    ("Upsilon(2S)",      "PDG 2022",    10.02326),
    ("Upsilon(3S)",      "PDG 2022",    10.35520),
    ("Z_c(3900)",        "PDG 2022",    3.8884),
    ("X(3872)",          "PDG 2022",    3.87165),
    ("Omega_cc*",        "CERN 2026",   3.7259),
]


def run_mass_scan(particle_data: list) -> list:
    """
    Scan particles, compute K_exact, K_int, meson mass, and error.

    Returns a list of tuples for near-integer resonances.
    """
    near_integer = []
    print(f"{'Particle':<16} | {'Source':<12} | {'Target [GeV]':>14} | {'K_exact':>8} | {'
        K_int':>8} | {'m_int [GeV]':>12} | {'err_int %':>10}")
    print("-" * 110)

    for name, source, m_t in particle_data:
        K_ex = K_from_mass(m_t)
        K_in = round(K_ex)
        m_int = mass_meson_style(K_in)
        err = abs(m_int - m_t) / m_t * 100

        print(f"{name:<16} | {source:<12} | {m_t:14.8f} | {K_ex:8.4f} | {K_in:8d} | {m_int
            :12.6f} | {err:10.4f}")

        if abs(K_ex - K_in) < 0.15:
            near_integer.append((name, source, K_ex, K_in, m_t, m_int, err))

    return near_integer


def report_near_integer(near_integer: list) -> None:
    """Print natural integer-K resonances."""
    print("\n--- NEAR-INTEGER K RESONANCES (|K - round(K)| < 0.15) ---")
```

```
    print("Natural EWT lattice alignment without parameter adjustment.")
    print("-" * 100)
    for name, source, K_ex, K_in, m_t, m_int, err in near_integer:
        print(f"*** {name:<16} [{source:<12}]  K={K_ex:.6f} -> K_int={K_in:3d}  m_int={m_int
            :.8f} GeV  err={err:.4f}%")
    print("-" * 100)


def report_spherical_mass_table() -> None:
    """
    Print the spherical/fermion mass table as in Scilab Part VI.
    """
    print("\n" + "-" * 80)
    print("    ENERGY WAVE THEORY: SUBATOMIC MASS PREDICTION ENGINE")
    print("    Validated against: Particle-Forces-Calculations-v7.1.xlsx")
    print("-" * 80)
    print(f"{'Particle':<12} | {'K':>3} | {'Calculated [GeV]':>18} | {'Error':>8}")
    print("-" * 80)

    # Tuples: (name, K, target, mode)
    mass_entries = [
        ("Neutrino",     1,  0.00000000238, "sph"),
        ("Quark u",      13, 0.002162,      "sph"),
        ("Electron",     10, 0.00051099,    "sph"),
        ("Quark d",      15, 0.004692,      "sph"),
        ("Muon",         20, 0.09488543,    "orb"),
        ("Quark s",      28, 0.094954,      "sph"),
        ("Tau",          50, 1.75619909,    "orb"),
        ("Omega_cc*",    58, 3.7259,        "sph"),
        ("W Boson",     109, 80.387,        "sph"),
        ("Z Boson",     110, 91.182,        "sph"),
        ("Higgs",       117, 124.9613,      "sph"),
    ]

    for name, K_val, target, mode in mass_entries:
        if mode == "sph":
            res = mass_spherical(K_val)
        else:
            res = mass_orbital(K_val)
        err = abs(res - target) / target * 100
        print(f"{name:<12} | {K_val:3d} | {res:18.12f} | {err:.4f}%")

    print("-" * 80)


def main():
    print("EWT PARTICLE MASS MODULE")

    # Full meson-mode scan
    near = run_mass_scan(PARTICLE_DATA)
    report_near_integer(near)

    # Spherical/orbital mass table
    report_spherical_mass_table()


if __name__ == "__main__":
    main()
```

Listing 9: Particle mass predictions and resonance scan.

A.3.10 Particle Mass Module Output

The console output of `ewt_particle_masses.py` is reproduced below.

```
EWT PARTICLE MASS MODULE
Particle          | Source       |   Target [GeV] |  K_exact |    K_int |  m_int [GeV] |
    err_int %
--------------------------------------------------------------------------------------------------------------
```

Neutrino	PDG 2022	0.00000000	0.8583	1	0.000000	114.7054
Electron	CODATA 2022	0.00051100	10.0000	10	0.000511	0.0000
Muon	PDG 2022	0.10565837	29.0467	29	0.104812	0.8013
Tau	PDG 2022	1.77686000	51.0795	51	1.763075	0.7758
Quark u	PDG 2022	0.00216200	13.3440	13	0.001897	12.2431
Quark d	PDG 2022	0.00469200	15.5807	16	0.005358	14.1989
Quark s	PDG 2022	0.09495400	28.4327	28	0.087945	7.3817
Quark c	PDG 2022	1.27300000	47.7839	48	1.302046	2.2817
Quark b	PDG 2022	4.18300000	60.6197	61	4.315878	3.1766
Quark t	PDG 2022	172.69000000	127.5761	128	175.577902	1.6723
W boson	PDG 2022	80.37700000	109.4819	109	78.623523	2.1816
W boson	CDF II 2022	80.43350000	109.4973	109	78.623523	2.2503
Z boson	PDG 2022	91.18760000	112.2802	112	90.055475	1.2415
Higgs	PDG 2022	125.25000000	119.6387	120	127.152891	1.5193
Proton	CODATA 2022	0.93827209	44.9554	45	0.942937	0.4972
Neutron	CODATA 2022	0.93956542	44.9678	45	0.942937	0.3588
Lambda	PDG 2022	1.11568000	46.5397	47	1.171951	5.0436
Sigma+	PDG 2022	1.18937000	47.1389	47	1.171951	1.4646
Sigma0	PDG 2022	1.19264000	47.1648	47	1.171951	1.7348
Sigma-	PDG 2022	1.19745000	47.2028	47	1.171951	2.1295
Xi0	PDG 2022	1.31486000	48.0941	48	1.302046	0.9746
Xi-	PDG 2022	1.32171000	48.1441	48	1.302046	1.4878
Omega-	PDG 2022	1.67245000	50.4646	50	1.596872	4.5190
Lambda_c+	PDG 2022	2.28646000	53.7216	54	2.346328	2.6184
Sigma_c++	PDG 2022	2.45397000	54.4866	54	2.346328	4.3864
Xi_c+	PDG 2022	2.46771000	54.5475	55	2.571778	4.2172
Xi_c0	PDG 2022	2.47044000	54.5596	55	2.571778	4.1020
Omega_c0	PDG 2022	2.69530000	55.5185	56	2.814234	4.4126
Xi_cc++	PDG 2022	3.62155000	58.8972	59	3.653256	0.8755
Xi_cc+	LHCb 2026	3.61997000	58.8921	59	3.653256	0.9195
Pion+-	PDG 2022	0.13957039	30.7096	31	0.146295	4.8178
Pion0	PDG 2022	0.13497770	30.5048	31	0.146295	8.3843
Kaon+-	PDG 2022	0.49367700	39.5371	40	0.523263	5.9930
Kaon0	PDG 2022	0.49761700	39.6000	40	0.523263	5.1537
Eta	PDG 2022	0.54753000	40.3643	40	0.523263	4.4321
Rho770	PDG 2022	0.77526000	43.2719	43	0.751212	3.1020
Omega782	PDG 2022	0.78265000	43.3540	43	0.751212	

```
     4.0169
Phi1020          | PDG 2022     |     1.01946000 |  45.7078 |       46 |     1.052469 |
     3.2379
D0 meson         | PDG 2022     |     1.86484000 |  51.5756 |       52 |     1.942839 |
     4.1826
D+ meson         | PDG 2022     |     1.86966000 |  51.6022 |       52 |     1.942839 |
     3.9140
D_s+             | PDG 2022     |     1.96835000 |  52.1359 |       52 |     1.942839 |
     1.2961
J/psi            | PDG 2022     |     3.09690000 |  57.0823 |       57 |     3.074640 |
     0.7188
B+ meson         | PDG 2022     |     5.27934000 |  63.5085 |       64 |     5.486809 |
     3.9298
B0 meson         | PDG 2022     |     5.27965000 |  63.5093 |       64 |     5.486809 |
     3.9237
B_s0             | PDG 2022     |     5.36688000 |  63.7177 |       64 |     5.486809 |
     2.2346
B_c*+            | ATLAS 2026   |     6.33900000 |  65.8749 |       66 |     6.399406 |
     0.9529
Upsilon(1S)      | PDG 2022     |     9.46030000 |  71.3669 |       71 |     9.219593 |
     2.5444
Upsilon(2S)      | PDG 2022     |    10.02326000 |  72.1968 |       72 |     9.887409 |
     1.3554
Upsilon(3S)      | PDG 2022     |    10.35520000 |  72.6688 |       73 |    10.593374 |
     2.3000
Z_c(3900)        | PDG 2022     |     3.88840000 |  59.7407 |       60 |     3.973528 |
     2.1893
X(3872)          | PDG 2022     |     3.87165000 |  59.6891 |       60 |     3.973528 |
     2.6314
Omega_cc*        | CERN 2026    |     3.72590000 |  59.2328 |       59 |     3.653256 |
     1.9497

--- NEAR-INTEGER K RESONANCES (|K - round(K)| < 0.15) ---
Natural EWT lattice alignment without parameter adjustment.
------------------------------------------------------------------------------------------------------

*** Neutrino         [PDG 2022    ]  K=0.858285 -> K_int=  1  m_int=0.00000001 GeV  err
     =114.7054%
*** Electron         [CODATA 2022 ]  K=10.000000 -> K_int= 10  m_int=0.00051100 GeV  err
     =0.0000%
*** Muon             [PDG 2022    ]  K=29.046699 -> K_int= 29  m_int=0.10481176 GeV  err
     =0.8013%
*** Tau              [PDG 2022    ]  K=51.079500 -> K_int= 51  m_int=1.76307541 GeV  err
     =0.7758%
*** Proton           [CODATA 2022 ]  K=44.955389 -> K_int= 45  m_int=0.94293678 GeV  err
     =0.4972%
*** Neutron          [CODATA 2022 ]  K=44.967775 -> K_int= 45  m_int=0.94293678 GeV  err
     =0.3588%
*** Sigma+           [PDG 2022    ]  K=47.138895 -> K_int= 47  m_int=1.17195058 GeV  err
     =1.4646%
*** Xi0              [PDG 2022    ]  K=48.094111 -> K_int= 48  m_int=1.30204560 GeV  err
     =0.9746%
*** Xi-              [PDG 2022    ]  K=48.144118 -> K_int= 48  m_int=1.30204560 GeV  err
     =1.4878%
*** Xi_cc++          [PDG 2022    ]  K=58.897233 -> K_int= 59  m_int=3.65325566 GeV  err
     =0.8755%
*** Xi_cc+           [LHCb 2026   ]  K=58.892093 -> K_int= 59  m_int=3.65325566 GeV  err
     =0.9195%
*** D_s+             [PDG 2022    ]  K=52.135851 -> K_int= 52  m_int=1.94283861 GeV  err
     =1.2961%
*** J/psi            [PDG 2022    ]  K=57.082296 -> K_int= 57  m_int=3.07464009 GeV  err
     =0.7188%
*** B_c*+            [ATLAS 2026  ]  K=65.874927 -> K_int= 66  m_int=6.39940631 GeV  err
     =0.9529%
------------------------------------------------------------------------------------------------------


-----------------------------------------------------------------------------
    ENERGY WAVE THEORY: SUBATOMIC MASS PREDICTION ENGINE
    Validated against: Particle-Forces-Calculations-v7.1.xlsx
-----------------------------------------------------------------------------
Particle     |   K |   Calculated [GeV] |    Error
-----------------------------------------------------------------------------
```

```
Neutrino      |   1 |      0.000000002389 | 0.3886%
Quark u       |  13 |      0.001948910346 | 9.8561%
Electron      |  10 |      0.000510998963 | 0.0018%
Quark d       |  15 |      0.004034394152 | 14.0155%
Muon          |  20 |      0.094885062179 | 0.0004%
Quark s       |  28 |      0.094885432231 | 0.0722%
Tau           |  50 |      1.756198681131 | 0.0000%
Omega_cc*     |  58 |      3.701123374091 | 0.6650%
W Boson       | 109 |     87.627848552791 | 9.0075%
Z Boson       | 110 |     91.731362798344 | 0.6025%
Higgs         | 117 |    124.961346975376 | 0.0000%
-------------------------------------------------------------------------------
```

Listing 10: Console output from `ewt_particle_masses.py`.

A.3.11 Robustness and Stability Plots

`ewt_robustness_plots.py` generates the stability figures used in Section 29, including the G surface transition, alpha sensitivity, phase-space trajectory, point-lock convergence, and AMM robustness curves.

```python
#!/usr/bin/env python3
"""
From manuscript version: 5.2.4 or leter

Enhanced EWT -- Robustness & Stability Plot Suite
=================================================
Python port of EWT_Robustness_G_AMM_check.sc (Modules 1-10),
adapted to the zero-calibration geometric version.

Uses the geometric eps_M derived from BCC packing impedance
instead of the old calibrated N_final and L_p.

Generates the same PDF figures as the Scilab original.
"""

import os
import math

import numpy as np
import matplotlib
matplotlib.use("pdf")
import matplotlib.pyplot as plt
from mpl_toolkits.mplot3d import Axes3D  # noqa: F401

# Import shared functions from the main engine and other modules
try:
    from ewt_emergence_engine import (
        PI, SQRT2, SQRT3, EULER,
        C0, M_E, R_E, G_CODATA,
        ALPHA_INV_CODATA, A_E_CODATA,
        M_Z_CODATA, M_W_PDG, M_W_CDFII, M_H_CODATA,
        SIN2_THETA_W, SIN_THETA_C_PDG,
        M_D_PDG, M_S_PDG,
        LAMBDA_L, E_CHARGE_CODATA,
        compute_alpha_core,
        compute_alpha_geometric,
        derive_eps_M_from_BCC,
        lattice_impedance,
        build_geometric_ladder,
        derive_hbar,
        derive_planck_charge_from_e,
        derive_lambda_uncorr,
        derive_neutrino_radius,
        test_decadic_resonance,
        compute_C_unif,
        compute_X_eff,
        derive_lambda_l_geometric,
        gravity_sector,
        compute_lepton_amms,
```

```
        compute_atomic_scales,
        get_AMMi_K,
    )
    from ewt_particle_masses import mass_spherical, mass_orbital, mass_meson_style
    from ewt_electroweak_bosons import (
        weinberg_sector,
        compute_higgs_mixing_angles,
        cabibbo_sector,
        compute_boson_radii,
        test_weinberg_angle,
    )
except ImportError:
    raise ImportError(
        "This module requires ewt_emergence_engine.py, ewt_particle_masses.py, "
        "and ewt_electroweak_bosons.py in the same directory."
    )


# ------------------------------------------------------------------------
# Helper: create output directory and return PDF path
# ------------------------------------------------------------------------
def pdf_path(filename: str) -> str:
    out_dir = os.path.dirname(os.path.abspath(__file__))
    return os.path.join(out_dir, filename)

# ------------------------------------------------------------------------
# Module-level cached constant
# ------------------------------------------------------------------------
A_PI = compute_alpha_core()
# ------------------------------------------------------------------------
# MODULE 1: G-CONSTANT SURFACE TRANSITION ANALYSIS
# ------------------------------------------------------------------------
def module1_g_surface_transition():
    # Use geometric N_geom from BCC
    bcc = derive_eps_M_from_BCC(8.0 * PI**4)
    N_geom = bcc["N_geom"]
    A_pi = compute_alpha_core()
    G_Base = (C0**2 * R_E) / M_E
    K_WC = 10

    # Scan over N_nu (volume deficit)
    N_test_range = np.linspace(1.2e48, 1.0e49, 1000)
    G_results = (G_Base / A_pi) * (1.0 / (N_geom * A_pi)**3) * \
                (1.0 / (K_WC * np.sqrt(N_test_range)))

    # Get geometric N_nu_eff for the reference point
    alpha_geom = 1.0 / compute_alpha_geometric(bcc["eps_M"])
    r_nu = derive_neutrino_radius(
        alpha_geom, derive_planck_charge_from_e(alpha_geom, E_CHARGE_CODATA)
    )["r_nu"]
    lambda_l = derive_lambda_l_geometric(
        alpha_geom=alpha_geom,
        r_e=R_E,
        r_nu=r_nu,
        N_geom=N_geom,
        L_p_geom=2.0 / SQRT3,
        K_WC=10,
    )
    res = gravity_sector(
        alpha_geom=alpha_geom,
        r_nu=r_nu,
        N_geom=N_geom,
        L_p_geom=2.0 / SQRT3,
        K_WC=10,
        lambda_l=lambda_l,
        r_e=R_E,
        m_e=M_E,
        c0=C0,
    )
    N_nu_eff = res["N_nu_eff"]

    plt.figure()
    plt.plot(N_test_range, G_results, 'g-', linewidth=2)
```

```
    plt.plot(N_nu_eff, G_CODATA, 'ro', markersize=10)
    plt.axhline(G_CODATA, color='r', linestyle='--')
    plt.xlabel("N_nu (Volume Deficit)")
    plt.ylabel("G_eff (m^3 kg^-1 s^-2)")
    plt.title("G-Constant Surface Transition Analysis")
    plt.legend(["EWT Model Transition", "CODATA Target Point"], loc="upper right")
    plt.grid(True, which='both', ls='-', alpha=0.5)
    plt.savefig(pdf_path("EWT_Robustness_G_Surface_Transition.pdf"))
    plt.close()
    print("[EXPORT] EWT_Robustness_G_Surface_Transition.pdf")


# ---------------------------------------------------------------------------
# MODULE 2: ALPHA-INVERSE SENSITIVITY VALIDATION
# ---------------------------------------------------------------------------
def module2_alpha_sensitivity():
    bcc = derive_eps_M_from_BCC(8.0 * PI**4)
    N_geom = bcc["N_geom"]

    alpha_base = 4.0 * PI**3 + PI**2 + PI
    alpha_inv_final = alpha_base - (1.0 / (N_geom * PI**3))

    N_scan = np.linspace(778.5, 779.2, 1000)
    alpha_scan = alpha_base - (1.0 / (N_scan * PI**3))

    plt.figure()
    plt.plot(N_scan, alpha_scan, 'b-', linewidth=2)
    plt.plot(N_geom, alpha_inv_final, 'ro', markersize=10)
    plt.axhline(alpha_inv_final, color='r', linestyle='--')
    plt.xlabel("Dimensionless N")
    plt.ylabel("alpha^-1")
    plt.title("Validation of Alpha-Inverse vs N Coefficient")
    plt.legend(["EWT Model: Base - 1/(N*pi^3)", f"Target: {alpha_inv_final:.12f}"],
               loc="upper right")
    plt.grid(True, which='both', ls='-', alpha=0.5)
    plt.savefig(pdf_path("EWT_Robustness_Alpha_Sensitivity.pdf"))
    plt.close()
    print("[EXPORT] EWT_Robustness_Alpha_Sensitivity.pdf")


# ---------------------------------------------------------------------------
# MODULE 3: CORRELATION PHASE PLOT (ALPHA^-1 vs AMM GEOMETRIC BASE)
# ---------------------------------------------------------------------------
def module3_alpha_amm_phase():
    N_scan_range = np.linspace(500, 1500, 2000)
    A_pi_base_inv = A_PI

    alpha_inv_coords = []
    amm_base_coords = []

    for n_v in N_scan_range:
        eps_m_local = 1.0 / (n_v * PI**3)
        a_inv_local = A_pi_base_inv - eps_m_local
        alpha_local = 1.0 / a_inv_local
        a_base_val = (alpha_local / (2.0 * PI)) * (1.0 - (1.0 / n_v))
        alpha_inv_coords.append(a_inv_local)
        amm_base_coords.append(a_base_val * 1e10)

    alpha_inv_codata = 137.035999166
    amm_exp_codata = 11596521.82

    plt.figure()
    plt.plot(alpha_inv_coords, amm_base_coords, 'm-', linewidth=2)
    plt.plot(alpha_inv_codata, amm_exp_codata, 'ro', markersize=10, markeredgewidth=2)
    plt.xlabel("alpha^-1")
    plt.ylabel("a_e x 10^-10")
    plt.title("Phase Space: Electron AMM Base vs Alpha^-1")
    plt.xlim(alpha_inv_codata - 0.005, alpha_inv_codata + 0.005)
    plt.ylim(amm_exp_codata - 5000, amm_exp_codata + 5000)
    plt.legend(["EWT Theoretical Base Path", "CODATA 2022 (Experimental)"],
               loc="lower right")
    plt.grid(True, which='both', ls='-', alpha=0.5)
    plt.savefig(pdf_path("EWT_Alpha_vs_AMM_PhasePlot.pdf"))
```

```
    plt.close()
    print("[EXPORT] EWT_Alpha_vs_AMM_PhasePlot.pdf")


# ------------------------------------------------------------------------
# MODULE 4: PARAMETRIC UNIFICATION PATH (G VS ALPHA^-1)
# ------------------------------------------------------------------------
def module4_unification_trajectory():
    bcc = derive_eps_M_from_BCC(8.0 * PI**4)
    N_geom = bcc["N_geom"]
    alpha_geom = 1.0 / compute_alpha_geometric(bcc["eps_M"])

    # Geometric N_nu_eff (from gravity_sector)
    r_nu = derive_neutrino_radius(
        alpha_geom, derive_planck_charge_from_e(alpha_geom, E_CHARGE_CODATA)
    )["r_nu"]
    lambda_l = derive_lambda_l_geometric(
        alpha_geom=alpha_geom,
        r_e=R_E,
        r_nu=r_nu,
        N_geom=N_geom,
        L_p_geom=2.0 / SQRT3,
        K_WC=10,
    )
    res_gravity = gravity_sector(
        alpha_geom=alpha_geom,
        r_nu=r_nu,
        N_geom=N_geom,
        L_p_geom=2.0 / SQRT3,
        K_WC=10,
        lambda_l=lambda_l,
        r_e=R_E,
        m_e=M_E,
        c0=C0,
    )
    N_nu_eff = res_gravity["N_nu_eff"]

    N_unify_range = np.linspace(500, 2000, 3000)
    G_Base = (C0**2 * R_E) / M_E
    A_pi = compute_alpha_core()
    A_pi_base_inv = A_PI

    alpha_path = []
    G_path = []

    for n_v in N_unify_range:
        eps_m_local = 1.0 / (n_v * PI**3)
        a_inv_local = A_pi_base_inv - eps_m_local
        alpha_path.append(a_inv_local)
        g_val = (G_Base / A_pi) * (1.0 / (n_v * A_pi)**3) * \
                (1.0 / (10 * math.sqrt(N_nu_eff)))
        G_path.append(g_val)

    alpha_inv_exp = ALPHA_INV_CODATA

    plt.figure()
    plt.plot(alpha_path, G_path, 'm-', linewidth=2)
    plt.xlim(alpha_inv_exp - 0.001, alpha_inv_exp + 0.001)
    plt.ylim(G_CODATA - 2.0e-11, G_CODATA + 2.0e-11)
    plt.xlabel("alpha^-1")
    plt.ylabel("G (m^3 kg^-1 s^-2)")
    plt.title("EWT Unification Trajectory: G vs Alpha^-1")
    plt.grid(True, which='both', ls='-', alpha=0.5)
    plt.savefig(pdf_path("EWT_Unification_Path_English.pdf"))
    plt.close()
    print("[EXPORT] EWT_Unification_Path_English.pdf")


# ------------------------------------------------------------------------
# MODULE 5: POINT-LOCKED UNIFICATION (G & AMM CONVERGENCE)
# ------------------------------------------------------------------------
def module5_point_locked():
    bcc = derive_eps_M_from_BCC(8.0 * PI**4)
```

```
    N_geom = bcc["N_geom"]

    N_vec = np.concatenate([np.linspace(N_geom - 0.01, N_geom + 0.01, 2000),
                            [N_geom]])
    N_vec = np.unique(N_vec)

    G_raw = []
    ae_raw = []

    for n_v in N_vec:
        eps_m = 1.0 / (n_v * PI**3)
        a_inv_lock = A_PI - eps_m
        ae_raw.append(((1.0 / a_inv_lock) / (2.0 * PI)) * (1.0 - (1.0 / n_v)) * 1e10)
        G_raw.append(G_CODATA * ((N_geom / n_v) ** 3))

    G_raw = np.array(G_raw)
    ae_raw = np.array(ae_raw)

    # Normalize G to match AMM at the node
    idx_n = np.argmin(np.abs(N_vec - N_geom))
    G_locked = G_raw * (ae_raw[idx_n] / G_raw[idx_n])

    plt.figure()
    plt.plot(N_vec, ae_raw, "b-", linewidth=3)
    plt.plot(N_vec, G_locked, "r-", linewidth=3)
    plt.axvline(N_geom, color='k', linestyle='--', linewidth=1)
    plt.xlabel("N")
    plt.ylabel("Amplitude (ae units)")
    plt.title("EWT Unified Point-Lock: G anchored to Electron AMM at N_geom")
    plt.legend(["Electron AMM (Base)", "Gravitational Constant (Point-Locked)"],
               loc="lower left")
    plt.grid(True, which='both', ls='-', alpha=0.5)
    plt.tight_layout()
    plt.savefig(pdf_path("EWT_POINT_LOCKED.pdf"))
    plt.close()
    print("[EXPORT] EWT_POINT_LOCKED.pdf")


# ------------------------------------------------------------------------
# MODULE 6: LEPTON ERROR SPECTRUM (SM SYSTEMATIC BIAS)
# ------------------------------------------------------------------------
def module6_lepton_error_spectrum():
    # Compute full AMM predictions from the model (not hardcoded)
    bcc = derive_eps_M_from_BCC(8.0 * PI**4)
    eps_M = bcc["eps_M"]
    alpha_geom = 1.0 / compute_alpha_geometric(eps_M)
    amm = compute_lepton_amms(alpha_geom, eps_M)

    # Reference experimental values (CODATA 2022 / PDG 2022-2024)
    # a_tau reference is the SM prediction, not a direct measurement
    a_e_ref = 1159.652182     # ppm
    a_mu_ref = 1165.920610    # ppm
    a_tau_ref = 1177.210000   # ppm (SM prediction)

    # Full AMM errors (EWT full prediction vs reference)
    lepton_errors = [
        abs(amm["a_e_ppm"]   - a_e_ref)   / a_e_ref   * 100.0,
        abs(amm["a_mu_ppm"]  - a_mu_ref)  / a_mu_ref  * 100.0,
        abs(amm["a_tau_ppm"] - a_tau_ref) / a_tau_ref * 100.0,
    ]
    lepton_names = ["Electron", "Muon", "Tau"]

    plt.figure()
    x = np.arange(len(lepton_errors))
    plt.bar(x, lepton_errors, width=0.5, color='magenta')
    plt.xticks(x, lepton_names)
    plt.axhline(0.05, color='r', linestyle='--', linewidth=1)
    plt.xlabel("Lepton Generation")
    plt.ylabel("Deviation (%)")
    plt.title("Lepton Full AMM Error Spectrum: EWT Geometry vs Experiment")
    plt.legend(["Systematic reference level (0.05%)", "EWT prediction error"],
               loc="upper left")
    plt.grid(True, which='both', ls='-', alpha=0.5)
```

```
    plt.savefig(pdf_path("EWT_Lepton_Error_Spectrum.pdf"))
    plt.close()
    print("[EXPORT] EWT_Lepton_Error_Spectrum.pdf")


# ------------------------------------------------------------------------
# MODULE 7: STRUCTURAL ROBUSTNESS OF STATUTORY DENSITY (N_nu_stat)
# ------------------------------------------------------------------------
def module7_statutory_robustness():
    bcc = derive_eps_M_from_BCC(8.0 * PI**4)
    alpha_geom = 1.0 / compute_alpha_geometric(bcc["eps_M"])
    r_nu = derive_neutrino_radius(
        alpha_geom, derive_planck_charge_from_e(alpha_geom, E_CHARGE_CODATA)
    )["r_nu"]
    lambda_l = derive_lambda_l_geometric(
        alpha_geom=alpha_geom,
        r_e=R_E,
        r_nu=r_nu,
        N_geom=bcc["N_geom"],
        L_p_geom=2.0 / SQRT3,
        K_WC=10,
    )

    N_nu_stat_base = (r_nu / (2.0 * lambda_l * EULER)) ** 3

    dr_ratio = np.linspace(-0.3, 0.3, 200)

    compliance_r_nu = ((r_nu * (1 + dr_ratio)) / (2.0 * lambda_l * EULER)) ** 3
    compliance_r_nu /= N_nu_stat_base

    compliance_r_emc = (r_nu / (2.0 * (lambda_l * (1 + dr_ratio)) * EULER)) ** 3
    compliance_r_emc /= N_nu_stat_base

    plt.figure()
    plt.plot(dr_ratio, np.ones_like(dr_ratio) * 1.3, 'r:', linewidth=1)
    plt.plot(dr_ratio, np.ones_like(dr_ratio) * 0.7, 'r:', linewidth=1)
    plt.plot(dr_ratio, np.ones_like(dr_ratio), 'k--', linewidth=2)
    plt.plot(dr_ratio, compliance_r_nu, 'b-', linewidth=2)
    plt.plot(dr_ratio, compliance_r_emc, 'r-', linewidth=2)

    plt.xlim(-0.3, 0.3)
    plt.ylim(0.6, 1.5)
    plt.xticks([-0.3, -0.15, 0, 0.15, 0.3],
               ['-30%', '-15%', '0%', '15%', '30%'])
    plt.xlabel("Relative Lattice Fluctuation (dr/r)")
    plt.ylabel("Normalized N_stat Stability Response")
    plt.title("Structural Robustness of Statutory Density N_nu_stat")
    plt.legend(["Upper Bound (1.3)", "Lower Bound (0.7)", "Statutory Lock-in (1.0)",
                "Fluctuation by r_nu", "Fluctuation by lambda_l"],
               loc="upper left")
    plt.grid(True, which='both', ls='-', alpha=0.5)
    plt.tight_layout()
    plt.savefig(pdf_path("EWT_Robustness_Analysis_DataDriven.pdf"))
    plt.close()
    print("[EXPORT] EWT_Robustness_Analysis_DataDriven.pdf")


# ------------------------------------------------------------------------
# MODULE 8: STIFFNESS-VOLUME STABILITY (FINAL PRECISION VERSION)
# ------------------------------------------------------------------------
def module8_stiffness_equilibrium():
    # Hierarchical gap computed from gravity_sector (not hardcoded)
    bcc = derive_eps_M_from_BCC(8.0 * PI**4)
    N_geom = bcc["N_geom"]
    alpha_geom = 1.0 / compute_alpha_geometric(bcc["eps_M"])
    r_nu = derive_neutrino_radius(
        alpha_geom, derive_planck_charge_from_e(alpha_geom, E_CHARGE_CODATA)
    )["r_nu"]
    lambda_l = derive_lambda_l_geometric(
        alpha_geom=alpha_geom,
        r_e=R_E,
        r_nu=r_nu,
        N_geom=N_geom,
```

```
        L_p_geom=2.0 / SQRT3,
        K_WC=10,
    )
    res_gravity = gravity_sector(
        alpha_geom=alpha_geom,
        r_nu=r_nu,
        N_geom=N_geom,
        L_p_geom=2.0 / SQRT3,
        K_WC=10,
        lambda_l=lambda_l,
        r_e=R_E,
        m_e=M_E,
        c0=C0,
    )
    N_nu_stat = res_gravity["N_nu_statutory"]
    N_nu_eff = res_gravity["N_nu_eff"]
    ratio_stat = N_nu_stat / N_nu_eff

    stiffness_exponent = -1.0 / 6.0

    dr_range = np.linspace(-0.25, 0.25, 200)
    N_nu_norm = (1 + dr_range) ** 3
    N_req_norm = (1 + dr_range) ** (3 * stiffness_exponent)

    plt.figure(figsize=(9, 7))
    plt.plot(dr_range, N_nu_norm, "b-", linewidth=3)
    plt.plot(dr_range, N_req_norm, "r-", linewidth=3)
    plt.plot(0, 1.0, "ko", markersize=12, markeredgewidth=2)

    plt.xlim(-0.25, 0.25)
    plt.ylim(0.4, 2.0)
    plt.xlabel("Relative Radius Fluctuation (dr/r)")
    plt.ylabel("Normalized Response (Value / N_eff)")
    plt.title("Gravity Stability: Nodal Stiffness vs. Soliton Volume")
    plt.legend(["Soliton Vol. Response (r^3)",
                "Nodal Stiffness (r^-0.5 from N_eff^-1/6)",
                "Effective Lock-in Point"], loc="upper center")
    plt.grid(True, which='both', ls='-', alpha=0.5)
    plt.tight_layout()
    plt.savefig(pdf_path("EWT_Stiffness_Equilibrium.pdf"))
    plt.close()
    print("[EXPORT] EWT_Stiffness_Equilibrium.pdf")
    print(f"       [HIERARCHY] Stat/Eff Density Gap: {ratio_stat:.2e}")


# ------------------------------------------------------------------------
# MODULE 9: LEPTODYNAMICS STABILITY & AMM ROBUSTNESS ANALYSIS
# ------------------------------------------------------------------------
def module9_amm_robustness():
    # Real sensitivity of a_mu and a_tau to eps_M perturbation
    # (computed from the model, not synthetic linear placeholders)
    bcc = derive_eps_M_from_BCC(8.0 * PI**4)
    eps_M = bcc["eps_M"]

    # Baseline predictions at the model eps_M
    alpha_geom_base = 1.0 / compute_alpha_geometric(eps_M)
    amm_base = compute_lepton_amms(alpha_geom_base, eps_M)
    a_mu_base = amm_base["a_mu_ppm"]
    a_tau_base = amm_base["a_tau_ppm"]

    dr_range = np.linspace(-0.05, 0.05, 100)
    a_mu_norm = []
    a_tau_norm = []

    for dr in dr_range:
        eps_M_pert = eps_M * (1.0 + dr)
        alpha_geom_pert = 1.0 / compute_alpha_geometric(eps_M_pert)
        amm_pert = compute_lepton_amms(alpha_geom_pert, eps_M_pert)
        a_mu_norm.append(amm_pert["a_mu_ppm"] / a_mu_base)
        a_tau_norm.append(amm_pert["a_tau_ppm"] / a_tau_base)

    a_mu_norm = np.array(a_mu_norm)
    a_tau_norm = np.array(a_tau_norm)
```

```
    plt.figure()
    plt.plot(dr_range * 100, a_mu_norm, "g-", linewidth=2)
    plt.plot(dr_range * 100, a_tau_norm, "m-", linewidth=2)
    plt.plot(0, 1.0, "ro", markersize=10)

    plt.xlabel("eps_M fluctuation (d(eps_M)/eps_M, %)")
    plt.ylabel("Normalized AMM Response (a_i / a_i^base)")
    plt.title("Real Sensitivity: Lepton AMM vs eps_M Perturbation")
    plt.legend(["Muon AMM (2D planar shell, O_mu-suppressed)",
                "Tau AMM (3D volumetric shell, O_tau = 1)",
                "Baseline at N_geom"], loc="upper left")
    plt.grid(True, which='both', ls='-', alpha=0.5)
    plt.savefig(pdf_path("EWT_AMM_Robustness_Leptons.pdf"))
    plt.close()
    print("[EXPORT] EWT_AMM_Robustness_Leptons.pdf")


# ------------------------------------------------------------------------
# MODULE 10: WEINBERG & CABIBBO (Shift-Normalized)
# ------------------------------------------------------------------------
def module10_weinberg_cabibbo():
    bcc = derive_eps_M_from_BCC(8.0 * PI**4)
    N_geom = bcc["N_geom"]

    # Physical constants
    M_Z_exp = 91.1876
    sin2W_target = 0.23122

    eps_M_final = 1.0 / (N_geom * PI**3)
    C_local_final = eps_M_final / (2.0 * math.sqrt(2.0))
    C_gap_final = 1.0 + (PI**6) * C_local_final

    M_W_Ideal = M_Z_exp * math.sqrt((1.0 - sin2W_target) * C_gap_final)
    sin2W_at_Nfinal = 1.0 - ((M_W_Ideal / M_Z_exp) ** 2) * (1.0 / C_gap_final)

    N_scan_W = np.linspace(778.5, 779.2, 1000)
    N_scan_W = np.unique(np.concatenate([N_scan_W, [N_geom]]))

    sin2W_results = []
    for n_v in N_scan_W:
        eps_m_local = 1.0 / (n_v * PI**3)
        C_local = eps_m_local / (2.0 * math.sqrt(2.0))
        C_gap = 1.0 + (PI**6) * C_local
        sin2W_results.append(1.0 - ((M_W_Ideal / M_Z_exp) ** 2) * (1.0 / C_gap))

    # Cabibbo using EWT spherical-mode quark masses
    m_d_ewt = mass_spherical(15)
    m_s_ewt = mass_spherical(28)

    sinC_results = []
    for n_v in N_scan_W:
        # eps_flavor(N) = 1/(8*pi^5*(1-zeta)) = 1/(pi*N)
        eps_flavor_local = 1.0 / (PI * n_v)
        sinC_results.append(
            math.sqrt(m_d_ewt / m_s_ewt) * (1.0 + eps_flavor_local)
        )

    eps_flavor_final = 1.0 / (PI * N_geom)
    sinC_final = math.sqrt(m_d_ewt / m_s_ewt) * (1.0 + eps_flavor_final)

    shift_C = sin2W_at_Nfinal - sinC_final
    sinC_shifted = np.array(sinC_results) + shift_C

    plt.figure(figsize=(9, 7))
    plt.plot(N_scan_W, sin2W_results, 'c-', linewidth=3)
    plt.plot(N_geom, sin2W_at_Nfinal, 'ro', markersize=10)
    plt.plot(N_scan_W, sinC_shifted, 'g-', linewidth=3)

    shift_label = f"sin(theta_C) + shift (shift = {shift_C:.10f})"
    plt.xlabel("N Coefficient")
    plt.ylabel("Angle Value (Shifted)")
    plt.title("Shift-Normalized Robustness: Weinberg & Cabibbo Angles")
```

```
    plt.legend(["sin^2(theta_W)", "N_geom Lock-in", shift_label,
                "Cabibbo Lock-in (shifted)"], loc="lower right")

    y_min = min(min(sin2W_results), min(sinC_shifted))
    y_max = max(max(sin2W_results), max(sinC_shifted))
    padding = (y_max - y_min) * 0.15
    plt.ylim(y_min - padding, y_max + padding)

    plt.grid(True, which='both', ls='-', alpha=0.5)
    plt.tight_layout()
    plt.savefig(pdf_path("EWT_Weinberg_Cabibbo_Robustness.pdf"))
    plt.close()
    print("[EXPORT] EWT_Weinberg_Cabibbo_Robustness.pdf")


# ----------------------------------------------------------------------
# Main
# ----------------------------------------------------------------------
def main():
    print("Enhanced EWT -- Robustness Plot Suite (Geometric)")

    module1_g_surface_transition()
    module2_alpha_sensitivity()
    module3_alpha_amm_phase()
    module4_unification_trajectory()
    module5_point_locked()
    module6_lepton_error_spectrum()
    module7_statutory_robustness()
    module8_stiffness_equilibrium()
    module9_amm_robustness()
    module10_weinberg_cabibbo()

    print("All figures exported.")


if __name__ == "__main__":
    main()
```

Listing 11: Robustness and stability plot generator.

A.3.12 Robustness and Stability Plots Output

The console output of `ewt_robustness_plots.py` is reproduced below.

```
Enhanced EWT -- Robustness Plot Suite (Geometric)
[EXPORT] EWT_Robustness_G_Surface_Transition.pdf
[EXPORT] EWT_Robustness_Alpha_Sensitivity.pdf
[EXPORT] EWT_Alpha_vs_AMM_PhasePlot.pdf
[EXPORT] EWT_Unification_Path_English.pdf
[EXPORT] EWT_POINT_LOCKED.pdf
[EXPORT] EWT_Lepton_Error_Spectrum.pdf
[EXPORT] EWT_Robustness_Analysis_DataDriven.pdf
[EXPORT] EWT_Stiffness_Equilibrium.pdf
       [HIERARCHY] Stat/Eff Density Gap: 5.28e+03
[EXPORT] EWT_AMM_Robustness_Leptons.pdf
[EXPORT] EWT_Weinberg_Cabibbo_Robustness.pdf
All figures exported.
```

Listing 12: Console output from `ewt_robustness_plots.py`.

Statements and Declarations

Funding: The author declares that no funds, grants, or other support were received during the preparation of this manuscript.

Competing Interests: The author has no relevant financial or non-financial interests to disclose.

Author Contributions: Ł. Smoliński is the sole author of this manuscript, responsible for the conceptualization, methodology, formal analysis, and writing.

Data and Code Availability: The complete repository for Project MagnetismGravity is openly accessible to support scientific reproducibility:

- **Datasets:** The EWT core dataset is available on Hugging Face: `https://huggingface.co/datasets/luksmol/EWT-Vacuum-Lattice-Unified-Physics/tree/main`.
- **Interactive Auditor:** The web-based EWT Auditor application is deployed at: `https://huggingface.co/spaces/luksmol/EWTAuditor`.
- **Version Control:** The active development source code is maintained on GitHub: `https://github.com/lsmolinski/MagnetismGravity`.

References

[1] Yee, J. (2020). The Relationship of the Speed of Light to Aether Density. Preprint, ResearchGate. DOI: 10.13140/RG.2.2.23961.98404.

[2] Wheeler, J. A. (1962). *Geometrodynamics*. Academic Press.

[3] Wheeler, J. A. (1957). On the Nature of Quantum Geometrodynamics. *Annals of Physics*, **2**(6), 604–614. DOI: 10.1016/0003-4916(57)90050-7.

[4] Wheeler, J. A. (1968). Superspace and the Nature of Quantum Geometrodynamics. In C. DeWitt & J. A. Wheeler (Eds.), *Battelle Rencontres: 1967 Lectures in Mathematics and Physics* (pp. 242–307). W. A. Benjamin.

[5] DeWitt, B. S. (1967). Quantum Theory of Gravity. I. The Canonical Theory. *Physical Review*, **160**(5), 1113–1148. DOI: 10.1103/PhysRev.160.1113.

[6] Misner, C. W., Thorne, K. S., & Wheeler, J. A. (1973). *Gravitation*. W. H. Freeman. ISBN: 978-0-7167-0344-0.

[7] Wheeler, J. A. (1964). Geometrodynamics and the Issue of the Final State. In C. DeWitt & B. DeWitt (Eds.), *Relativity, Groups and Topology*. Gordon and Breach.

[8] Wheeler, J. A. (1980). Pregeometry: Motivations and Prospects. In A. R. Marlow (Ed.), *Quantum Theory and Gravitation* (pp. 1–11). Academic Press.

[9] Wheeler, J. A. (1990). Information, Physics, Quantum: The Search for Links. In W. H. Zurek (Ed.), *Complexity, Entropy, and the Physics of Information* (pp. 3–28). Addison-Wesley.

[10] Yee, J. (2019). *The Relationship of the Fine Structure Constant and Pi* (Version 2) Preprint, ResearchGate. DOI: 10.13140/RG.2.2.30832.92162.

[11] Yee, J. and Gardi, L. (2019). The Geometry of Spacetime and the Unification of the Electromagnetic, Gravitational and Strong Forces. DOI: 10.13140/RG.2.2.23094.24642.

[12] Yee, J. i Gardi, L. (2020). The Physics of Subatomic Particles and their Behavior Modeled with Classical Laws. Preprint, ResearchGate. DOI: 10.13140/RG.2.2.27250.25283.

[13] Yee, J. i Gardi, L. (2020). Forces: Explained and Derived by Energy Wave Equations. URL:Forces Explained and Derived by Energy Wave Equations.

[14] Yee, J. (2020). The Geometry of Particle Standing Waves. DOI: 10.13140/RG.2.2.27401.88169.

[15] Yee, J. (2021). The Relation of Relativistic Energy to Particle Wavelength. Technical Report / Preprint. URL: 10.13140/RG.2.2.28388.889668.

[16] Smoliński, Ł. and Yee, J. (2025). Bridging Geometry and Measurement: The Geometric Correction Terms ϵ_M and ϵ_G in the Derivation of the Fine-Structure Constant (α) Within the Energy Wave Theory (EWT) Model. DOI: 10.5281/zenodo.17397260.

[17] Yee, J. and Smoliński, Ł. (2025). The Geometric Black Hole: The Role of ϵ_G in Extreme Wave Geometries. DOI: 10.5281/zenodo.17397981.

[18] Mohr, P. J., Newell, D. B., Taylor, B. N., and Tiesinga, E. (2022). *CODATA Recommended Values of the Fundamental Physical Constants: 2022*. Rev. Mod. Phys. **97**, 025002. DOI: 10.1103/RevModPhys.97.025002.

[19] Padmanabhan, T. (2010). Thermodynamics of Spacetime: The Einstein Equation and the Principle of Equipartition. *Journal of Physics: Conference Series*, **235**, 012015. DOI: 10.1088/1742-6596/235/1/012015.

[20] Verlinde, E. (2017). Emergent Gravity and the Dark Universe. *SciPost Physics*, **2**, 016. DOI: 10.21468/SciPostPhys.2.3.016.

[21] Visser, M. (2020). Sakharov's induced gravity: a modern perspective. *Annalen der Physik*, **532**, 2000287. DOI: 10.48550/arXiv.gr-qc/0204062.

[22] Fukuda, Y., et al. (Super-Kamiokande Collaboration). (1998). Evidence for Oscillation of Atmospheric Neutrinos. *Physical Review Letters*, **81**, 1562. DOI: 10.1103/PhysRevLett.81.1562.

[23] Ahmad, Q. R., et al. (SNO Collaboration). (2002). Direct Evidence for Neutrino Flavor Transformation from Neutral-Current Interactions in the Sudbury Neutrino Observatory. *Physical Review Letters*, **89**, 011301. DOI: 10.1103/PhysRevLett.89.011301.

[24] Beda, A. G., et al. (GEMMA Collaboration). (2012). Results of Search for the Neutrino Magnetic Moment in GEMMA Experiment. *Physics of Particles and Nuclei Letters*, **9**, 143–149. DOI: 10.1155/2012/350150.

[25] Studenikin, A. I. (2017). Neutrino Magnetic Moment: A Review. *Modern Physics Letters A*, **32**, 1730001. DOI: 10.1142/S0217732305017482.

[26] Yee, J. (2019). The Geometry of Particles and the Explanation of their Creation and Decay. ResearchGate Preprint. DOI: 10.13140/RG.2.2.14966.14401.

[27] Yee, J., Zhu, Y., Zhou, G. (2017). Atomic Orbitals: Explained with Classical Mechanics. Vixra.org. URL: 10.13140/RG.2.2.19156.88963.

[28] Yee, J., Zhu, Y., & Zhou, G. F. (2017). The Relation of Particle Sequence to Atomic Sequence. *Journal of Physical Mathematics*, **8**, 247. DOI: 10.4172/2090-0902.1000247.

[29] Yee, J., Gardi, L. (2019). The Relationship of Mass and Charge. Technical Paper. URL: 10.13140/RG.2.2.12645.45289.

[30] Sakharov, A. D. (1968). Vacuum Quantum Fluctuations in Curved Space and the Theory of Gravitation. *Doklady Akademii Nauk SSSR*, **177**, 70–71.

[31] Jacobson, T. (1995). Thermodynamics of Spacetime: The Einstein Equation of State. *Physical Review Letters*, **75**, 1260–1263. DOI: 10.1103/PhysRevLett.75.1260.

[32] Verlinde, E. P. (2011). On the Origin of Gravity and the Laws of Newton. *Journal of High Energy Physics*, **04**, 029. DOI: 10.1007/JHEP04(2011)029.

[33] Padmanabhan, T. (2004). Gravity and the Thermodynamics of Horizons. *General Relativity and Gravitation*, **36**, 2271–2276. DOI: 10.48550/arXiv.gr-qc/0311036.

[34] Caligiuri, L. M., & Sorli, A. (2014). Gravity Originates from Variable Energy Density of Quantum Vacuum. *American Journal of Modern Physics*, **3**(3), 118–128. DOI: 10.11648/j.ajmp.20140303.11.

[35] Daywitt, W. C. (2015). The Trouble with the Equations of Modern Fundamental Physics. *American Journal of Modern Physics*, **5**(1-1), 23–32. DOI: 10.11648/j.ajmp.s.2016050101.14 [DOI is broken]. URL: https://www.sciencepublishinggroup.com/article/10.11648.j.ajmp.s.2016050101.14

[36] Schwinger, J. (1948). On Quantum-Electrodynamics and the Magnetic Moment of the Electron. *Physical Review*, **73**, 416–417. DOI: 10.1103/PhysRev.73.416.

[37] Yee, J. (2014). Explaining the Mass of Leptons with Wave Structure Matter. Technical Paper, Third Draft (Corrected May 18, 2021). URL: https://www.rxiv.org/pdf/1302.0147v4.pdf.

[38] Yee, J., Zhu, Y., and Zhou, G.F. (2024). The Relation of Particle Sequence to Atomic Sequence. URL: https://doi.org/10.4172/2090-0902.1000247.

[39] Aoyama, T., Kinoshita, T., Nio, M. (2018). Revised and improved value of the QED tenth-order electron anomalous magnetic moment. *Phys. Rev. D* **97**, 036001. DOI: 10.1103/PhysRevD.97.036001.

[40] Visser, M. (2002). Sakharov's induced gravity: a modern perspective. *Modern Physics Letters A*, **17**, 977–991. DOI: 10.1142/S0217732302006886.

[41] Eidelman, S., & Passera, M. (2007). Theory of the tau lepton anomalous magnetic moment. *Modern Physics Letters A*, **22**, 159. DOI: 10.1142/S0217732307022694.

[42] Yee, J. (2018). The Periodic Table of Subatomic Particles. URL: 10.13140/RG.2.2.33610.62401.

[43] Tiesinga, E., Mohr, P. J., Newell, D. B., and Taylor, B. N. (2021). CODATA recommended values of the fundamental physical constants: 2018. *Reviews of Modern Physics*, 93(2), 025010. URL: 10.1103/RevModPhys.93.025010.

[44] Navas, S., et al. (Particle Data Group). (2024). Review of Particle Physics. *Physical Review D*, 110, 030001. URL: 10.1103/PhysRevD.110.030001.

[45] Aoyama, T., et al. (2020). The anomalous magnetic moment of the muon in the Standard Model. *Physics Reports*, 887, 1-166. URL: 10.1016/j.physrep.2020.07.006.

[46] Yee, J. (2020). Fundamental Physical Constants: Explained and Derived by Energy Wave Equations (vB). URL: https://vixra.org/pdf/1606.0113vB.pdf.

[47] Han, S. on behalf of the LHCb collaboration. (2026). Search for the doubly charmed baryon Ξ_{cc}^{+} with the upgraded LHCb detector. *60th Rencontres de Moriond*, LHCb-PAPER-2026-009 (in preparation).

[48] E. Di Valentino, J. Levi Said, et al. (The CosmoVerse Network). *The CosmoVerse White Paper: Addressing observational tensions in cosmology with systematics and fundamental physics*. URL: https://arxiv.org/abs/2504.01669.

[49] ATLAS Collaboration. Observation of a B_c^{*+} meson with the ATLAS detector. URL: https://arxiv.org/abs/2605.16228.

[50] LHCb Collaboration (2026). *Observation of a new doubly charmed baryon* Ω_{cc}^{*}. CERN Seminar, 24 June 2026. URL: https://indico.cern.ch/event/1595459/.

[51] LaFrenière, G. (2000). The Michelson-Morley Experiment: A physical explanation of the null result. URL: https://mildred.github.io/glafreniere/sa_Michelson.htm.

[52] Lorentz, H. A. (1904). Electromagnetic phenomena in a system moving with any velocity smaller than that of light. *Proceedings of the Royal Netherlands Academy of Arts and Sciences*, **6**, 809–831. Reprint DOI: 10.1007/978-94-015-3445-1_5.

[53] FitzGerald, G. F. (1889). The Ether and the Earth's Atmosphere. *Science*, **13**(328), 390. Full text available at: Wikisource:The_Ether_and_the_Earth%27s_Atmosphere.

[54] OpenWave Collaboration (2026). *M4.11 — N as the Internal Lock-In Point: Drift, Resolution, Grouping*. OpenWave research repository. URL: `https://github.com/openwave-labs/openwave/blob/main/openwave/xperiments/m4_ewt/research/scripts/m4_11_lock_in_n_scan.py`.

[55] OpenWave Collaboration (2026). *M4.3–M4.9 — Local Metric Phenomena and Emergent Encoding Validation Suite*. OpenWave research repository. URL: `https://github.com/openwave-labs/openwave/tree/main/openwave/xperiments/m4_ewt/research/scripts`.

[56] OpenWave Collaboration (2026). *M4.10 and M4.12 — Newtonian Force from EMC Field Overlap*. OpenWave research repository. URL: `https://github.com/openwave-labs/openwave/tree/main/openwave/xperiments/m4_ewt/research/scripts`.

[57] M. Thomson, *Model-Independent Measurement of the $e^+e^- \to HZ$ Cross Section at a Future e^+e^- Linear Collider using Hadronic Z Decays*, arXiv:1509.02853 [hep-ex] (2015). Available at: https://arxiv.org/abs/1509.02853.

[58] K. Fujii *et al.*, *Physics Case for the 250 GeV Stage of the International Linear Collider*, arXiv:1710.07621 [hep-ex] (2017). Available at: https://arxiv.org/abs/1710.07621.

[59] Z. Chen, Y. Yang, M. Ruan, D. Wang, G. Li, S. Jin, and Y. Ban, *Cross Section and Higgs Mass Measurement with Higgsstrahlung at the CEPC*, arXiv:1601.05352 [hep-ex] (2016). Available at: https://arxiv.org/abs/1601.05352.

[60] Snowmass 2021 Higgs Physics Topical Group, *Report of the Topical Group on Higgs Physics for Snowmass 2021: The Case for Precision Higgs Physics*, arXiv:2209.07510 [hep-ph] (2022). Available at: https://arxiv.org/abs/2209.07510.

[61] N. Craig, J. Gu, Z. Liu, and K. Wang, *Beyond Higgs Couplings: Probing the Higgs with Angular Observables at Future e^+e^- Colliders*, *Journal of High Energy Physics* **03** (2016) 050. doi:10.1007/JHEP03(2016)050.

[62] Z. Liu, *Probing the Higgs with Angular Observables at Future e^+e^- Colliders*, *International Journal of Modern Physics A* **31**, 1644005 (2016). doi:10.1142/S0217751X1644005X.

[63] S.-F. Ge, H.-J. He, and R.-Q. Xiao, *Probing New Physics Scales from Higgs and Electroweak Observables at e^+e^- Higgs Factory*, *Journal of High Energy Physics* **10** (2016) 007. doi:10.1007/JHEP10(2016)007.

[64] R. Gatto, G. Sartori, and M. Tonin, *Weak self-masses, Cabibbo angle, and broken $SU(2) \times SU(2)$*, Physics Letters B **28**, 128–130 (1968). DOI: 10.1016/0370-2693(68)90150-0.

[65] H. Kleinert, *Gravity as a Theory of Defects in a Crystal with Only Second Gradient Elasticity*, Annalen der Physik **499**, 117–119 (1987). DOI: 10.1002/andp.19874990206.

[66] N. Manton and P. Sutcliffe, *Topological Solitons*, Cambridge University Press, Cambridge (2004). DOI: 10.1017/CBO9780511617034.

[67] Planck Collaboration, Aghanim, N., Akrami, Y., et al. (2020). *Planck 2018 results. VI. Cosmological parameters.* Astronomy & Astrophysics, **641**, A6. DOI: 10.1051/0004-6361/201833910.

[68] DESI Collaboration, Adame, A. G., Aguilar, J., et al. (2024). *DESI 2024 VI: Cosmological Constraints from the Measurements of Baryon Acoustic Oscillations.* The Astronomical Journal, **168**, 58. DOI: 10.3847/1538-3881/ad0b08.

[69] Riess, A. G., Yuan, W., Macri, L. M., et al. (2022). *A Comprehensive Measurement of the Local Value of the Hubble Constant with 1 $km\,s^{-1}\,Mpc^{-1}$ Uncertainty from the Hubble Space Telescope and the SH0ES Team.* The Astrophysical Journal Letters, **934**, L7. DOI: 10.3847/2041-8213/ac5c5b.

[70] Fixsen, D. J. (2009). *The Temperature of the Cosmic Microwave Background.* The Astrophysical Journal, **707**, 916–920. DOI: 10.1088/0004-637X/707/2/916.

[71] Haug, E. G. (2022). *Progress in the Composite View of the Newton Gravitational Constant and Its Link to the Planck Scale.* Universe, **8**(9), 454. DOI: 10.3390/universe8090454.

[72] Haug, E. G., & Tatum, E. T. (2024). *The Hawking Hubble Temperature as a Minimum Temperature, the Planck Temperature as a Maximum Temperature and the CMB Temperature as Their Geometric Mean Temperature.* Journal of Applied Mathematics and Physics, **12**, 3328–3340. DOI: 10.4236/jamp.2024.1210198.

[73] Haug, E. G. (2020). *Collision Space-Time: Unified Quantum Gravity.* Physics Essays, **33**(1), 46–78. DOI: 10.4006/0836-1398-33.1.46.

[74] Haug, E. G. (2023). *Not Relying on the Newton Gravitational Constant Gives More Accurate Gravitational Predictions.* Journal of Applied Mathematics and Physics, **11**, 3124–3158. DOI: 10.4236/jamp.2023.1110205.

[75] Haug, E. G. (2024). *CMB, Hawking, Planck, and Hubble Scale Relations Consistent with Recent Quantization of General Relativity Theory.* International Journal of Theoretical Physics, **63**, 57. DOI: 10.1007/s10773-024-05570-6.

[76] Haug, E. G. (2026). *Thermodynamic Cosmology from the CMB Temperature and the Planck Temperature and the Relation to Quantum Cosmology.* Astronomy, **5**(4), 19. DOI: 10.3390/astronomy5040019.

[77] Haug, E. G. (2020). *Finding the Planck Length Multiplied by the Speed of Light Without Any Knowledge of G, c, or ħ, Using a Newton Force Spring.* Journal of Physics Communications, **4**(7), 075001. DOI: 10.1088/2399-6528/ab9dd7.

Final Remark

The history of physics teaches us that the path to unification often lies within the most fundamental constituents of nature. Just as the quest to fully understand the electron once held the promise of unraveling quantum electrodynamics, the evidence presented in this geometric, unified model points to a more profound truth. By demonstrating that the neutrino's inherent wave geometry is the common source of the gravitational constant **G**, the fine-structure constant α, and the anomalous magnetic moments, the picture is also completed by the discovery of the Weinberg and Cabibbo angles as the structural resonance limits of the vacuum lattice. The conclusion is inescapable: **It would have been enough to understand the neutrino's geometric anchor.**