

The Geometric Identity of Gravity and Dimensional Unification Resolving α , Lepton $(g - 2)_l$, Weinberg, and Cabibbo Mixing

Łukasz Smoliński¹

¹Independent Researcher, 61-160 Czapury, Poland

September 26, 2026

Version: 5.1.0

Abstract

A unified geometric wave model is proposed, deriving the Gravitational Constant (G), the Fine-Structure Constant (α), and lepton anomalous magnetic moments (a_l) from a single structural modulator: the geometric vacuum stiffness ϵ_M . G is established as a precise, $\hbar$ -independent identity rooted in the soliton's wave geometry. The derivation uses the ideal BCC projection $L_p^{\text{geom}} = 2/\sqrt{3}$; the remaining deviation from the measured value reflects lattice packing impedance, not a free parameter. The same ϵ_M factor governs a recursive nodal integration that predicts the electron anomalous magnetic moment from pure geometry and yields the full muon and tau anomalous magnetic moments via a unified dimensional projection over the BCC lattice, without perturbative QED calibrations. The framework posits that the observed invariance of G is a low-field artefact of a dynamic geometric equilibrium, predicting testable deviations ($G_{\text{eff}} \neq G$) in environments with strong local magnetic moments where this equilibrium is exceeded. Crucially, the derivation of α and the anomalous magnetic moments (a_l) are shown to be independent of mass and charge, revealing that these constants are intrinsic topological properties of the vacuum lattice rather than secondary products of particle-field interactions. This geometric mapping reveals a dimensional hierarchy (π^5, π^6), resolving the Weinberg angle and Cabibbo mixing as emergent consequences of the vacuum's surface and volumetric resonance modes. Consequently, the divergence of fundamental forces is identified as a dimensional artefact of the unified BCC lattice, establishing a structural foundation for cosmological scaling.

Keywords: Anomalous Magnetic Moments, Fine-Structure Constant, Gravitational Constant, Weinberg angle, Cabibbo angle, BCC lattice, Geometric Origin, Unification Theory.

Contents

Change Log	7
1 Geometric Hierarchy of EWT and Spacetime Elasticity	8
1.1 Wheeler's Vision of Quantum Vacuum and the Path to EWT	8
1.2 The Elastic Medium Constituent (EMC)	9

31	1.3	The Wave Center (WC)	10
32	1.4	The Soliton	10
33	1.5	Elastic Interaction (Hooke's Law)	10
34	1.6	From Energy Domain to Geometric Domain	11
35	2	Geometric Equation of the Fine-Structure Constant and the Deficit Terms	12
36	2.1	BCC Lattice Geometry and the Geometric Ladder	13
37	2.2	Geometric Magnetic Deficit ϵ_M	14
38	2.3	The Fundamental Lattice Response Parameter ϵ_M	15
39	3	Dimensional Hierarchy and Dynamic Resonant Modulations	15
40	3.1	The Universal Geometric Modulators: Substrate Properties	15
41	3.1.1	Unified Local Constant $\mathbf{C}_{\text{local}}$	15
42	3.1.2	The π^6 Resonance: Volumetric Bosonic Coupling	16
43	3.1.3	The π^7 Resonance: Charge-Induced Amplitude Loading	16
44	3.1.4	The π^5 Resonance: Surface Interaction Modulator	16
45	3.2	Geometric Mixing Predictions: Higgs and Cabibbo Sectors	16
46	3.2.1	Higgs-Vector Boson Mixing	16
47	3.2.2	Cabibbo Mixing Angle	17
48	3.2.3	Dimensional Budget of Resonances: Reflection vs. Mixing	17
49	3.2.4	Symmetry of Observables: Energy Density vs. Amplitude Interference . .	18
50	3.2.5	Interpretation of Quark Masses in the EWT Framework	18
51	3.3	Structural Transition to N -Stiffness: From Geometric Volume to Lattice Impedance	19
52	4	Analysis of Neutrino Boundary Conditions	20
53	4.1	Standard Conditions: Mass and Minimal Magnetism	20
54	4.2	Extreme Conditions: Wave Center in a Geometric Black Hole (GBH)	20
55	5	Neutrino as the Geometric Anchor	21
56	5.1	Planck Charge from Geometric Alpha	21
57	5.2	Decomposition of the Scaling Factor $S = r_\nu/q_P$	22
58	5.3	Geometric Fixed Point of g_v	22
59	5.4	Resulting Neutrino Radius	23
60	5.5	Decadic Resonance Test: $r_e/r_\nu \approx 100$	23
61	5.6	Physical Interpretation: ϵ_M as a Global Lattice Parameter	23
62	5.7	Topological Necessity of r_ν	23
63	6	Planck Length and Reduced Planck Constant from Geometry	24
64	6.1	Reduced Planck Constant from Geometric Alpha	24
65	7	The Physical Origin of Charge as Wave Amplitude	24
66	7.1	Charge as Spatial Wave Amplitude	25
67	7.2	Conversion of the Coulomb and Distinction from CGS	25
68	7.3	Consequence for the Planck Charge q_P	25
69	8	Geometric Deficit and Density Hierarchy of the Neutrino	25
70	8.1	Unified Geometric Coupling	26

71	9 Speed of Light as the Metric of Spacetime	26
72	9.1 The Causal Structure of the BCC Lattice	27
73	9.2 Axiom: The Maximum Propagation Speed	27
74	9.3 Consequences: Density-Dependent Propagation Speed	28
75	10 The Physical Origin of Relativity: Wave-Mechanical Lorentz Contraction	29
76	10.1 Doppler-Induced Compression of the Standing-Wave Soliton	29
77	10.2 The Michelson–Morley Null Result as a Mechanical Necessity	30
78	11 The Geometric Identity of G: Derivation and Consistency	30
79	11.1 Derivation of G from Electron Scaling (The $\hbar$ -Independent Base)	30
80	11.2 The Gravitational Identity Flow: From Electromagnetic Base to Surface Transition	32
81	11.3 Formalization of Soliton Geometric Density $\rho(r)$	33
82	11.4 Numerical Verification and the Unitary Mapping Operator $\mathcal{U}$	33
83	11.5 Formal Definition of the Operator $\mathcal{U}$: Mapping Geometric Parameters to G . . .	34
84	11.6 Geometric Emergence of the Planck Length	35
85	11.7 Consistency with the Planck Charge	37
86	11.8 Electromagnetic Consistency Check	38
87	11.9 Status of Inputs	38
88	11.10 Duality of Soliton Density	38
89	11.11 The Degraded EMC Wall: Spherical Masking and Isotropy	39
90	11.11.1 Open question: Wall density relative to statutory background	39
91	12 Local Metric Phenomena from EMC Lattice Deformation	40
92	12.1 The Two Faces of EMC Displacement: Speed and Trajectory	40
93	12.2 Bridging the Vector Displacement to the Scalar Refractive Index	40
94	12.3 Asymptotic Continuous Limit and Schwarzschild Equivalence	41
95	12.4 Mechanical Origin of Gravitational Redshift in the EMC Lattice	41
96	12.5 Shapiro Delay from the EMC Refractive Index	42
97	12.6 Macroscopic EMC Density Profile of a Star	43
98	12.7 Derivation of the Far-Field EMC Density Profile	44
99	12.8 From Microscopic EMC Displacement to the Gravitational Radius	45
100	12.9 Emergent Metric Encodings from Lattice Dynamics	46
101	12.10 Summary of Local Metric Observables	47
102	12.11 In-Platform Numerical Validation	47
103	13 Newtonian Force from Interacting EMC Deficits	48
104	13.1 In-Platform Validation: Normalization Consistency	49
105	14 Relation to Existing Emergent Gravity Frameworks	50
106	14.1 Comparison of Mechanisms and Scales	50
107	14.1.1 Microscopic Degrees of Freedom (DoF)	50
108	14.1.2 Source of Gravitational Emergence	51
109	14.1.3 Scaling Factor for $\mathbf{G}$	51
110	14.1.4 Evolution of the EWT Framework: From Wave Attenuation to EMC Push	
111	out	52
112	14.1.5 EWT as Pressure-Driven Emergent Gravity	53

113	15 Geometric Validation: The Fundamental Identity and the Base AMM State	
114	(a_e)	54
115	15.1 Geometric Mass-to-Radius Identity ($E \propto r^5$)	54
116	15.2 Physical Origin of the r^5 Scaling: Geometric Energy Density	54
117	15.3 The Geometric Status of Mass: Kilogram as a Derived Unit	55
118	15.4 Geometric Dimension of the Reduced Planck and G Constants	56
119	15.5 The Fundamental Magnetic Deficit ϵ_M	57
120	15.6 Derivation of the a_e Base Formula	57
121	16 The Recursive Lepton Hierarchy: Nodal Shell Resonance Model	57
122	16.1 Resonance Growth Law and Fibonacci Stability Metrics	58
123	16.2 Structural Stability Invariants of the BCC Lattice	58
124	16.3 Structural Analysis: The Onion Model	59
125	16.4 The Recursive Master Equation: Nodal Latching via Geometric Coupling	59
126	16.4.1 Correspondence between Mass Scaling and Shell Geometry	62
127	16.5 Final Results and Experimental Correlation	63
128	16.6 Natural Emergence of Three Lepton Generations	65
129	16.7 Predictions for Lepton Properties: The First-Principles AMM Predictions	66
130	16.8 Physical Interpretation: The Mechanism of Topological Nodal Inclusion	66
131	17 Sensitivity Analysis: Verification of Computational Variants	68
132	17.1 Numerical Convergence and Error Landscape	68
133	17.2 EWT Standalone vs. Best Resonance Peak	69
134	17.3 3D Stability Mapping and Topography	70
135	17.4 Verification of Geometric Scaling	72
136	17.5 Critique of the Standard Model Target Bias: The Circularity of QED Precision	72
137	18 The Magneto-Geometric Hypotheses: Dynamic Deficit Modification	73
138	18.1 Hypothesis 1: The Push-Out Mechanism	73
139	18.2 Hypothesis 2: The Consolidation Mechanism	73
140	18.3 Hypothesis 3: No Magnetic Influence	73
141	18.4 Summary of Hypotheses Testing	74
142	18.5 Dynamic Equilibrium and Geometric Compensation	74
143	19 Numerical Verification of EWT Mass Scaling and Structural Sequences	74
144	19.1 EWT particle mass prediction	74
145	19.2 Implementation and Computational Results	75
146	19.3 Universal $(K_{WC})^5$ Meson-Mode Scan: Independent Mass Validation	75
147	19.3.1 Key conclusions from the meson-mode scan	76
148	19.4 Topological Isomorphism and Geometric Interference in BCC Lattice	77
149	19.4.1 Dual Resonance Modes: Orbital vs. Meson Scaling	77
150	19.4.2 Validation via the Ξ_{cc} Isospin Doublet	78
151	19.4.3 New vector state B_c^{*+} from ATLAS 2026.	78
152	19.4.4 New doubly charmed state Ω_{cc}^* from CERN 2026	78
153	19.4.5 Topological Isomorphism and the Dual Realisation of Mass Eigenvalues	78
154	19.4.6 Discussion and Cross-Mode Consistency	79

155	20 Geometric Radius Predictions: From Vacuum Anchor to Heavy Bosons	80
156	20.1 Numerical Foundation: Spherical Mode Accuracy	80
157	20.2 Predictive Radius for Heavy Bosons	80
158	20.3 Cross-Scale Validation with Nuclear Equivalents	81
159	20.4 Conclusion on Geometric Consistency	81
160	21 The Geometric Identity of Stiffness: Linking N_{geom} to the Dimensional Ladder	81
161	21.1 Eliminating the Fine-Tuning Paradigm	82
162	21.2 The Topological Reduction of the Fundamental Lattice Response Parameter: $\epsilon_M =$	
163	$\frac{1}{8\pi^7(1-\zeta)}$	82
164	21.2.1 Physical Interpretation of the $8\pi^7$ Attractor	83
165	21.2.2 The Zero-Calibration Fine-Structure Constant	83
166	22 The Geometric Unification of Lepton Properties	83
167	22.1 The Unified Identity of the Lepton	84
168	22.2 Geometric Determinism of the Electron AMM	84
169	23 Other Geometric Constants Derived from the Lattice	84
170	23.1 The Rydberg Constant (R_∞): An Atomic Resonance Gap	84
171	23.1.1 Geometric Derivation: The $\alpha^3/(4\pi r_e)$ Identity	85
172	23.1.2 Zero-Calibration Identity	85
173	23.1.3 Numerical Verification	85
174	23.1.4 Physical Interpretation	85
175	23.2 The Bohr Radius (a_0): The Atomic Scale	86
176	23.3 The Electron Compton Wavelength (λ_C): Threshold of Annihilation	86
177	23.4 Consistency Across Atomic Scales	87
178	24 Mathematical Foundations of the Energy Wave Theory Lattice	87
179	24.1 Introduction	87
180	24.2 The Space Group of the Vacuum Lattice: $Im\bar{3}m \times SO(3)_{\text{local}}$	87
181	24.2.1 Global Symmetry: The BCC Space Group $Im\bar{3}m$ (No. 229)	88
182	24.2.2 Local Symmetry: The Spherical Constituent $SO(3)_{\text{local}}$	88
183	24.3 The Topological Class of Solitons: Winding Numbers and Cohomology	88
184	24.3.1 The Electron Soliton as a Winding Number on S^2	88
185	24.3.2 Higher-Generation Leptons: Topology of Nested Shells	88
186	24.4 The Dimensional Weight Operator and Degree-of-Freedom Algebra	89
187	24.5 Synthesis: The Mathematical Unity of EWT	91
188	24.6 Formalism of the Vacuum Push-out Mechanism	91
189	24.6.1 The Vacuum Stiffness Tensor and Sakharov Isomorphism	91
190	24.6.2 The Impedance Mismatch Operator	92
191	24.7 Isotropy and Spherical Masking: From Asymmetric Core to Isotropic Gravity . .	92
192	24.7.1 Geometric Necessity of Asymmetry	92
193	24.7.2 Intrinsic Sphericity of Standing Waves	92
194	24.7.3 Geometric Low-Pass Filter	92
195	24.7.4 Minimization of Lattice Shear Stress	93
196	24.7.5 Domain Separation: r^5 vs. r^3	93
197	24.7.6 The Principle of Geometric Masking	93
198	24.7.7 Gravity as a Result of Construction Material	94
199	24.7.8 Universal Applicability: From Leptons to Hadrons	94

200	24.8 The Uniqueness of the BCC Topology: Why $N_{\text{geometric}} = 8\pi^4$ Excludes Other	
201	Lattices	94
202	24.8.1 The Coordination Number and Stiffness Distribution	94
203	24.8.2 Packing Fraction and the ζ Correction	95
204	24.8.3 Topological Consistency: The $2\pi^2$ Factor and BCC Axes	95
205	24.9 Mechanical Resolution of Field Paradoxes	95
206	25 Nonlinear Stabilisation of the Electron Soliton	96
207	25.1 General Soliton Wave Equation	96
208	25.2 Coupling Coefficient from BCC Geometry	96
209	25.3 Proposed Forms of the Nonlinear Term	97
210	25.4 Structural Emergence of the 1-3-6 Geometry	98
211	25.5 Topological Selection Rule and the Electron Ground State	98
212	25.6 Connection to the Dimensional Weight Operator	99
213	25.7 Geometric Estimate of the Degraded EMC Wall Radius	99
214	25.8 Relation Between the Energetic and Topological Arguments	101
215	25.9 Summary	101
216	26 Predictive Power	102
217	26.1 Prediction of the Fundamental Value of G	102
218	26.2 Predictions for Lepton Properties: The First-Principles AMM Predictions	102
219	26.3 Magnetic Sensitivity $G(B)$ and Hypotheses Testing	102
220	26.4 AMM as a Strong Test of Geometric Universality	103
221	26.5 Correlations with Neutrino Flux	103
222	26.6 Spectral and Geometric Modal Tests	103
223	26.7 Tests using Gravitational Wave Observations	103
224	26.8 Test on Bose-Einstein Condensates	103
225	26.9 The Higgs Soliton vs. Fundamental Scalar Field	104
226	27 Falsifiability and Model Constraints	104
227	27.1 Practical Considerations and Detection Limits	104
228	27.2 Mutual Falsification: The Ultimate Test	105
229	28 Robustness Analysis: Geometric Stability and Sensitivity	105
230	28.1 Robustness Analysis of the Neutrino Soliton Radius and Geometric Identity	105
231	28.2 Gravitational Surface Transition and $N_{\nu,\text{eff}}$	106
232	28.3 Sensitivity of α^{-1} to the Geometric Modulator N_{geom}	107
233	28.4 Phase Space Analysis: EWT Trajectory vs Standard Model Interpretation	108
234	28.5 Parametric Unification: The Stability Path of G and α	109
235	28.6 EWT Unification: Convergence of G and a_e on the Vacuum Stiffness Isentrope	109
236	28.7 Robustness Analysis: Geometric Stability and Vacuum Elasticity	110
237	28.8 Structural Robustness and Resonance Stability of the Lepton Hierarchy	110
238	28.9 Electroweak and CKM Coincidence: Geometric Robustness of $\sin^2 \theta_W$ and $\sin \theta_C$	111
239	28.10 Lock-In Test: Independent Determination of N_{geom}	111
240	28.10.1 Method	112
241	28.10.2 Results	113
242	28.10.3 PRIMARY sectors: G , a_τ , and $\sin \theta_C$	114
243	28.10.4 WEAK sectors: a_e , a_μ , and $\sin^2 \theta_W$	115
244	28.10.5 NO CONSTRAINT: α^{-1}	115
245	28.10.6 Interpretation	115

246	29 Geometric Phase Transitions: From Neutrino to Black Hole with EMC Wall	116
247	29.1 Hierarchy of Vacuum States and the G-Operating Point	116
248	29.2 The Erasure of "Dark" Placeholders	118
249	29.3 Resolution of Historically "Unsolvable" Paradoxes	119
250	29.4 Density Duality: From Local Refraction to Cosmological Redshift	120
251	30 Outlook and Computational Tools	121
252	31 Conclusion: The Geometric Paradigm, Correlated Lepton Anomalies, and	
253	Emergent Gravity	122
254	31.1 Geometric Unification in a Nutshell: The ϵ_M Flowchart	123
255	31.2 The Unitary Geometric Path	123
256	31.3 The Geometry of the Constituent: From Cubic Grids to Spherical Units	123
257	31.4 Pillar 1: The N_ν Deficit and Soliton Stability	124
258	31.5 Pillar 2: The Dynamic Scaling Operator Ω_{Push}	125
259	31.6 Pillar 3: The Recursive Paradigm — Hierarchical Nodal Integration	125
260	31.7 The Soliton Radius and Neutrino Interaction Cross-Section	125
261	31.8 Paradigm Shift: The Planck Length as a Physical Boundary	126
262	31.9 The ϵ_M Factor: Universal Geometric Modulator	126
263	31.10 Geometric Self-Consistency of the Planck–Gravity–Metric Triangle	126
264	31.11 Final Synthesis	127
265	A Appendix	128
266	A.1 List of Model Parameters and Physical Constants	128
267	A.2 Computational Scripts (Python)	129
268	A.2.1 Emergence Engine	129
269	A.2.2 Emergence Engine Output	146
270	A.2.3 Extended Lepton AMM	148
271	A.2.4 Extended Lepton AMM Output	150
272	A.2.5 AMM Resonance Scanner	151
273	A.2.6 AMM Resonance Scanner Output	154
274	A.2.7 Electroweak and Boson Module	154
275	A.2.8 Electroweak and Boson Module Output	158
276	A.2.9 Particle Mass Module	158
277	A.2.10 Particle Mass Module Output	162
278	A.2.11 Robustness and Stability Plots	164
279	A.2.12 Robustness and Stability Plots Output	172
280		
281		

Change Log

Version	Changes
5.1.0	Corrected the r^5 scaling label (“Resonance Mode Count” instead of “Resonance Frequency”); added the Lock-In Test and in-platform validation subsections; added OpenWave M4 references; minor typographical corrections.

286 1 Geometric Hierarchy of EWT and Spacetime Elasticity

287 Energy Wave Theory (EWT) is based on a quantum medium and establishes a clear hierarchy
288 of entities, starting from the Planck scale. This geometric structure is crucial for the elasticity
289 of spacetime and maintaining the constancy of the speed of light [1].

290 1.1 Wheeler’s Vision of Quantum Vacuum and the Path to EWT

291 The search for a discrete, pregeometric structure underlying spacetime has a rich intellectual
292 history, with John Archibald Wheeler as one of its most profound and visionary architects.
293 Wheeler’s work spanned several interlocking themes, each of which resonates – sometimes sup-
294 portively, sometimes critically – with the Energy Wave Theory (EWT) presented here.

295 Geometrodynamics and the 3-Geometry

296 Wheeler championed the idea that physics could be reduced to pure geometry [2]. In his ge-
297 ometrodynamic program, particles such as electrons and photons were to be understood as
298 **geons** – gravitational and electromagnetic fields trapped by their own curvature, embodying the
299 principle of “mass without mass” [6]. The Wheeler–DeWitt equation [5] attempted to quantise
300 this 3-geometry, laying the foundation for canonical quantum gravity.

301 EWT shares the conviction that geometry is primary, but it replaces Wheeler’s continu-
302 ous 3-geometry with a **discrete body-centered cubic (BCC) lattice** of Elastic Medium
303 Constituents (EMCs). The gravitational constant G emerges not from quantised continuum cur-
304 vature, but from the volumetric packing deficit of these spherical units – a concrete, calculable
305 pregeometry.

306 Quantum Foam and Pregeometry

307 Perhaps Wheeler’s most radical proposal was that at the Planck scale, spacetime is no longer
308 smooth but becomes a chaotic, fluctuating **quantum foam** [3] – a topological tapestry of worm-
309 holes and virtual geometries. He further argued that such a foam must be preceded by an
310 even more fundamental structure: **pregeometry** [8], the information-theoretic or combinatorial
311 substrate from which geometry itself crystallises.

312 In EWT, the quantum foam is replaced by a **static, ordered BCC crystal** with a well-defined
313 packing fraction ($\eta \approx 0.68$) and a residual impedance ζ . This is not a foam but a **mechanical**
314 **lattice**, whose elasticity (encoded in the effective geometric stiffness $N_{\text{geom}} = 8\pi^4(1 - \zeta)$) gives
315 rise to both gravitational and electromagnetic constants. Thus EWT provides a **concrete real-**
316 **isation of Wheeler’s pregeometry**, replacing speculative foam with an engineerable lattice.

317 It from Bit – and the Alternative of Geometric Realism

318 Wheeler’s famous aphorism “**it from bit**” [9] asserted that every physical entity (“it”) derives
319 from yes/no questions (“bits”), elevating information to the ontologically primitive level.

EWT takes a different stance. Here, the primitive is **geometric**: the BCC lattice and its spherical EMCs are not made of bits; they are the actual fabric of the vacuum. Information is an emergent property of geometric configurations, not their foundation. Consequently, EWT aligns more closely with a **geometric realism** than with Wheeler’s informational idealism – a crucial philosophical fork that distinguishes the present work from the Wheelerian mainstream.

Evolution of Wheeler’s Ideas in EWT

The table below summarises how each major Wheelerian concept has been transformed in the EWT framework:

Wheeler concept	Original meaning	EWT implementation / stance
Geometrodynamics	Continuous 3-geometry, geons	Discrete BCC lattice, EMC units, $N_{\text{final}} = 8\pi^4$
Quantum foam	Chaotic Planck-scale topology	Ordered BCC crystal, static packing deficit
Pregeometry	Abstract information substrate	Concrete BCC lattice with calculable elasticity
It from bit	Information ontological primacy	Superseded by Geometric Realism (Geometry precedes information)

Thus, while EWT draws inspiration from Wheeler’s insistence on a pregeometric foundation, it replaces the fluid, information-theoretic speculations with a **rigid, calculable lattice** – a shift from “foam” to “crystal” and from “bit” to “geometric element”. This evolution allows EWT to compute G , α , and the lepton anomalous magnetic moments from first principles, fulfilling Wheeler’s dream of a pregeometry while discarding the elements that proved non-predictive.

1.2 The Elastic Medium Constituent (EMC)

The **Elastic Medium Constituent (EMC)** is the smallest, fundamental geometric entity, which serves as the physical quantization limit of the proposed medium. Its size and spacing are directly linked to the requirement for the medium to propagate the electromagnetic wave at the speed of light c .

- **Geometric Radius (r_{EMC}):** In EWT, the EMC radius is typically defined as a magnitude on the order of 10^{-35} m, closely related to the **Planck Length** ($\lambda_l \approx 1.616 \times 10^{-35}$ m). This radius is the fundamental unit of length. Assuming the approximate value derived from the Planck scale:

$$r_{\text{EMC}} \approx 1.0 \times 10^{-35} \text{ m} \quad (1)$$

- **Inter-Constituent Distance (d_{EMC}) and Wavelength Quantum (λ_l):** The distance between the centers of adjacent EMCs defines the minimum wavelength, which directly influences the wave propagation speed c in the medium [1]. This distance is equal to the **Planck Length**:

$$d_{\text{EMC}} = \lambda_l \approx 1.616 \times 10^{-35} \text{ m} \quad (2)$$

- **Geometric Reference and Quantization:** The radius r_{EMC} , the distance d_{EMC} define the **absolute limit of spatial quantization** within the medium. The EMC is utilized as a geometric reference point for all larger structures.

1.3 The Wave Center (WC)

The Wave Center (WC) is defined as the focal point of oscillation in the elastic medium. The continuous interference of waves focused on this point establishes a standing wave structure (Soliton), which constitutes the localized oscillating volume of the medium. A single WC represents the fundamental unit of stored energy in the medium. Complex particles, such as the electron, are modeled as structures composed of multiple WCs. The variable $\mathbf{K}_{\text{WC}}$ represents the **number of constituent Wave Centers** in a given composite structure. For example, in the derivation of the electron's mass and charge, K_{WC} is a crucial summation factor [26].

1.4 The Soliton

A **Soliton** (or particle) in EWT is a stable structure of standing waves created by one or more WCs.

- **Standing Wave Boundary:** The Soliton is defined by the boundary where the generated standing waves transition into traveling waves. This transition point defines the geometric radius of the particle.
- **Energy and Mass Relation:** The energy stored in the standing waves of the Soliton is the source of the particle's rest mass, linking the wave structure directly to the mass-energy equivalence $E = mc^2$ [26].

1.5 Elastic Interaction (Hooke's Law)

The stability of the geometric structure hinges upon the nature of the medium's internal forces. Interactions between the Elastic Medium Constituents are modeled as ****elastic compression interactions (repulsion)****, a mechanism essential for Soliton stability and the propagation of energy.

- **Interaction Nature (Hooke's Law):** The force of repulsion between adjacent EMCs, when displaced from their equilibrium position (x), strictly follows **Hooke's Law** [11, 12]. This principle is interpreted as the fundamental **elasticity of the quantum medium**, which is necessary to **counteract enforce structural symmetry**:

$$\vec{F}_{\text{Hooke}} = -k\vec{x} \quad (3)$$

where k is the **elastic constant** specific to the medium.

- **Significance for Solitons and Volume Symmetry:** This elastic interaction ensures that the medium prevents unlimited compression of the Constituents.

The macroscopic stiffness parameter N (as defined in Eq. 14) used in the derivation of the gravitational constant and anomalous magnetic moments is the emergent aggregate of the microscopic elastic constants k of the BCC lattice substrate. While k defines the fundamental repulsive interaction between individual EMCs following Hooke's Law (3), N represents the global resistance of the vacuum to volumetric displacement. This connection bridges the localized elastic forces shown in Fig. 2 with the large-scale stability required for the precise determination of leptonic anomalies.




Figure 1: **The Elastic Medium Constituent (EMC) and Hooke's Law.** The EMC is the minimal geometric unit (defined by $r_{\text{EMC}} \sim l_p$) where the fundamental elasticity ($\mathbf{k}$) of the medium manifests as a repulsive force $\mathbf{F} = -k\mathbf{x}$, ensuring the Soliton's stability against geometric collapse. (Note: l_p denotes the Planck length, establishing the geometric scale for r_{EMC} .)




Figure 2: **Elastic Interaction between two EMCs.** The geometric overlap (compression $\vec{x}$) between adjacent Constituents generates the repulsive Hooke's Force $\mathbf{F} = -k\vec{x}$. This mechanism is the source of Soliton stability, balancing the internal geometric deficit ($\epsilon_{\mathbf{G}}$).

1.6 From Energy Domain to Geometric Domain

The Energy Wave Theory (EWT), as developed by Yee [46], successfully derived 23 fundamental physical constants from five wave constants: longitudinal amplitude A_l , longitudinal wavelength λ_l , aether density ρ , wave speed c , and the electron's wave center count $K_{WC} = 10$. In that framework, charge was already reinterpreted as wave amplitude, measured in meters, hinting at an underlying geometric reality. The present work extends this program by identifying the physical substrate of these waves: a Body-Centered Cubic (BCC) lattice of spherical Elastic Medium Constituents (EMCs). The wave constants are thus replaced by geometric invariants of this lattice: the statutory neutrino radius r_ν (related to λ_l and K_{WC}), the effective nodal stiffness $N_{\text{geom}} = 8\pi^4(1-\zeta)$ (derived from the BCC coordination number and packing impedance), and the

396 magnetic deficit $\epsilon_M = 1/(N_{\text{geom}}\pi^3)$ (encoding the 7-dimensional weak interaction scale). This
 397 transition from the energy domain to the geometric domain not only preserves the predictive
 398 power of EWT but also unifies gravity, electromagnetism, and the weak force under a single
 399 topological framework.

400 2 Geometric Equation of the Fine-Structure Constant and 401 the Deficit Terms

402 The **fine-structure constant** (α) is defined within Energy Wave Theory (EWT) as a geometric
 403 ratio describing the relative strengths of fundamental forces—charge energy, magnetism, and
 404 gravity [10, 16].

405 The derivation of the inverse fine-structure constant ($1/\alpha$) is realized as the combination of a
 406 purely geometric core and a small topological correction. The core is obtained from the surface
 407 geometry of the soliton, while the correction is dictated by the BCC lattice structure.

408 This approach redefines fundamental particle properties:

- 409 • **Charge as Amplitude:** Electric charge is interpreted not as an abstract quantity, but as
 410 the physical **amplitude** (x) of a standing wave oscillation within a quantum medium.
- 411 • **Geometric Ratio for α :** The fine-structure constant is defined as the ratio of squared
 412 amplitudes, which is equated to a geometric ratio involving the total surface area of energy
 413 propagation (S):

$$\alpha = \frac{A_1^2}{A_0^2} = \frac{x^2}{S} \quad (4)$$

414 Geometric Soliton Core ($4\pi^3 + \pi^2 + \pi$)

415 The derivation of the pure geometric constant $\mathbf{A}_\pi$ was first proposed by Jeff Yee [10] within the
 416 framework of the Energy Wave Theory. The core postulate involves equating the fine-structure
 417 constant to a specific ratio of surface areas in the quantum background, and the derivation
 418 proceeds in the following steps:

419 **Step 1: Defining the Total Surface Area S** The total surface area S is the sum of the sphere
 420 surface area and the total cone surface area:

$$S = \underbrace{4\pi l^2}_{\text{Sphere Area}} + \underbrace{(\pi r l + \pi r^2)}_{\text{Total Cone Area}} \quad (5)$$

421 **Step 2: Substitution of Geometric Relations** ($r = x$, $l = \pi x$) Substituting $r = x$ and $l = \pi x$
 422 expresses S solely in terms of π and x :

$$S = 4\pi(\pi x)^2 + (\pi(x)(\pi x) + \pi x^2) \quad (6)$$

423 **Step 3: Simplification** The expression is simplified by factoring out x^2 :

$$S = 4\pi^3 x^2 + \pi^2 x^2 + \pi x^2 = x^2(4\pi^3 + \pi^2 + \pi) \quad (7)$$

424 **Step 4: Final Geometric Fine-Structure Constant** Substituting the simplified S into $\alpha =$
 425 x^2/S , the x^2 terms cancel, yielding α as a pure function of π :

$$\mathbf{A}_\pi^{-1} = \frac{1}{4\pi^3 + \pi^2 + \pi} \quad (8)$$

Kinematic Origin of the π Factor The relation $l = \pi x$ defines the propagation distance in terms of the source amplitude. It is not an arbitrary geometric choice but a consequence of the harmonic motion of the central EMC in an elastic medium with constant propagation speed c . In one complete oscillation the central granule travels a total longitudinal displacement of $2x$. During the same time interval, the disturbance emitted by this motion propagates outward at the lattice speed c . Because the central granule follows a sinusoidal path, its corresponding longitudinal phase advance is exactly πx , whereas the undisturbed wavefront travels the longer distance $l = \pi x$. The factor π therefore enters through the kinematics of harmonic oscillation, not as a fitted parameter. This mechanical origin of l is what ultimately generates the characteristic powers π^3, π^2, π in the fine-structure geometry.

“In the time that it takes for the center granule to vibrate, traveling a distance (r), and then returning (total of $2r$), the granules that it affects will have traveled a distance of length (l). The harmonic motion of the center granule over time can be represented by a sine wave, where the travel in one direction and a return to equilibrium is a half-wavelength, or π .”

“Assuming constant velocity for granules (expected to be the speed of light), the distance that the granules will have traveled is π times x . In other words, while the center granule travels a total distance of $2x$ during vibration, the straight-line motion of the granules it effects travels $\sim 3.14x$ because it doesn’t slow down to account for a turn in direction.”

— Yee, J. (2019). *The Relationship of the Fine Structure Constant and Pi*, p. 3–4. [10]

2.1 BCC Lattice Geometry and the Geometric Ladder

The vacuum lattice is a discrete body-centred cubic (BCC) lattice of Elastic Medium Constituents (EMCs). The BCC lattice is characterised by:

- 8 nearest neighbours,
- packing fraction

$$\eta_{\text{BCC}} = \frac{\sqrt{3}\pi}{8}, \quad (9)$$

- ideal projection factor

$$L_p^{\text{geom}} = \frac{2}{\sqrt{3}}, \quad (10)$$

The factor L_p^{geom} is the inverse of the dimensionless nearest-neighbour distance in the BCC lattice. It is not an adjustable constant but a direct consequence of the $Im\bar{3}m$ space group: the nearest neighbour lies at $d/a = \sqrt{3}/2$, and its inverse gives the geometric projection factor $2/\sqrt{3}$.

The BCC geometry introduces the constants π and $\sqrt{3}$ directly through the packing fraction. In later derivations, the face-diagonal factor $\sqrt{2}$ appears in the lattice projection operators and in the unified coupling bridge, while Euler’s number e enters through the Eulerian dilution of the statutory vacuum density. Neither of these constants is fitted; each has a well-defined geometric role in the corresponding section.

The geometry of the lattice imposes a natural dimensional hierarchy on the soliton’s wave structure. Each additional active degree of freedom contributes one factor of π , giving rise to the following geometric ladder:

The factor π appearing in every rung of this ladder is not introduced by hand. Its origin is the same as in the geometric soliton core: the kinematic relation $l = \pi x$, which reflects one full harmonic phase advance of the central EMC during one oscillation. Each additional degree of freedom therefore multiplies the phase-space budget by one further factor of π .

Table 1: The Geometric Ladder: Dimensional Interaction Topology

Scale	Interaction	Budget Components	Dim.
π^7	Charged Weak	$3D + A + f + r + Q$	7
π^6	Neutral Weak	$3D + A + f + r$	6
π^5	Flavor Mixing	$3D + A + f$	5
π^4	Int. Binding	$3D + A$	4
π^3	Spatial Substrate	$3D$	3

Legend: A = amplitude, f = frequency, r = radius, Q = charge.

The π^7 rung is particularly important: it represents the charged weak sector, in which the soliton engages seven independent geometric degrees of freedom. The BCC coordination number is 8. The magnetic deficit ϵ_M introduced below is therefore naturally associated with the inverse of the product $8\pi^7$, with a small correction due to the non-ideal spherical packing of the lattice.

2.2 Geometric Magnetic Deficit ϵ_M

The ideal stiffness of the BCC lattice follows from its eight nearest neighbours and the four-dimensional saturation budget:

$$N_{\text{ideal}} = 8\pi^4. \quad (11)$$

The non-ideal spherical packing of the EMCs gives rise to a small lattice impedance

$$\zeta = \frac{1 - \eta_{\text{BCC}}}{\eta_{\text{BCC}} N_{\text{ideal}}}. \quad (12)$$

Applying this impedance to the ideal stiffness yields the effective geometric stiffness:

$$N_{\text{geom}} = N_{\text{ideal}}(1 - \zeta). \quad (13)$$

The magnetic deficit is then defined by the key relation

$$\epsilon_M = \frac{1}{N_{\text{geom}} \pi^3} = \frac{1}{8\pi^7(1 - \zeta)} \quad (14)$$

In the ideal continuum limit $\zeta \rightarrow 0$, this reduces to

$$\epsilon_M^{(\text{ideal})} = \frac{1}{8\pi^7}. \quad (15)$$

The complete geometric inverse fine-structure constant therefore has the form

$$\frac{1}{\alpha} = A_\pi - \epsilon_M - \sum \epsilon_G, \quad (16)$$

where $\sum \epsilon_G$ is a lower-order gravitational deficit term. For the electron this term is negligible compared with ϵ_M , so the working expression used throughout this work is

$$\frac{1}{\alpha_{\text{geom}}} = A_\pi - \epsilon_M. \quad (17)$$

The omitted lower-order term $\sum \epsilon_G$ is retained as a formal remainder; it is not fitted and may become relevant when the lower-order gravitational correction is no longer negligible.

This expression contains no calibrated parameters. All quantities on the right-hand side are fixed by the BCC lattice geometry and the mathematical constants π , e , and $\sqrt{2}$.

2.3 The Fundamental Lattice Response Parameter ϵ_M

A central role in the Enhanced EWT framework is played by the parameter ϵ_M , traditionally referred to as the *magnetic deficit factor*. Initially, this parameter was identified as a functional heuristic—a "magnetic deficit" required to reconcile the fine-structure constant α within the energy wave equations. However, as the framework evolved, it became evident that ϵ_M was not a localized adjustment, but a fundamental property defining the very "stiffness" of the 3D space occupied by matter.

Within the unified geometric context of this work, ϵ_M is more fundamentally defined as the **Global Lattice Impedance Constant**. It represents the intrinsic structural resistance of the BCC vacuum substrate to any deviation from ideal spherical wave symmetry. This realization transformed it from a specialized electromagnetic factor into a cornerstone of Enhanced EWT, acting as a universal "scaling bridge" across different physical sectors.

By treating ϵ_M as a singular topological property, the model achieves numerical convergence across disparate scales—from the gravitational constant G to the anomalous magnetic moments of leptons—without the need for independent empirical constants. The structural coherence of this approach is formally validated in Section 28, where it is shown that this single value functions as the primary scaling anchor for the entire theory.

Ultimately, this work reveals that ϵ_M is not an arbitrary input, but a direct consequence of the BCC lattice coordination. Its explicit geometric form is given in Eq. (14).

3 Dimensional Hierarchy and Dynamic Resonant Modulations

This section translates the geometric ladder introduced in Section 2 into concrete lattice operators for the bosonic and fermionic sectors. The high-precision **2022 CDF II** measurement of M_W serves as an experimental anchor, revealing that the reported Standard Model mass tensions arise from the omission of volumetric lattice impedance inherent to the BCC substrate.

3.1 The Universal Geometric Modulators: Substrate Properties

The formation of mixing angles within the EWT framework is governed by a hierarchy of modulators derived from the global magnetic deficit ϵ_M . These factors account for the intrinsic lattice impedance and the resonant displacement required to maintain equilibrium within the BCC vacuum substrate. Rather than being empirical parameters, these modulators represent the structural response of the lattice to different interaction topologies (volumetric vs. surface).

3.1.1 Unified Local Constant C_{local}

The **Unified Local Constant** is the structural bridge between the global lattice magnetic deficit (ϵ_M) and the local chiral projection ($2\sqrt{2}$):

$$C_{\text{local}} \equiv \frac{\epsilon_M}{2\sqrt{2}} \quad (18)$$

519 3.1.2 The π^6 Resonance: Volumetric Bosonic Coupling

520 Vector bosons (W, Z, H) are high-energy excitations involving the full phase space. The modu-
521 lator $\mathbf{C}_{\text{gap}}$ accounts for the volumetric resistance of the lattice:

$$C_{\text{gap}} = 1 + \pi^6 \cdot C_{\text{local}} \quad (19)$$

522 The Weinberg angle emerges as a structural equilibrium point between the 6D volumetric
523 resonance (π^6) and the boson mass ratio:

$$\sin^2 \theta_W = 1 - \left(\frac{M_W}{M_Z} \right)^2 \cdot \frac{1}{C_{\text{gap}}} \quad (20)$$

524 By utilizing this structural constant, the framework identifies the ideal mass attractor for the
525 W boson, accounting for the 6D volumetric resonance:

$$M_W^{\text{EWT, Ideal}} = M_Z^{\text{CODATA}} \cdot \sqrt{(1 - \sin^2 \theta_W^{\text{Target}}) \cdot C_{\text{gap}}} \quad (21)$$

526 3.1.3 The π^7 Resonance: Charge-Induced Amplitude Loading

527 The highest rung of the geometric ladder, π^7 , is associated with the charged weak sector ($W^\pm$).
528 Unlike the neutral Z^0 and H^0 solitons, the W boson carries non-compensated wave amplitude
529 (charge), which acts as an additional **7th degree of freedom** within the BCC substrate.

530 In the EWT framework, the W boson mass is not an independent input but a **geometric**
531 **attractor**: the vacuum must sustain a mass of approximately 80.5141 GeV to maintain structural
532 equilibrium against the charge-induced displacement.

533 3.1.4 The π^5 Resonance: Surface Interaction Modulator

534 For the quark sector and flavour mixing, the interaction topology shifts from the volume to the
535 **soliton boundary**. The modulator $\mathbf{C}_{\text{fermion}}$ represents a **5D holographic projection** (3D
536 momentum + 2D spherical surface):

$$\mathbf{C}_{\text{fermion}} \equiv (1 + \pi^5 \cdot \mathbf{C}_{\text{local}})^2 \quad (22)$$

537 The quadratic form reflects the bi-local nature of mixing between two quark states within the
538 lattice substrate.

539 3.2 Geometric Mixing Predictions: Higgs and Cabibbo Sectors

540 3.2.1 Higgs-Vector Boson Mixing

541 The Higgs boson interactions are governed by the π^6 volumetric laws mapped onto the Higgs
542 mass scale ($M_H^{\text{CODATA}} = 125.25$ GeV), utilising the geometric $M_W^{\text{EWT, Ideal}}$:

$$\sin^2 \theta_{ZH}^{\text{EWT}} = \mathbf{0.4686} \quad (23)$$

543 **Falsification of the Standard Model (VEV Mechanism):** The stability of Eq. (23)
544 serves as a critical test. Confirming this discrete value would prove the Higgs is a composite
545 soliton, thereby **falsifying the Standard Model**, as such a resonant state is incompatible with
546 the vacuum-expectation-value (VEV) mechanism.

Table 2: Geometric Mixing Angles for Higgs-Vector Boson Pairs

Mixing Interaction	Calculation Formula	EWT Prediction
$\sin^2 \theta_{ZH}$	$1 - \left(\frac{M_Z^{\text{EWT}}}{M_H^{\text{EWT}}} \right)^2 \cdot \frac{1}{C_{\text{gap}}}$	0.4686
$\sin^2 \theta_{WH}$	$1 - \left(\frac{M_W^{\text{EWT, Ideal}}}{M_H^{\text{EWT}}} \right)^2 \cdot \frac{1}{C_{\text{gap}}}$	0.5906

3.2.2 Cabibbo Mixing Angle

Using the π^5 surface modulator and the EWT-derived masses for d and s quarks, obtained from the spherical mode at wave centre counts $K_{WC} = 15$ and $K_{WC} = 28$:

$$\sin \theta_C = \sqrt{\frac{M_d}{M_s}} \cdot C_{\text{fermion}} \quad (24)$$

where

$$C_{\text{fermion}} = \left(1 + \pi^5 C_{\text{local}} \right)^2, \quad (25)$$

and the masses calculated from the spherical mode are:

$$M_d^{\text{EWT}} = 0.004034 \text{ GeV}, \quad (26)$$

$$M_s^{\text{EWT}} = 0.094885 \text{ GeV}. \quad (27)$$

The experimental value (PDG 2022) is $\sin \theta_C \approx 0.2243$. The EWT-derived masses lead to a relative difference of about 7.24% (Variant A in Table 3).

To isolate the precision of the geometric mixing mechanism itself, the calculation is repeated using PDG 2022 experimental quark masses as input:

$$M_d^{\text{PDG}} = 0.004692 \text{ GeV}, \quad (28)$$

$$M_s^{\text{PDG}} = 0.094954 \text{ GeV}. \quad (29)$$

Substituting these values into the same operator yields a deviation of only 0.0055% (Variant B), confirming that the π^5 surface resonance operator C_{fermion} correctly captures the physics of flavour mixing at the level of numerical precision.

Table 3: Cabibbo angle prediction: sensitivity to quark mass input.

Variant	M_d [GeV]	M_s [GeV]	$\sin \theta_C$	Error
A: EWT spherical mode	0.004034	0.094885	0.20805	7.24%
B: PDG 2022 targets	0.004692	0.094954	0.22429	0.0055%
PDG 2022 (reference)	—	—	0.22430	—

3.2.3 Dimensional Budget of Resonances: Reflection vs. Mixing

The scaling of the modulators is determined by the “dimensional budget” of the resonance involved in the interaction with the BCC lattice:

- 562 • **Bosons (π^6 and π^7):** These are volumetric excitations where the action occurs in 3D
563 space, involving 1D amplitude, 1D frequency, and 1D radius (r). For charged bosons, an
564 additional 1D of freedom (charge) expands the budget to π^7 . These processes represent
565 a unified “reflection” of the energy packet off the lattice, resulting in a linear correction
566 (C_{gap}).
- 567 • **Fermions (π^5):** In the fermionic sector, the interaction is localised on the phase-surface.
568 The budget includes 3D space, 1D amplitude, and 1D frequency, but excludes the volu-
569 metric radius (r). This π^5 resonance describes the surface integrity of the soliton.
- 570 • **The Square (Interference):** Unlike the singular reflection of bosons, the Cabibbo angle
571 represents the **mixing** of two such π^5 states. Consequently, the lattice modulation must
572 be applied to both participating objects, leading to the quadratic form $C_{\text{fermion}} = (1 +$
573 $\pi^5 C_{\text{local}})^2$.

574 3.2.4 Symmetry of Observables: Energy Density vs. Amplitude Interference

575 The contrasting mathematical forms of the key observables— $\sin^2 \theta_W$ for the bosonic sector and
576 $\sin \theta_C \propto \sqrt{M_d/M_s}$ for the fermionic sector—are not coincidental. They reflect a fundamental
577 transition in the nature of the interaction within the BCC lattice, rooted in the dimensional
578 hierarchy of the Geometric Ladder.

- 579 • **Bosonic Sector (Volumetric Energy Density):** The Weinberg angle, $\sin^2 \theta_W$, emerges
580 from the ratio of the W and Z boson masses. In the EWT framework, mass is a measure
581 of volumetric energy density, which at the π^6 and π^7 levels involves the full set of degrees
582 of freedom: 3D space, amplitude A , frequency f , radius r , and (for charged bosons) charge
583 Q . The squared form of the observable directly mirrors the squared relation between mass
584 and the underlying wave amplitude ($E \propto A^2$), making $\sin^2 \theta_W$ the natural expression for
585 a volumetric, energy-density-based coupling.
- 586 • **Fermionic Sector (Surface Amplitude Interference):** In contrast, the Cabibbo angle,
587 $\sin \theta_C$, describes the mixing of two boundary-localised fermions (d and s quarks). This is a
588 bi-local interference process occurring on the π^5 phase-surface, where the relevant physical
589 quantity is the wave amplitude A itself, not the volume-integrated energy. At the π^5 level,
590 the soliton possesses 3D space, amplitude A , and frequency f —the minimal set required
591 for a massive surface excitation. Its mass scales as $M \propto A^2$, as energy is quadratic in
592 amplitude. To access the amplitude A directly, one must therefore take the square root
593 of the mass. This operation effectively projects the energy-based observable from the π^5
594 level down to the π^4 level of the Geometric Ladder, where the fundamental amplitude A
595 resides in 3D space as the static structural skeleton of the particle (3D + A). The linear
596 form $\sin \theta_C \propto \sqrt{M_d/M_s}$ is thus the direct signature of amplitude interference at the soliton
597 boundary.

598 This dichotomy— $\sin^2$ for volumetric energy ratios versus $\sin$ for surface amplitude interference—
599 is a direct and testable consequence of the dimensional hierarchy. It confirms that the EWT
600 framework does not merely fit parameters, but derives the very structure of physical laws from
601 the topology of the vacuum substrate.

602 3.2.5 Interpretation of Quark Masses in the EWT Framework

603 While the Standard Model operates on the abstraction of point-particles within a passive vacuum,
604 the Enhanced EWT model restores the physical necessity of the medium. Matter cannot exist in

space without being of space. The π^4 Quark Stability budget formalises this necessity: the 3D spatial substrate (π^3) is not merely a background, but the indispensable structural foundation for any dynamic amplitude (π^1) and/or frequency (π^1) to manifest as a stable particle.

Remark on Quark Masses and the Nature of Confinement:

In the EWT framework, quarks are not fundamental particles but topological excitations confined within hadronic solitons. Their “masses” are not intrinsic properties of free entities, but effective measures of the binding energy within the collective π^5 phase-surface. The permanent isolation of a quark is geometrically impossible: it would require severing the π^4 amplitude-space core from the π^5 surface, leaving a residual frequency without a spatial substrate.

3.3 Structural Transition to N -Stiffness: From Geometric Volume to Lattice Impedance

To unify the model, the correction parameters C_{gap} and C_{fermion} are expressed directly through the geometric vacuum stiffness N_{geom} introduced in Section 2. Using the identity $C_{\text{local}} = \epsilon_M/(2\sqrt{2})$ and $\epsilon_M = 1/(N_{\text{geom}}\pi^3)$, the geometric structure reveals the scaling hierarchy:

1. Direct N -Representation

$$C_{\text{gap}} = 1 + \pi^6 \cdot C_{\text{local}} = 1 + \frac{\pi^3}{2\sqrt{2} \cdot N_{\text{geom}}} \quad (30)$$

$$C_{\text{fermion}} = (1 + \pi^5 \cdot C_{\text{local}})^2 = \left(1 + \frac{\pi^2}{2\sqrt{2} \cdot N_{\text{geom}}}\right)^2 \quad (31)$$

2. Physical Interpretation: The Pi-Power Scaling

- **C_{gap} (Volumetric Coupling):** The π^3/N_{geom} term identifies the magnetic gap as a 3D volumetric resonance. The π^3 factor represents the interaction of the spherical soliton with the BCC lattice stiffness.
- **C_{fermion} (Surface Scaling):** The reduction to π^2/N_{geom} within the squared operator identifies the fermion correction as a 2D surface effect. The square accounts for the bi-directional (spin-lattice) coupling required for mass stabilisation.
- **The N_{geom} -Anchor:** Both constants are locked to the same geometric stiffness parameter, proving that the difference between magnetic and mass-surface anomalies is purely a matter of geometric dimensionality (π^3 vs π^2).

The transition to the N_{geom} -anchor reveals a fundamental split in the vacuum’s response: the bosonic sector is governed by 3D volumetric resonance, while the fermionic sector is dictated by 2D surface-coupling dynamics.

4 Analysis of Neutrino Boundary Conditions

Before proceeding to the detailed analysis of the neutrino's boundary conditions, it is crucial to place the Geometric Gravity mechanism of EWT—derived from the Gravitational Deficit ϵ_G —within the broader context of modern theoretical physics. The Energy Wave Theory's postulate that gravity arises from a deficit in the **Elastic Medium Constituent** (EMC) components conceptually aligns with the **Emergent Gravity** paradigm. This framework suggests that gravity is not a fundamental force, but rather an effective phenomenon arising from the microstructure or thermodynamics of spacetime [19]. Key proponents, such as Padmanabhan and Verlinde, develop theories where gravity emerges from spacetime thermodynamics or the information encoded on a holographic screen [20]. While such models face significant theoretical and empirical challenges [21], the EWT approach offers a concrete geometric mechanism linking a specific wave-centre structure to a macroscopic gravitational effect, without invoking entropic or thermodynamic assumptions. This links a specific wave-centre structure to a macroscopic gravitational effect, providing a unique, finite, and quantifiable emergent model.

The stability of the Neutrino as a Soliton is dependent on the precise balancing of internal geometric forces. These forces are represented by the **Constant Geometric Deficit** (ϵ_G), the **global Magnetic Deficit** (ϵ_M), and the **Geometric Soliton Core**.

4.1 Standard Conditions: Mass and Minimal Magnetism

Under standard conditions, the Neutrino is confirmed to possess non-zero mass (empirically validated by oscillation experiments [22, 23]) and a negligible, but potentially non-zero, magnetic moment [24].

- **Stability (Hooke's Force):** Non-zero mass is understood to imply that the Neutrino possesses an internal **Constant Geometric Deficit** ϵ_G . The internal geometric pressure, caused by ϵ_G and other factors, is **balanced** by the repulsion force resulting from **Hooke's Law** (Elastic Interaction) between the constituents, which keeps the Soliton in a stable state.
- **Charge:** The Soliton's charge is **considered effectively zero** (≈ 0).
- **Magnetism and ϵ_M :** The global magnetic deficit ϵ_M is fixed by the BCC lattice geometry and does not vanish for the neutrino. However, because the neutrino carries no charge and no spin, it does not generate the local magnetic torque that compresses charged leptons. As a result, the neutrino exhibits only a negligible local magnetic response, even though a tiny moment $\mu_\nu \neq 0$ is not excluded.

4.2 Extreme Conditions: Wave Center in a Geometric Black Hole (GBH)

In the case of a Geometric Black Hole (GBH) [17], extreme geometric compression and the dominance of the accumulated gravitational field ($\sum \epsilon_G$) impose severe conditions.

- **Dominance of ϵ_G and Stability (Hooke's):** At the Critical Distance (d_{crit}) of the GBH, geometric forces are so extreme that **all geometric terms other than gravity are suppressed**. However, the **Elastic Interaction** (F_{Hooke}) is an invariant property and **is still active**, guaranteeing that the Soliton's radius does not fall below r_ν (the limit of minimal Soliton stability), and the Neutrino remains a stable WC.

- 667 • **Empirical Validation (Supernovae):** The observation that **99% of supernova energy**
668 **is released as neutrinos** constitutes empirical confirmation that the collapse of matter
669 under extreme gravity leads to the conversion of the dominant mass into the **minimal**
670 **geometric state** of the Neutrino (r_ν) at the point d_{crit} . This fact justifies the adoption
671 of the Neutrino as the fundamental WC for the GBH model.
- 672 • **Local Magnetic Suppression:** Under conditions of extreme compression, the soliton's
673 local spin and magnetic degrees of freedom are geometrically frozen. This does not mean
674 that the global lattice parameter ϵ_M changes; rather, the local contribution of the spin
675 sector to the soliton's energy balance is suppressed.
- 676 • **Pure WC (Boundary Condition):** At the critical point (d_{crit}), the geometric boundary
677 conditions of the GBH force the complete elimination of charge and the local magnetic
678 response. The Neutrino is converted into a **pure Wave Center (WC)** with the single
679 dominant characteristic (ϵ_G):

$$\text{Charge} = 0 \quad \text{and the local magnetic contribution is suppressed.} \quad (32)$$

680 Hypothesis of Spin Recovery Beyond the GBH Horizon

681 Within the EWT framework, the geometric structure of the WC (and thus its local spin/magnetic
682 properties) may be **partially recovered or enhanced** outside the GBH's critical boundary
683 d_{crit} . This phenomenon is driven by the reduced compression and resulting non-linear elasticity
684 of the spacetime medium away from the core, aligning conceptually with information preservation
685 hypotheses near the horizon.

686 5 Neutrino as the Geometric Anchor

687 The statutory neutrino radius r_ν sets the fundamental length scale of the BCC lattice. In this
688 section, r_ν is not treated as an independent input. Instead, it emerges from the geometric
689 fine-structure constant α_{geom} , the elementary charge amplitude e , and the topology of the BCC
690 lattice.

691 5.1 Planck Charge from Geometric Alpha

692 The Planck charge in EWT is interpreted as the fundamental geometric amplitude. It is not
693 introduced as a separate empirical constant; it follows from the relation between the elementary
694 charge amplitude e and the geometric fine-structure constant:

$$q_P = \frac{e}{\sqrt{\alpha_{geom}}}, \quad (33)$$

695 where $e = 1.602176634 \times 10^{-19}$ m is the CODATA elementary charge amplitude, and α_{geom} is
696 the purely geometric fine-structure constant derived in Section 2.

697 The resulting ratio

$$\frac{q_P}{e} = \frac{1}{\sqrt{\alpha_{geom}}} \approx 11.70624886 \quad (34)$$

698 is therefore a geometric prediction, not a fitted input.

5.2 Decomposition of the Scaling Factor $S = r_\nu/q_P$

The dimensionless scaling factor $S = r_\nu/q_P$ is obtained by combining three physically distinct mechanisms that act in the undisturbed BCC vacuum lattice.

1. **Static lattice projection.** The soliton potential, described by the inverse geometric fine-structure constant $\alpha_{\text{inv, geom}}$, is distributed over the eight BCC coordination nodes and the spherical wave front:

$$S_{\text{proj}} = \frac{\alpha_{\text{inv, geom}}}{8 + \pi}. \quad (35)$$

2. **Dynamic wave expansion.** The natural exponential decay of the standing-wave amplitude from the centre outward introduces Euler's number e as an integrated factor:

$$S_{\text{exp}} = e. \quad (36)$$

3. **Discrete lattice impedance.** The discrete BCC lattice and wave propagation along the face diagonals add a small correction:

$$\delta_{\text{imp}} = (1 - g_v)(\sqrt{2} - 1). \quad (37)$$

Here g_v is the geometric correction factor associated with the neutrino ground state. The factor $\sqrt{2} - 1$ measures the excess diagonal path length, while $1 - g_v$ quantifies the deviation of the relaxed neutrino state from an ideal continuum response.

The total scaling factor is the sum of these components:

$$S_{\text{tot}} = S_{\text{proj}} + S_{\text{exp}} + \delta_{\text{imp}}. \quad (38)$$

5.3 Geometric Fixed Point of g_v

The dynamic relation for the neutrino radius is

$$r_\nu = \frac{2q_P e^2}{g_v}. \quad (39)$$

In terms of the scaling factor, this becomes

$$S_{\text{tot}} = \frac{2e^2}{g_v}. \quad (40)$$

Equating this with Eq. (38) gives

$$\frac{2e^2}{g_v} = \frac{\alpha_{\text{inv, geom}}}{8 + \pi} + e + (1 - g_v)(\sqrt{2} - 1). \quad (41)$$

Rearranging leads to the quadratic equation

$$(\sqrt{2} - 1)g_v^2 - \left(\frac{\alpha_{\text{inv, geom}}}{8 + \pi} + e + \sqrt{2} - 1 \right) g_v + 2e^2 = 0. \quad (42)$$

This equation has two roots:

$$g_v \approx 0.983594 \quad \text{and} \quad g_v \approx 36.27. \quad (43)$$

Only the first root satisfies $0 < g_v < 1$, which is required for stable wave propagation and a physical soliton. Thus

$$g_v \approx 0.983594 \quad (44)$$

is the unique geometric fixed point of the BCC lattice. It is not an adjustable parameter, but a direct consequence of the lattice topology.

5.4 Resulting Neutrino Radius

Using the fixed-point value of g_v and the geometric Planck charge q_P , the neutrino radius is

$$r_\nu = q_P S_{\text{tot}} \approx 2.817935 \times 10^{-17} \text{ m.} \quad (45)$$

This value is in close agreement with the earlier dynamical expression

$$r_\nu = \frac{2q_P e^2}{g_v}, \quad (46)$$

and the two approaches differ only at the 10^{-6} level. This confirms that r_ν is derived from the same BCC geometry as α_{geom} and ϵ_M .

5.5 Decadic Resonance Test: $r_e/r_\nu \approx 100$

An immediate test of the geometric anchor is provided by the ratio of the classical electron radius to the neutrino radius:

$$\frac{r_e}{r_\nu} \approx 100.00017. \quad (47)$$

The corresponding fifth power

$$\left(\frac{r_e}{r_\nu}\right)^5 \approx 1.0000087 \times 10^{10} \quad (48)$$

reproduces the 10^{10} scaling expected from the difference between the electron wave-centre count $K_{WC} = 10$ and the neutrino count $K_{WC} = 1$. This is a direct consequence of the geometric mass-to-radius identity, Eq. (163), discussed in Section 15.1. It provides a non-trivial consistency test linking the geometric radius to the r^5 energy scaling used throughout the model.

5.6 Physical Interpretation: ϵ_M as a Global Lattice Parameter

The magnetic deficit ϵ_M is a global topological property of the BCC lattice and does not vanish for any soliton. What differs between the neutrino and charged leptons is the local magnetic response.

The neutrino carries no charge and no spin; it therefore generates no local magnetic torque. Consequently, its soliton relaxes to the natural radius r_ν given by Eq. (45).

For the electron, the local magnetic response compresses the soliton below this relaxed state. The factor g_v measures exactly this difference: it is the ratio between the relaxed neutrino radius and the compressed wavelength that would follow from the simple wave constants. The global ϵ_M remains unchanged; it is the local magnetic compression that differs between particles.

Thus the neutrino is not a “small electron”, but the fundamental, torque-free ground state of the BCC lattice. The electron is a compressed and locally magnetised excitation of this same ground state.

5.7 Topological Necessity of r_ν

The decomposition of S_{tot} into

$$\frac{\alpha_{\text{inv, geom}}}{8 + \pi}, \quad e, \quad (1 - g_v)(\sqrt{2} - 1) \quad (49)$$

shows that the neutrino radius is fixed by three simultaneous geometric constraints:

- 752 • the static soliton potential distributed over the eight BCC directions and the spherical wave
753 front,
- 754 • the natural exponential decay encoded by Euler’s number e ,
- 755 • the discrete lattice correction along the face diagonals.

756 No free parameter is introduced. All quantities are determined by π , e , $\sqrt{2}$, the BCC coor-
757 dination number, and the geometric fine-structure constant.

758 This completes the emergence of the neutrino as a topological anchor of the BCC lattice. No
759 fitted parameter is introduced in this derivation.

760 6 Planck Length and Reduced Planck Constant from Ge- 761 ometry

762 The geometric fine-structure constant α_{geom} , the electron radius r_e , and the electron mass m_e
763 determine the reduced Planck constant without invoking $\hbar$ as an independent input. The same
764 geometric machinery then yields the Planck length λ_l as a derived quantity.

765 6.1 Reduced Planck Constant from Geometric Alpha

766 In EWT, the classical electron radius is not defined through $\hbar$ but is treated as a geometric scale
767 fixed by the electron soliton. The reduced Planck constant follows from the relation

$$\hbar_{\text{geom}} = \frac{m_e c r_e}{\alpha_{\text{geom}}}, \quad (50)$$

768 where c is the metric conversion factor defined in Section 9.

769 Using the CODATA values

$$m_e = 9.1093837015 \times 10^{-31} \text{ kg}, \quad r_e = 2.8179403262 \times 10^{-15} \text{ m}, \quad (51)$$

770 and the geometric fine-structure constant α_{geom} , Eq. (50) gives

$$\hbar_{\text{geom}} \approx 1.0545738 \times 10^{-34} \text{ kg m}^2 \text{ s}^{-1}, \quad (52)$$

771 in agreement with the CODATA value at the level of the underlying experimental inputs. No
772 new free parameter is introduced.

773 7 The Physical Origin of Charge as Wave Amplitude

774 The identification of charge with wave amplitude is a foundational postulate of Energy Wave
775 Theory, established in the primary EWT literature [29]. The present section makes the dimen-
776 sional reduction explicit, in the same way that the kilogram was reduced to a geometric unit in
777 Section 15.3.

7.1 Charge as Spatial Wave Amplitude

In EWT, the elementary charge e of a single electron represents the physical wave displacement amplitude of the lattice constituents at the first wavelength. Because displacement in a spatial lattice is fundamentally a geometric length, charge carries the dimension of length ($[Q] = [L]$).

In the International System of Units (SI), where the fundamental unit of spatial length is the metre, the elementary charge amplitude is given by:

$$e_{\text{geom}} = 1.602176634 \times 10^{-19} \text{ m}. \quad (53)$$

This is a direct dimensional mapping: the numerical value of the CODATA elementary charge is preserved, while the dimension changes from coulombs (C) to metres (m).

7.2 Conversion of the Coulomb and Distinction from CGS

From the definition $1 \text{ C} = 1 \text{ A s}$ and the geometric representation of charge as amplitude (m), the coulomb is recognized as a derived unit of spatial displacement:

$$e_{\text{SI}} = 1.602176634 \times 10^{-19} \text{ C} \iff e_{\text{geom}} = 1.602176634 \times 10^{-19} \text{ m}. \quad (54)$$

Resolution of the CGS / Statcoulomb Ambiguity:

It is crucial to emphasize that e_{geom} is a spatial displacement ($[L]$), not a force-derived quantity. In Gaussian CGS units, the statcoulomb is defined via Coulomb's Law as $\text{g}^{1/2} \text{ cm}^{3/2} \text{ s}^{-1}$, which assigns charge a composite mechanical dimension rather than a pure spatial length.

Transforming e_{geom} to CGS spatial units simply corresponds to expressing the displacement in centimetres ($1.602176634 \times 10^{-17} \text{ cm}$), rather than substituting the numerical value of the statcoulomb (4.8032×10^{-10}). The numerical equality $1 \text{ C} \rightarrow 1 \text{ m}$ is anchored specifically to the SI convention, where spatial displacement is measured in metres.

7.3 Consequence for the Planck Charge q_P

The Planck charge in EWT represents the maximum fundamental geometric amplitude of the BCC lattice:

$$q_P = \frac{e_{\text{geom}}}{\sqrt{\alpha_{\text{geom}}}}. \quad (55)$$

Evaluating this relation with $e_{\text{geom}} = 1.602176634 \times 10^{-19} \text{ m}$ yields:

$$q_P = 1.8755478 \times 10^{-18} \text{ m}. \quad (56)$$

The resulting quantity q_P is therefore an absolute spatial length of the lattice. The zero-calibration derivation of the statutory neutrino radius $r_\nu = q_P \cdot S_{\text{tot}}$ is a purely geometric relation mapping this amplitude scale to the lattice step.

8 Geometric Deficit and Density Hierarchy of the Neutrino

The neutrino soliton serves as the reference wave centre for the BCC vacuum. Its geometric radius r_ν , derived in Section 5, sets the fundamental length scale from which the EMC density hierarchy follows.

At this stage the EMC spacing λ_l is treated as a formal geometric parameter. Its numerical value is fixed later by the self-consistent gravitational identity of Section 11.6, where the Planck-length condition and the geometric expression for G form a closed algebraic system. The density hierarchy below therefore uses λ_l symbolically before that value is substituted.

This geometric saturation limit and the associated density hierarchy follow from the EWT granular density model developed by Yee [11].

1. Absolute Maximum Capacity ($N_{\nu,\max}$)

$$N_{\nu,\max} = \left(\frac{r_\nu}{\lambda_l}\right)^3 \approx 5.3004 \times 10^{54} \quad (57)$$

2. Statutory Background Density ($N_{\nu,\text{stat}}$)

$$N_{\nu,\text{stat}} = \left(\frac{r_\nu}{2\lambda_{le}}\right)^3 \approx 3.2986 \times 10^{52} \quad (58)$$

3. Effective Gravitational Density ($N_{\nu,\text{eff}}$)

The effective density inside the electron soliton is obtained from the statutory density and the geometric dilution factor X_{eff} :

$$N_{\nu,\text{eff}} = \frac{N_{\nu,\text{stat}}}{X_{\text{eff}}}, \quad (59)$$

where X_{eff} is the geometric dilution factor defined in Section 11.6.

8.1 Unified Geometric Coupling

The transition from the statutory background to the effective density is governed by the unified coupling operator C_{unif} and the geometric dilution factor X_{eff} , already introduced in Section 6. Both quantities are fixed by the BCC lattice geometry and the electron wave-centre count $K_{WC} = 10$.

The resulting effective gravitational density is

$$N_{\nu,\text{eff}} = \frac{N_{\nu,\text{stat}}}{X_{\text{eff}}} \approx 6.247 \times 10^{48}. \quad (60)$$

This value follows from the geometric dilution of the statutory vacuum by the soliton wave structure. The concept of such a density deficit and its role in gravitational emergence originates from the EWT volume-deficit models of Yee [11, 1]. In the present formulation the deficit is not fitted; it is a direct consequence of the same BCC geometry that determines α_{geom} , r_ν , and λ_l .

The extreme dilution of approximately 99.98% relative to $N_{\nu,\text{stat}}$ explains the weakness of gravity as a pressure deficit within the high-density medium.

9 Speed of Light as the Metric of Spacetime

Before deriving the gravitational constant, the role of the speed of light c must be addressed. In the conventional formulation, c is treated as a fundamental constant, empirically determined and independent of any underlying structure. Within the EWT framework, this status is revised: c is not an independent parameter but the intrinsic metric conversion factor between the spatial and temporal dimensions of the BCC lattice, determined entirely by the discrete geometry of the elastic medium.

9.1 The Causal Structure of the BCC Lattice

The BCC spacetime lattice is characterised by three distinct density levels that define its structural states:

- **Statutory density** $N_{\nu,\text{stat}}$: the equilibrium density of the undisturbed vacuum, derived in Eq. (58).
- **Effective density** $N_{\nu,\text{eff}}$: the reduced density inside a soliton (matter-occupied region), derived in Eq. (79).
- **Maximum density** $N_{\nu,\text{max}}$: the absolute saturation limit of the lattice, given by Eq. (57).

The lattice is characterised by two fundamental geometric scales:

- **The spatial step**: the inter-constituent distance λ_l , which follows from the packing geometry of the Elastic Medium Constituents (EMCs) and is identified with the Planck length. Its value is fixed by the statutory neutrino radius r_ν and the Eulerian dilution factor $2e$ through the relation

$$\lambda_l = \frac{r_\nu}{2e N_{\nu,\text{stat}}^{1/3}}, \quad (61)$$

where $N_{\nu,\text{stat}}$ is the statutory background density. Equivalently, it appears in the uncorrected neutrino wavelength

$$\lambda_{\text{uncorr}} = 2q_P e^2, \quad (62)$$

which, after the geometric correction g_v , yields the statutory radius $r_\nu = \lambda_{\text{uncorr}}/g_v$. The spatial step λ_l is thus a derived property of the BCC lattice, not an independent input.

- **The temporal step**: the fundamental cycle time t_p , which represents the characteristic oscillation period of an EMC in the lattice. In the spring-mass representation of the BCC medium, this period is determined by the effective elastic constant of the lattice and the Planck mass, both of which are themselves consequences of the lattice geometry (Sec. 31.8). The formal derivation of t_p from first principles is a natural direction for future research; for the present purposes, it suffices to note that t_p is a structural invariant of the lattice, defined by the discrete causal structure itself.

The Planck charge q_P is not an independent parameter but the fundamental amplitude scale of the BCC lattice. It is related to the elementary charge e through the fine-structure constant:

$$e^2 = \alpha q_P^2. \quad (63)$$

This relation follows directly from the definitions of α and q_P , and it holds identically in both SI units (where $q_P = \sqrt{4\pi\epsilon_0\hbar c}$) and in the EWT amplitude representation (where both charges are measured in meters). It expresses the fact that the elementary charge is the lattice amplitude reduced by the coupling factor α .

9.2 Axiom: The Maximum Propagation Speed

The following fundamental axiom is posited:

In the BCC lattice at the statutory density $N_{\nu,\text{stat}}$, there exists a maximum propagation speed c for any disturbance. This speed is a structural property of the lattice, determined by its density and elastic response, and is independent of the amplitude or frequency of the disturbance.

This axiom is physically motivated: in any elastic medium with finite density and stiffness, there exists a speed of propagation (the speed of sound in the medium). The BCC lattice, as a discrete elastic medium, is no exception. The speed c is therefore not an empirical input but a derived property of the lattice, on the same footing as the speed of sound in a solid.

In any discrete causal manifold, the maximum propagation speed is fixed by the ratio of the spatial step to the temporal step:

$$c \equiv \frac{\lambda_l}{t_p}. \quad (64)$$

Because the speed of light is the fixed metric conversion $c \equiv \lambda_l/t_p$, specifying λ_l therefore fixes the temporal step $t_p = \lambda_l/c$. In the natural units of the lattice ($c = 1$) the spatial and temporal steps are identical.

This relation is not a postulate nor an empirical input; it is the definition of the causal structure of the lattice, expressing the unique conversion factor between the temporal and spatial coordinates in the metric:

$$ds^2 = -c^2 dt^2 + dx^2 + dy^2 + dz^2. \quad (65)$$

9.3 Consequences: Density-Dependent Propagation Speed

The axiom immediately implies that the propagation speed in the lattice is a function of the local EMC density:

$$v(N_\nu) = c \cdot f\left(\frac{N_\nu}{N_{\nu,\text{stat}}}\right), \quad (66)$$

where the function f encodes the elastic response of the lattice. The three density regimes yield distinct physical consequences:

- **In vacuum** ($N = N_{\nu,\text{stat}}$), the speed is exactly c , explaining the constancy of the speed of light in free space. The vacuum is defined as the region where the lattice maintains its statutory density.
- **In matter** ($N = N_{\nu,\text{eff}} < N_{\nu,\text{stat}}$), the speed is reduced, providing a fundamental mechanism for refraction and the slowing of light in media. This is the origin of the refractive index $n = c/v > 1$.
- **In the extreme EMC packaging density limit** ($N \rightarrow N_{\nu,\text{max}}$), the lattice approaches its elastic saturation limit. The propagation properties of this regime — relevant to early-universe cosmology — lie beyond the scope of the present analysis and are reserved for future work.

The ratio of the spatial step to the temporal step in the lattice gives the maximum propagation speed in the statutory vacuum as defined in eq. 64.

Remark on the status of t_p in natural units. The preceding analysis noted that the formal derivation of the temporal step t_p from the elastic constants of the BCC medium remains an open objective in SI units. Within the natural units of the lattice ($c = 1$), however, this problem does not arise: the defining relation $c \equiv \lambda_l/t_p$ immediately yields

$$t_p = \frac{\lambda_l}{c} = \lambda_l, \quad (67)$$

so the lattice possesses a *single* fundamental scale λ_l , and t_p is not an independent quantity but an alternative labelling of the same geometric step in the temporal direction. The open objective

of deriving the *numerical value* of c in SI units (metres per second) is therefore reframed as a question about the external convention that separates the metre from the second — not a question about the internal structure of the lattice. Within EWT, the lattice is characterised by one geometric primitive: λ_l .

Remark on the Dimensional Status of c :

The speed of light c does not appear in any of the dimensionless predictions of the EWT framework (the fine-structure constant α , the anomalous magnetic moments a_e , a_μ , a_τ , or the lepton mass ratios). It enters only when expressing dimensional SI quantities such as G or m_e , where it plays the role of a *metric conversion factor* between the spatial step λ_l and the temporal step t_p of the BCC lattice ($c \equiv \lambda_l/t_p$). In natural units ($c = 1$), the EWT framework is parameter-free: all physical predictions reduce to dimensionless ratios determined by BCC lattice topology alone. The appearance of c in SI expressions reflects the unit system's convention for separating spatial and temporal dimensions, not an independent physical input to the theory. Its numerical value in SI units is therefore a consequence of the arbitrary definitions of the metre and the second, not a fundamental property of the vacuum lattice. This status is consistent with the SI 2019 definition, in which c is an exact definitional constant rather than a measured quantity.

With this geometric interpretation of the speed of light, the base gravitational scaling may now be derived.

10 The Physical Origin of Relativity: Wave-Mechanical Lorentz Contraction

As established in Sec. 9, the speed of light c is the structural conversion factor between the spatial and temporal steps of the BCC lattice. The present section demonstrates that this same value is measured by all inertial observers, not as a primary postulate, but as a mechanical consequence of the wave-structure of matter. The derivation follows the Doppler-based approach developed by Gabriel LaFrenière [51] and historically proposed by H.A. Lorentz [52] and G.F. FitzGerald [53] as a physical explanation of the Michelson–Morley null result.

10.1 Doppler-Induced Compression of the Standing-Wave Soliton

In EWT, a particle is a standing wave (soliton) formed by the interference of incoming and outgoing waves in the BCC medium. When such a soliton moves relative to the lattice with velocity $v = \beta c$, the underlying waves undergo an anisotropic Doppler shift.

Let λ_0 be the wavelength of the soliton at rest. For a soliton moving at speed v , the forward-propagating wave is compressed to λ_f and the backward wave is expanded to λ_b :

$$\lambda_f = \lambda_0(1 - \beta), \quad \lambda_b = \lambda_0(1 + \beta). \quad (68)$$

The resulting standing-wave wavelength in the direction of motion, λ' , is the geometric mean of the two, modulated by the requirement of phase synchronization across the soliton's diameter:

$$\lambda' = \sqrt{\lambda_f \cdot \lambda_b} = \lambda_0 \sqrt{1 - \beta^2} = \frac{\lambda_0}{\gamma}, \quad (69)$$

where $\gamma = (1 - \beta^2)^{-1/2}$ is the Lorentz factor. This result shows that the physical distance between the wave-center nodes — the “size” of the particle — must contract.

10.2 The Michelson–Morley Null Result as a Mechanical Necessity

The invariance of the measured speed of light in the Michelson–Morley experiment is usually interpreted as a property of spacetime. In EWT, it is a property of the **measuring apparatus**. Because all matter (including the interferometer arms) consists of waves in the BCC lattice, the apparatus itself undergoes a physical contraction:

1. **Longitudinal arm ($L_{\parallel}$):** Contracts by exactly $1/\gamma$ owing to the wave-node compression derived above.

2. **Transverse arm ($L_{\perp}$):** Remains unchanged in length, but the wave path is elongated by γ because of the diagonal trajectory through the medium.

The total round-trip times for the longitudinal path ($t_{\parallel}$) and transverse path ($t_{\perp}$) become identical:

$$t_{\parallel} = \frac{L_0/\gamma}{c-v} + \frac{L_0/\gamma}{c+v} = \frac{2L_0}{c} \gamma, \quad t_{\perp} = \frac{2L_0 \sqrt{1 + (v\gamma/c)^2}}{c} = \frac{2L_0}{c} \gamma. \quad (70)$$

Thus $t_{\parallel} = t_{\perp}$. The null result is not due to the absence of a medium, but because the wave-mechanical contraction of the soliton perfectly masks the motion through the lattice.

Remark on time dilation: Substituting $c = 1$ (the natural units of the BCC lattice, Sec. 9) into the Michelson–Morley round-trip times yields $t_{\parallel} = t_{\perp} = 2L_0\gamma$ for both arms. In the lattice rest frame the round-trip time is therefore $t = 2L_0\gamma$. A clock carried by the moving interferometer, whose rate is dilated by $1/\gamma$, measures

$$t' = \frac{t}{\gamma} = 2L_0,$$

the same elapsed time as an identical apparatus at rest in the BCC lattice. The oscillation period of the soliton therefore dilates to $T' = \gamma T_0$, with the observed tick rate scaling as $f' = f_0/\gamma$ — the temporal counterpart of the spatial contraction derived above.

11 The Geometric Identity of G : Derivation and Consistency

The core hypothesis of the Energy Wave Theory (EWT) is that the gravitational constant $\mathbf{G}$ is not a fundamental constant but an **emergent, calculated geometric identity** derived from the electron’s geometric parameters and the BCC lattice topology. This section derives a precise $\hbar$ -independent formula for $\mathbf{G}$ and explains the physical meaning of its scaling terms.

11.1 Derivation of G from Electron Scaling (The $\hbar$ -Independent Base)

Traditional definitions of $\mathbf{G}$ rely on Planck units, which inherently contain the reduced Planck constant ($\hbar$). EWT posits that the gravitational constant is derived from the scaling of the electron’s properties ($\mathbf{m}_e, \mathbf{r}_e$), which are geometric wave-based quantities.

958 The Base Gravitational Scaling ($\mathbf{G}_{\text{Base}}$)

959 The fundamental scaling constant for gravity is established by the ratio of energy density to
960 mass, using the speed of light (c), the classical electron radius ($\mathbf{r}_e$), and the electron mass ($\mathbf{m}_e$):

$$\mathbf{G}_{\text{Base}} = \frac{c^2 \mathbf{r}_e}{\mathbf{m}_e} \approx 2.780252 \times 10^{32} \text{ m}^3 \text{kg}^{-1} \text{s}^{-2} \quad (71)$$

961 This term is the sole source of the required physical units for the gravitational constant:

$$\left[\frac{c^2 \mathbf{r}_e}{\mathbf{m}_e} \right] = \frac{(\text{m} \cdot \text{s}^{-1})^2 \cdot \text{m}}{\text{kg}} = \text{m}^3 \cdot \text{kg}^{-1} \cdot \text{s}^{-2} \quad (72)$$

962 All subsequent geometric factors in the final identity for $\mathbf{G}$ are therefore dimensionless.

963 The $\hbar$ -Independence of $\mathbf{G}_{\text{Base}}$

964 The form of $\mathbf{G}_{\text{Base}}$ is inherently independent of $\hbar$. This is demonstrated by substituting the
965 standard definition of the classical electron radius ($\mathbf{r}_e = \alpha \frac{\hbar}{\mathbf{m}_e c}$) into the conventional expression
966 for the gravitational constant based on particle properties ($\mathbf{G}_{\text{Base}} = \alpha \frac{\hbar c}{\mathbf{m}_e^2}$):

$$\mathbf{G}_{\text{Base}} = \alpha \frac{\hbar c}{\mathbf{m}_e^2} = \alpha \frac{\left(\frac{\mathbf{r}_e \mathbf{m}_e c}{\alpha} \right) c}{\mathbf{m}_e^2} = \frac{\mathbf{r}_e \mathbf{m}_e c^2}{\mathbf{m}_e^2} = \frac{c^2 \mathbf{r}_e}{\mathbf{m}_e} \quad (73)$$

967 The cancellation of $\hbar$ confirms that the base gravitational scale is a classical geometric prop-
968 erty of the electron soliton ($\mathbf{m}_e, \mathbf{r}_e, c$), rather than a purely quantum effect.

969 Status of m_e and r_e in the Present Work

970 In the current formulation, m_e , r_e , and c are the metric and structural inputs used to anchor the
971 electron soliton. The radius r_e is related to the geometric neutrino anchor by

$$r_e = 100 r_\nu, \quad (74)$$

972 while m_e is constrained by the $E \propto r^5$ scaling law discussed in Section 15.1. Their full derivation
973 from the BCC lattice alone is a central open objective.

974 The subsequent derivation of G therefore contains no free dimensional parameters: the only
975 inputs are r_e , m_e , c , and the elementary charge amplitude e . In natural units ($c = 1$), the
976 number of independent inputs reduces further, and the gravitational constant becomes a purely
977 geometric prediction.

978 The Final Geometric Identity for G

979 The observed gravitational constant $\mathbf{G}$ is obtained by scaling $\mathbf{G}_{\text{Base}}$ with the total dimensionless
980 geometric weakness factor. In the zero-calibration formulation, this factor is built solely from
981 the geometric soliton core A_π , the effective BCC stiffness N_{geom} , the electron wave-centre count
982 $K_{WC} = 10$, and the effective volume deficit $N_{\nu, \text{eff}}$.

983 The geometric identity for $\mathbf{G}$ is therefore:

$$\mathbf{G} \equiv \mathbf{G}_{\text{Base}} \cdot \frac{1}{A_\pi} \cdot \left(\frac{1}{N_{\text{geom}} A_\pi} \right)^3 \cdot \frac{1}{K_{WC} \sqrt{N_{\nu, \text{eff}}}} \quad (75)$$

984 The only dimensional factor in this expression is $\mathbf{G}_{\text{Base}}$. Every other factor is a pure geometric
985 number determined by the BCC lattice and the electron soliton.

Geometric Dependence on λ_l

The effective volume deficit $N_{\nu,\text{eff}}$ depends on the EMC spacing λ_l through the statutory density:

$$N_{\nu,\text{eff}} = \frac{N_{\nu,\text{stat}}}{X_{\text{eff}}}, \quad N_{\nu,\text{stat}} = \left(\frac{r_\nu}{2e\lambda_l} \right)^3. \quad (76)$$

Thus Eq. (75) defines G as a function of the formal parameter λ_l . This is not a fault of the model; it is the expected self-consistent coupling between gravity and the fundamental lattice spacing. The value of λ_l will be fixed in Section 11.6 by combining the geometric G with the Planck-length definition.

Compact ϵ_M Form

Using the geometric magnetic deficit

$$\epsilon_M = \frac{1}{N_{\text{geom}}\pi^3}, \quad (77)$$

Eq. (75) can also be written as

$$\mathbf{G} = \frac{\mathbf{G}_{\text{Base}}}{A_\pi} \left(\frac{\epsilon_M \pi^3}{A_\pi} \right)^3 \frac{1}{K_{WC} \sqrt{N_{\nu,\text{eff}}}}. \quad (78)$$

This form shows that gravity and the magnetic deficit share the same geometric origin. It also makes clear that the weakness of gravity is a consequence of the cubic suppression by ϵ_M/A_π , followed by the surface projection $N_{\nu,\text{eff}}^{-1/2}$.

11.2 The Gravitational Identity Flow: From Electromagnetic Base to Surface Transition

The derivation of the Gravitational Constant $\mathbf{G}$ follows a systematic, multi-step geometric flow. This process resolves the scale disparity between electromagnetic and gravitational interactions by accounting for the displacement of the medium's constituents within the Soliton structure.

Step 1: Establishment of the EM Base ($\mathbf{G}_{\text{Base}}$): The process is initiated by defining $\mathbf{G}_{\text{Base}}$, which links the classical electron radius (r_e) and its mass (m_e). This term represents the maximum potential coupling scale before geometric dilution is applied.

Step 2: Geometric Stability Factor (A_π^{-1}): The first scaling step introduces the normalization to the static wavelength-to-radius ratio of the Soliton.

Step 3: Nodal Push-Out Potential (Ω_{Push}): This phase incorporates the volumetric redistribution of energy. By applying the cubic factor $(N_{\text{geom}} A_\pi)^{-3}$, the model quantifies the structural dilution of the soliton in the BCC lattice.

Step 4: Geometric Surface Transition ($\mathbf{T}_{\text{Surface}}$): The conversion of the internal displacement into the observable gravitational force is governed by the factor $(K_{WC} \cdot \sqrt{N_{\nu,\text{eff}}})^{-1}$. Here, $K_{WC} = 10$ Wave Centers act as the primary operators of the Push-Out mechanism, evacuating the medium from its statutory density to the effective density of the matter-occupied region.

1016 The square root ($N_{\nu,\text{eff}}^{-1/2}$) serves as the surface transition operator, projecting the three-
 1017 dimensional internal packing deficit onto the two-dimensional effective surface of the Soli-
 1018 ton. Gravity is thus revealed as an emergent, pressure-driven phenomenon acting on this
 1019 surface.

1020 **Step 5: Final Gravitational Identity:** The integration of these modular scales yields the geo-
 1021 metric identity for $\mathbf{G}$ given in Eq. (75).

1022 Physical Significance

1023 The extreme weakness of gravity is a direct consequence of this hierarchical dilution. While the
 1024 Soliton's mass is defined by high-frequency energy localization, the gravitational constant $\mathbf{G}$ is
 1025 a measure of the effective displacement deficit. The transition from the statutory vacuum to the
 1026 diluted interior creates a pressure gradient, whereby the universal medium strives for equilibrium
 1027 by filling the geometric void.

1028 The Holographic Nature and Experimental Implications

1029 The presence of the square root ($\sqrt{N_{\nu,\text{eff}}}$) in the final identity reinforces the principle that
 1030 gravity is an emergent surface phenomenon. By projecting the three-dimensional volumetric
 1031 deficit onto a two-dimensional boundary, the model aligns with holographic and thermodynamic
 1032 interpretations of gravitational flux.

1033 11.3 Formalization of Soliton Geometric Density $\rho(r)$

1034 To avoid ambiguity between energy density ρ_E and the packing density, the internal density
 1035 function $\rho(r)$ is introduced to represent the local distribution of Elastic Medium Constituents
 1036 (EMC) within the Soliton. This profile is physically important, but the numerical value of the
 1037 effective deficit is fixed algebraically by the unified coupling described in Section 8:

$$N_{\nu,\text{eff}} = \frac{N_{\nu,\text{stat}}}{X_{\text{eff}}}, \quad (79)$$

1038 where X_{eff} is the geometric dilution factor from Section 6. The radial profile $\rho(r)$ is compatible
 1039 with this value, and the structural parameters of any assumed profile do not enter as free inputs
 1040 into the gravitational identity.

1041 11.4 Numerical Verification and the Unitary Mapping Operator $\mathcal{U}$

1042 Numerical Consistency

1043 Using the geometric value N_{geom} and the self-consistent λ_l from Eq. (94), the geometric identity
 1044 Eq. (75) gives

$$G_{\text{EWT}} \approx 6.67752 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}. \quad (80)$$

1045 The corresponding CODATA 2022 value is

$$G_{\text{CODATA}} = 6.674305 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}, \quad (81)$$

1046 so the relative deviation is approximately 0.048%.

1047 This is a genuine geometric prediction.

1048 The Unitary Mapping Operator $\mathcal{U}_{\text{Geom} \rightarrow G}$

1049 The final step in the geometric formalization is the introduction of the **Unitary Mapping Op-**
 1050 **erator** $\mathcal{U}$, which symbolically represents the non-linear transformation of the Soliton's internal
 1051 geometry into the macroscopic gravitational constant:

$$\mathcal{U}_{\text{Geom} \rightarrow G} : A_\pi, N_{\text{geom}}, N_{\nu, \text{eff}} \mapsto G \equiv \frac{c^2 r_e}{m_e} \cdot \Omega_G. \quad (82)$$

1052 This operator formalizes that G is not a standalone fundamental constant but a direct manifes-
 1053 tation of the geometric scaling applied to the electron's $\hbar$ -independent base.

1054 11.5 Formal Definition of the Operator $\mathcal{U}$: Mapping Geometric Pa- 1055 rameters to G

1056 To codify the micro-macro transformation in a rigorous mathematical manner, a nonlinear func-
 1057 tional operator $\mathcal{U}$ is defined:

$$\mathcal{U} : \mathcal{D} \subset \mathbb{R}^n \rightarrow \mathbb{R}, \quad \mathcal{U}(\mathbf{P}) \mapsto G \quad (83)$$

1058 where the parameter vector $\mathbf{P}$ contains all relevant microscopic variables of the model. This
 1059 functional approach expresses the gravitational constant G as a product of the **Base Soliton**
 1060 **Identity** ($\mathbf{G}_{\text{Base}}$) and a **Geometric Scaling Functional** (F_{geom}):

$$\mathcal{U}(\mathbf{P}) = \mathbf{G}_{\text{Base}} \cdot F_{\text{geom}}(\mathbf{P}), \quad \mathbf{G}_{\text{Base}} = \frac{c^2 \mathbf{r}_e}{\mathbf{m}_e}. \quad (84)$$

1061 The explicit geometric form used in this work is

$$\mathcal{U}(\mathbf{P}) = \frac{c^2 r_e}{m_e} \cdot \frac{1}{A_\pi} \cdot \left(\frac{1}{N_{\text{geom}} A_\pi} \right)^3 \cdot \frac{1}{K_{WC} \sqrt{N_{\nu, \text{eff}}}} \quad (85)$$

1062 where A_π is the core geometric factor and $N_{\nu, \text{eff}}$ is the effective volume deficit derived from the
 1063 unified coupling C_{unif} .

1064 Properties of the Operator $\mathcal{U}$

- 1065 • **Non-linearity:** The mapping is characterized by cubic scaling of the stiffness factor and
 1066 an inverse square-root dependency on the effective constituent count.
- 1067 • **Monotonicity:** For a fixed statutory background, any increase in $N_{\nu, \text{eff}}$ results in a de-
 1068 terministic decrease in the emergent value of G .
- 1069 • **Zero calibration:** The operator contains only geometric and structural inputs: $A_\pi, N_{\text{geom}},$
 1070 $K_{WC}, N_{\nu, \text{eff}}$.

1071 the G implementation result showed in table 4.

Table 4: Numerical Verification of the Gravitational Constant G within the Geometric EWT Framework

Parameter / Result	Numerical Value	Source and Physical Context
I. Base Soliton Identity	$2.7802522591 \times 10^{32}$	$G_{\text{Base}} = \frac{c^2 r_e}{m_e}$. Fundamental Soliton base.
II. Raw Geometry Value	$6.6804369615 \times 10^{-11}$	$G_{\text{EWT, raw}}$. Pure $K + 1$ projection; error 0.09187% relative to CODATA.
III. Geometric Result	$6.6775199755 \times 10^{-11}$	Zero-calibration geometric G . Uses L_p^{geom} and N_{geom} ; error 0.04817%.
IV. CODATA Target	$6.6743050000 \times 10^{-11}$	Experimental reference value (CODATA 2022).

Note: All values are expressed in units of $m^3 kg^{-1} s^{-2}$.

11.6 Geometric Emergence of the Planck Length

The Planck length in EWT is identified with the fundamental EMC spacing λ_l . In the zero-calibration formulation, λ_l is not taken from experiment; it follows from the geometric identity for the gravitational constant and the Planck-length definition.

Before the effective dilution factor X_{eff} can be written down, the unified coupling operator C_{unif} that governs it must be derived. The operator bridges the raw geometric projection of the wave centres onto the BCC lattice with the electromagnetic response of the medium, and its three terms each have a distinct structural origin.

The Fundamental Geometric Ratio: C_{Raw}

Prior to any electromagnetic coupling, the model identifies the pure interaction ratio between the soliton's wave centres and the background lattice. This factor,

$$C_{\text{Raw}} = 1 + \frac{1}{K_{WC}}, \quad (86)$$

is the “ideal” geometric scaffold of the displacement. The term $1/K_{WC}$ is the harmonic reciprocal of the $K_{WC} = 10$ wave centres; it normalises the global volume deficit to the individual node's displacement capacity. The additive $+1$ is the statutory equilibrium of the medium, the unperturbed vacuum state that the correction is applied on top of. For $K_{WC} = 10$ this gives $C_{\text{Raw}} = 1.1$, the baseline efficiency of the vacuum displacement before any elastic or fine-structure corrections.

The Lattice Projection Factor L_p^{geom}

The natural geometric starting point for the interfacial correction is the ideal BCC projection factor

$$L_p^{\text{geom}} = \frac{2}{\sqrt{3}} \approx 1.154700538, \quad (87)$$

which arises as the inverse of the dimensionless nearest-neighbour distance ($d_{NN}/a = \sqrt{3}/2$) in the $Im\bar{3}m$ lattice. Its combination $\pi L_p^{\text{geom}} = 2\pi/\sqrt{3}$ is the dimensionless fundamental wavevector

for a standing wave along the [111] body diagonal, capturing the projection of the 3D lattice onto the 2D soliton interface.

The Unified Bridge: C_{unif}

The full operator adds the electromagnetic coupling across the lattice interface:

$$C_{\text{unif}} = \frac{1}{K_{WC}} + 1 + \frac{\alpha_{\text{geom}}}{\pi L_p^{\text{geom}}}. \quad (88)$$

The three terms correspond to the three distinct contributions of the medium's response:

- **Harmonic reciprocal** $1/K_{WC}$: the specific contribution of the $K_{WC} = 10$ wave centres, ensuring that the global deficit is normalised to the individual node's displacement capacity.
- **Unitary baseline** $+1$: the statutory equilibrium of the medium, the unperturbed vacuum state. It ensures the correction is additive to the existing background density $N_{\nu,\text{stat}}$, maintaining continuity of the elastic field.
- **Interfacial tension bridge** $\alpha_{\text{geom}}/(\pi L_p^{\text{geom}})$: the electromagnetic strain coupled through the lattice projection factor and the geometric π . This term scales the large-scale volumetric deficit down to the interfacial tension at the soliton boundary.

With C_{unif} established, the effective dilution factor follows directly:

$$X_{\text{eff}} = \frac{A_\pi 3 K_{WC} \sqrt{2}}{C_{\text{unif}}}, \quad (89)$$

where $A_\pi = 4\pi^3 + \pi^2 + \pi$ is the geometric core of the soliton. The numerator counts the active geometric budget of the soliton (the core geometry A_π , the three spatial dimensions, the K_{WC} wave centres, and the $\sqrt{2}$ face-diagonal coupling); the denominator C_{unif} is the structural impedance of the medium through which that budget is projected. The ratio is the geometric dilution of the statutory background density inside the soliton.

The self-consistent derivation of λ_l starts from the Planck-length definition,

$$\lambda_l = \sqrt{\frac{\hbar_{\text{geom}} G_{\text{geom}}}{c^3}}, \quad (90)$$

in which $\hbar_{\text{geom}}$ is given by Eq. (50), while G_{geom} is the geometric gravitational constant whose explicit form is derived in Section 11. In the present section it is sufficient to use its geometric structure,

$$G_{\text{geom}} = \frac{G_{\text{Base}}}{A_\pi} \left(\frac{1}{N_{\text{geom}} A_\pi} \right)^3 \frac{1}{K_{WC} \sqrt{N_{\nu,\text{eff}}}}, \quad (91)$$

where

$$G_{\text{Base}} = \frac{c^2 r_e}{m_e}. \quad (92)$$

The effective density $N_{\nu,\text{eff}}$ is related to the statutory density $N_{\nu,\text{stat}}$ by

$$N_{\nu,\text{eff}} = \frac{N_{\nu,\text{stat}}}{X_{\text{eff}}}, \quad N_{\nu,\text{stat}} = \left(\frac{r_\nu}{2e\lambda_l} \right)^3. \quad (93)$$

Substituting these expressions into the Planck-length definition cancels m_e , c , and the common A_π factors. Solving the resulting algebraic equation for λ_l gives

$$\lambda_l = \left[\frac{r_e^2 \sqrt{X_{\text{eff}}}}{\alpha_{\text{geom}} A_\pi^4 N_{\text{geom}}^3 K_{WC} \left(\frac{r_\nu}{2e} \right)^{3/2}} \right]^2, \quad (94)$$

where e is Euler's number. Normalized form is:

$$\lambda_l = \left[\frac{r_e^2 \sqrt{\frac{3\sqrt{2} A_\pi}{C_{\text{unif}}}}}{\alpha_{\text{geom}} A_\pi^4 N_{\text{geom}}^3 \sqrt{K_{WC}} \left(\frac{r_\nu}{2e} \right)^{3/2}} \right]^2. \quad (95)$$

Equation (95) substitutes the definition of X_{eff} directly, so that the wave-centre count enters once through $\sqrt{K_{WC}}$ rather than twice. The fine-structure constant appears twice in the expression, but in two distinct roles: the leading α_{geom} is the electromagnetic coupling of the soliton itself, while the α_{geom} inside C_{unif} is the electromagnetic strain across the soliton–vacuum interface. The two appearances are not a redundancy; they reflect the two surfaces through which the same coupling is projected.

Planck introduced this length as a formal dimensional combination of $\hbar$, G , and c , without reference to a microscopic substrate. In the present framework, the same quantity appears as the self-consistent solution of the geometric BCC equations, thereby providing a physical origin for the scale that Planck treated only as a formal unit.

Equations (94) and (95) are therefore not an independent ansatz. They are the direct consequence of combining the geometric fine-structure constant, the geometric gravitational scaling, and the Planck-length definition into a single closed algebraic constraint. Here e is Euler's number and r_ν is the geometric neutrino radius derived in Section 5.

Numerically,

$$\lambda_l \approx 1.6166464 \times 10^{-35} \text{ m}. \quad (96)$$

The CODATA value is

$$\lambda_{l,\text{CODATA}} = 1.6162 \times 10^{-35} \text{ m}, \quad (97)$$

so the relative deviation is approximately 0.0276%. This is a genuine geometric prediction.

X_{eff} is not a free parameter; it is the geometric bridge that connects the statutory EMC background to the effective soliton deficit. It encapsulates how $K_{WC} = 10$, the ideal BCC projection, and the geometric core A_π jointly dilute the vacuum density inside matter.

11.7 Consistency with the Planck Charge

The geometric Planck charge was derived earlier as

$$q_P = \frac{e}{\sqrt{\alpha_{\text{geom}}}}. \quad (98)$$

With the above values,

$$q_P \approx 1.8755478 \times 10^{-18} \text{ m}, \quad (99)$$

which matches the expected amplitude scale.

The fact that λ_l , q_P , and $\hbar_{\text{geom}}$ all emerge from the same BCC geometry confirms that the model does not treat the Planck scale as an external boundary but as an internal consequence of the lattice structure.

11.8 Electromagnetic Consistency Check

In SI units, the relation between charge, $\hbar$, and α is

$$e^2 = 4\pi\epsilon_0\hbar c\alpha. \quad (100)$$

Substituting the geometric expression for $\hbar$, Eq. (50), cancels α_{geom} identically and yields

$$e^2 = 4\pi\epsilon_0 m_e c^2 r_e. \quad (101)$$

In the natural units of the BCC lattice ($c = 1$, $4\pi\epsilon_0 = 1$; see Section 9), this reduces to the purely geometric identity

$$e^2 = m_e r_e. \quad (102)$$

This is not an independent derivation of e , since m_e and r_e are inputs, but it is a powerful internal consistency test: it shows that the electromagnetic scale (e) is rigidly tied to the soliton's mass and radius, with no residual dependence on $\hbar$ or α .

11.9 Status of Inputs

The only experimental inputs in model are

$$r_e, \quad m_e, \quad c, \quad e. \quad (103)$$

In natural units ($c = 1$; see Section 9), c is a unit conversion and m_e is geometrically tied to r_e through the r^5 scaling law. In SI units the relation between charge, $\hbar$, and α is $e^2 = 4\pi\epsilon_0\hbar c\alpha$. In the EWT framework ϵ_0 is not an independent input: together with μ_0 it is fixed by the metric condition $c^2 = 1/(\mu_0\epsilon_0)$ and therefore carries no new geometric information. No calibrated parameters appear in this section.

The quantities G_{geom} , λ_l , $\hbar_{\text{geom}}$, and α_{geom} are not independent predictions. They form a single closed algebraic system fixed by the BCC lattice geometry, the electron soliton inputs r_e, m_e, c, e , and the geometric self-consistency condition

$$\lambda_l = \sqrt{\frac{\hbar_{\text{geom}} G_{\text{geom}}}{c^3}}. \quad (104)$$

Solving this system is the geometric self-consistency of the model, not a stepwise derivation of one constant from another.

In the standard framework the Planck-length relation links three independent empirical constants. In the present model the same relation emerges as an internal geometric condition, because both $\hbar_{\text{geom}}$ and G_{geom} are already determined by the BCC lattice. The Planck scale is therefore not imported from the Standard Model; it is reconstructed as a consequence of the underlying geometry.

11.10 Duality of Soliton Density

Presented model inherently describes a fundamental duality of densities within the Soliton. The localized energy density ρ_E inside the Soliton must be higher than the surrounding medium, due to localized standing wave energy, to account for its mass $E = mc^2$. Conversely, the geometric packing density function $\rho(r)$ must be lower than the uniform density of the surrounding space, creating the measurable packing deficit $N_{\nu, \text{eff}}$.

This dichotomy means the Soliton is a region of high energy but low geometric packing efficiency. The dynamic processes described in models unifying mass and charge as wave energy

[26, 29, 15] can be hypothesized to be the cause of the packing density deficit of the EMC, by continuously forcing the Soliton's wave components outward, defining the effective volume and maintaining its stability.

The accompanying constant geometric term A_π^{-1} is specifically interpreted here as the Geometric Stability Factor. This factor quantifies the fixed boundary ratio resulting from the continuous dynamic process of outward forcing on the Soliton's internal wave structure.

This definition of density introduces a central principle of the Enhanced EWT Model: gravity is an emergent, pressure-driven phenomenon that acts on the Soliton's effective surface, as the universal medium strives for equilibrium by filling this geometric deficit. This provides a basis for interpreting gravity as a push force induced by a vacuum pressure gradient [34].

11.11 The Degraded EMC Wall: Spherical Masking and Isotropy

A fundamental puzzle in any geometric theory of matter is how an asymmetric, rotating not perfect spherical soliton can generate a perfectly isotropic gravitational field. The resolution lies in a structural boundary layer that surrounds every soliton – the **Degraded EMC Wall**.

The Degraded EMC Wall is a spherical shell of finite thickness located at a radius r_{mask} where the packing density of Elastic Medium Constituents (EMCs) transitions from the diluted interior of the soliton toward the statutory background density $N_{\nu, \text{stat}}$. The wall is a region of **elevated EMC density** caused by the active push-out mechanism: as the soliton's standing waves displace EMCs from the interior, these constituents accumulate just outside the soliton boundary, creating a local density peak:

$$\rho_{\text{EMC}}(r_{\text{mask}}^-) < \rho_{\text{EMC}}(r_{\text{wall}}), \quad (105)$$

where r_{wall} denotes a point located strictly inside the Degraded EMC Wall.

The exact value of this peak relative to $N_{\nu, \text{stat}}$ depends on the balance between the internal push-out force and the external statutory pressure of the surrounding BCC lattice. In ordinary solitons this equilibrium yields a moderate peak, while in extreme gravitational environments the peak can become very large.

The Degraded EMC Wall acts as a mandatory transducer between the asymmetric core and the exterior. Because the individual EMCs are spherical units, the BCC lattice can only achieve a stable, static interface with the statutory background by adopting a spherical shape. The wall therefore enforces a spherical boundary condition on the energy density distribution emerging from the asymmetric core. The effect can be quantified by the Isotropy Operator $\mathcal{I}$ (introduced in Section 24.7).

11.11.1 Open question: Wall density relative to statutory background

The precise value of the peak EMC density within the Degraded EMC Wall, ρ_{wall} , relative to the statutory background $N_{\nu, \text{stat}}$, remains undetermined. Two scenarios are physically permissible:

- **Scenario A (asymptotic approach):** $\rho_{\text{wall}} < N_{\nu, \text{stat}}$, with the density increasing monotonically from the soliton interior toward the statutory value.
- **Scenario B (local maximum):** $\rho_{\text{wall}} > N_{\nu, \text{stat}}$, corresponding to a genuine local density peak caused by the active push-out.

Both scenarios are compatible with the integral determination of $N_{\nu, \text{eff}}$ and with the derivation of G . Future high-resolution simulations may discriminate between them. For the purpose of the present work, we treat the wall density ratio as an open parameter $\eta = \rho_{\text{wall}}/N_{\nu, \text{stat}}$.

12 Local Metric Phenomena from EMC Lattice Deformation

12.1 The Two Faces of EMC Displacement: Speed and Trajectory

In the Enhanced EWT framework, light propagation involves two distinct, though coupled, consequences of the EMC lattice strain field. They are frequently conflated in continuous medium optics, so it is necessary to state them with structural precision.

1. **Coordinate phase velocity via lattice strain.** In natural lattice units ($c \equiv 1$), the statutory speed of light is defined by the undisturbed cell ratio $c = \lambda_l/t_p = 1$. Around a matter soliton, the radial displacement of EMC nodes creates a spatial strain $\epsilon(r) = \nabla \cdot \vec{u}(r)$. This strain stretches the effective coordinate cell spacing $\lambda_l(r)$, so that a wave packet requires more fundamental time ticks t_p to traverse a unit coordinate distance. This produces an effective slowdown in coordinate speed $v(r) < 1$, encoded by an isotropic scalar refractive index $n_\gamma(r) = 1/v(r) > 1$.
2. **Trajectory deflection via restoring displacement.** The actual bending of a wave packet trajectory is driven by the vector displacement field $\vec{u}(r)$, rather than by a scalar density deficit alone. In response to the core deficit of a matter soliton, the surrounding EMC lattice develops a directed restoring displacement field:

$$\vec{u}(r) = -u_r(r) \hat{r}, \quad (106)$$

where the dimensionless radial magnitude $u_r(r)$ is the cumulative structural response to the normalized density gradient:

$$u_r(r) = \int_r^\infty \left| \nabla \left(\frac{N_\nu(r')}{N_{\text{stat}}} \right) \right| dr'. \quad (107)$$

The negative sign in Eq. (106) signifies structural displacement *toward* the central deficit. Light rays propagating through this strained lattice continuously realign along the principal axes of the displacement field, deflecting the wavefront toward the mass center.

12.2 Bridging the Vector Displacement to the Scalar Refractive Index

To connect the vector kinematics of Eq. (106) to standard ray-tracing integrals, we evaluate the scalar refractive index $n_\gamma(r)$ as the local isotropic dilatation factor induced by the dimensionless radial displacement field $u_r(r)$.

A 3D standing-wave soliton exhibits an asymptotic radial energy density deficit decaying as $1/r$. In the simplest spherical weak-field model, the normalized local EMC density takes the form

$$\frac{N_\nu(r)}{N_{\text{stat}}} = 1 - \frac{r_s}{r}, \quad (108)$$

where $r_s = 2GM$ is the gravitational radius in natural units ($c = 1$). The exact normalized gradient is therefore

$$\left| \nabla \left(\frac{N_\nu(r)}{N_{\text{stat}}} \right) \right| = \frac{r_s}{r^2}. \quad (109)$$

Substituting this gradient into Eq. (107) yields exactly

$$u_r(r) = \int_r^\infty \frac{r_s}{r'^2} dr' = \frac{r_s}{r}. \quad (110)$$

1255 The scalar refractive index $n_\gamma(r)$ represents the local isotropic dilatation factor induced by
 1256 this displacement field:

$$n_\gamma(r) = 1 + u_r(r) = 1 + \frac{r_s}{r} = 1 + 2\Phi_N(r), \quad (111)$$

1257 where $\Phi_N(r) = GM/r = r_s/(2r)$ is the dimensionless Newtonian gravitational potential per unit
 1258 energy.

1259 12.3 Asymptotic Continuous Limit and Schwarzschild Equivalence

1260 Eq. (111) corresponds to the first-order weak-field expansion of the full continuous expression:

$$n_\gamma(r) = \left(1 - \frac{2r_s}{r}\right)^{-1/2}. \quad (112)$$

1261 It is important to clarify the theoretical status of Eq. (112): in Enhanced EWT, this function
 1262 does not represent an external metric tensor imported from General Relativity, nor is it an ad
 1263 hoc optical ansatz. Rather, Eq. (112) is the *asymptotic continuous limit* of the discrete BCC
 1264 lattice deformation field around a spherical soliton core.

1265 Formally, $n_\gamma(r)$ matches the effective optical metric factor of the Schwarzschild geometry.
 1266 In EWT, however, this exact mathematical form arises physically from the nonlinear elastic
 1267 response of the EMC medium under the $1/r$ gravitational potential load, without requiring a
 1268 curved spacetime manifold.

1269 Expanding Eq. (112) binomially:

$$n_\gamma(r) = \left(1 - \frac{2r_s}{r}\right)^{-1/2} \approx 1 + \frac{r_s}{r} + \mathcal{O}\left(\left(\frac{r_s}{r}\right)^2\right), \quad (113)$$

1270 recovers the exact factor of 2 in front of $\Phi_N(r)$, providing the structural lattice foundation for
 1271 the observed $1.75''$ solar limb deflection. In the undisturbed statutory vacuum ($r_s/r \rightarrow 0$),
 1272 $n_\gamma(r) \rightarrow 1$, restoring the fundamental lattice ratio $c = \lambda_l/t_p = 1$.

1273 12.4 Mechanical Origin of Gravitational Redshift in the EMC Lattice

1274 In Enhanced EWT, both electromagnetic propagation and gravitational time dilation arise from
 1275 a single microscopic mechanism: the spatial deformation of the EMC node separation $\lambda_l(r)$
 1276 around a matter soliton.

1277 In the natural units of the BCC lattice, the speed of light is the fundamental metric conversion
 1278 factor

$$c \equiv \frac{\lambda_l}{t_p} = 1.$$

1279 Consequently, length and time share the same dimension, $[m] = [s]$, and the gravitational poten-
 1280 tial is dimensionless without the explicit c^2 factor that appears only in the SI-unit convention.
 1281 In these natural units, the Newtonian potential is simply

$$\Phi_N(r) = \frac{GM}{r}, \quad (114)$$

1282 and the gravitational radius of the source is

$$r_s = 2GM. \quad (115)$$

1283 Define the dimensionless local EMC lattice density ratio as:

$$\eta(r) \equiv \frac{N_\nu(r)}{N_{\text{stat}}} = 1 - \frac{r_s}{r} = 1 - 2\Phi_N(r), \quad (116)$$

1284 where $\Phi_N(r) = GM/r = r_s/(2r)$ in natural units.

1285 A physical clock is modeled as a standing-wave soliton whose fundamental period $T(r)$ is
 1286 determined by the internal round-trip time of a longitudinal EMC lattice wave traversing the
 1287 soliton core. Because the internal wave relies on node-to-node propagation across the same
 1288 strained lattice spacing $\lambda_l(r)$ that dictates external optics, its effective propagation rate scales
 1289 with the square root of the local node availability $\eta(r)$:

$$v_{\text{clock}}(r) = \sqrt{\eta(r)} = \sqrt{1 - 2\Phi_N(r)} \approx 1 - \Phi_N(r). \quad (117)$$

1290 For a rest-frame internal path length $2L_0$, the clock period in the gravitational potential field
 1291 becomes:

$$T(r) = \frac{2L_0}{v_{\text{clock}}(r)} = \frac{T_0}{\sqrt{1 - 2\Phi_N(r)}} \approx T_0 (1 + \Phi_N(r)), \quad T_0 = 2L_0. \quad (118)$$

1292 The fractional frequency shift reproduces the standard gravitational redshift:

$$\frac{\Delta f}{f} = \frac{f(r) - f_0}{f_0} = -\Phi_N(r) = -\frac{GM}{r} = -\frac{r_s}{2r}. \quad (119)$$

1293 **Structural Unity of Metric Phenomena.** This completes the mechanical equivalence be-
 1294 tween General Relativity metric components and EWT lattice strain:

- 1295 • **Temporal metric component** ($\sqrt{g_{00}}$): Encoded by the internal clock wave speed $v_{\text{clock}}(r) =$
 1296 $\eta(r)^{1/2}$, producing gravitational time dilation.
- 1297 • **Effective optical metric factor:** Encoded by the optical refractive index $n_\gamma(r) = 1 +$
 1298 $r_s/r = 1 + 2\Phi_N(r)$, whose weak-field form is the first-order expansion of $(1 - 2r_s/r)^{-1/2}$,
 1299 producing solar light deflection.
- 1300 In standard GR language, this effective index combines the contributions of both g_{00} and
 1301 g_{rr} . In EWT it arises directly from the integrated EMC displacement u_r .

1302 Both fundamental phenomena are physically unified as dual manifestations of the same local
 1303 EMC density profile $\eta(r) = 1 - r_s/r$, but they project onto different observables: the clock
 1304 samples the local node availability through $\sqrt{\eta}$, while the light ray samples the integrated radial
 1305 displacement $u_r = 1 - \eta = r_s/r$.

1306 12.5 Shapiro Delay from the EMC Refractive Index

1307 The third local metric observable is the Shapiro delay: a light signal passing near a massive
 1308 body takes longer to cross the region than it would in the undisturbed statutory vacuum. In the
 1309 Enhanced EWT framework, this delay is not an independent postulate but a direct consequence
 1310 of the same EMC lattice strain that produces light bending and gravitational redshift.

1311 In natural lattice units ($c \equiv 1$), the coordinate speed of a photon in the strained EMC
 1312 geometry is

$$v_\gamma(r) = \frac{1}{n_\gamma(r)} = \left(1 - \frac{2r_s}{r}\right)^{1/2}, \quad (120)$$

where $n_\gamma(r)$ is the optical refractive index introduced in Sec. 12.3. The coordinate time required for a signal to travel along a path $\mathcal{C}$ is therefore

$$t = \int_{\mathcal{C}} \frac{ds}{v_\gamma(r)} = \int_{\mathcal{C}} n_\gamma(r) ds. \quad (121)$$

In the weak-field limit,

$$n_\gamma(r) - 1 = \frac{r_s}{r} = 2\Phi_N(r), \quad \Phi_N(r) = \frac{GM}{r}, \quad (122)$$

so the excess delay relative to the statutory vacuum along a one-way path is

$$\Delta t_{\text{one-way}} = \int_{\mathcal{C}} (n_\gamma(r) - 1) ds = 2 \int_{\mathcal{C}} \Phi_N(r) ds. \quad (123)$$

For a round-trip radar signal grazing the solar limb, the dominant contribution comes from the near-Sun segment of the trajectory. Approximating the path as a straight line with impact parameter $b \simeq R_\odot$, the integral over the two-way path gives the logarithmic form

$$\Delta t_{\text{round-trip}} \approx 4GM \ln\left(\frac{4R_ER_P}{b^2}\right) \quad (c = 1), \quad (124)$$

where R_E and R_P are the heliocentric distances of the emitter and receiver. Restoring the speed of light c yields the standard SI expression,

$$\Delta t_{\text{round-trip}} \approx \frac{4GM}{c^3} \ln\left(\frac{4R_ER_P}{b^2}\right) \quad (\text{SI}), \quad (125)$$

which matches the classic relativistic Shapiro delay.

The Shapiro delay constitutes the third independent projection of the single EMC lattice deformation surrounding a matter soliton.

12.6 Macroscopic EMC Density Profile of a Star

The EMC density profile around a macroscopic body such as the Sun is not a simple scaled copy of an individual lepton's internal profile. A star is a vast collection of overlapping matter solitons, each contributing a local displacement of the EMC medium. The resulting effective density field is therefore a collective, smoothed macroscopic structure rather than a single soliton boundary wall.

For a static, spherically symmetric body of total mass M , the normalised EMC density ratio,

$$\eta(r) \equiv \frac{N_\nu(r)}{N_{\text{stat}}}, \quad (126)$$

can be naturally partitioned into three characteristic regions:

1. **Interior deficit region** ($r < R_\odot$). Inside the body, the conserved energy density of the constitutive matter solitons forces a local reduction of the EMC lattice density. This deficit is finite and, for an ordinary star, remarkably shallow compared with the extreme depletion expected in a compact remnant or black hole configuration. Its exact functional shape depends on the internal equation of state and is not required for external metric tests.

1339 **2. Stellar transition layer** ($r \approx R_\odot$). At the physical boundary of the body, the EMC
 1340 density transitions from the interior deficit to the statutory background. For a star, this
 1341 layer is broad and diffuse relative to the subatomic Degraded EMC Wall of a single electron,
 1342 arising from the macroscopic superposition of countless individual soliton boundaries.

1343 **3. Far-field tail** ($r \gg R_\odot$). Far outside the body, the EMC density relaxes toward the
 1344 statutory value N_{stat} according to

$$\eta(r) = 1 - \frac{r_s}{r}, \quad r_s = 2GM \quad (c \equiv 1). \quad (127)$$

1345 This represents the weak-field, spherically symmetric solution consistent with the long-
 1346 range lattice strain generated by the enclosed mass M .

1347 The local metric phenomena evaluated in this work — solar light bending, gravitational
 1348 redshift, and Shapiro delay — are strictly insensitive to the detailed structure of the interior
 1349 and transition regions. They are entirely governed by the far-field tail $\eta(r) = 1 - r_s/r$, which is
 1350 universal for any static, spherically symmetric EMC deficit of total strain magnitude r_s .

1351 Thus, the solar EMC profile used in our analytical and numerical validations represents the
 1352 *macroscopic far-field tail* of the collective density field, distinct from the microscopic Degraded
 1353 EMC Wall of an isolated lepton.

1354 12.7 Derivation of the Far-Field EMC Density Profile

1355 The far-field density profile, $\eta(r) = 1 - r_s/r$, is derived from the static continuum equilibrium of
 1356 the EMC lattice rather than selected as an ad hoc fitting ansatz.

1357 In the long-wavelength limit, $r \gg \lambda_l$, the discrete BCC lattice is well approximated by an
 1358 isotropic elastic continuum. Outside a bounded source region, where external volume forces
 1359 vanish, the static response of such a medium is governed by the Laplace equation for the scalar
 1360 strain field. Writing the normalized local EMC density as

$$\eta(r) = 1 + \delta\eta(r), \quad |\delta\eta(r)| \ll 1, \quad (128)$$

1361 the linearised equilibrium condition takes the form

$$\nabla^2 \delta\eta(r) = 0, \quad r > R, \quad (129)$$

1362 where R is the effective radius of the central matter distribution.

1363 Under spherical symmetry and the boundary condition that the perturbation vanishes at
 1364 spatial infinity ($\delta\eta \rightarrow 0$ as $r \rightarrow \infty$), the general solution to Eq. (129) is uniquely constrained to
 1365 the harmonic radial tail:

$$\delta\eta(r) = -\frac{A}{r}, \quad A > 0. \quad (130)$$

1366 By analogy with electrostatics, the coefficient A plays the role of a monopole charge for the
 1367 strain field: it measures the total strength of the deformation induced by the central source. The
 1368 integration constant A is fixed by applying Gauss's theorem to the strain flux across a closed
 1369 surface surrounding the mass M . In natural units ($c \equiv 1$), this gives

$$A = r_s = 2GM. \quad (131)$$

1370 Consequently, the far-field profile is uniquely determined as:

$$\eta(r) = 1 - \frac{r_s}{r}, \quad r \gg R. \quad (132)$$

1371 Thus, the profile contains no free functional parameters and no empirical tuning coefficients.
 1372 The $1/r$ form is required by the harmonic equilibrium of the lattice, and its amplitude is fixed
 1373 by the enclosed gravitational radius r_s .

12.8 From Microscopic EMC Displacement to the Gravitational Radius

The far-field EMC density profile

$$\eta(r) = 1 - \frac{r_s}{r} \quad (133)$$

contains a single characteristic length scale, r_s . In the preceding numerical implementations, this amplitude was identified with the gravitational radius of the source. This section demonstrates that this identification is a necessary consequence of the additive microscopic EMC displacements of constituent matter solitons, rather than an independent boundary input.

Consider a static, spherically symmetric body of total mass $M = \sum_i m_i$, composed of elementary matter solitons. In the long-wavelength, weak-field regime, the lattice continuum behaves as a linear elastic medium, ensuring that the total strain field is the linear superposition of individual soliton perturbations.

For any elementary soliton i of mass m_i , the fundamental geometric identity of EWT yields the individual monopole strain amplitude:

$$a_i = 2 G_{\text{EWT, geo}} m_i, \quad (134)$$

where $G_{\text{EWT, geo}}$ is the uncalibrated geometric constant derived purely from the ideal BCC lattice projection parameter $L_p^{\text{geom}} = 2/\sqrt{3}$. Owing to field linearity in the far zone, the total monopole amplitude A of the macroscopic body is the sum of the constituent amplitudes:

$$A = \sum_i a_i = 2 G_{\text{EWT, geo}} \sum_i m_i = 2 G_{\text{EWT, geo}} M. \quad (135)$$

Equation (135) establishes that the macroscopic strain amplitude is strictly proportional to the total mass enclosed.

Applying Gauss's divergence theorem to the radial displacement field $\delta\eta(r) = \eta(r) - 1$, the outward strain flux through a sphere of radius r is given by:

$$\oint_{\partial V} \nabla(\delta\eta) \cdot d\mathbf{S} = 4\pi r^2 \frac{d\delta\eta}{dr} = 4\pi A. \quad (136)$$

For the fundamental harmonic solution $\delta\eta(r) = -A/r$, the radial derivative evaluates to:

$$\frac{d\delta\eta}{dr} = \frac{A}{r^2}. \quad (137)$$

Consequently, the Neumann boundary condition enforced at the source radius R in the numerical implementation is fully derived from first principles:

$$\left. \frac{d\delta\eta}{dr} \right|_R = \frac{2 G_{\text{EWT, geo}} M}{R^2}. \quad (138)$$

Integration yields the unique far-field density profile:

$$\eta(r) = 1 - \frac{2 G_{\text{EWT, geo}} M}{r}, \quad (139)$$

from which the gravitational radius naturally emerges as a derived effective quantity:

$$r_s \equiv 2 G_{\text{EWT, geo}} M. \quad (140)$$

1399 No empirical value of G is injected into this derivation. The minor numerical discrepancy
 1400 (~ 5 ppm) between $G_{\text{EWT, geo}}$ and the CODATA value lies within the experimental uncertainty
 1401 of Newton's constant and is fully accounted for by the non-ideal spherical packing impedance
 1402 ζ (Sec. 24.8.2). This completes the logical closure: the far-field profile and metric artifacts are
 1403 generated strictly from microscopic lattice parameters and the enclosed mass.

1404 12.9 Emergent Metric Encodings from Lattice Dynamics

1405 The elastic constant k introduced in Sec. 1.5 can be derived from a microscopic pair potential
 1406 $V(r)$ between neighbouring EMCs. In the continuum limit of a one-dimensional lattice, the
 1407 equilibrium spacing is

$$a(\eta) \equiv \frac{1}{\eta}, \quad (141)$$

1408 where η is the normalised EMC density.

1409 In this one-dimensional chain, the natural contact interaction is modelled by the repulsive
 1410 potential

$$V(r) = \frac{V_0}{r}, \quad (142)$$

1411 whose first derivative gives the force between neighbouring sites and whose second derivative
 1412 evaluated at the equilibrium spacing defines the local spring stiffness:

$$k(\eta) = \left. \frac{d^2 V}{dr^2} \right|_{r=1/\eta} \propto \eta^3. \quad (143)$$

1413 The mass per lattice site is fixed, since each site corresponds to one EMC. The mass per unit
 1414 length is therefore

$$\rho = \frac{m_0}{a(\eta)} = m_0 \eta. \quad (144)$$

1415 The long-wavelength speed of sound in a one-dimensional lattice is not $\sqrt{k/m_0}$, but rather

$$v(\eta) = a(\eta) \sqrt{\frac{k(\eta)}{m_0}}. \quad (145)$$

1416 Using the one-dimensional modulus

$$E(\eta) = k(\eta) a(\eta) \propto \eta^2, \quad (146)$$

1417 this can be written in the standard continuum form

$$v(\eta) = \sqrt{\frac{E(\eta)}{\rho(\eta)}} \propto \sqrt{\frac{\eta^2}{\eta}} = \sqrt{\eta}. \quad (147)$$

1418 This is exactly the clock-speed encoding

$$v_{\text{clock}} \propto \sqrt{\eta}, \quad (148)$$

1419 now obtained as an emergent property of the microscopic lattice dynamics.

1420 The corresponding effective refractive index follows from the same wave speed:

$$n_\gamma(\eta) = \frac{c_0}{v(\eta)} \propto \eta^{-1/2}. \quad (149)$$

1421 A physical clock in this medium is a standing-wave soliton of fixed linear dimension L . Its
 1422 fundamental frequency is therefore

$$f(\eta) \propto \frac{v(\eta)}{L} \propto \sqrt{\eta}. \quad (150)$$

1423 Thus the two exponents previously introduced as geometric encodings follow directly from
 1424 the lattice dynamics. They do not require an independent optical or clock hypothesis.

1425 **Dimensional reduction and 3D extension:** The one-dimensional chain represents the
 1426 effective longitudinal propagation along a high-symmetry axis of the three-dimensional BCC
 1427 lattice. In a full 3D lattice, the equilibrium spacing scales as $a(\eta) \propto \eta^{-1/3}$, while the volumetric
 1428 mass density scales as $\rho_{3D} \propto \eta$. The same wave-speed scaling $v(\eta) = \sqrt{E/\rho} \propto \sqrt{\eta}$ is then
 1429 recovered from a microscopic inverse-cube pair potential $V(r) \propto r^{-3}$.

1430 **Note on the separation of gravitational and electromagnetic domains:** In the EWT
 1431 framework, electromagnetic phenomena operate in the *soliton-energy domain*. Charge and radia-
 1432 tion are carried by the amplitude and phase of standing and travelling waves. Consequently, two
 1433 counter-propagating electromagnetic waves superpose linearly in free space: their interference
 1434 does not produce a leading-order change in the EMC packing density $\rho(r)$. Gravity, by contrast,
 1435 operates in the *EMC density domain*. It is the macroscopic response of the lattice to a localized
 1436 deficit in the packing density $N_{\nu,\text{eff}}$. This distinction explains why gravity is universal, acting on
 1437 all forms of energy that displace the medium, while remaining exceptionally weak compared to
 1438 electromagnetic effects, which directly involve the energy amplitudes of the solitons.

1439 12.10 Summary of Local Metric Observables

1440 Thus, the single normalized EMC density profile $\eta(r) = 1 - r_s/r$ unifies all three classic local
 1441 metric observables without introducing any free parameters:

- 1442 • **Solar light bending** through the spatial gradient of $n_\gamma(r)$,
- 1443 • **Gravitational redshift** through the internal clock speed $v_{\text{clock}} = \sqrt{\eta}$,
- 1444 • **Shapiro delay** through the integrated excess of $n_\gamma(r) - 1$ along the propagation path.

1445 The functional form of this underlying profile is uniquely determined by the harmonic equi-
 1446 librium of the EMC lattice, with its amplitude strictly fixed by the enclosed gravitational radius
 1447 r_s .

1448 12.11 In-Platform Numerical Validation

1449 The analytical derivations presented in this section are validated numerically in the OpenWave
 1450 research suite. The scripts implement the same EMC density profile $\eta(r) = 1 - r_s/r$ and the
 1451 same encodings used in the analytical treatment, and compare the results against established
 1452 reference values. The numerical outcomes are summarised in Table 5.

1453 The numerical results reproduce the analytical predictions of this section. The scripts are
 1454 part of the OpenWave research repository [55].

1455 For M4.3, M4.4, and M4.5 the encoding of the EMC density profile into the refractive index
 1456 and the clock speed is taken from this manuscript; the scripts perform the resulting integrations
 1457 and confirm the agreement with the general-relativistic reference values.

1458 For M4.6, the radial Laplace equation is solved numerically with the monopole flux condition
 1459 at the solar surface and the asymptotic Robin boundary condition at the outer boundary. The
 1460 resulting profile matches the analytic form to 5.7×10^{-14} of η .

Table 5: Numerical validation of the local metric phenomena. All scripts implement the same EMC density profile and encodings derived analytically in this section.

Artifact	Quantity	EWT result	Reference	Difference
M4.3	Solar-limb light bending	1.751728 arcsec	1.7517 arcsec (GR)	0.0016%
M4.4	Gravitational redshift $\Delta f/f$	-2.123135×10^{-6}	GR exact factor	0 (identity)
M4.5	Shapiro round-trip delay	239.014 μ s	239.014 μ s	0.000075%
M4.6	Far-field profile $\eta(r)$	RK4 solution	analytic $1 - r_s/r$	5.7×10^{-14}
M4.8	Monopole amplitude A	2.954320×10^3 m	CODATA G	65.65 ppm
M4.9	Stiffness exponent α	3.0	expected 3.0	< 0.05
M4.9	Wave-speed exponent β	0.5	expected 0.5	< 0.05

For M4.8, the monopole amplitude is computed from the geometric G chain of the emergence engine with the packing impedance $\zeta_{\text{est}} = (1 - \eta_{\text{BCC}})/(\eta_{\text{BCC}} N_{\text{ideal}})$, without calibrating to the measured fine-structure constant. The resulting amplitude differs from the CODATA value by 65.65 ppm. The pure geometric limit ($\zeta = 0$) gives 1743.57 ppm, establishing the packing correction as the relevant physical input.

For M4.9, the one-dimensional lattice dynamics with a repulsive pair potential $V(r) = V_0/r$ yields the exponents $\alpha = 3$ (stiffness scaling $k \propto \eta^\alpha$) and $\beta = 0.5$ (wave-speed scaling $v \propto \eta^\beta$), which are the values used in the encodings of the clock speed and the refractive index.

13 Newtonian Force from Interacting EMC Deficits

The same far-field density deficit that reproduces the local metric observables also gives rise to the Newtonian force law.

Consider two matter solitons of masses M_1 and M_2 , separated by a distance R . Each soliton sources a monopole density deficit

$$\delta\eta_i(r) = -\frac{A_i}{r}, \quad A_i = \frac{2G_{\text{EWT, geo}}M_i}{c^2}. \quad (151)$$

The static interaction energy arises from the integrated overlap of the two density gradients in the surrounding medium,

$$U_{\text{int}}(R) = -K_{\text{emc}} \int \nabla(\delta\eta_1) \cdot \nabla(\delta\eta_2) dV, \quad (152)$$

The minus sign follows from the negative-definite energy functional adopted for the EMC field; this functional is the geometric counterpart of the Newtonian field energy $-\frac{|\nabla\Phi|^2}{8\pi G}$, and the normalization constant is

$$K_{\text{emc}} = \frac{c^4}{16\pi G_{\text{EWT, geo}}}. \quad (153)$$

In spherical coordinates centred on the first source, the scalar product of the two gradients is

$$\nabla(\delta\eta_1) \cdot \nabla(\delta\eta_2) = A_1 A_2 \frac{r^2 - rR \cos \theta}{r^3(r^2 + R^2 - 2rR \cos \theta)^{3/2}}. \quad (154)$$

Performing the angular integration over the spherical shell at radius r yields

$$\int_0^{2\pi} \int_0^\pi [\nabla(\delta\eta_1) \cdot \nabla(\delta\eta_2)] \sin \theta d\theta d\phi = \begin{cases} 0, & r < R, \\ \frac{4\pi A_1 A_2}{r^4}, & r \geq R, \end{cases} \quad (155)$$

1481 which is the exact field-theoretic analogue of Newton’s shell theorem.

1482 Integrating over the volume element $dV = r^2 \sin \theta \, dr \, d\theta \, d\phi$ for the non-vanishing region $r \geq R$
 1483 gives

$$U_{\text{int}}(R) = -\frac{c^4}{16\pi G_{\text{EWT, geo}}} \cdot 4\pi A_1 A_2 \int_R^\infty \frac{dr}{r^2} = -\frac{c^4 A_1 A_2}{4G_{\text{EWT, geo}} R}. \quad (156)$$

1484 The attractive force directed along the line connecting the centers is

$$F = -\frac{dU_{\text{int}}}{dR} = -\frac{c^4 A_1 A_2}{4G_{\text{EWT, geo}} R^2}. \quad (157)$$

1485 Substituting the monopole amplitudes $A_i = 2G_{\text{EWT, geo}} M_i / c^2$ yields precisely

$$F = -\frac{G_{\text{EWT, geo}} M_1 M_2}{R^2}. \quad (158)$$

1486 No additional fundamental force is postulated. The inverse-square law is the static conse-
 1487 quence of two EMC density deficits interacting through vacuum push-out pressure. The 4π factor
 1488 emerges dynamically from the 3D angular integration, while the 16π factor represents the field
 1489 stress-energy normalization.

1490 13.1 In-Platform Validation: Normalization Consistency

1491 The derivation of the inverse-square force law from EMC field overlap is validated numerically
 1492 in the OpenWave research suite. The baseline gate compares the field-overlap force F_{EMC} with
 1493 the Newtonian force F_{Newton} , both built from the same geometric G_{geom} derived from the BCC
 1494 lattice. The result is summarised in Table 6.

Table 6: Baseline gate for the Newtonian force from EMC field overlap. Both F_{EMC} and F_{Newton} are built with $G_{\text{geom}} = 6.677519975508460 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$.

Quantity	Value
F_{EMC}	$3.544205553218 \times 10^{22} \text{ N}$
F_{Newton}	$3.544205979545 \times 10^{22} \text{ N}$
Relative difference	$1.203 \times 10^{-5}\%$

1495 Because G_{geom} enters both the monopole amplitude $A_i = 2G_{\text{geom}} M_i / c^2$ and the coupling
 1496 $K_{\text{emc}} = c^4 / (16\pi G_{\text{geom}})$, it cancels identically in $F_{\text{EMC}} \equiv F_{\text{Newton}}$. The gate therefore passes for
 1497 any value of G_{geom} and remains a normalization-consistency check on A , K_{emc} , and the angular
 1498 factor 4π . Mutating any single normalization factor breaks the agreement, as shown in Table 7.

Table 7: Mutation test of the normalization-consistency gate. Each mutation changes a single normalization factor while keeping the rest of the derivation unchanged.

Mutation	Observed relative difference	Gate result
Baseline	$1.203 \times 10^{-5}\%$	PASS
Angular factor ($4\pi \rightarrow 3\pi$)	25.0%	FAIL
Monopole amplitude ($2GM/c^2 \rightarrow GM/c^2$)	75.0%	FAIL
Coupling denominator ($16\pi \rightarrow 8\pi$)	100.0%	FAIL

1499 The amplitude can also be written without the symbol G_{geom} , directly in the chain factors
 1500 of the emergence engine:

$$A = \frac{2Mr_e}{m_e} \cdot \frac{\sqrt{X_{\text{eff}}}}{A_\pi^4 N_{\text{geom}}^3 K_{WC} \sqrt{N_{\nu,\text{stat}}}}, \quad X_{\text{eff}} = \frac{N_{\nu,\text{stat}}}{N_{\nu,\text{eff}}}. \quad (159)$$

1501 Every factor on the right-hand side is geometric: $A_\pi = 4\pi^3 + \pi^2 + \pi$, $N_{\text{geom}} = 8\pi^4(1 - \zeta)$,
 1502 $K_{WC} = 10$, and X_{eff} , $N_{\nu,\text{stat}}$ from the geometric chain. The anchors r_e, m_e are the measured
 1503 dimensional quantities. The product of Eq. (159) is the engine's G_{EWT} formula multiplied by
 1504 $2M/c^2$, term for term, so $A = 2G_{\text{geom}}M/c^2$ by value. The geometric form agrees with the
 1505 G -based form to machine precision (1.5×10^{-16} for the solar amplitude, 2.0×10^{-16} for the
 1506 terrestrial amplitude): the two are the same expression, so the agreement is an identity, not a
 1507 confirmation.

1508 The force gate does not depend on the value of G_{geom} . Making it a test of G_{geom} requires
 1509 an amplitude whose value is fixed without passing through the G_{EWT} formula; this remains an
 1510 open clause in the research program.

1511 **Strength clause.** The two halves of the strength clause for the gravitational sector are stated
 1512 as follows.

- 1513 • **Circularity: discharged.** G_{geom} enters the force test directly, derived from the BCC
 1514 geometry, without re-entering through an input.
- 1515 • **Accuracy: stated, not met.** The G_{geom} residual against CODATA is 0.048169%, or
 1516 $21.9\times$ the CODATA 2022 relative uncertainty on G (22 ppm). At that distance the derived
 1517 value does not agree with the measured G within its uncertainty. The dimensional anchors
 1518 r_e, m_e, c are the measured quantities used to build the dimensionless geometric ratio; the
 1519 derivation produces G_{geom} as a dimensionless ratio against $c^2 r_e / m_e$, not G from nothing.

1520 The scripts are part of the OpenWave research repository [56].

1521 14 Relation to Existing Emergent Gravity Frameworks

1522 The Geometric Identity Model (EWT) is fundamentally aligned with the concept of Emergent
 1523 Gravity (EG), where gravity is viewed as a collective, induced, or thermodynamic phenomenon
 1524 [31, 32, 30]. The derivation of G from explicit microscopic geometry and its dependence on the
 1525 vacuum's effective volume deficit places this work within the general realm of EG approaches.
 1526 However, EWT offers a unique, explicit microscopic realization that is independent of purely
 1527 thermodynamic or entropic principles.

1528 14.1 Comparison of Mechanisms and Scales

1529 The EWT model differs from established EG frameworks in three critical aspects: the nature
 1530 of the microscopic degrees of freedom, the source of gravitational emergence, and the explicit
 1531 scaling factor for the gravitational constant G .

1532 14.1.1 Microscopic Degrees of Freedom (DoF)

- 1533 • **Thermodynamic/Entropic (Jacobson, Verlinde):** The DoF are typically macroscopic
 1534 or informational, related to the ****area elements of a holographic screen**** or ****entanglement****
 1535 ****between quantum fields in the vacuum.**

- 1536 • **Induced (Sakharov):** The DoF are the **quantum fields themselves**, with gravity
1537 arising from the zero-point energy of the vacuum.
- 1538 • **EWT (Geometric Soliton):** The DoF are **explicitly geometric** and local: the internal
1539 structure and energy density profile $\rho(\mathbf{r})$ of the Soliton, and its internal magnetic coherence
1540 $\epsilon_{\mathbf{M}}$. This provides a tangible, particle-scale physical realization of the underlying DoF.

1541 14.1.2 Source of Gravitational Emergence

1542 The EWT model proposes a **dual mechanism** for the strength of gravity, distinguishing the
1543 source of the mass/energy from the source of its weakness, which is not present in standard EG
1544 theories:

- 1545 • **Thermodynamic/Entropic:** The source is the **thermodynamic state** (e.g., temper-
1546 ature T) or the **information content** of spacetime.
- 1547 • **Induced (Sakharov):** The source is the **bulk change in vacuum energy** caused by
1548 spacetime curvature.
- 1549 • **EWT (Geometric Soliton):**
 - 1550 a) **Primary Source ($\mathbf{G}_{\text{Base}}$):** The gravitational force originates from the fundamental
1551 **quantity of EMC** (energy-mass-coherence) within the Soliton ($\mathbf{m}_e$).
 - 1552 b) **Emergence/Weakness:** The vast weakness of gravity (scaling from $\mathbf{G}_{\text{Base}}$ to G) is
1553 caused by the **Geometric Deficit**—the ratio between the Soliton’s compact volume
1554 and the much larger effective volume it influences in the surrounding EWT medium.

1555 14.1.3 Scaling Factor for $\mathbf{G}$

1556 The frameworks rely on fundamentally different scaling parameters to define the observed value
1557 of G :

- 1558 • **Thermodynamic/Entropic:** G is typically scaled by $\hbar$ and parameters related to the
1559 **temperature of the horizon ($\mathbf{T}$)**, e.g., $\mathbf{G} \propto 1/(\mathbf{S} \cdot \mathbf{T})$.
- 1560 • **Induced (Sakharov):** G is inversely proportional to the **fourth power of the quantum**
1561 **field theory cutoff scale (Λ)**, $\mathbf{G} \propto 1/\Lambda^4$.
- 1562 • **EWT (Geometric Soliton):** G is scaled by the complete, derived, dimensionless scaling
1563 factor $\Omega_{\mathbf{G}}$:

$$\mathbf{G} = \mathbf{G}_{\text{Base}} \cdot \Omega_{\mathbf{G}}, \quad \text{where } \Omega_{\mathbf{G}} = \frac{1}{\mathbf{A}_{\pi}} \cdot \left(\frac{1}{\mathbf{N}_{\text{final}} \mathbf{A}_{\pi}} \right)^3 \cdot \frac{1}{\mathbf{K}_{\mathbf{WC}} \cdot \sqrt{\mathbf{N}_{\nu, \text{effective}}}} \quad (160)$$

1564 The EWT scaling factor $\Omega_{\mathbf{G}}$ is **explicitly constructed** from the Soliton’s geometric pa-
1565 rameters ($\mathbf{A}_{\pi}$, $\mathbf{N}_{\text{final}}$) and the unified volume deficit ($\mathbf{N}_{\nu, \text{effective}}$), linking G to the Soliton’s
1566 internal dynamics rather than external thermodynamic quantities.

1567 The EWT model thus acts as a bridge, providing the explicit, geometric and non-thermodynamic
1568 identity $\mathcal{U}(\mathbf{P})$ required to link fundamental particle parameters to the macroscopic gravitational
1569 constant G , while remaining compatible with the holographic and entropic concepts that underlie
1570 the vacuum structure.

Comparison with Induced Gravity (Sakharov)

The EWT scaling factor Ω_G exhibits a profound structural isomorphism with the concept of **Induced Gravity** proposed by Andrei Sakharov [30]. In Sakharov's framework, gravity is not a fundamental force but an emergent "elasticity of the vacuum," where the gravitational constant G scales inversely with the fourth power of a quantum field theory cutoff, $G \propto 1/\Lambda^4$.

In the EWT model, this "cutoff" is replaced by the geometric coupling constant A_π . The total suppression factor Ω_G incorporates the term A_π^{-4} (derived from the linear and volumetric saturation: $A_\pi^{-1} \cdot A_\pi^{-3}$), providing a **purely geometric derivation** for the quartic attenuation observed in emergent gravity theories.

While Sakharov required vacuum fluctuations to define the scale, EWT identifies this scale as the physical phase-saturation point of the BCC lattice. This suggests that the 10^{42} weakness of gravity is the result of the push-out effective mechanism modulated by "stiffness" coupled with the holographic projection of nodal energy ($\sqrt{N_{\nu,\text{eff}}}$).

Gravity is scaled by the total energy density of the space occupied by matter, where the attenuation follows $G \propto 1/A_\pi^4$, further diluted by the holographic transition to the soliton's surface.

This perspective implies that within a certain range, the gravitational strength is determined by the lattice's volumetric resistance, while the final 10^{42} weakness is dominated by the geometric projection:

$$G = \underbrace{\frac{G_{\text{Base}}}{A_\pi^4 \cdot N_{\text{final}}^3}}_{\text{Volumetric Energy Density (push-out)}} \cdot \underbrace{\frac{1}{K_{WC} \cdot \sqrt{N_{\nu,\text{eff}}}}}_{\text{Surface Dilution}} \quad (161)$$

14.1.4 Evolution of the EWT Framework: From Wave Attenuation to EMC Push out

The introduction of the Push out Mechanism marks a fundamental shift in the interpretation of gravitational forces within the Energy Wave Theory (EWT). While the baseline EWT model [11] treats the BCC lattice primarily as a passive *wave carrier*—where gravity is attributed to a minute loss of wave energy (shadow effect)—the current framework redefines the BCC lattice as an **active actor**.

In this refined model, the gravitational constant is no longer dependent on an arbitrary wave attenuation coefficient. Instead, the extreme weakness of gravity ($\sim 10^{-42}$) is a direct consequence of **holographic energy dilution**. The energy density, concentrated within the volumetric saturation of the lattice (A_π^4), is projected onto the 3D surface of the matter soliton and further corrected by the effective EMC density $N_{\nu,\text{eff}}$.

This transition is governed by the fundamental geometric derivative of the vacuum state, where the volumetric saturation A_π^4 yields the **push-out force density**:

$$y = A_\pi^4 \implies y' = 4A_\pi^3 \quad (162)$$

- **Baseline EWT:** Relies on a nearly infinitesimal energy loss during wave propagation to account for the weakness of gravity, which lacks a direct structural justification.
- **Push-out Model (Enhanced EWT):** Identifies gravity as a "blurred image" of an extremely strong lattice interaction. The 10^{-42} factor emerges naturally from the geometric ratio between the 4D saturation base (A_π^4) and the holographic surface projection of the soliton.

By replacing "leaking wave energy" with the **structural stiffness** and **statutory density** ($N_{\nu,\text{stat}}$) of the lattice, the model provides a deterministic explanation for gravitational emergence. Gravity is thus revealed as the mechanical reaction of the BCC medium to a localized EMC deficit, governed by the derivative of the vacuum's volumetric impedance.

Structural Range: Continuity vs. Wave Dissipation

A significant challenge for the baseline EWT model is explaining the immense, theoretically infinite range of gravity. In a framework based on infinitesimal wave attenuation, a signal as weak as 10^{-42} would realistically be dissipated by the vacuum's stochastic noise over long distances.

The **EMC Push-out Mechanism** provides a structural solution to this problem:

- **Geometric Continuity:** Unlike a wave signal that requires constant energy propagation, the push-out mechanism creates a **static structural deformation** of the BCC lattice. Since the medium is a continuous mechanical entity, the gradient between $N_{\nu,\text{stat}}$ and $N_{\nu,\text{eff}}$ must propagate throughout the entire network to maintain topological equilibrium.
- **Infinite Reach:** The $1/r^2$ scaling is revealed as a geometric requirement of 3D projection from the 4D saturation base (A_π^4), not a result of "fading waves." This explains why gravity remains stable across cosmological scales—it is not a "message" being sent, but a permanent "tilt" in the lattice geometry.

By shifting the focus from "energy loss" to **lattice elasticity**, the Enhanced EWT accounts for both the extreme weakness of the force (via holographic dilution) and its universal reach (via structural integrity), identifying gravity as the deterministic mechanical response of the vacuum's impedance.

14.1.5 EWT as Pressure-Driven Emergent Gravity

The fundamental mechanism proposed by EWT—where mass creates a **Geometric Deficit** ($N_{\nu,\text{effective}}$) which is then compensated by the surrounding EWT medium—aligns the model with modern interpretations of gravity as a **Push Force** or a vacuum pressure effect.

This interpretation contrasts sharply with the classic Newtonian view (attraction) and even with General Relativity (spacetime curvature), instead focusing on local density gradients:

- **Density Deficit and Energy Conservation:** The Soliton structure (the particle) constitutes a localized region of **lower packing** (a geometric deficit N_ν) compared to the undisturbed background of the EWT medium (the vacuum). Simultaneously, due to continuous internal wave interference, the Soliton **conserves** a large, localized amount of **energy** (mass equivalent).
- **Equilibrium Drive (Gravity):** The resulting force (gravity) is the expression of the **EWT medium's fundamental drive to restore equilibrium** by moving into the region of lower packing. This motion generates a net **pressure gradient**, which pushes all matter towards the center of the deficit.

This conceptualization places EWT in close affinity with models advocating that gravity is generated by the **pressure of a Quantum Vacuum** (or: *Medium*) flowing from higher to lower energy density areas [35], offering a coherent, picture of the gravitational interaction that is both emergent and pressure-driven.

This interpretation provides direct justification for the appearance of the square root term ($N_{\nu,\text{effective}}^{-1/2}$) in the scaling factor Ω_G . This term represents the **geometric transition** from

the underlying three-dimensional volume deficit ($\mathbf{N}_{\nu,\text{effective}}$) to the **two-dimensional surface effect** (pressure) exerted by the EWT medium, aligning EWT with the surface-based scaling principles of holographic gravity models.

15 Geometric Validation: The Fundamental Identity and the Base AMM State (a_e)

The internal consistency of the Energy Wave Theory (EWT) is verified through the geometric relationship between a soliton's energy and its spatial extent. In this framework, the anomalous magnetic moment (AMM) is treated as a deterministic consequence of the vacuum's elastic properties.

15.1 Geometric Mass-to-Radius Identity ($E \propto r^5$)

The core principle of EWT [37, 29] dictates that the energy E of a soliton scales with the fifth power of its geometric radius r ($E \propto r^5$). This identity provides the deterministic link between a particle's measured mass and its geometric size:

$$\frac{E_x}{E_e} = \left(\frac{r_x}{r_e}\right)^5 \implies r_x = r_e \left(\frac{E_x}{E_e}\right)^{1/5} \quad (163)$$

This relationship ensures that the geometric radius r_f is not an arbitrary parameter but is directly derivable from the particle's measured energy. This identity confirms that the soliton's scale is governed by the Wave Count hierarchy, establishing a unified set of parameters for both mass and magnetic anomaly. Radius scaling is directly related to the mass scaling model presented in section 19.3.

15.2 Physical Origin of the r^5 Scaling: Geometric Energy Density

The quintic relationship between energy and radius ($E \propto r^5$), as identified in the foundational EWT framework [37, 29], is here demonstrated to be a rigorous requirement of energy conservation within the vacuum lattice. While the r^5 scaling provides a powerful predictive tool, its physical origin lies in the convergence of three fundamental geometric and dynamic factors of the soliton:

1. **Volumetric Occupancy (r^3):** The three-dimensional spatial extent of the wave center within the BCC lattice.
2. **Amplitude Scaling (r^1):** Since charge is expressed as wave amplitude A in meters [29], the stability of the soliton requires the amplitude to be geometrically coupled with the radius ($A \propto r$) to maintain the structural integrity.
3. **Resonance Mode Count (r^1):** The number of standing-wave periods N that fit within the soliton scales linearly with its radius ($N \propto r$). A larger soliton supports more resonant modes, each oscillating at the characteristic lattice frequency $f \propto 1/r$.

The product of these factors ($r^3 \times r^1 \times r^1 = r^5$) confirms that mass is a measure of "dynamic information density". However, the divergence between this quintic energy surge and the cubic geometric compensation capacity ($\propto r^3$) explains the precision limits encountered in earlier models (e.g., the 10-16% error for the muon and tau in [37]).

1688 The three contributions combine into two scaling laws that are the formal basis of the Onion
1689 Model. The first is the energy–radius identity,

$$\boxed{E \propto r^5} \quad (164)$$

1690 which follows directly from the product of the three factors above and is identical to the mass-
1691 to-radius relation in Eq. (163). The second is the volumetric capacity of the three-dimensional
1692 lattice,

$$\boxed{V \propto r^3} \quad (165)$$

1693 which expresses the geometric compensation available in real space as the soliton expands.
1694 The mismatch between the quintic energy surge, Eq. (164), and the cubic geometric compensa-
1695 tion, Eq. (165), is the source of the $r^5/r^3 = r^2$ disparity that forces recursive shell formation in
1696 the Onion Model described in 16.3.

1697 15.3 The Geometric Status of Mass: Kilogram as a Derived Unit

1698 The $E \propto r^5$ scaling law acquires its full dimensional significance when combined with the single
1699 structural constant that defines the causal structure of the BCC lattice: the maximum propaga-
1700 tion speed c .

1701 **Step 1 — Metric unification via c .** The axiom $c \equiv \lambda_l/t_p$ (Section 9) identifies c as the
1702 intrinsic conversion factor between the spatial and temporal steps of the BCC lattice. Adopting
1703 $c = 1$ (the natural units of the lattice) unifies the dimensions of space and time:

$$c = 1 \implies [s] = [m]. \quad (166)$$

1704 **Step 2 — Mass from the r^5 geometric scaling law.** The fundamental mass-radius
1705 relation of EWT (Section 15.1) states that the energy of a soliton is entirely defined by its
1706 geometric extent, scaling as the fifth power of its radius:

$$E \propto r^5. \quad (167)$$

1707 Assuming the proportionality constant is a dimensionless topological feature of the BCC lattice,
1708 the geometric dimension of energy is identically $[r^5] = [m^5]$.

1709 In classical SI units, energy is defined by $E = mc^2$. Equating the geometric and classical
1710 dimensions yields the dimensional structure of mass:

$$[m \cdot c^2] = [r^5] \implies [\text{kg}] \cdot [\text{m}^2 \cdot \text{s}^{-2}] = [\text{m}^5]. \quad (168)$$

1711 Solving for the kilogram reveals its true derived nature in the EWT framework:

$$[\text{kg}] = [\text{m}^3 \cdot \text{s}^2]. \quad (169)$$

1712 When evaluated in the natural units of the lattice ($c = 1 \implies [s] = [m]$), the dimensional
1713 conversion becomes completely spatial:

$$[\text{kg}] = [\text{m}^3 \cdot \text{m}^2] = [\text{m}^5]. \quad (170)$$

1714 **Step 3 — Mass hierarchy from r^5 .** With the mass scale anchored to r_e through the r^5
1715 law, the entire lepton mass hierarchy follows from the topological integer sequence K_{WC} :

$$\frac{m_x}{m_e} = \left(\frac{r_x}{r_e}\right)^5 = \left(\frac{K_x}{K_e}\right)^5, \quad \text{dimensionless.} \quad (171)$$

1716 **Step 4 — Gravitational consistency.** The independently derived gravitational identity
 1717 (Section 11) provides a non-trivial consistency check. The base gravitational scale is defined as
 1718 $G_{\text{Base}} = c^2 r_e / m_e$. Substituting the dimensional relation $[\text{kg}] = [\text{m}^3 \cdot \text{s}^2]$ from Eq. (170) gives:

$$[G_{\text{Base}}] = \frac{[\text{m}^2 \cdot \text{s}^{-2}] \cdot [\text{m}]}{[\text{m}^3 \cdot \text{s}^2]} = [\text{s}^{-4}]. \quad (172)$$

1719 Adopting $c = 1$ ($[\text{s}] = [\text{m}]$) reduces this to $[G_{\text{Base}}] = [\text{m}^{-4}]$, which is exactly the dimension of the
 1720 ratio r_e / m_e when mass carries the geometric dimension $[\text{m}^5]$:

$$[r_e / m_e] = \frac{[\text{m}]}{[\text{m}^5]} = [\text{m}^{-4}]. \quad (173)$$

1721
 1722 The gravitational constant therefore requires no independent mass scale: its dimension, and
 1723 ultimately its value, is entirely determined by the geometry of the BCC lattice.

1724 In this framework, the SI kilogram occupies the same logical status as π in Euclidean ge-
 1725 ometry: it is not a free parameter, but a structural consequence of the underlying geometric
 1726 medium. The reduction of all three SI mechanical base units to a single geometric primitive (the
 1727 BCC lattice spacing λ_l) is the dimensional expression of the claim that EWT is a parameter-free
 1728 geometric theory of matter.

1729 15.4 Geometric Dimension of the Reduced Planck and G Constants

1730 The reduction of mass to a geometric quantity has an immediate consequence for the reduced
 1731 Planck constant. In SI units $\hbar$ has dimension

$$[\hbar] = [\text{kg}][\text{m}]^2[\text{s}]^{-1}. \quad (174)$$

1732 Using the natural-unit identifications $[\text{s}] = [\text{m}]$ and $[\text{kg}] = [\text{m}]^5$, this becomes

$$[\hbar] = [\text{m}]^5 \cdot [\text{m}]^2 \cdot [\text{m}]^{-1} = [\text{m}]^6. \quad (175)$$

1733 In the geometric formulation of EWT, $\hbar$ is therefore not an independent quantum input. It is a
 1734 derived geometric quantity of dimension length to the sixth power.

1735 The same reduction applies to the gravitational constant. From the defining relation $[G] =$
 1736 $[\text{m}]^3 / ([\text{kg}][\text{s}]^2)$, the natural-unit substitutions give

$$[G] = \frac{[\text{m}]^3}{[\text{m}]^5 [\text{m}]^2} = [\text{m}]^{-4}. \quad (176)$$

1737 This is consistent with the earlier derivation of the base gravitational scale from the electron
 1738 soliton.

1739 As a result, the Planck length

$$l_{\text{P}} = \sqrt{\frac{\hbar G}{c^3}} \quad (177)$$

1740 reduces at $c = 1$ to the geometric dimensional identity

$$[l_{\text{P}}] = \sqrt{[\hbar][G]} = \sqrt{[\text{m}]^6 [\text{m}]^{-4}} = [\text{m}]. \quad (178)$$

1741 Thus l_{P} is dimensionally identical to the primary EMC lattice step λ_l . In the geometric EWT
 1742 framework, λ_l is not constructed from $\hbar$ and G . The single lattice length λ_l fixes the geometric

units from which $\hbar$ and G inherit their SI dimensions when the system is expressed in laboratory units.

This geometric dimension admits a direct interpretation within the Geometric Ladder of the EWT framework. The sixth degree of freedom is associated with the volumetric bosonic budget $3D + A + f + r$, which is represented by the π^6 rung (Section 2.1). The reduced Planck constant is therefore the geometric measure of the six-dimensional phase-space volume available to a soliton resonance. The dimensional exponent 6 is not accidental: it is the same 6 that appears in the π^6 volumetric coupling operator for neutral weak interactions.

Using the Dimensional Weight Operator formalism, the six factors of π forming the π^6 rung are precisely those that define the six-dimensional phase-space cell whose quantum is $\hbar$.

15.5 The Fundamental Magnetic Deficit ϵ_M

In EWT, the magnetic moment correction is a purely geometric phenomenon resulting from the non-ideal stiffness of the elastic medium. The fundamental magnetic deficit constant, ϵ_M , is defined by the effective geometric stiffness modulus (N_{geom}) and the volumetric factor (π^3):

$$\epsilon_M \equiv \frac{1}{N_{\text{geom}} \cdot \pi^3} \quad (179)$$

The total stiffness deficit (Loss Factor) of the elastic medium is thus expressed as $\epsilon_M \cdot \pi^3 = 1/N_{\text{geom}}$.

15.6 Derivation of the a_e Base Formula

The anomalous magnetic moment of the electron (a_e) is derived as the product of the Ideal Geometric Moment (the Schwinger Term) and the Stiffness Retention Factor:

$$a_{\text{Base}}^{\text{Geometric}} = \frac{\alpha_{\text{geom}}}{2\pi} \cdot (1 - \epsilon_M \cdot \pi^3) \quad (180)$$

Numerical verification, using the geometric α_{geom} and ϵ_M from the BCC lattice, yields $a_{\text{Base}}^{\text{Geometric}} \approx 1159.916228$ ppm. This result aligns with the experimental value of the electron ($a_e^{\text{exp}} \approx 1159.652182$ ppm) with a relative error of approximately 0.02277%, establishing the electron as the fundamental geometric ground state.

16 The Recursive Lepton Hierarchy: Nodal Shell Resonance Model

Heavier leptons — the Muon (Ψ_μ) and the Tau (Ψ_τ) — emerge through **Recursive Nodal Inclusion** (the "Onion Model"). Each generation is treated as a cumulative wave-packing excitation within the BCC vacuum lattice, where the nodal structure is governed by a characteristic geometric factor $2\pi^2$ (the surface area of a torus, though other topologies yielding the same factor are not excluded).

The numerical values quoted in this section are obtained directly from the geometric Python implementation described in the Appendix.

Methodological Framework: Nodal Metrics and Geometric Determinism

To maintain theoretical rigor, a distinction is established between the discrete lattice and the resonant excitation. While the **Wave Center (Soliton)** (K_{WC}) is the fundamental physical and energy-conserving entity, its internal dynamics involve a level of complexity that makes direct continuous analysis less efficient. In contrast, the **Node** provides a highly effective **topological metric** for the EWT framework. By utilizing the discrete points of the BCC lattice as a reference for **phase space occupancy**, the "Onion Model" derives lepton generations through the precise counting of nodal density (K).

In this approach, nodes do not function as a rigid measurement of spatial distance, but rather as a **quantized gauge of resonant stability**. When the energy surge ($\propto r^5$) exceeds the volumetric capacity ($\propto r^3$) of the base geometry, the system's adaptation is captured by its "latching" onto additional nodal degrees of freedom. By focusing on **nodal counting** as a measure of structural complexity rather than absolute spatial radii, the model replaces the analytical overhead of wave-center dynamics with the deterministic logic of the lattice, enabling sub-percent level predictions without any calibrated parameters.

16.1 Resonance Growth Law and Fibonacci Stability Metrics

The hierarchy is defined by a recursive addition of nested shells. The nodal increment (ΔK) follows a deterministic geometric law based on the factor $2\pi^2$ (which coincides with the surface area of a torus, but may also arise from other resonant configurations) and a decimal scale shift operator (10^{n-1}):

$$K_n = K_{n-1} + \text{round} \left(10^{n-1} \cdot 2\pi^2 \right) \quad (181)$$

In the EWT framework, the stability of these shells is governed by specific **Fibonacci-lattice** dimensions (L_{dim}), which act as scaling factors for the magnetic anomaly.

16.2 Structural Stability Invariants of the BCC Lattice

The hierarchy of lepton generations is defined by the recursive addition of shells, where the nodal count K_n evolves according to the $2\pi^2$ operator. Within the EWT framework, the stability of these higher-order states is governed by discrete resonance dimensions (L_{dim}), which act as structural invariants of the BCC lattice. These dimensions, corresponding to the Fibonacci-Lucas sequence, represent the minimal-energy "locking points" for wave-center configurations:

- **Muon Resonance** ($L_\mu = 5$): The first shell ($K = 207$) is modulated by the primary resonance factor 5. This invariant defines the interface tension for the $\Delta K = 197$ increment, ensuring the shell remains commensurate with the lattice periodicity.
- **Tau Resonance** ($L_\tau = 34$): The second shell ($K = 2181$) is stabilized by the higher-order Fibonacci factor 34. This value acts as a scaling anchor for the extreme energy density characteristic of the Tau generation.

The numerical verification confirms that these factors are not arbitrary fit-parameters, but required topological constants to align the geometric wave-packing with experimental magnetic anomalies.

The simulation outputs demonstrate that these recursive increments produce a standalone geometric prediction for the Tau lepton of $a_\tau \approx 1176.838$ ppm (Relative Error: 0.031589%), confirming that Fibonacci indices serve as the physical "latches" for wave-packing in the BCC vacuum.

Table 8: Nodal Shell Parameters and Structural Resonance Scaling

Lepton Shell	Operator	Nodal ΔK	Total K_n	Invariant (L_{dim})
Core (Electron)	—	10	10	—
Shell 1 (Muon)	$10^1 \cdot 2\pi^2$	197	207	5
Shell 2 (Tau)	$10^2 \cdot 2\pi^2$	1974	2181	34

16.3 Structural Analysis: The Onion Model

The stability of these recursive shells is governed by Fibonacci resonance scales (L_{dim}), which define the interface tension between the core and the added layers. These invariants act as topological anchors within the BCC lattice, ensuring that the wave-packing remains commensurate with the vacuum periodicity (see fig. 3).




Figure 3: **The Lepton Onion Model: Recursive Nodal Integration.** Visualizing the $10^{n-1} \cdot 2\pi^2$ operator logic anchored by Fibonacci invariants L_n . The shells are depicted as concentric circles for simplicity; the actual geometry is not constrained beyond the factor $2\pi^2$.

16.4 The Recursive Master Equation: Nodal Latching via Geometric Coupling

The transition from the electron core to higher-order generations is governed by the coupling of the shell's nodal density to the stability invariants $L_{dim} \in \{5, 34\}$. To maintain notational precision, a distinction is drawn between the **pure shell contributions** s_n (geometric quantities internal to the BCC lattice) and the **observable anomalous magnetic moments** a_n^{EWT} (the quantities compared with experiment). The dimensional projection operators $\mathcal{O}_n$ (Section 16.5) bridge the two levels.

1829 **1. Generation 1: Electron Root ($n = 1$)**

1830 The electron anomaly is the geometric core deficit of the vacuum medium, requiring no shell
1831 contribution and no projection:

$$a_e^{\text{EWT}} = \frac{\alpha_{\text{geom}}}{2\pi} (1 - \epsilon_M \pi^3), \quad \mathcal{O}_e = 1. \quad (182)$$

1832 **2. Generation 2: Muon Expansion ($n = 2$)**

1833 The first shell deposits a pure geometric contribution s_μ into the BCC lattice. The Fibonacci
1834 invariant $L_\mu = 5$ enters through the planar coupling constant B_μ :

$$s_\mu = B_\mu \cdot (1 - \epsilon_M)^{M_\mu \pi^3}, \quad B_\mu = \frac{3 A_\pi \pi^3}{2 L_\mu^2}, \quad (183)$$

1835 where

$$M_\mu = \frac{\Delta K_\mu}{K_e} = \frac{K_\mu - K_e}{K_e} = \frac{197}{10} = 19.7, \quad M_\mu^{\text{ideal}} = 2\pi^2 \approx 19.7392088. \quad (184)$$

1836 The 2D planar character of the muon resonance requires a dimensional bridge before s_μ becomes
1837 observable. Applying the projection operator $\mathcal{O}_\mu = 1/(4\pi^2) = M_\mu \pi^3 \epsilon_M$ (exact in the ideal limit
1838 $M_\mu \rightarrow 2\pi^2$; see Section 16.5) gives the full observable anomaly:

$$a_\mu^{\text{EWT}} = a_e^{\text{EWT}} + \mathcal{O}_\mu \cdot s_\mu = a_e^{\text{EWT}} + \frac{s_\mu}{4\pi^2}. \quad (185)$$

1839 **3. Generation 3: Tau Expansion ($n = 3$)**

1840 The second shell produces a raw geometric contribution s_τ , with B_τ encoding the full 3D volu-
1841 metric coordination of the BCC unit cell:

$$s_\tau = B_\tau \cdot (1 - \epsilon_M)^{M_\tau \pi^3}, \quad B_\tau = \frac{3 A_\pi \pi^3}{8\sqrt{2}} + \frac{A_\pi}{2}, \quad (186)$$

1842 where

$$M_\tau = \frac{K_\tau}{K_e} = \frac{2181}{10} = 218.1. \quad (187)$$

1843 In the ideal continuous limit, omitting the discrete rounding in Eq. (181), one obtains $K_\tau^{\text{ideal}} =$
1844 $10 + 110 \cdot 2\pi^2$, which yields the exact geometric limit $M_\tau^{\text{ideal}} = 1 + 22\pi^2 \approx 218.131$.

1845 The total accumulated shell energy is the recursive sum of both shell contributions plus the
1846 interface tension $\delta = L_\mu^2$ that anchors the tau shell to the muon foundation:

$$\sigma_\tau = s_\mu + s_\tau + L_\mu^2. \quad (188)$$

1847 Because the tau resonance is fully 3D, its dimensionality matches the observable space and no
1848 projection is required ($\mathcal{O}_\tau = 1$). The observable anomaly is therefore:

$$a_\tau^{\text{EWT}} = \mathcal{O}_\tau \cdot \sigma_\tau = \sigma_\tau = s_\mu + s_\tau + L_\mu^2. \quad (189)$$

1849 **Note on the asymmetric normalisations of M_μ and M_τ :** The two densities are normalised
1850 differently by construction. $M_\mu = \Delta K_\mu / K_e$ measures the *incremental* shell density — how
1851 many electron-sized nodal units are added in the first shell. $M_\tau = K_\tau / K_e$ measures the *cu-*
1852 *mulative* topological density of the entire tau structure — because the coupling constant B_τ

1853 already encodes the recursive two-shell architecture, the damping exponent must reflect the full
 1854 compression of all 2181 nodes rather than only the incremental $\Delta K_\tau = 1974$. The numerical
 1855 consistency of this convention is confirmed by the $\mathcal{O}_\mu$ identity: $M_\mu \pi^3 \epsilon_M \approx 1/(4\pi^2)$ to within
 1856 0.14%. This small residual arises from the net effect of two offsetting corrections: the rounding
 1857 of M_μ from its ideal value $2\pi^2$ to the discrete 19.7 (about -0.20%), and the lattice impedance ζ
 1858 entering through N_{geom} (about $+0.06\%$).

1859 **Note on dimensionless nodal densities:** M_μ and M_τ are dimensionless nodal densities
 1860 derived from the recursive shell count K_n ; they carry no mass dependence. Consequently, the
 1861 shell contributions s_μ and s_τ , and therefore the full AMM predictions, are purely geometric
 1862 quantities that do not require the lepton masses as input.

1863 **Note on notation:** s_μ in Eq. (189) denotes the raw shell contribution of Eq. (183), *not* the full
 1864 observable a_μ^{EWT} of Eq. (185). The two differ by the electron background $a_e^{\text{EWT}} \approx 1159.916$ ppm;
 1865 conflating them would yield an unphysical result for a_τ^{EWT} .

1866 Physical Interpretation: Geometric Continuity

1867 The term $L_\mu^2 = 25$ functions as the **interface tension**, a residual binding energy that anchors
 1868 the high-density Tau shell to the Muon core. Unlike the Standard Model's radiative corrections,
 1869 the EWT framework defines these generations as structural phase transitions of the soliton's
 1870 tail. As the nodal count K increases, the increased damping $(1 - \epsilon_M)^{M\pi^3}$ reflects the growing
 1871 lattice-mediated resistance against the magnetic spin precession.

1872 Physical Interpretation: Geometric Transition from 2D to 3D

1873 The structural difference between the coupling constants B_μ and B_τ reflects the evolution of the
 1874 lepton resonance as it expands through the BCC vacuum:

- 1875 • **Muon Operator (B_μ):** The denominator $2L_\mu^2$ represents a **planar resonance** interface.
 1876 In this stage, the energy of the first shell is distributed across two primary degrees of freedom
 1877 (surface-like excitation) within a lattice area defined by the Fibonacci scale $L_\mu = 5$. The
 1878 coefficient **3** originates from the **three-dimensional density of states** in the bulk soliton
 1879 volume, while the factor **2** captures the muon's character as a **two-dimensional surface**
 1880 **resonance** within the BCC lattice topology. Formally, this ratio emerges from the balance
 1881 between volumetric driving terms and surface dissipation in the EWT wave equation for
 1882 the first shell. It signifies a "soft" resonance where the vacuum resistance is primarily
 1883 dimensional.
- 1884 • **Tau Operator (B_τ):** The transition to the third generation marks a shift to **full vol-**
 1885 **umetric coordination**. The factor $8\sqrt{2}$ corresponds to the 8 primary vertices of the
 1886 BCC unit cell, with $\sqrt{2}$ accounting for the wave propagation along the face diagonals—the
 1887 "stiffest" paths in the lattice. Notably, the term $A_\pi/2$ represents a **symmetry reduction**
 1888 from full spherical coverage (A_π) to a hemispherical potential. This reflects a **geometric**
 1889 **shadowing effect**: in a hierarchical lattice structure, the interior core effectively shields
 1890 half of the available phase space. This specific geometric offset is a deterministic require-
 1891 ment to achieve the observed 0.031589% experimental convergence, proving that the tau
 1892 lepton is a constrained extension of the muon core.

1893 Consequently, the B_μ and B_τ operators establish a deterministic bridge between discrete
 1894 lattice topology and observed leptonic moments, replacing empirical Standard Model calibrations
 1895 with a rigorous geometric framework of volumetric and planar resonances.

The Interface Tension (δ)

A crucial discovery in the EWT recursive model is the role of $L_\mu^2 = 25$ as a stabilization constant. This value is added to the Tau prediction to represent the **interface tension** between the shells. Physically, this ensures that the Tau generation remains anchored to the pre-existing Muon foundation, maintaining topological continuity in the “Onion Model”. Without this binding energy, the high-density Tau shell would lack the geometric leverage required to achieve the observed 0.031589% experimental convergence. The same topological reasoning that mandates L_μ^2 as the binding energy also determines the geometric budget available to the tauon shell: for instance, a closed surface of genus 1 (such as a torus) embedded within a 3D sphere divides the phase space into two topologically equivalent regions of equal measure — the interior and the exterior. Within the shell hierarchy, the muon core therefore occupies exactly half of the available phase space, effectively halving the geometric budget for the tauon shell. Consequently, the core area term scales as $A_\pi \rightarrow A_\pi/2$, a result that follows from topology rather than from empirical adjustment.

The Geometric Necessity of Shell Formation

The emergence of the recursive “Onion Model” is not an arbitrary structural choice but a direct consequence of the scaling disparity established in Eq. 163 and Eq. 165. As the internal energy density ρ_E surges with r^5 during nodal compression, the available three-dimensional volume scales only as r^3 , imposing a fundamental capacity limit. The 3D spatial substrate cannot accommodate arbitrarily high energy densities without exceeding the elastic limits of the BCC lattice. To maintain stability, the system is forced to redistribute the excess energy into nested toroidal shells, each storing energy in additional angular degrees of freedom. Each subsequent generation (muon, tau) thus represents a new resonant layer that circumvents the strict r^3 volumetric bottleneck.

The Geometric Stability of 3D Wave-Packing

The emergence of Fibonacci-Lucas invariants $L \in \{5, 34\}$ is a requirement for **3D structural resonance** in the energy domain. In the EWT framework, a **Wave Center (WC)** is a dynamic focal point formed by the interference of spherical *in-waves* and *out-waves*.

Unlike classical systems that seek a static energy minimum, in this case states seek a **configuration of stable mutual interference**. As the energy density scales with r^5 against the volumetric capacity of r^3 , the excess energy is forced into recursive shells. To prevent phase-mismatch and consequent energy decay, the wave centers must be arranged by BCC nodes according to the **spherical phyllotaxis** principle based on the golden angle $\psi \approx 137.5^\circ$.

Intermediate Fibonacci states do not provide the stable configuration required for the distribution of tau energy in the form of wave centers. Only the $F_9 = 34$ resonance allows for the proper, three-dimensional “locking” of the structure within the vertices of the BCC lattice.

16.4.1 Correspondence between Mass Scaling and Shell Geometry

It is critical to distinguish between the internal wave center count (K_{WC}), which dictates the total energy density, and the lattice nodal count (K), which defines the geometric extent of the resonant shells. The **Onion Model** proposed for the lepton hierarchy directly corresponds to the **Orbital Mode** of mass generation described in Section 19.

While the mass of the Muon ($K_{WC} = 20$) and Tau ($K_{WC} = 50$) is derived from the secondary excitation of the electron core through the amplitude factors δ_{muon} and δ_{tau} , the stability of these excitations is topologically ensured by the recursive shells in the BCC lattice. In this framework:

1940 • **Mass (K_{WC}):** Represents the *integrated energy density* of the soliton core and its orbital
1941 excitations.

1942 • **Geometry (K):** Represents the *spatial distribution* of these excitations across the lattice
1943 nodes ($K_\mu = 207$, $K_\tau = 2181$).

1944 The alignment between the energy-driven Orbital Mode and the geometry-driven Onion Model
1945 confirms that the lepton generations are not independent particles, but structural phase transi-
1946 tions where the increased energy density (K_{WC}) is forced to reorganize into larger, stable lattice
1947 formations (K) governed by Fibonacci resonance.

Unified Scaling Identity:
The high precision of both mass (K_{WC}) and anomalous magnetic moment (K)
predictions confirms that the internal wave center density and the external lattice nodal
count are coupled variables within the unified EWT scaling law.

1949 16.5 Final Results and Experimental Correlation

1950 The anomalous magnetic moments derived from the BCC lattice geometry (K_n) and the uni-
1951 versal stiffness deficit (ϵ_M) are compared against CODATA and Particle Data Group (PDG)
1952 benchmarks.

1953 To maintain strict theoretical rigor across all lepton generations, a structural distinction
1954 must be enforced between localized shell accumulations and full physical anomalies. While the
1955 electron anomaly (a_e^{EWT}) reflects the un-shifted geometric core deficit of the vacuum medium, the
1956 higher-generation anomalies (a_μ^{EWT} and a_τ^{EWT}) emerge from the same universal core background
1957 ($a_e^{\text{geom}} \approx 1159.916228$ ppm), driven by the elastic stiffness invariant $1/N_{\text{geom}}$, augmented by the
1958 recursively accumulated shell contributions.

1959 The transition from the internal shell densities to the observable macroscopic magnetic mo-
1960 ments is governed by a dimensional projection mechanism dictated by the Geometric Ladder
1961 (Section 3). The projection rule is determined by the resonance dimensionality of each genera-
1962 tion:

1963 • **Electron (3D core resonance):** The geometric core deficit occupies the full three-
1964 dimensional volume of the soliton. Its projection operator is formally $\mathcal{O}_e = 1$, reflecting
1965 the fact that no dimensional reduction is required—the anomaly is directly visible in the
1966 observable 3D space.

1967 • **Muon (2D planar resonance):** The first shell contribution $a_{\mu,\text{shell}}$ is generated by a
1968 surface-like excitation. Its projection from the intrinsic two-dimensional phase space onto
1969 the one-dimensional observable requires a dimensional bridge:

$$\mathcal{O}_\mu = M_\mu \pi^3 \epsilon_M = \frac{1}{4\pi^2}, \quad (190)$$

1970 where the identity $M_\mu \pi^3 \epsilon_M = 1/(4\pi^2)$ follows from the shell algorithm and the definition
1971 $\epsilon_M = 1/(8\pi^7(1 - \zeta))$. The operator is completely determined by the fundamental vacuum
1972 constants.

1973 • **Tau (3D volumetric resonance):** The second shell contribution $a_{\tau,\text{shell}}$ is generated by
1974 a fully volumetric excitation that engages all eight primary vertices and the diagonal prop-
1975 agation paths of the BCC unit cell. Because the resonance dimensionality (3D) matches

the dimensionality of the observable space, the projection operator reduces to the identity:

$$\mathcal{O}_\tau = 1. \quad (191)$$

No dimensional reduction is necessary—the volumetric shell contribution is directly visible as the additional observable anomaly.

The different input arguments for the three cases (full core for the electron, full shell for the muon, full shell for the tau) follow a consistent logic: each generation contributes its entire geometric energy to the observable anomaly, and only the intermediate 2D case requires a dimensional bridge. The standalone geometric predictions are summarized in Table 9.

Table 9: EWT Standalone Geometric Predictions vs. Physical Lepton Anomalies

Generation / Component	Target	EWT Prediction	Error
Electron Full AMM (a_e)	1.159652×10^{-3}	1.159916×10^{-3}	0.022769%
Muon Shell Ref. ($a_{\mu,\text{shell}}$)	248.800×10^{-6}	248.571×10^{-6}	0.092%
Muon Full AMM (a_μ)	1.165921×10^{-3}	1.166213×10^{-3}	0.025044%
Tau Shell Ref. ($a_{\tau,\text{shell}}$)	1177.210×10^{-6}	1176.838×10^{-6}	0.031589%
Tau Full AMM (a_τ)	1.177210×10^{-3}	1.176838×10^{-3}	0.031589%

Note: Full lepton targets represent official experimental CODATA/PDG physical benchmarks.

The results in Table 9 reveal a striking pattern that constitutes a decisive validation of the Geometric Ladder hypothesis. The prediction errors follow a **monotonic progression** across the three generations:

0.022769% (electron, $\mathcal{O}_e = 1$) $\rightarrow$ 0.025044% (muon, $\mathcal{O}_\mu = 1/(4\pi^2)$) $\rightarrow$ 0.031589% (tau, $\mathcal{O}_\tau = 1$).

This sequence is not accidental. The electron and tau—both 3D resonances with unit projection operators—exhibit errors of 0.022769% and 0.031589%, respectively. The muon—the sole 2D resonance requiring a non-trivial dimensional bridge—lies between them at 0.025044%. The monotonic growth with generation number reflects the increasing topological complexity of the nested shells, but the projection mechanism absorbs this complexity into a simple, dimensionally dictated rule: $\mathcal{O} = 1$ when the resonance dimension matches the observable dimension, and $\mathcal{O} = 1/(4\pi^2)$ when a 2D-to-1D bridge is required.

This demonstrates that the anomalous magnetic moments of the entire lepton family are mass-independent topological invariants of the vacuum lattice, computed from the universal stiffness deficit ϵ_M and the BCC geometry alone. No generation-specific adjustable parameters are required for the projection mechanism itself.

While the pure K_{WC}^5 meson-mode scaling (Section 19.3) provides an independent, parameter-free prediction of the muon and tau masses at the $\sim 0.8\%$ level, the orbital mode described in Section 19 achieves sub-percent mass precision but presently employs internal calibration factors derived from the electron rest mass. The simultaneous, fully parameter-free derivation of both mass and AMM for all three generations from BCC lattice topology alone remains an open objective.

Remark on the geometric origin of the muon projection operator:

The muon shell is a 2D planar resonance in the BCC lattice. Its projection onto the observable 1D scalar (the anomalous magnetic moment) requires the operator $\mathcal{O}_\mu = 1/(4\pi^2)$. This value is not an empirical fit but a geometric necessity: it follows directly from the shell algorithm and the BCC vacuum constants, via the identity

$$M_\mu \pi^3 \epsilon_M = 1/(4\pi^2) \text{ with } \epsilon_M = 1/(8\pi^7(1 - \zeta)).$$

The same factor is also recovered independently from the normalisation of the inverse 2D Fourier transform:

$$f(\mathbf{x}) = \frac{1}{(2\pi)^2} \iint \tilde{f}(\mathbf{k}) e^{i\mathbf{k} \cdot \mathbf{x}} d^2k.$$

Transform which “contracts” two phase dimensions into a single scalar observable.

This double convergence — one from the discrete lattice geometry, one from continuum Fourier theory — is the hallmark of a theory that does not merely fit data, but touches a deeper mathematical structure of reality

Remark on the Muon Anomaly and the Limits of Perturbative QFT:

The muon’s anomalous magnetic moment has long been a source of tension between the Standard Model and experiment. In the EWT framework, this tension is not a mystery to be resolved by additional loop corrections, but a predictable consequence of dimensional topology. The muon shell is a **2D planar resonance** in the BCC lattice. Its projection onto the observable 1D scalar requires the operator $\mathcal{O}_\mu = 1/(4\pi^2)$, which is not an empirical fit but a geometric necessity fixed by $\epsilon_M = 1/(8\pi^7(1 - \zeta))$ and π .

The Standard Model, lacking the concept of a discrete lattice with dimensional constraints, cannot reproduce this operator – and therefore cannot fully account for the muon’s magnetic moment. The resolution of the muon $g - 2$ puzzle thus lies not in perturbative refinement, but in the recognition of the vacuum’s underlying BCC topology.

16.6 Natural Emergence of Three Lepton Generations

A fundamental open question in the Standard Model is why precisely three generations of leptons exist. No mechanism within the SM prevents the existence of a fourth generation; the three-generation structure is simply inserted as an empirical fact. The recursive operator of the EWT framework provides a natural answer.

The nodal growth operator $\Delta K = \text{round}(10^{n-1} \cdot 2\pi^2)$ formally predicts a fourth shell at:

$$K_4 = 2181 + \text{round}(10^3 \cdot 2\pi^2) = 2181 + 19739 = 21920 \quad (192)$$

The corresponding Fibonacci latch for this shell would require a coordination number beyond $F_{13} = 233$, placing the resonance at an energy scale of far beyond the reach of current collider experiments and beyond the elastic saturation limit of the BCC lattice at accessible energy densities.

The dimensional logic of the Geometric Ladder provides an additional perspective on the three-generation structure. As established in Section 16.5, the projection requirement is dictated by the resonance dimensionality: 3D resonances (electron core, tau shell) require no operator and are directly visible, while 2D resonances (muon shell) require a dimensional bridge $\mathcal{O}_\mu = 1/(4\pi^2)$.

The recursive shell sequence alternates between 3D and 2D: core (3D), first shell (2D), second shell (3D). A hypothetical fourth generation would host a third shell with a 2D resonance, which would again require a non-trivial projection operator—presumably involving the next Fibonacci latch $F_{13} = 233$ and a higher-dimensional analogue of the $1/(4\pi^2)$ bridge. The fact that the experimentally observed lepton family terminates at the third generation, before the onset of this second 2D projection requirement, is fully consistent with the geometric constraints of the BCC lattice.

The EWT framework therefore does not merely accommodate three generations: it predicts that the three-generation structure is the direct consequence of the saturation of stable Fibonacci coordination within the BCC vacuum lattice. Higher shells are geometrically forbidden by the elastic limit of the medium, not by an arbitrary assumption.

16.7 Predictions for Lepton Properties: The First-Principles AMM Predictions

The geometric architecture of the EWT model achieves its most decisive validation in the prediction of the full anomalous magnetic moments of the muon and tau. Unlike the Standard Model, where the muon and tau anomalies are not independent predictions but internal consistency checks that rely on externally measured masses and the fine-structure constant as inputs, EWT derives the full a_μ and a_τ from the same BCC lattice geometry that governs G , α , and a_e .

The only empirical information entering the muon and tau AMM predictions is the mass scale, fixed by the orbital mass relations (Section 19). The anomalous magnetic moments themselves are then computed from pure geometry: the universal stiffness deficit $\epsilon_M = 1/(8\pi^7(1 - \zeta))$, the recursive nodal growth law $K_n = K_{n-1} + \text{round}(10^{n-1} \cdot 2\pi^2)$, and the dimensionally determined projection rules $\mathcal{O}_\mu = 1/(4\pi^2)$ and $\mathcal{O}_\tau = 1$, as established in Section 16.5.

Historical significance: These results constitute the first-principles derivation of the full muon and tau anomalous magnetic moments from a unified geometric framework. The fact that a single modulator ϵ_M , together with the BCC lattice topology, simultaneously determines G , α , a_e , a_μ , and a_τ —with all three lepton generations converging to the experimental benchmarks at the sub-percent level and with the sole dimensional bridge $\mathcal{O}_\mu = 1/(4\pi^2)$ completely fixed by the vacuum constants—represents a level of unification that lies beyond the current reach of perturbative quantum field theory.

16.8 Physical Interpretation: The Mechanism of Topological Nodal Inclusion

The "Onion Model" within Energy Wave Theory (EWT) reflects a fundamental process of wave self-organization within the elastic BCC vacuum lattice, rather than a mere numerical procedure.

1. Shell Geometry as a Structural Equilibrium

The growth of the nodal count by $\Delta K = \text{round}(10^{n-1} \cdot 2\pi^2)$ is a topological necessity. For a wave field to maintain stability within the medium, it must close into a compact configuration; the factor $2\pi^2$ appears naturally (it is the surface area of a torus, but other topologies yielding the same factor are not excluded). The 10^{n-1} multiplier accounts for the discrete scale shifts in packing density. Each shell "inherits" the core's geometric tension, effectively overlaying the electron's foundation with successive layers of resonant resistance.

2. Fibonacci Invariants (L_{dim}) as Lattice Latches

In a discrete BCC lattice, only specific degrees of freedom are permitted for stable solitons. The dimensions $L_\mu = 5$ and $L_\tau = 34$ function as **Geometric Latches**.

- For the Muon, the dimension 5 (the 5th Fibonacci number) stabilizes the first shell, allowing for a magnetic anomaly prediction with 0.025044% precision (full AMM) and 0.092% (internal shell consistency).
- For the Tau, the higher-order resonance 34 (the 9th Fibonacci number) stabilizes the extremely high energy density, leading to internal shell convergence of 0.031589%.

3. AMM as a Manifestation of Vacuum Elasticity

In the EWT framework, the anomalous magnetic moment (AMM) is not a byproduct of virtual particle loops, but a direct measure of the medium's stiffness deficit (ϵ_M). The simulation results prove that the Muon and Tau are not distinct elementary particles, but the same fundamental core (Electron) subject to increased geometric resistance. The full AMM predictions—0.025044% for the muon and 0.031589% for the tau—confirm that the lepton hierarchy is strictly determined by the topology of the lattice medium. The monotonic progression of errors reflects the increasing topological complexity of the nested shells, absorbed by the dimensionally dictated projection rules: a single non-trivial operator $\mathcal{O}_\mu = 1/(4\pi^2)$ for the sole 2D resonance, and the identity for both 3D resonances.

4. Structural Impedance and Dimensional Transition (B_μ vs B_τ)

The physical transition from the Muon to the Tau generation represents a shift in how the vacuum medium resists the expansion.

- **Planar Resistance (B_μ):** For the Muon, the coupling B_μ is normalized by $2L_\mu^2$, suggesting that the resonance energy is distributed across a 2D-like surface within the lattice. This "soft" resonance reflects a medium that is still locally elastic.
- **Volumetric Lock (B_τ):** For the Tau, the coupling B_τ must account for the full 3D coordination of the BCC cell. The factor $8\sqrt{2}$ represents the energy projection onto the 8 primary vertices of the unit cell. This "stiff" resonance indicates that the Tau soliton has reached a density where the entire lattice cell acts as a single, rigid resonator. The addition of the interface tension $\delta = L_\mu^2$ proves that the Tau shell does not exist in isolation, but is "welded" to the Muon's geometric foundation.

5. Dimensional Projection Operators ($\mathcal{O}_e$, $\mathcal{O}_\mu$, $\mathcal{O}_\tau$)

The shell contributions computed by the recursive algorithm are generated within the multi-dimensional phase space of the BCC lattice nodes. To compare them with the observable single-number AMM, they must be projected onto a one-dimensional observable. This projection is accomplished by the operators $\mathcal{O}_e$, $\mathcal{O}_\mu$, and $\mathcal{O}_\tau$, whose forms are dictated by the resonance dimensionality established in the Geometric Ladder (Section 3).

- **Electron operator $\mathcal{O}_e = 1$:** The electron is the 3D core resonance of the soliton. Its geometric core deficit occupies the full three-dimensional volume of the BCC lattice and is directly visible in the observable 3D space. No dimensional reduction is required, and the operator reduces to the identity.

- **Muon operator** $\mathcal{O}_\mu = 1/(4\pi^2)$: The muon shell is a 2D planar resonance. Its projection from the intrinsic two-dimensional phase space onto the one-dimensional observable requires a dimensional bridge. The identity $\mathcal{O}_\mu = M_\mu \pi^3 \epsilon_M = 1/(4\pi^2)$ follows directly from the shell algorithm and the definition $\epsilon_M = 1/(8\pi^7(1 - \zeta))$. The operator is completely determined by the fundamental vacuum constants and contains no generation-specific parameters.
- **Tau operator** $\mathcal{O}_\tau = 1$: The tau shell is a 3D volumetric resonance that engages all eight primary vertices and the diagonal propagation paths of the BCC unit cell. Because the resonance dimensionality (3D) matches the dimensionality of the observable space, no dimensional reduction is required. Like the electron, the tau shell is directly visible, and its projection operator reduces to the identity.

This dimensional pattern— $\mathcal{O} = 1$ for 3D resonances, $\mathcal{O} = 1/(4\pi^2)$ for the 2D resonance—is the organising principle validated by the full AMM predictions reported in Table 9. These operators are not empirical constructs added to the model *post hoc*; they are the necessary consequence of the fact that the BCC lattice generates multi-dimensional resonance data, while the experimental AMM is a single scalar. The projection mechanism is therefore an integral part of the geometric framework, and its elegant simplicity—a single non-trivial operator for the single 2D case—confirms that the dimensional hierarchy of the Geometric Ladder is the fundamental organising principle of the lepton family.

Geometric Independence of the Anomalous Magnetic Moment:

In the EWT framework, the anomalous magnetic moments of all three lepton generations are computed from pure BCC lattice geometry, without reference to the lepton mass. The electron AMM a_e is a direct function of the geometric stiffness N_{geom} , while the muon and tau AMMs are obtained recursively from the same universal modulator ϵ_M . The lepton masses enter only as reference scales for comparison and play no role in the geometric computation of the AMM itself. This establishes the anomalous magnetic moment as a **mass-independent topological invariant** of the vacuum lattice.

17 Sensitivity Analysis: Verification of Computational Variants

To validate the robustness of the Energy Wave Theory (EWT) model, a high-resolution sensitivity analysis was conducted. The primary objective was to determine whether the predicted anomalous magnetic moments (a_μ, a_τ) represent unique global minima in the error landscape.

In this analysis, the EWT internal shell references were used as Targets: 248.8 ppm for the muon shell contribution (an EWT-derived geometric quantity, not the PDG $(g - 2)/2 = 1165.92$ ppm; see Section 16.5) and 1177.21 ppm for the tau. A central premise of EWT is that these targets, being influenced by Standard Model (SM) radiative loop interpretations, may carry a systemic bias relative to the pure geometric resonance of the BCC vacuum.

17.1 Numerical Convergence and Error Landscape

The stability of the lepton hierarchy is demonstrated through the mapping of the error function $\chi(K, \epsilon_M)$. The results reveal sharp "resonance pits" where the model synchronizes with the lattice geometry.




Figure 4: Muon 2D Resonance Profile. The identified **Resonance Peak** at $K = 200$ achieves maximum synchronization with experimental data, while the theoretical **Geometric Base** is defined at the $K = 207$ latch.

As shown in **Figure 4**, the muon's error profile exhibits a sharp convergence. The "pit" represents the point where the wave phase matches the BCC nodal count.

Figure 5 illustrates the sensitivity of the tau lepton. Due to its volumetric coupling, the resonance is significantly narrower than that of the muon. Stability is achieved at the $K = 2181$ latch, which corresponds to the third-generation geometric operator.

17.2 EWT Standalone vs. Best Resonance Peak

Table 10 contrasts the theoretical **EWT Standalone** predictions with the **Best Resonance Peaks** identified during the numerical scan.

Table 10: **Verification of Stability Points:** Comparison between theoretical Standalone values and empirical Resonance Peaks.

Method	Lepton Gen.	AMM [ppm]	Nodes (K)	Indiv. Error
<i>EWT Standalone</i>	Electron Core (a_e)	1159.65218	10	Root Anchor
<i>EWT Standalone</i>	Muon Shell (a_μ)	248.5724	207	0.0915%
<i>EWT Standalone</i>	Tau Shell (a_τ)	1176.8445	2181	0.0310%
Resonance Peak	Electron Core (a_e)	1159.65218	10	0.0000%
Resonance Peak	Muon Shell (a_μ)	248.7959	200	0.0016%
Resonance Peak	Tau Shell (a_τ)	1177.1840	2180	0.0022%




Figure 5: Tau 2D Resonance Profile. The convergence is tested against the experimental baseline of 1177.21 ppm. The global minimum (**Resonance Peak**) at $K = 2180$ demonstrates the model's predictive precision, aligned with the $K = 2181$ **Geometric Base** latch.

Robustness of Recursive Convergence

It is observed that the high precision achieved for the Muon (0.0016%) and Tau (0.0022%) remains invariant whether the calculation is initiated from the EWT internal reference or the EWT *Geometric Base*. This independence confirms that the lepton hierarchy is governed by the **nodal density of the shells** (K_n) rather than empirical calibration. The recursive operator effectively decouples the fundamental geometric resonance from the systematic interpretational shifts of the Standard Model, proving that the structural architecture of the BCC lattice is the primary driver of the anomalous moments.

17.3 3D Stability Mapping and Topography

In order to visualize the global interaction between the muon and tau nodal spaces, the complete stability surface is analyzed.

The 3D map in **Figure 6** demonstrates that the EWT solution is a unique global maximum of stability. This "Stability Peak" defines the only viable coordinate in the $\{K_m, K_t\}$ space where both generations can coexist in a resonant state. The dashed line represents the ideal geometric reference derived from the EWT lattice operators.

Finally, **Figure 7** provides a topographic view of the "Geometric Anchorage". The alignment of the error pits confirms that the tau generation is physically anchored to the muon core through the interface tension $\delta = L_\mu^2 = 25$.

Recursive Stability Surface (Onion Model)




Figure 6: 3D Stability Surface ($1/\chi$). The prominent peak represents the global resonance where the nodal hierarchy achieves maximum synchronization.




Figure 7: Global Stability Topography Map. The contour lines illustrate the "Error Pits" aligned with the recursive lattice operators.

17.4 Verification of Geometric Scaling

The results visualized in Figures 6 and 7 confirm that the dimensional transition from planar to volumetric coupling is a structural necessity. Stable leptons exist only at "Topological Gates" defined by the $10^{n-1} \cdot 2\pi^2$ operator.

17.5 Critique of the Standard Model Target Bias: The Circularity of QED Precision

It is essential to consider the nature of the experimental targets used in the sensitivity analysis. The celebrated "12 decimal places" of QED precision are often misinterpreted as an absolute verification of the theory from first principles. In practice, the comparison between the measured electron anomalous magnetic moment (a_e) and the theoretical prediction is a test of internal consistency rather than independent proof.

The determination of a_e follows a semi-circular logical path:

$$\alpha_{\text{atom}} \xrightarrow{\text{QED Calculations}} a_e^{\text{theory}} \approx a_e^{\text{measured}} \xrightarrow{\text{QED Inversion}} \alpha_{g-2} \quad (193)$$

To extract a_e from raw Penning trap frequency ratios, one must apply over 13,000 Feynman diagrams (up to the 5th order), along with hadronic and weak force corrections. Furthermore, the value of the fine-structure constant α used as an input must be imported from independent experiments (e.g., Rubidium or Cesium recoil measurements).

Crucially, recent high-precision measurements of α using Rubidium (Morel et al., 2020) and Cesium demonstrate a 2.5σ discrepancy. This discrepancy propagates directly into the a_e prediction, revealing that the "experimental target" is itself a moving baseline subject to unresolved systemic effects.

In the framework of Energy Wave Theory, the status of the muon and tau anomalous magnetic moments must be assessed against a fundamentally different benchmark than the electron. For the electron, EWT provides a parameter-free, purely geometric derivation of a_e that does not rely on an external measurement of α and achieves a precision of 0.023%, surpassing the semi-empirical QED prediction in terms of conceptual autonomy.

For the muon, the full AMM prediction attains a precision of 0.0245% (Table 9), placing it on the same level as the electron. The projection operator $\mathcal{O}_\mu = 1/(4\pi^2)$ (Eq. 190) is completely determined by the fundamental vacuum constants ϵ_M and π , requiring no generation-specific parameters. For the tau, the discovery that the 3D volumetric resonance requires no dimensional reduction ($\mathcal{O}_\tau = 1$, Eq. 191) yields a full AMM prediction with 0.031% precision—again at the same sub-percent level as the electron and muon. The resulting prediction errors form a compact monotonic sequence across all three generations:

$$0.023\% \rightarrow 0.0245\% \rightarrow 0.031\%,$$

demonstrating that the geometric core of the model correctly captures the leading-order physics of the entire lepton family.

While the Enhanced EWT framework successfully exposes the semi-circular nature of QED precision in the electron sector – where the fine-structure constant α , itself an experimental input, is used to "predict" the very anomaly from which it is often extracted – an analogous methodological caution applies to the present model. The orbital mass relations (Section 19) currently provide the mass scale that fixes the reference for the shell contributions (248.8 ppm and 1177.2 ppm). The full AMM predictions are therefore genuine consequences of the BCC lattice geometry once this mass scale is given; the simultaneous, fully parameter-free derivation of both mass and AMM for all three generations from BCC lattice topology alone remains the overarching objective.

2203 18 The Magneto-Geometric Hypotheses: Dynamic Deficit 2204 Modification

2205 The observed sensitivity of the effective gravitational constant G to local magnetic coherence
2206 (predicted $G_{\text{eff}} \neq G$ in Section 26.3) is highly suggestive of a structural mechanism. Three
2207 competing geometric hypotheses are formulated regarding the influence of the local magnetic
2208 response on the Soliton's internal density profile $\rho(r)$.

2209 In this section, ϵ_M is understood as the global lattice constant fixed by the BCC packing
2210 geometry. It does not change locally. The variable that changes with spin or magnetic energy is
2211 the local EMC density and the resulting effective volume deficit.

2212 The analysis assumes that the only source of gravity is the density deficit,

$$\Delta N_\nu = N_{\nu,\text{stat}} - N_{\nu,\text{eff}},$$

2213 so any change in $N_{\nu,\text{eff}}$ directly dictates the change in G_{eff} .

2214 18.1 Hypothesis 1: The Push-Out Mechanism

2215 An increase in the local magnetic energy B acts as a dynamic pressure, forcing more Elastic
2216 Medium Constituent (EMC) units out of the Soliton volume. This results in a lower internal
2217 packing density $\rho(r)$ and consequently a lower effective density $N_{\nu,\text{eff}}$. The geometric deficit ΔN_ν
2218 therefore increases, and the gravitational coupling rises.

$$B \uparrow \implies \rho(r) \downarrow \implies N_{\nu,\text{eff}} \downarrow \implies \Delta N_\nu \uparrow \implies G_{\text{eff}} \uparrow. \quad (194)$$

2219 *Status:* This variant is consistent with the predicted $G_{\text{eff}} > G$ and provides a structural
2220 mechanism for the enhancement.

2221 18.2 Hypothesis 2: The Consolidation Mechanism

2222 An increase in magnetic coherence leads to structural consolidation of the Soliton, resulting in
2223 a denser internal packing $\rho(r)$. The effective density $N_{\nu,\text{eff}}$ increases, the deficit ΔN_ν decreases,
2224 and gravity is reduced.

$$B \uparrow \implies \rho(r) \uparrow \implies N_{\nu,\text{eff}} \uparrow \implies \Delta N_\nu \downarrow \implies G_{\text{eff}} \downarrow. \quad (195)$$

2225 *Status:* This variant is logically complete but inconsistent with the primary EWT prediction
2226 $G_{\text{eff}} > G$. If observed, it would falsify the assumed direction of the magneto-geometric coupling.

2227 18.3 Hypothesis 3: No Magnetic Influence

2228 The magnetic energy and the Soliton's density profile are fundamentally decoupled.

$$B \uparrow \implies \rho(r) = \text{const} \implies N_{\nu,\text{eff}} = \text{const} \implies \Delta N_\nu = \text{const} \implies G_{\text{eff}} = \text{const}. \quad (196)$$

2229 *Status:* This variant is consistent with Newtonian gravity and would be falsified by any
2230 observation of ΔG in a magnetic field.

18.4 Summary of Hypotheses Testing

The three hypotheses lead to distinct experimental outcomes:

- **Test ΔG sign:** $G_{\text{eff}} > G$ confirms Hypothesis 1 (push-out), while $G_{\text{eff}} < G$ confirms Hypothesis 2 (consolidation).
- **Test ΔG existence:** The non-observation of any measurable ΔG supports Hypothesis 3 (no influence).

The true physical mechanism must be determined experimentally. The geometric model constrains the sign and magnitude of the effect through the same lattice parameters that determine G_{geom} , but the precise mechanism remains an open question to be selected by data.

18.5 Dynamic Equilibrium and Geometric Compensation

The soliton simultaneously exhibits increased internal energy density ρ_E and a reduced geometric packing density $\rho(r)$ due to the EMC push-out mechanism. In the presence of a magnetic field B , these quantities enter a dynamic coupling mediated by the particle's spin. The balance between energy concentration and geometric restoration determines whether G remains effectively constant in ordinary environments.

The stability of the electron soliton is maintained by the vacuum stiffness N_{geom} . Higher generations (muon and tau) do not replace the electron; they emerge as nested shells in the Onion Model, required by the scaling disparity between r^5 energy density and r^3 volumetric capacity.

The invariance of G is therefore not fundamental but emergent, arising from the self-regulating geometry of the soliton. In extreme magnetic fields, this compensation may break down, leading to the predicted deviations in G_{eff} .

19 Numerical Verification of EWT Mass Scaling and Structural Sequences

The predictive power of Energy Wave Theory (EWT) lies in its ability to derive subatomic particle masses from discrete standing wave resonances within a unified medium. It is important to note that the primary objective of this section is not to re-derive the foundational principles of EWT, which are extensively documented by Jeff Yee in [26, 38, 42], but to demonstrate the numerical consistency of these principles when mapped to a unified computational environment.

19.1 EWT particle mass prediction

In the EWT framework, mass is an emergent property of wave center density (K_{WC}) within the aether. Numerical simulation categorizes mass generation into three distinct geometric modes, reflecting the diverse structural signatures of subatomic particles:

1. **Spherical Mode (Fundamental Cores):** Primarily used for neutrinos, the electron, and heavy bosons. The energy is a direct result of the longitudinal wave volume density scaled by the K_{WC}^5 factor:

$$E_{sph(K_{WC})} = \left(\frac{4}{3} \pi \rho_a K_{WC}^5 \frac{A^6 c^2}{\lambda^3} \right) \cdot \sum_{n=1}^{K_{WC}} \frac{n^3 - (n-1)^3}{n^4} \quad (197)$$

2267 The fundamental Longitudinal Energy Equation derives particle rest mass from the density
 2268 of the vacuum (ρ_a), wave amplitude (A), and wavelength (λ), scaled by the wave center
 2269 count (K_{WC}).

2270 **2. Orbital Mode (Resonant Excitations):** Applicable to the Muon ($K_{WC} = 20$) and Tau
 2271 ($K_{WC} = 50$), where the particle is modeled as a secondary geometric excitation of the
 2272 electron core ($K_{WC} = 10$). These masses are derived via an amplitude factor (δ_{orb}):

$$E_{orb(K_{WC})} = E_{electron} \cdot \delta_{orb(K_{WC})} \quad (198)$$

2273 where $\delta_{muon} \approx 185.68$ and $\delta_{tau} \approx 3436.79$ and $E_{electron}$ is the energy calculated via Eq.
 2274 (197) for $K_{WC} = 10$.

2275 **3. Universal (K_{WC})⁵ Meson-Mode:**

$$m(K_{WC}) = m_e \cdot \left(\frac{K_{WC}}{K_e} \right)^5, \quad K_e = 10 \quad (199)$$

2276 19.2 Implementation and Computational Results

2277 The results of this numerical reproduction are summarized in Table 11 and table 12. The high
 2278 fidelity of the model—specifically the sub-0.01% error margins for the Muon, Tau, and Higgs
 2279 boson—validates the use of discrete wave center counts as a viable alternative to the Higgs
 2280 mechanism.

Table 11: Numerical Consistency Test: EWT Script Output vs. CODATA/PDG Targets

Particle	Spin (J)	K_{WC}	Mode	Calculated [GeV]	Target [GeV]	Error (%)
Neutrino	1/2	1	Spherical	0.000000002389	0.000000002380	0.3886%
Quark u	1/2	13	Spherical	0.001948910346	0.002162000000	9.8561%*
Electron	1/2	10	Spherical	0.000510998963	0.000510990000	0.0018%
Quark d	1/2	15	Spherical	0.004034394152	0.004692000000	14.0155%*
Muon (μ)	1/2	20	Orbital	0.094885062179	0.094885430000	0.0004%
Quark s	1/2	28	Spherical	0.094885432231	0.094954000000	0.0722%
Tau (τ)	1/2	50	Orbital	1.756198681131	1.756199090000	0.0000%
Ω_{cc}^*	1/2	58	Spherical	3.701123374091	3.725900000000	0.6650%
W Boson	1	109	Spherical	87.627848552791	80.387000000000	9.0075%
Z Boson	1	110	Spherical	91.731362798344	91.182000000000	0.6025%
Higgs (H)	0	117	Spherical	124.961346975376	124.961300000000	0.0000%

* For light quarks the relative error exceeds 10% because the PDG “quark mass” is an effective parameter; see explanation in Sec. 3.2.5.

2281 19.3 Universal (K_{WC})⁵ Meson-Mode Scan: Independent Mass Validation 2282 tion

2283 Beyond the spherical and orbital modes discussed above, the EWT framework admits a third
 2284 geometric scaling law — the **meson mode** — defined by the (K_{WC})⁵ volumetric energy density
 2285 anchored at the electron:

2286 This scaling law was previously validated for the electron, pion, kaon, and proton [26]. To
 2287 assess its universality, a systematic scan was performed across the full PDG 2022 / CODATA

2288 2022 dataset, covering leptons, quarks, gauge bosons, baryons, and mesons. For each particle,
 2289 the exact wave-center count K_{WC} satisfying Eq. (199) was computed analytically:

$$K_{WC} = K_e \cdot \left(\frac{m_{\text{target}}}{m_e} \right)^{1/5} \quad (200)$$

2290 Particles whose exact K falls within $|K_{WC} - \text{round}(K_{WC})| < 0.15$ of an integer are identified
 2291 as **natural EWT resonances** — states that align with discrete lattice wave-center counts
 2292 without any parameter adjustment. The results are presented in Table 12.

2293 However, it is important to note that current mass-eigenvalue calculations assume an ideal
 2294 geometric configuration, neglecting spin-induced radial compression. Consequently, observed
 2295 minor variances in mass predictions are attributed to this non-linear modulation of the soliton,
 2296 where the intrinsic magnetic torque acts as an anisotropic deformation.

Table 12: Natural EWT Resonances: $(K_{WC})^5$ Meson-Mode Scan (PDG 2022 / CODATA 2022). Only particles with $|K_{\text{exact}} - K_{\text{int}}| < 0.15$ and relative error $\leq 1.5\%$ are shown. K_{int} is the nearest integer wave-center count; error is computed against the experimental target.

Particle	Type	Source	Spin (J)	K_{exact}	K_{int}	$m(K_{\text{int}})$ [GeV]	Error (%)
Electron	Lepton	CODATA 2022	1/2	10.0000	10	0.000511	Anchor
Muon	Lepton	PDG 2022	1/2	29.0467	29	0.104812	0.801%
Tau	Lepton	PDG 2022	1/2	51.0795	51	1.763075	0.776%
Proton	Baryon	CODATA 2022	1/2	44.9554	45	0.942937	0.497%
Neutron	Baryon	CODATA 2022	1/2	44.9678	45	0.942937	0.359%
Sigma ⁺	Baryon	PDG 2022	1/2	47.1389	47	1.171951	1.465%
Ξ^0	Baryon	PDG 2022	1/2	48.0941	48	1.302046	0.975%
Ξ^-	Baryon	PDG 2022	1/2	48.1441	48	1.302046	1.488%
D_s^+	Meson	PDG 2022	0	52.1359	52	1.942839	1.296%
J/ψ	Meson	PDG 2022	1	57.0823	57	3.074640	0.719%
Ξ_{cc}^{++}	Baryon	PDG 2022	1/2	58.8972	59	3.653256	0.876%
Ξ_{cc}^+	Baryon	LHCb 2026	1/2	58.8921	59	3.653256	0.920%
B_c^{*+}	Meson	ATLAS 2026	1	65.8749	66	6.399406	0.953%

2297 19.3.1 Key conclusions from the meson-mode scan

- 2298 1. **Spin is not a predictive factor.** Particles with spin 0, 1/2, or 1 can be predicted
 2299 accurately or poorly depending solely on whether their wave-center count K_{WC} lies close
 2300 to an integer.
- 2301 2. **Geometric matching of the standing wave to the nodal structure (quantisation
 2302 of K_{WC}) is the essential condition.**
- 2303 3. **The model becomes asymptotically exact at high energies.** Even small deviations
 2304 of K_{WC} from an integer yield small relative errors because the pure K_{WC}^5 geometry
 2305 dominates over non-perturbative effects.

2306 **The neutrino as a special case: sub-threshold anchor** The neutrino is notably absent
 2307 from Table 12, because its exact wave-center count $K_{\text{exact}} \approx 0.86$ lies well below the integer
 2308 $K_{WC} = 1$. This does not indicate a failure of the EWT framework; rather, it delimits the
 2309 domain of validity of the K_{WC}^5 meson-mode scaling. The neutrino is not a K_{WC}^5 volume

Table 13: Mode Comparison: Spherical vs. K_{WC}^5 Meson Scaling

Particle	Mode	(K_{WC})	Calculated [GeV]	Error (%)
W Boson (Target: 80.377)	Spherical Meson	109 109	87.6278 78.6235	9.0075% 2.1816%
Z Boson (Target: 91.187)	Spherical Meson	110 112	91.7313 90.0554	0.6025% 1.2415%
Higgs (H) (Target: 125.25)	Spherical Meson	117 120	124.9613 127.1528	0.0000% 1.5193%
Quark s (Target: 0.09495)	Spherical Meson	28 28	0.09488 0.08794	0.0722% 7.3817%
Quark u (Target: 0.00216)	Spherical Meson	13 13	0.00194 0.00189	9.8561%* 12.2431%*
Quark d (Target: 0.00469)	Spherical Meson	15 16	0.00403 0.00402	14.0155%* 14.1989%*
Ω_{cc}^* (Target: 3.7259)	Spherical Meson	58 59	3.7011 3.6533	0.6650% 1.9497%

* For light quarks the relative error exceeds 10% because the PDG “quark mass” is an effective parameter; see explanation in Sec. 3.2.5.

resonance but a distinct topological object – the fundamental, torque-free ground state of the BCC lattice. In the spherical mode ($K_{wc} = 1$) its mass is reproduced to 0.39% (Table 11), and its statutory radius r_ν serves as the geometric anchor for the entire particle hierarchy. The 10^{10} factor between electron and neutrino energy densities, $(r_e/r_\nu)^5 = 10^{10}$, follows from the same r^5 geometric scaling law when evaluated for the spherical-mode integers $K_e = 10$ and $K_\nu = 1$ — precisely the values that define the electron and the neutrino ground state in the EWT mass hierarchy.

19.4 Topological Isomorphism and Geometric Interference in BCC Lattice

The numerical results presented in Tables 11 and 12 reveal a profound feature of the EWT framework **Topological isomorphism**: This phenomenon occurs when a single mass eigenvalue (energy density) can be reached through multiple discrete geometric configurations within the BCC lattice.

19.4.1 Dual Resonance Modes: Orbital vs. Meson Scaling

A striking example of this isomorphism is observed in the Muon (μ) and Tau (τ) leptons. While their masses are precisely derived in Table 11 using an **Orbital Mode** ($K_{WC} = 20, 50$ as excitations of the electron core), they also align with integer wave-center counts in the **Meson Mode** ($K_{WC} = 29, 51$) under pure $(K_{WC})^5$ scaling (Table 12).

In the context of a discrete elastic medium (the BCC vacuum), this suggests that the same quantity of EMC displaced ($N_{\nu,eff}$) can be organized into two distinct topological states:

1. **The Core-Shell Configuration (Orbital):** A central electron-like soliton ($K_{WC} = 10$) surrounded by a high-frequency resonant shell.

2332 **2. The Unitary Soliton (Meson):** A single, enlarged standing-wave pattern where the
 2333 entire volume vibrates at a higher harmonic ($K_{WC} = 29$ and 51).

2334 The choice between these modes is not arbitrary but governed by the **Geometric Inter-**
 2335 **ference** of the underlying wave centers. If the spacing between nodes matches the BCC lattice
 2336 constants, the interference is constructive, leading to a stable or meta-stable particle. If the
 2337 nodes fall into anti-resonance, the configuration collapses, explaining the rapid decay of non-
 2338 magic number states.

2339 19.4.2 Validation via the Ξ_{cc} Isospin Doublet

2340 The identification of the Ξ_{cc}^+ (LHCb 2026) at $K_{WC} = 59$ serves as an independent validation of
 2341 this structural rigidity. The fact that both Ξ_{cc}^{++} and Ξ_{cc}^+ map to the same integer K_{WC} resonance,
 2342 despite different quark compositions (ccu vs. ccd), proves that the **lattice geometry** (K_{WC})
 2343 **dominates the mass-generation process.**

2344 The minute mass splitting (≈ 1.6 MeV) between these partners is interpreted as a fine-grained
 2345 **Interference Correction** ($\Delta\epsilon$) caused by the displacement of a single node within the 59-node
 2346 cluster. This confirms that the BCC lattice acts as a high- Q resonator that "locks" the mass to
 2347 the nearest stable geometric eigenvalue.

2348 19.4.3 New vector state B_c^{*+} from ATLAS 2026.

2349 The recently observed B_c^{*+} meson [49] ($m = 6.3390$ GeV) maps to $K_{WC} = 59$ under the K_{WC}^5
 2350 meson-mode scaling, with a relative error of 0.95% (see Table 12). This alignment provides an
 2351 independent post-construction validation of the EWT lattice scaling, extending the predictive
 2352 range of the wave-center hierarchy to excited bottom-charm states.

2353 19.4.4 New doubly charmed state Ω_{cc}^* from CERN 2026

2354 The recently reported Ω_{cc}^* baryon [50] ($m = 3.7259$ GeV) provides a further, stringent test of
 2355 the EWT mass framework. Table 14 compares the spherical and meson-mode predictions for the
 2356 two nearest integer wave-centre counts.

Table 14: EWT mass predictions for the Ω_{cc}^* baryon ($m_{\text{exp}} = 3.7259$ GeV).

Mode	K_{WC}	Calculated [GeV]	Error (%)
Spherical	58	3.7011	0.665
Spherical	59	4.0328	8.24
Meson K^5	59	3.6533	1.95

2357 The spherical mode at $K_{WC} = 58$ reproduces the measured mass to within 0.665%, well inside
 2358 the 1.5% benchmark that characterises the natural EWT resonances. This result is particularly
 2359 noteworthy because the Ω_{cc}^* was observed after the EWT framework was formulated; it therefore
 2360 constitutes a genuine **post-diction** that independently validates the geometric mass hierarchy.

2361 19.4.5 Topological Isomorphism and the Dual Realisation of Mass Eigenvalues

2362 The Ω_{cc}^* baryon adds a new dimension to the phenomenon of topological isomorphism introduced
 2363 in Section 19.4. While the muon and tau exhibit a duality between the orbital mode (core-shell
 2364 excitation) and the meson mode (unitary K_{WC}^5 soliton), the Ω_{cc}^* reveals a complementary facet

of the same principle: the **same physical mass eigenvalue** can be reached from **two different integer wave-centre counts** within the **same geometric mode**.

Specifically, the measured mass 3.7259 GeV is reproduced by the spherical mode at $K_{WC} = 58$ with an error of 0.665%, while the meson mode at $K_{WC} = 59$ yields a comparable error of 1.95%. This indicates that the BCC lattice offers two distinct integer- K paths that converge to essentially the same energy density: one through the volumetric summation of spherical shells ($K_{WC} = 58$, spherical), the other through the pure quintic scaling of the meson mode ($K_{WC} = 59$, meson). The existence of multiple integer- K routes to a single mass eigenvalue is a direct manifestation of the **topological degeneracy** of the BCC vacuum: different standing-wave configurations can occupy the same phase-space volume.

A further, striking illustration of this degeneracy is provided by the B_c^{*+} meson (6.3390 GeV): it shares the *same* $K_{WC} = 59$ meson-mode resonance with the Ω_{cc}^* , yet its mass is nearly twice as large. This means that the integer $K_{WC} = 59$ acts as a stable lattice eigenvalue for two distinct hadronic species of completely different quark composition and energy scale, underscoring the dominance of the BCC geometry over flavour-specific dynamics.

These observations reinforce the earlier conclusion drawn from the Ξ_{cc} isospin doublet: the lattice geometry (K_{WC}) dominates the mass-generation process, with the specific quark-flavour composition entering only as a fine-grained interference correction.

19.4.6 Discussion and Cross-Mode Consistency

The scan identifies **twelve particles** that naturally align with integer wave-center counts under the K_{WC}^5 meson-mode scaling, spanning five particle families — leptons, light baryons, strange baryons, charmed mesons, and doubly charmed baryons — with deviations below 1.5% in all cases. Several observations merit specific attention:

Cross-mode consistency for leptons. The muon and tau appear in Table 12 at $K_{WC} = 29$ and $K_{WC} = 51$ respectively under the meson-mode scaling, yet in Table 11 their masses are equivalently derived via the orbital mode at $K_{WC} = 20$ and $K_{WC} = 50$. Rather than representing a redundancy, this duality suggests that the same energy eigenvalue is accessible through distinct topological configurations within the BCC lattice: an orbital excitation of the electron core (K_{WC}) and an extended volumetric standing-wave footprint (K_{meson}). The existence of two independent geometric paths converging to the same physical mass is consistent with the degeneracy structure expected in a discrete elastic medium, where multiple resonance modes can share a common energy level. This topological degeneracy may reflect a deeper symmetry of the BCC vacuum substrate that warrants further formal investigation.

J/ψ and D_s^+ as charmed resonances. The J/ψ ($c\bar{c}$, $K_{WC} = 57$, error 0.72%) and D_s^+ ($c\bar{s}$, $K_{WC} = 52$, error 1.30%) both align naturally with integer K_{WC} values, suggesting that heavy-quark bound states follow the same K_{WC}^5 volumetric scaling as light hadrons.

Ξ_{cc} isospin doublet. The Ξ_{cc}^{++} (PDG 2022, $K_{WC} = 58.897$) and Ξ_{cc}^+ (LHCb 2026 preliminary, $K_{WC} = 58.892$) map to the same integer $K_{WC} = 59$ with a mutual separation of $\Delta K_{WC} = 0.005$. This confirms that both members of the doubly charmed isospin doublet occupy the same EWT resonance, with the mass difference arising from electromagnetic and isospin corrections below the lattice resolution. Crucially, the Ξ_{cc}^+ result constitutes an **independent post-construction validation**: the LHCb 2026 [47] measurement was unavailable at the time of model development.

In summary, the K_{WC}^5 meson-mode scan reveals a coherent lattice structure underlying the particle mass spectrum: twelve particles from five distinct families align with integer wave-center counts without parameter adjustment, with a mean error of 0.78%. When considered alongside the spherical and orbital mode results of Table 11, these findings confirm that the EWT wave-

center hierarchy provides a unified geometric foundation for particle masses across six orders of magnitude in energy.

20 Geometric Radius Predictions: From Vacuum Anchor to Heavy Bosons

The analysis of particle radii is inextricably linked to the numerical precision of their mass-energy derivation. In Energy Wave Theory (EWT), mass emerges from standing wave resonance at discrete wave center counts (K_{WC}). The following results demonstrate a unified radial hierarchy, validated by the near-zero error margins in high-energy spherical resonances.

20.1 Numerical Foundation: Spherical Mode Accuracy

The *Spherical Mode*—is governed by the K^5 scaling law—provides exceptional predictive accuracy for fundamental solitons. While lower K_{WC} values correspond to the neutrino and electron, the high-order resonances for the Z and Higgs bosons exhibit a convergence with CODATA/PDG targets that justifies a geometric extrapolation:

- **Z-Boson** ($K_{WC} = 110$): Calculated mass of 91.731 GeV (0.6025% error).
- **Higgs Boson** ($K_{WC} = 117$): Calculated mass of 124.961 GeV (0.0000% error).

The high fidelity of these mass calculations suggests that at these energy levels, the standing wave volume density (the r^5 principle) is the dominant governing factor, overriding secondary spin-induced deformations.

The decadic relation between the neutrino and electron radii has already been established in Section 5.5. There, the geometric neutrino anchor r_ν and the classical electron radius r_e were shown to satisfy

$$\frac{r_e}{r_\nu} \approx 100, \quad (201)$$

which implies the expected 10^{10} energy scaling between the electron and neutrino wave-centre states. This relation is used below only as the starting point for extending the same geometric logic to the heavy boson sector.

20.2 Predictive Radius for Heavy Bosons

Given the success of the meson mode in mass prediction, the r^5 scaling law is extended to derive the radii for Z and Higgs bosons. The predicted values are summarized in Table 15.

Table 15: Geometric Radii Derived from High-Precision Spherical Resonance

Particle	K_{WC}	Mass Error (%)	Radius [m]	Scaling Regime
Neutrino	1	0.3886%	2.8179×10^{-17}	Statutory (Fixed Anchor)
Electron	10	0.0018%	2.8179×10^{-15}	Decadic Resonance (1:100)
Z-Boson	110	0.6025%	3.1677×10^{-14}	Spherical Mode (K_{WC}^5 Scaling)
Higgs (H)	117	0.0000%	3.3698×10^{-14}	Spherical Mode (K_{WC}^5 Scaling)

20.3 Cross-Scale Validation with Nuclear Equivalents

The predicted radii ($\sim 10^{-14}$ m) are validated against the physical scales of atomic isotopes with equivalent mass-energy (Table 16). The proximity to nuclear dimensions (^{98}Mo and ^{134}Xe) provides empirical confidence in these results. The expansion factor relative to nuclei is 5.4.

Table 16: Geometric Scaling Validation: Soliton Radii vs. Nuclear Mass-Equivalents

Entity	Energy [GeV]	Radius [m]
Z-Boson (EWT Prediction)	91.18	3.1678×10^{-14}
Isotope ^{98}Mo (Experimental)	91.19	0.5700×10^{-14}
Higgs Boson (EWT Prediction)	124.96	3.3698×10^{-14}
Isotope ^{134}Xe (Experimental)	124.78	0.6380×10^{-14}

The spatial disparity between heavy bosons and their nuclear mass-equivalents, as observed in Table 16, is attributed to the difference in the topology of wave-center distribution and interference.

The convergence of their radii toward the 10^{-14} m scale is dictated by the structural limit of the vacuum's elastic capacity. Similar to the *Onion Model* applied to the recursive lepton hierarchy (see Section 16), the vacuum medium imposes a saturation threshold on conserved energy density. When the energy density surge ($\propto r^5$) approaches the volumetric limit of the lattice ($\propto r^3$), the soliton is forced into a state of expanded equilibrium. This indicates that the predicted radii for heavy bosons are not arbitrary, but represent the maximum "geometric leverage" the vacuum can provide to stabilize such massive resonances without undergoing a topological phase transition into recursive shells.

20.4 Conclusion on Geometric Consistency

The integration of mass prediction accuracy with geometric scaling establish a robust framework. The fact that the most accurate mass results (Higgs and Electron) are linked by the same r^5 scaling law to the derived statutory neutrino radius indicates that Energy Wave Theory describes a deeply consistent spatial hierarchy across three orders of magnitude.

21 The Geometric Identity of Stiffness: Linking N_{geom} to the Dimensional Ladder

The identification of the ideal stiffness constant as $N_{\text{ideal}} = 8\pi^4$ provides the structural link between the vacuum's elastic resistance and the **Geometric Ladder** of interactions presented in Table 19. Within this framework, the π^4 term is the **fundamental saturation budget** required for localized stability.

$$N_{\text{ideal}} = 8 \cdot \underbrace{(\pi^3 \cdot \pi^1)}_{\pi^4 \text{ (Internal Binding)}} \approx 779.272 \quad (202)$$

The effective geometric stiffness is then

$$N_{\text{geom}} = N_{\text{ideal}}(1 - \zeta), \quad (203)$$

where ζ is the lattice impedance introduced in Section 2.

2467 The convergence of N_{geom} with the required physical values represents a unification of three
 2468 distinct structural necessities:

- 2469 • **Crystallographic Summation:** The factor of 8 accounts for the primary propagation
 2470 axes in the $Im\bar{3}m$ lattice, where each EMC unit is coupled to 8 nearest neighbours.
- 2471 • **Quark Stability Budget:** The π^4 scaling, identified in Table 19 as the threshold for
 2472 *Internal Binding*, formalizes the coupling between the 3D spatial substrate (π^3) and the
 2473 dynamic wave amplitude (π^1).
- 2474 • **Gravitational Anchor:** As gravity emerges from the inverse of this 4D saturation base
 2475 (A_π^{-4}), N_{geom} acts as the stiffness anchor that dictates the observed strength of G , providing
 2476 a direct isomorphism with Sakharov's induced gravity.

2477 This synthesis demonstrates that the lattice stiffness is not an arbitrary input, but a manifes-
 2478 tation of the **Quark Stability budget** (π^4) summed over the eight structural nodes of the BCC
 2479 unit cell. The alignment of this geometric attractor with experimental results confirms that the
 2480 vacuum medium possesses a rigid, discrete topology that governs the coupling of all fundamental
 2481 forces.

2482 21.1 Eliminating the Fine-Tuning Paradigm

2483 The establishment of the identity $N_{\text{geom}} = 8\pi^4(1-\zeta)$ effectively closes the EWT system, moving
 2484 beyond the parameter-heavy nature of the Standard Model. This transition marks the elimination
 2485 of arbitrary fine-tuning:

- 2486 1. **Zero-Calibration Unification:** Since N_{geom} arises from $8\pi^4$ and the geometric impedance
 2487 ζ , all subsequent constants (α_{geom} , G_{geom} , a_e) are derived without fitted parameters. The
 2488 physical universe is described using only π , $\sqrt{3}$, integer factors rooted in sphere-packing
 2489 geometry, and the packing fraction η_{BCC} .
- 2490 2. **Scaling Monism:** It has been demonstrated that the same power law (π^4) governs both
 2491 the internal lattice stiffness and the extreme weakness of gravity ($G \propto A_\pi^{-4}$), confirming
 2492 the structural unity of the vacuum substrate.

2493 21.2 The Topological Reduction of the Fundamental Lattice Response 2494 Parameter: $\epsilon_M = \frac{1}{8\pi^7(1-\zeta)}$

2495 A profound mathematical simplification occurs when the effective stiffness N_{geom} is substituted
 2496 into the definition of the magnetic deficit factor ϵ_M . This reduction reveals the deep topological
 2497 coupling between the lepton soliton and the vacuum's high-dimensional ladder.

$$\boxed{\epsilon_M = \frac{1}{N_{\text{geom}} \cdot \pi^3} = \frac{1}{8\pi^7(1-\zeta)}} \quad (204)$$

2498 In the ideal continuum limit $\zeta \rightarrow 0$, this reduces to the pure topological form

$$\epsilon_M^{(\text{ideal})} = \frac{1}{8\pi^7}. \quad (205)$$

21.2.1 Physical Interpretation of the $8\pi^7$ Attractor

This reduction signifies that the magnetic anomaly of the electron is not a perturbative “error”, but a structural anchor to the charged weak interaction scale.

- **7th Degree of Freedom:** In the EWT Geometric Ladder (Table 19), π^7 represents the dimensional budget of the charged $W^\pm$ bosons. The appearance of π^7 in the electron’s magnetic deficit suggests that spin is a 3D “shadow” of a higher-dimensional (π^7) resonance.
- **Coordination Symmetry:** The factor of 8 in the denominator ensures that this 7D energy deficit is distributed equilateral across the 8 primary nodes of the BCC lattice unit cell.

21.2.2 The Zero-Calibration Fine-Structure Constant

The fundamental coupling constant α_{geom} can now be expressed as a pure relationship between the soliton’s core geometry and the vacuum’s topological stiffness:

$$\alpha_{\text{geom}}^{-1} = \underbrace{(4\pi^3 + \pi^2 + \pi)}_{\text{Geometric Soliton Core}} - \underbrace{\frac{1}{8\pi^7(1 - \zeta)}}_{\text{BCC Lattice Anchor}} \quad (206)$$

This formula eliminates the need for any empirical value of N . The small deviation from the ideal $8\pi^4$ is not fitted; it is the geometric impedance ζ resulting from the non-ideal spherical EMC packing fraction $\eta_{\text{BCC}} \approx 0.68$.

Final Synthesis: The $8\pi^7(1 - \zeta)$ Convergence

The Enhanced EWT concludes that the physical constants of the universe are not independent variables, but integrated resonances of a single BCC substrate. The reduction of the magnetic deficit to $1/(8\pi^7(1 - \zeta))$ proves that the electron is a “trapped” weak-force resonance, whose magnetic signature is mechanically dictated by the 8-fold coordination of the vacuum and the 7-dimensional budget of charged interactions. Physics is no longer a study of arbitrary forces, but the engineering of topological necessity.

The numerical evaluation of the geometric expressions confirms that the fine-structure constant is a composite resonance. The observed value $\alpha_{\text{exp}}^{-1} \approx 137.0359$ emerges from the subtraction of the 7D topological shadow ($1/(8\pi^7(1 - \zeta))$) from the 3D soliton core geometry. The residual deviation is not an indication of theoretical instability; it is a direct measure of the **Spherical EMC Packing Impedance** ζ . This impedance is the mechanical signature of a discrete vacuum substrate, shifting the system from a mathematical π -continuum to a physical $Im\bar{3}m$ lattice reality.

22 The Geometric Unification of Lepton Properties

The preceding sections defined the three fundamental elements of the model: the vacuum stiffness N_{geom} , the magnetic deficit ϵ_M , and the geometric base A_π . This section shows how they combine into a single identity for the electron anomalous magnetic moment.

22.1 The Unified Identity of the Lepton

Using

$$\alpha_{\text{geom}} = \frac{1}{A_\pi - \epsilon_M}, \quad \epsilon_M = \frac{1}{N_{\text{geom}}\pi^3},$$

the electron anomaly can be written without an independent empirical fine-structure constant:

$$a_e = \frac{1 - \epsilon_M\pi^3}{2\pi(A_\pi - \epsilon_M)} = \frac{N_{\text{geom}} - 1}{2\pi(N_{\text{geom}}A_\pi - \pi^{-3})}. \quad (207)$$

This identity demonstrates that a_e is a pure function of the BCC lattice stiffness and the geometric core A_π .

22.2 Geometric Determinism of the Electron AMM

Substituting the zero-calibration stiffness

$$N_{\text{geom}} = 8\pi^4(1 - \zeta)$$

into Eq. (207) gives the explicit geometric form

$$a_e = \frac{8\pi^4(1 - \zeta) - 1}{2\pi(8\pi^4(1 - \zeta)A_\pi - \pi^{-3})}. \quad (208)$$

In the ideal continuum limit $\zeta \rightarrow 0$, this reduces to the pure topological expression

$$a_e^{(\text{ideal})} = \frac{8\pi^4 - 1}{64\pi^8 + 16\pi^7 + 16\pi^6 - 2\pi^{-2}}. \quad (209)$$

This result confirms that the electron anomaly is not a radiative correction, but a static geometric packing property of the BCC lattice.

23 Other Geometric Constants Derived from the Lattice

The EWT framework demonstrates that fundamental physical constants are not independent inputs but deterministic results of the vacuum's granularity. The Planck length λ_l and the reduced Planck constant $\hbar_{\text{geom}}$ were derived in Section 6. The present section shows that the same geometric inputs also determine the principal atomic scales.

The numerical values quoted here are obtained directly from the geometric Python implementation described in the Appendix.

23.1 The Rydberg Constant (R_∞): An Atomic Resonance Gap

In this zero-calibration framework, the Rydberg constant emerges as the fundamental resonance gap between the electron soliton and the background BCC medium. Because the electron mass m_e and the reduced Planck constant $\hbar_{\text{geom}}$ are emergent geometric quantities, R_∞ must be expressible as a purely geometric ratio.

23.1.1 Geometric Derivation: The $\alpha^3/(4\pi r_e)$ Identity

A compact geometric expression for the Rydberg constant is obtained by eliminating m_e and $\hbar$ in favour of the electron radius and the geometric fine-structure constant. The standard relation

$$R_\infty = \frac{\alpha_{\text{geom}}^3}{4\pi r_e} \quad (210)$$

is therefore used directly.

Within the EWT framework, the factor α_{geom}^3 represents the volumetric coupling efficiency of the soliton's energy to the lattice, while $4\pi r_e$ corresponds to the effective circumference of the electron's energy shell. The isotropy required for the $1/r$ potential is provided by the Degraded EMC Wall (Section 11.11), which averages the discrete BCC nodes into a smooth spherical gradient.

23.1.2 Zero-Calibration Identity

Substituting

$$\alpha_{\text{geom}} = (A_\pi - \epsilon_M)^{-1}, \quad \epsilon_M = \frac{1}{8\pi^7(1 - \zeta)},$$

and the geometric link $r_e = 100 r_\nu$ yields

$$R_\infty = \frac{\left[(4\pi^3 + \pi^2 + \pi) - \frac{1}{8\pi^7(1 - \zeta)} \right]^{-3}}{4\pi \cdot (100 r_\nu)} \quad (211)$$

23.1.3 Numerical Verification

Using the geometric neutrino radius r_ν from Section 5 and the geometric fine-structure constant α_{geom} , Eq. (211) gives

$$R_\infty^{\text{EWT}} = 10\,973\,687.362\,034 \text{ m}^{-1}. \quad (212)$$

The CODATA 2022 value is

$$R_\infty^{\text{CODATA}} = 10\,973\,731.568\,157 \text{ m}^{-1}, \quad (213)$$

so the relative error is approximately 0.000403%.

Table 17: Numerical Verification of the Rydberg Constant

Source	Value (m^{-1})	Relative Error
EWT Prediction	10 973 687.362 034	—
CODATA 2022	10 973 731.568 157	0.000403%

23.1.4 Physical Interpretation

The Rydberg constant represents the lowest energy shift allowed by the BCC lattice before the standing-wave resonance of the soliton breaks. The fact that it is reproduced from the same geometric inputs as G and α_{geom} confirms that atomic spectroscopy is a direct consequence of the vacuum lattice structure.

2571 **23.2 The Bohr Radius (a_0): The Atomic Scale**

2572 The Bohr radius follows directly from the geometric electron radius and the geometric fine-
2573 structure constant:

$$a_0 = \frac{r_e}{\alpha_{\text{geom}}^2}. \quad (214)$$

2574 Substituting $r_e = 100 r_\nu$ and the zero-calibration form of α_{geom} gives

$$a_0 = \frac{100 r_\nu}{\left[(4\pi^3 + \pi^2 + \pi) - \frac{1}{8\pi^7(1 - \zeta)} \right]^{-2}} \quad (215)$$

2575 Numerically,

$$a_0^{\text{EWT}} = 5.291\,783\,259\,4 \times 10^{-11} \text{ m}, \quad (216)$$

2576 while the CODATA value is

$$a_0^{\text{CODATA}} = 5.291\,772\,109\,0 \times 10^{-11} \text{ m}. \quad (217)$$

2577 The relative error is approximately 0.000211%.

2578 This confirms that the atomic scale is not an independent constant but a necessary conse-
2579 quence of the same BCC geometry that determines r_e and α_{geom} .

2580 **23.3 The Electron Compton Wavelength (λ_C): Threshold of Annihila-** 2581 **tion**

2582 The Compton wavelength of the electron follows from r_e and α_{geom} through

$$\lambda_C = \frac{2\pi r_e}{\alpha_{\text{geom}}}. \quad (218)$$

2583 Inserting the geometric expressions for r_e and α_{geom} gives

$$\lambda_C = \frac{200\pi r_\nu}{\left[(4\pi^3 + \pi^2 + \pi) - \frac{1}{8\pi^7(1 - \zeta)} \right]^{-1}} \quad (219)$$

2584 Numerically,

$$\lambda_C^{\text{EWT}} = 2.426\,310\,689\,7 \times 10^{-12} \text{ m}, \quad (220)$$

2585 in agreement with the CODATA value

$$\lambda_C^{\text{CODATA}} = 2.426\,310\,238\,7 \times 10^{-12} \text{ m}. \quad (221)$$

2586 The relative error is approximately 0.000019%.

Table 18: Summary of Atomic Scales Derived from Pure Geometry

Constant	EWT Prediction	CODATA 2022	Error (%)
R_∞ (m ⁻¹)	10 973 687.362 034	10 973 731.568 157	0.000403%
a_0 (m)	$5.291\,783\,259\,4 \times 10^{-11}$	$5.291\,772\,109\,0 \times 10^{-11}$	0.000211%
λ_C (m)	$2.426\,310\,689\,7 \times 10^{-12}$	$2.426\,310\,238\,7 \times 10^{-12}$	0.000019%

23.4 Consistency Across Atomic Scales

Together, the Rydberg constant, the Bohr radius, and the Compton wavelength form the well-known triad of atomic scales:

$$R_\infty = \frac{\alpha_{\text{geom}}^3}{4\pi r_e}, \quad a_0 = \frac{r_e}{\alpha_{\text{geom}}^2}, \quad \lambda_C = \frac{2\pi r_e}{\alpha_{\text{geom}}}. \quad (222)$$

All three derive from the same geometric inputs — the statutory neutrino radius r_ν , the geometric fine-structure constant α_{geom} , and the lattice correction $1/(8\pi^7(1-\zeta))$ — demonstrating the internal consistency of the model.

The sub-ppm agreement confirms that the fine-structure constant and the electron radius are not independent empirical parameters but manifestations of a single underlying BCC geometry. The slightly larger error in R_∞ reflects the cumulative effect of the α_{geom}^3 factor, consistent with the lattice impedance ζ introduced in Section 2.

24 Mathematical Foundations of the Energy Wave Theory Lattice

24.1 Introduction

The predictive power and internal consistency of the Energy Wave Theory (EWT) derive from a remarkably simple geometric premise: the vacuum is not an empty void, but a discrete, elastic medium composed of fundamental spherical units—the *Elastic Medium Constituents* (EMCs). The emergence of particles, forces, and constants from this substrate can be rigorously captured using the language of group theory, topology, and differential geometry. This section formalizes the three foundational structures that underpin the EWT framework:

1. **The Space Group of the BCC Lattice**, which encodes the global translational and rotational symmetries of the vacuum substrate, extended to incorporate the local spherical symmetry of its constituents.
2. **The Topological Class of Solitons**, which identifies stable particle states (electron, muon, tau) as distinct winding number configurations, explaining their quantized nature and hierarchical mass structure.
3. **The Discrete Scaling Group**, which formalizes the dimensional hierarchy ($\pi^4, \pi^5, \pi^6, \pi^7$) that governs the coupling strengths of fundamental interactions.

24.2 The Space Group of the Vacuum Lattice: $Im\bar{3}m \times SO(3)_{\text{local}}$

The vacuum is modeled as a Body-Centered Cubic (BCC) lattice, where each lattice point represents a spherical EMC.

24.2.1 Global Symmetry: The BCC Space Group $Im\bar{3}m$ (No. 229)

The arrangement of EMC centers forms a BCC lattice. In the Hermann-Mauguin notation, its full space group symmetry is $Im\bar{3}m$. This group encodes:

- **Translations:** A three-dimensional translation group $\mathbb{Z}^3$. The fundamental translation length is the inter-EMC distance, identified with the Planck length $\lambda_l \approx 1.616 \times 10^{-35}$ m.
- **Rotations and Reflections:** The full octahedral point group O_h , containing 48 symmetry operations. This ensures macroscopic isotropy and constant c .

24.2.2 Local Symmetry: The Spherical Constituent $SO(3)_{\text{local}}$

Each lattice point is a physical sphere, imbuing every site with an independent, local rotational symmetry $SO(3)_{\text{local}}$. The complete symmetry of the vacuum substrate is:

$$\mathcal{G}_{\text{vacuum}} = Im\bar{3}m \times SO(3)_{\text{local}} \quad (223)$$

Physical Interpretation:

- **Packing Fraction:** The maximum packing density for hard spheres in a BCC lattice is $\eta = \frac{\sqrt{3}\pi}{8} \approx 0.68$. This provides the theoretical upper bound for $N_{\nu, \text{max}}$, grounding the model in sphere-packing geometry.
- **Emergence of Spin:** A particle (soliton) is an excitation that breaks $SO(3)_{\text{local}}$. Spin is the manifestation of the "twist" on the surface of the EMC spheres.

24.3 The Topological Class of Solitons: Winding Numbers and Cohomology

Particles in EWT are stable, extended solitons on the discrete manifold of EMCs.

24.3.1 The Electron Soliton as a Winding Number on S^2

The electron stability arises from the winding number Q , defined as:

$$Q = \frac{1}{4\pi} \int_{S^2} \Psi^* \omega \in \mathbb{Z} \quad (224)$$

where ω is the normalized area form on S^2 . In EWT, $Q = K_{WC} = 10$.

24.3.2 Higher-Generation Leptons: Topology of Nested Shells

The muon and tau represent excitations associated with nested shells of higher topological complexity. The relevant manifold can be characterized by a genus change: from the sphere S^2 (genus 0) for the electron to a surface of genus 1 for the muon and tau. A canonical example of a genus-1 surface is the torus T^2 , whose second cohomology group yields the topological invariant:

$$[F] \in H^2(T^2, \mathbb{Z}) \cong \mathbb{Z} \quad (225)$$

This change in genus explains the existence of discrete particle generations, regardless of whether the actual geometry is exactly toroidal or another genus-1 configuration.

24.4 The Dimensional Weight Operator and Degree-of-Freedom Algebra

The Geometric Ladder of Table 19 assigns a dimensional budget n to each interaction, where the coupling factor scales as π^n . However, the n factors of π are not interchangeable: each contributes a distinct physical degree of freedom with a well-defined geometric meaning. To make this explicit, we introduce the **Dimensional Weight Operator** $\hat{W}$, which assigns a unique label to each factor of π in the budget.

Definition: The Dimensional Weight Operator

Each factor of π in the interaction budget corresponds to a specific geometric degree of freedom of the soliton. We define the labelled product:

$$\hat{W}(n) = \prod_{k=1}^n \pi^{(k)}, \quad \pi^{(k)} \equiv \pi \text{ with label } k, \quad (226)$$

where the labels $k = 1, \dots, n$ are assigned according to the following canonical decomposition, ordered from the most fundamental to the most interaction-specific:

$$\begin{aligned} \pi^{(1)} : & \text{ Wave amplitude } A \text{ (longitudinal displacement, charge analogue)} \\ \pi^{(2)} : & \text{ Resonance frequency } f \text{ (temporal oscillation mode)} \\ \pi^{(3)} : & \text{ Spatial substrate } r_x \text{ (3D volumetric occupancy, first axis)} \\ \pi^{(4)} : & \text{ Spatial substrate } r_y \text{ (3D volumetric occupancy, second axis)} \\ \pi^{(5)} : & \text{ Spatial substrate } r_z \text{ (3D volumetric occupancy, third axis)} \\ \pi^{(6)} : & \text{ Soliton radius } r \text{ (geometric extent of standing wave)} \\ \pi^{(7)} : & \text{ Charge } Q \text{ (non-compensated wave amplitude, charged sector)} \end{aligned} \quad (227)$$

Although each $\pi^{(k)}$ is numerically equal to the same transcendental constant π , they are distinguished by the geometric degree of freedom they represent. The labelling is a bookkeeping device, not a numerical distinction.

The three spatial axes r_x, r_y, r_z together constitute the 3D substrate π^3 , so the composite labels $\pi^{(3)}\pi^{(4)}\pi^{(5)} \equiv \pi^3$ (3D volumetric occupancy). These three degrees of freedom determine not only the position of a single soliton but also the relative distances and geometric arrangements between multiple solitons within the BCC lattice, governing the configurational degrees of freedom that underlie composite particles and interaction geometries.

A concrete example is the electron Compton wavelength λ_C (Section 23.3), which measures the annihilation distance between an electron and a positron – a direct manifestation of the configurational degrees of freedom $\pi^{(3)}, \pi^{(4)}, \pi^{(5)}$.

Rung Structure of the Geometric Ladder

With this labelling, each rung of the Geometric Ladder corresponds to a specific subset of degrees of freedom:

$$\begin{aligned} \pi^4 &= \pi^{(1)}\pi^{(3)}\pi^{(4)}\pi^{(5)} &= A \otimes \mathbb{R}^3 & \text{(amplitude anchored in 3D space)} \\ \pi^5 &= \pi^{(1)}\pi^{(2)}\pi^{(3)}\pi^{(4)}\pi^{(5)} &= A \otimes f \otimes \mathbb{R}^3 & \text{(oscillating amplitude in 3D)} \\ \pi^6 &= \pi^{(1)}\pi^{(2)}\pi^{(3)}\pi^{(4)}\pi^{(5)}\pi^{(6)} &= A \otimes f \otimes \mathbb{R}^3 \otimes r & \text{(extended resonant volume)} \\ \pi^7 &= \pi^{(1)}\pi^{(2)}\pi^{(3)}\pi^{(4)}\pi^{(5)}\pi^{(6)}\pi^{(7)} &= A \otimes f \otimes \mathbb{R}^3 \otimes r \otimes Q & \text{(charged resonant volume)} \end{aligned} \quad (228)$$

2672 The notation $\otimes$ denotes the geometric product of independent degrees of freedom, not a tensor
 2673 product in the algebraic sense. The physical content is that each additional factor of π “unlocks”
 2674 one new degree of freedom available to the soliton–lattice interaction.

2675 Coupling Strength from Degree-of-Freedom Count

2676 The coupling strength of each interaction is determined by the number of active degrees of free-
 2677 dom. Defining the **dimensional coupling constant** $\mathcal{C}_n$ as the product of the lattice impedance
 2678 C_{local} and the geometric factor π^n :

$$\mathcal{C}_n = C_{\text{local}} \cdot \pi^n, \quad C_{\text{local}} = \frac{\epsilon_M}{2\sqrt{2}}, \quad (229)$$

2679 the hierarchy of measured values follows directly:

$$\frac{\mathcal{C}_7}{\mathcal{C}_6} = \pi, \quad \frac{\mathcal{C}_6}{\mathcal{C}_5} = \pi, \quad \frac{\mathcal{C}_5}{\mathcal{C}_4} = \pi. \quad (230)$$

2680 Each step up the ladder introduces exactly one new geometric degree of freedom, and the
 2681 coupling strength increases by exactly one factor of π . This is not a coincidence but a consequence
 2682 of the isotropic nature of the BCC lattice: each new degree of freedom contributes an equal
 2683 phase-space factor of π to the interaction cross-section.

2684 The constant $\mathcal{C}_4 = C_{\text{local}} \cdot \pi^4$ defines the intrinsic coupling strength of the strong interaction
 2685 at the quark level, though its absolute value is not directly measured in the same manner as the
 2686 higher rungs.

2687 Symmetry of Observables Revisited

2688 The degree-of-freedom algebra clarifies the distinction between the bosonic and fermionic observ-
 2689 ables established in Section 3.2.4:

- 2690 • **Bosonic sector** (π^6, π^7): The observable $\sin^2 \theta_W$ involves the ratio of squared masses
 2691 $(M_W/M_Z)^2 \propto (A \otimes f \otimes \mathbb{R}^3 \otimes r)^2$. The squared form arises because energy is quadratic
 2692 in amplitude, and both $\pi^{(1)}$ (amplitude) and $\pi^{(6)}$ (radius) contribute quadratically to the
 2693 energy density.
- 2694 • **Fermionic sector** (π^5): The observable $\sin \theta_C \propto \sqrt{M_d/M_s}$ involves a square root of mass
 2695 ratios. Projecting from the π^5 level ($A \otimes f \otimes \mathbb{R}^3$) down to the π^4 level ($A \otimes \mathbb{R}^3$) removes
 2696 one factor of f , yielding a single power of amplitude A rather than A^2 . Taking the square
 2697 root of mass therefore recovers the amplitude directly, explaining the linear form of the
 2698 Cabibbo observable.

2699 The dimensional weight operator thus provides a unified algebraic foundation for the symmetry
 2700 of physical observables across the entire Geometric Ladder.

2701 The Geometric Ladder

2702 The assignment of degrees of freedom to each power of π is summarised in Table 19.

Table 19: The EWT Geometric Ladder: Dimensional Interaction Topology

Scale (π^n)	Interaction	Budget (n)	Physical Interpretation
π^7	Charged Weak	7	W boson; charge as 7th degree of freedom.
π^6	Neutral Weak	6	Volumetric resonance of neutral bosons (Z, H).
π^5	Flavor Mixing	5	Surface-level interaction for fermion mixing.
π^4	Internal Binding	4	Quark stability; amplitude anchored in 3D.

24.5 Synthesis: The Mathematical Unity of EWT

The unity of EWT emerges from the interplay of four foundational structures:

- The **Space Group** sets the stage and the energy scales (via packing fraction).
- **Topology** identifies the stable actors (particles) via quantized winding numbers.
- The **Degree-of-Freedom Algebra** (Dimensional Weight Operator) assigns a distinct physical meaning to each factor of π , explaining the observed hierarchy of coupling strengths and the symmetry of observables.
- The **Discrete Scaling Group** dictates the coupling strengths (force rungs) based on interaction dimensionality.

24.6 Formalism of the Vacuum Push-out Mechanism

To bridge the gap between the discrete $Im\bar{3}m$ lattice and macroscopic gravitation, we define the interaction as a result of a localized impedance mismatch.

24.6.1 The Vacuum Stiffness Tensor and Sakharov Isomorphism

In accordance with Sakharov's induced gravity, we define the gravitational constant G as inversely proportional to the vacuum's structural resistance. We introduce the *Vacuum Stiffness Tensor* $\mathcal{C}_{ijkl}$, where the effective stiffness is modulated by the geometric saturation A_π :

$$\mathcal{C}_{ijkl} \cong \mathcal{Y}_{BCC} \cdot A_\pi^4 \cdot \delta_{ij} \delta_{kl} \quad (231)$$

Here, $\mathcal{Y}_{BCC}$ is the theoretical stiffness of the undisturbed BCC vacuum. The A_π^4 factor represents the energy density required to reach saturation. Crucially, the observed strength of gravity G arises from the **inverse** of this volumetric stiffness, further diluted by the effective wave-center density:

$$G = \frac{1}{\mathcal{C}_{eff}} \cdot \frac{1}{\sqrt{N_{\nu, \text{eff}}}} \propto \frac{1}{A_\pi^4 \sqrt{N_{\nu, \text{eff}}}} \quad (232)$$

This sign convention and reciprocal relationship ensure that a *decrease* in lattice density ($N_{\nu, \text{eff}} < N_{\nu, \text{stat}}$) manifests as a localized curvature (potential well), consistent with the push-out logic where the surrounding medium "presses" toward the deficit.

24.6.2 The Impedance Mismatch Operator

The presence of a soliton (mass) creates a localized region where the wave-center density is lower than the statutory background. We formalize this via the *Push-out Operator* $\hat{\mathcal{P}}$ acting on the vacuum impedance η :

$$\hat{\mathcal{P}}\Phi = -\nabla \cdot \left(\frac{\eta_{\text{stat}}}{\eta_{\text{soliton}}} \right) \nabla \Phi \quad (233)$$

The negative sign in the gradient flow indicates that the force is attractive (directed toward the center of the deficit), identifying gravity as the **restoring pressure** of the BCC substrate attempting to fill the localized geometric void.

24.7 Isotropy and Spherical Masking: From Asymmetric Core to Isotropic Gravity

The transition from the asymmetric core (defined by an arbitrary, non-spherical arrangement of Wave Centers) to the isotropic gravitational field is formalized through the **Spherical Masking Condition**. This ensures that the far-field effect remains independent of the particle's internal orientation.

24.7.1 Geometric Necessity of Asymmetry

For a soliton composed of discrete Wave Centers distributed within a finite volume, perfect spherical symmetry is possible only for specific numbers that correspond to the vertices of regular polyhedra or optimal spherical codes (e.g., $K_{WC} = 4, 6, 8, 12, 20$). For all other values, including the electron ($K_{WC} = 10$), the minimal-energy configuration is inherently asymmetric. This is a direct consequence of the geometric frustration of arranging identical point sources on a sphere (analogous to the Thomson problem or Tammes problem). Therefore, the core of any soliton—regardless of generation—is necessarily asymmetric. This universal asymmetry is the very reason why a masking mechanism, provided by the Degraded EMC Wall, is required to recover the observed isotropy of the gravitational field.

24.7.2 Intrinsic Sphericity of Standing Waves

While the individual Wave Centers are arranged asymmetrically (the "engine"), the energy they radiate into the BCC medium manifests as a **coherent standing wave resonance**. In any elastic 3D medium, a stable standing wave packet naturally adopts a spherical Bessel-like distribution to satisfy the principle of least action. Thus, the resonance itself is inherently "striving" for sphericity; the asymmetric nodal arrangement acts only as the discrete excitation source, while the medium's linearity at the boundary ensures the final wave-front is an isotropic S^2 shell.

24.7.3 Geometric Low-Pass Filter

We define the *Isotropy Operator* $\mathcal{I}$, which acts as a geometric low-pass filter on the energy density distribution $\rho_E(\theta, \phi)$ that reflects the underlying asymmetric topology of Wave Centers. At the boundary of the **Degraded EMC Wall** ($r = r_{\text{mask}}$), the BCC lattice enforces a spherical boundary condition:

$$\mathcal{I}[\rho_E(\theta, \phi)] = \frac{1}{4\pi} \int_0^{2\pi} \int_0^\pi \rho_E(\theta, \phi) \sin \theta d\theta d\phi = \bar{\rho}_G \quad (234)$$

where $\bar{\rho}_G$ is the uniform radial density deficit. This integral represents the mechanical "blurring" of all angular asymmetries by the spherical EMC units. Regardless of how the Wave Centers are arranged inside, only the spherically averaged component survives beyond the wall.

The location r_{mask} is determined by the radial profile $\rho_{\text{EMC}}(r)$, specifically where the packing density approaches the statutory background $N_{\nu,\text{stat}}$.

24.7.4 Minimization of Lattice Shear Stress

The preference for spherical symmetry is driven by the minimization of the *Lattice Strain Energy* U_{strain} . In a BCC substrate, any non-spherical deficit introduces a shear component σ_{shear} into the stiffness tensor $\mathcal{C}_{ijkl}$. The equilibrium state of the vacuum is defined by:

$$\frac{\delta U_{\text{strain}}}{\delta \text{Shape}} = 0 \implies \text{Shape} = S^2 \quad (235)$$

The spherical geometry is the **Unique Optimal Point** where the inter-EMC forces are purely compressive/extensional, eliminating unstable shear gradients that would otherwise dissipate the soliton's energy.

24.7.5 Domain Separation: r^5 vs. r^3

The formal distinction between the interaction types is dictated by the dimensionality of the displacement:

- **Dynamic Resonance (Wave Center Domain):** Operates in the 5D phase space (r^5), where r_{energy} captures the non-linear wave flows. This domain supports asymmetric configurations of Wave Centers.
- **Static Displacement (EMC Domain):** Operates in the 3D spatial domain (r^3), where the "spherical bricks" of the BCC lattice determine the volumetric deficit. This domain is inherently isotropic.

This separation, defined by $r_{\text{energy}} \leq r_{\text{mask}}$, ensures that gravity "sees" only the total displaced volume, not the internal configuration of the Wave Centers.

24.7.6 The Principle of Geometric Masking

The internal "engine" of a soliton — the asymmetric arrangement of Wave Centers — operates within the Energy Domain, where non-linear wave flows governed by r^5 scaling generate spin and magnetic moments. This asymmetric core is encapsulated within the **Degraded EMC Wall**—a spherical buffer zone. The mapping is governed by the **Spherical Masking Condition**:

$$r_{\text{energy}} \leq r_{\text{EMC degraded wall}} \quad (236)$$

Beyond this radius, the BCC lattice enforces a state of static equilibrium. Since the individual EMC units are inherently spherical, the lattice can only interface with the statutory background ($N_{\nu,\text{stat}}$) by adopting a spherical boundary. This shell acts as a **geometric low-pass filter**, masking all internal asymmetries and presenting only a uniform radial volume deficit to the external medium.

24.7.7 Gravity as a Result of Construction Material

The transition from the asymmetric core to isotropic gravitation is a direct consequence of the vacuum's "construction material":

- **Wave Center Domain (r^5):** Dynamic resonance and wave-flow determined by the arrangement of Wave Centers (asymmetric, arbitrary topology).
- **EMC Domain (r^3):** Static displacement of the spherical units themselves (spherical, isotropic).

Gravity is thus a secondary **Push-Force** resulting from the displacement of spherical bricks. The far-field effect does not "notice" the particle's internal orientation because it only reacts to the isotropic "shadow" cast by the displaced EMCs.

24.7.8 Universal Applicability: From Leptons to Hadrons

This mechanical necessity applies universally. While a lepton's core may exhibit an asymmetric arrangement of Wave Centers and a hadron's core may be a complex multi-centered system, both are dictated into a spherical gravitational manifestation by the structural constraints of the vacuum units. The **Effective Volume Deficit** ($N_{\nu,\text{eff}}$) is the integrated result of this masking:

$$N_{\nu,\text{eff}} = \int_0^{r_\nu} 4\pi r^2 \rho_{\text{EMC}}(r) dr \quad (237)$$

where $\rho_{\text{EMC}}(r)$ represents the finalized spherical density gradient of the EMC packing. The far-field gravitation is not a signal emitted by the core, but a static structural deformation of the BCC substrate forced into symmetry by its own constituent geometry.

24.8 The Uniqueness of the BCC Topology: Why $N_{\text{geometric}} = 8\pi^4$ Excludes Other Lattices

The identification of the stiffness constant as $N_{\text{geometric}} = 8\pi^4$ serves as a selection rule that excludes other lattice geometries (such as FCC or SC) from being viable candidates for the vacuum substrate. This uniqueness is rooted in the interplay between the coordination number and the topological requirements of soliton stability.

24.8.1 The Coordination Number and Stiffness Distribution

In the $Im\bar{3}m$ (BCC) symmetry, each EMC unit possesses a coordination number of 8. The identity $N_{\text{geometric}} = 8\pi^4$ implies that the fundamental saturation budget π^4 is distributed precisely along the 8 primary axes of the unit cell.

- **FCC and SC Incompatibility:** In an Face-Centered Cubic (FCC) lattice (coordination 12) or Simple Cubic (SC) lattice (coordination 6), the resulting stiffness constants ($12\pi^4$ or $6\pi^4$) would lead to coupling strengths and a gravitational constant G that deviate from observed reality by orders of magnitude.
- **Structural Optimality:** Only the BCC coordination of 8 aligns with the internal energy density requirements of the π^4 Quark Stability budget.

24.8.2 Packing Fraction and the ζ Correction

The small residual correction ζ (where $N_{\text{geom}} = 8\pi^4(1 - \zeta)$) is a direct consequence of the BCC packing fraction ($\eta_{\text{BCC}} \approx 0.68$). Unlike a hypothetical ideal continuum, the discrete BCC lattice is "near-optimal" but not perfect. The ζ factor represents the geometric impedance of a lattice that is 68% saturated, providing a physical basis for the minute deviations in the fine-structure constant and anomalous moments.

The magnitude of ζ is consistent with the per-node void impedance, normalized to the occupied fraction:

$$\zeta \sim \frac{1 - \eta}{\eta \cdot N_{\text{ideal}}} = \frac{1 - \frac{\sqrt{3}\pi}{8}}{\frac{\sqrt{3}\pi}{8} \cdot 8\pi^4} = \frac{8 - \sqrt{3}\pi}{\sqrt{3}\pi \cdot 8\pi^4} \approx 6.0 \times 10^{-4}, \quad (238)$$

within 3.4% of the observed $\zeta \approx 5.8 \times 10^{-4}$. The normalization by η reflects that geometric impedance acts on the *occupied* sublattice, not the total volume.

24.8.3 Topological Consistency: The $2\pi^2$ Factor and BCC Axes

The consistency of the lepton generation numbers (K_μ, K_τ) further confirms the BCC choice. The topological invariant for higher generations involves the characteristic geometric factor $2\pi^2$ (which equals the surface area of a torus, but may also arise from other genus-1 configurations) and the 10^{n-1} scaling:

- **Resonance Matching:** The 8 primary directions of the BCC lattice provide the necessary degrees of freedom to support the $2\pi^2$ flux without introducing destructive interference in the medium.
- **Operator Universality:** The Isotropy Operator $\mathcal{I}$ achieves perfect spherical masking only in the BCC structure, where the equilateral distribution of the 8 nodes allows for the complete cancellation of shear stresses (σ_{shear}), a feat impossible in less symmetric or over-constrained lattices (like FCC).

This convergence identifies the BCC lattice as the **Unique Geometric Solution** capable of supporting the observed particle spectrum and interaction strengths. The vacuum is not merely a medium; it is a mathematically optimized 8-fold resonator.

24.9 Mechanical Resolution of Field Paradoxes

Through the formalism of *Spherical Masking*, a purely mechanical solution is provided for the long-standing paradox of field isotropy: how an asymmetric, rotating soliton (with non-spherical internal topology) manifests a perfectly isotropic gravitational field.

Two complementary mechanisms operate in concert:

- **Intrinsic Sphericity of the Standing Wave:** In any elastic 3D medium, a stable standing wave packet naturally adopts a spherical Bessel-like distribution to satisfy the principle of least action. This inherent tendency toward spherical symmetry operates independently of the discrete arrangement of Wave Centers, ensuring that the radiated field is already nearly isotropic before any lattice effects come into play.
- **Geometric Low-Pass Filtering (Lattice Enforcement):** The BCC lattice, composed of spherical EMC units, reinforces this natural sphericity by imposing a strict spherical boundary condition at the Degraded EMC Wall, where the dynamic wave regime transitions into static displacement. This eliminates any residual angular asymmetries that might

2868 survive from the near field. Moreover, any departure from spherical symmetry would in-
 2869 troduce shear stresses (σ_{shear}) into the stiffness tensor C_{ijkl} . The spherical geometry is
 2870 the **Unique Optimal Operating Point** where the inter-EMC forces are purely compres-
 2871 sive/extensional, minimizing internal shear stress and avoiding unstable energy dissipation.

2872 • **Gravity as Restoring Pressure:** The derivation of gravity as a "Push-out" force is
 2873 finalized. The constant G is revealed not as an independent coupling, but as a manifestation
 2874 of the geometric density deficit, explaining its weakness as a holographic projection of 4D
 2875 saturation into 3D space.

2876 The separation between the asymmetric core and the isotropic gravitational field is formalized
 2877 by the **Spherical Masking Condition**:

$$r_{\text{energy}} \leq r_{\text{EMC degraded wall}} \quad (239)$$

2878 This condition states that the asymmetric energy core (r^5 domain) is contained within the
 2879 radius of the Degraded EMC Wall (r^3 domain), enforcing gravitational isotropy through the
 2880 structural constraints of the spherical vacuum units.

2881 25 Nonlinear Stabilisation of the Electron Soliton

2882 The geometric identities derived in Sections 11 and 2 determine the coupling constants of the
 2883 vacuum, but they do not by themselves constitute a dynamical stability proof. A stable soliton
 2884 requires a nonlinear term $\mathcal{F}$ in the wave equation that counteracts dispersion. The form of $\mathcal{F}$
 2885 is constrained by the BCC lattice geometry, the topological class of the electron, and the native
 2886 1-3-6 wave-centre arrangement inherited from the EWT standing-wave model.

2887 25.1 General Soliton Wave Equation

2888 The scalar amplitude $\Psi(\mathbf{r}, t)$ of the standing wave satisfies

$$\left(\frac{\partial^2}{\partial t^2} - c^2 \nabla^2 \right) \Psi(\mathbf{r}, t) + \mathcal{F}(\Psi) = 0. \quad (240)$$

2889 The first two terms describe free wave propagation in the elastic BCC medium; the third term
 2890 encodes the nonlinear self-interaction that makes the soliton possible.

2891 25.2 Coupling Coefficient from BCC Geometry

2892 The magnetic deficit ϵ_M is the fundamental lattice response parameter derived in Section 15.5:

$$\epsilon_M = \frac{1}{N_{\text{geom}} \pi^3} = \frac{1}{8\pi^7(1 - \zeta)}. \quad (241)$$

2893 In the ideal continuum limit $\zeta \rightarrow 0$, this reduces to $\epsilon_M = 1/(8\pi^7)$. The approximate equality
 2894 reflects the 0.058% lattice impedance δ between the calibrated stiffness N_{final} and the ideal
 2895 geometric limit $8\pi^4$, as discussed in Section 21.1. Physically, ϵ_M measures the fractional loss
 2896 of stiffness caused by the soliton's spin-induced magnetic torque. To eliminate any systemic
 2897 ambiguity with the standard wave vector k , the natural nonlinear coupling strength is designated
 2898 by the parameter γ , defined as the inverse of this deficit:

$$\gamma \equiv \frac{1}{\epsilon_M} = N_{\text{geom}} \pi^3 \approx 2.414 \times 10^4. \quad (242)$$

This coefficient is not a free parameter; it follows from the BCC coordination number (8) and the dimensional budget of the charged weak scale (π^7).

For numerical implementation, two values of γ are available:

- $\gamma_{\text{geo}} = 8\pi^7 \approx 2.4162 \times 10^4$ — the pure geometric limit corresponding to ideal BCC packing ($N_{\text{geo}} = 8\pi^4$);
- $\gamma_{\text{final}} = N_{\text{final}}\pi^3 \approx 2.4148 \times 10^4$ — the value calibrated to the measured fine-structure constant, differing from γ_{geo} by the lattice impedance $\delta \approx 0.058\%$ (see Section 21.1).

The choice between them is not arbitrary: the 0.058% difference encodes the non-ideal spherical packing fraction of the physical BCC lattice, and it may prove decisive for discriminating $K = 10$ from neighbouring configurations in a numerical simulation. The OpenWave implementation should initially use γ_{final} (anchored to the measured α) and, if necessary, explore the geometric limit as a consistency check.

25.3 Proposed Forms of the Nonlinear Term

Three complementary variants are proposed, ordered by increasing structural fidelity to the EWT 1-3-6 architecture.

Variant A — Pure cubic nonlinearity

The simplest form compatible with NLS-type soliton equations is the cubic (self-focusing) nonlinearity:

$$\mathcal{F}(\Psi) = \gamma \Psi^3 = \frac{1}{\epsilon_M} \Psi^3 = N_{\text{final}} \pi^3 \Psi^3. \quad (243)$$

A term proportional to Ψ^3 is the lowest-order odd nonlinearity that preserves the sign of the restoring force; it appears generically in Lagrangian derivations from a quartic potential $V(\Psi) = \frac{1}{4}\gamma\Psi^4$.

Variant B — Nonlinearity modulated by EMC packing density

In the EWT framework the nonlinear stiffness should be active only where the granular (EMC) density is depleted relative to the statutory vacuum background ρ_0 . Let $\rho(r)$ be the local EMC density inside the soliton; the deficit fraction is $(1 - \rho(r)/\rho_0)$. The nonlinear term then becomes

$$\mathcal{F}(\Psi, \rho) = \gamma \left(1 - \frac{\rho(r)}{\rho_0}\right) \Psi^3. \quad (244)$$

Equation (244) directly couples the wave equation to the push-out mechanism (Section 11.10): the nonlinearity is maximal in the soliton core where $\rho(r) \ll \rho_0$, and vanishes in the surrounding statutory vacuum. This provides a unified description of soliton stability and gravitational emergence from a single density deficit.

Variant C — Explicit 1-3-6 source-driven dynamics

In the native EWT model the wave centres (WCs) are not an abstract continuum but ten discrete, mobile sources arranged in the 1-3-6 configuration. The vector wave equation of M4 then takes the driven form

$$\left(\frac{\partial^2}{\partial t^2} - c^2 \nabla^2 \right) \Psi(\mathbf{r}, t) + \gamma \|\Psi\|^2 \Psi = \mathbf{J}_{1-3-6}(\mathbf{r}, t), \quad (245)$$

where the source operator $\mathbf{J}_{1-3-6}$ is the sum of ten vectorial centres partitioned into three topological classes:

$$\mathbf{J}_{1-3-6}(\mathbf{r}, t) = \mathbf{j}_{\text{core}}(\mathbf{r} - \mathbf{r}_0) + \sum_{i=1}^3 \mathbf{j}_{\text{inner}}(\mathbf{r} - \mathbf{r}_i(t)) + \sum_{j=1}^6 \mathbf{j}_{\text{outer}}(\mathbf{r} - \mathbf{r}_j(t)). \quad (246)$$

Each class carries a distinct phase offset, and the kinematic rule inherited from the Yee standing-wave model applies: every centre migrates toward the local minimum of the field amplitude $\|\Psi\|$. The phase asymmetry between the inner (3) and outer (6) groups forces a synchronised rotation of the vertices, generating the 720° spin characteristic of the electron.

25.4 Structural Emergence of the 1-3-6 Geometry

Why the 1-3-6 arrangement, and not another partition of ten? The answer follows from the interplay of the nonlinearity and the BCC lattice topology:

1. The **core** (1) anchors the soliton at a single BCC lattice node, providing the central phase reference.
2. The **inner shell** (3) forms an equilateral triangle. Three is the minimal number of vertices that can define a plane; the plane's normal coincides with the spin axis. Any other number would either fail to define a unique axis (1 or 2) or, it is proposed, introduce internal phase frustration (4 or more on a single shell at this radius) — a physical claim whose verification requires the same numerical proof as the rest of the stability argument.
3. The **outer shell** (6) completes the octahedral coordination of the BCC unit cell. Six wave centres placed at the face-centre-equivalent positions are proposed to saturate the remaining propagation axes, thereby guaranteeing isotropic far-field emission after spherical masking by the Degraded EMC Wall (Section 24.7) — a claim that, like the others in this section, requires numerical verification.

It is proposed that a configuration such as 2-3-5 or 2-2-6 would either misalign the spin axis or leave some BCC axes unmatched, producing a net multipole moment that the nonlinearity cannot cancel — a prediction to be tested in the OpenWave M4 simulation. The 1-3-6 triad is therefore the unique self-consistent candidate that simultaneously satisfies: (a) the topological winding number $Q = 10$, (b) the cubic nonlinearity with coefficient γ , and (c) the octahedral symmetry of the $Im\bar{3}m$ vacuum lattice. A rigorous proof that this configuration indeed minimises the energy functional constructed from $\gamma\Psi^3$ and the $Im\bar{3}m$ projection remains a task for the OpenWave M4 simulation.

25.5 Topological Selection Rule and the Electron Ground State

The nonlinear wave equation alone does not select a specific number of wave centres. The selection of $K_{\text{WC}} = 10$ is enforced by the topological class of the soliton (Section 19.4). On the spherical shell S^2 that encloses the soliton, the field configuration is characterised by an integer winding number (equation 224):

$$Q = \frac{1}{4\pi} \int_{S^2} \Psi^* \omega \in \mathbb{Z},$$

where ω is the normalised area form on S^2 . For the electron $Q = K_{\text{WC}} = 10$.

This topological winding number is not an empirical adjustment; it emerges from the discrete projection of the BCC lattice's $Im\bar{3}m$ space group symmetry onto the continuous spherical boundary S^2 . While the 8 primary propagation axes define the innermost volumetric kernel, the configuration of $Q = 10$ wave centres is a natural candidate for the lowest-energy, stable homotopy class capable of establishing a fully isotropic boundary condition on S^2 (a problem analogous to the global minimum of the spherical Thomson packing problem for discrete phase nodes). A rigorous proof that $Q = 10$ yields a deeper potential well than $Q = 8, 9, 11$ requires an explicit evaluation of the energy functional built from the nonlinear term $\gamma\Psi^3$ and the $Im\bar{3}m$ projection, which remains a task for the OpenWave M4 simulation.

Because Q is a topological invariant, it cannot change under continuous deformations of the wave field; a transition to $Q = 9$ or $Q = 8$ would require a discontinuous rupture of the elastic medium, requiring infinite energy. The soliton is therefore permanently protected against perturbations that would alter its internal multiplicity.

Equation (240) together with any of the three nonlinear variants and the topological condition $Q = 10$ constitutes a well-posed, falsifiable model of the electron core. The stability of the 1-3-6 configuration within this model is at present a postulate, not a proof; its numerical verification is the immediate objective of the ongoing OpenWave M4 implementation described in Section 30.

1. The **nonlinearity** (γ) creates a deep, narrow potential well that resists dispersion.
2. The **topological winding number** is proposed to fix the ground state at exactly ten wave centres, pending numerical confirmation that $Q = 10$ is energetically preferred over neighbouring integers $Q = 8, 9, 11$ within the $Im\bar{3}m$ projection.
3. The **BCC stiffness constant** N_{final} (or its geometric limit $8\pi^4$) fixes the numerical value of the coupling without adjustable parameters.

25.6 Connection to the Dimensional Weight Operator

The Dimensional Weight Operator $\hat{W}(n)$ (Section 24.4) assigns a distinct degree of freedom to each factor of π in the coupling budget. The nonlinear coefficient (242) can be written as

$$\gamma = \frac{1}{\epsilon_M} = N_{\text{final}} \pi^3 \approx 8\pi^7 = 8 \cdot \pi^{(1)}\pi^{(2)}\pi^{(3)}\pi^{(4)}\pi^{(5)}\pi^{(6)}\pi^{(7)}, \quad (247)$$

where the seven factors correspond to the seven degrees of freedom of the charged weak scale (Table 19). The approximate equality with $8\pi^7$ reflects the geometric limit γ_{geo} , while the exact factorisation in terms of labelled $\pi^{(k)}$ degrees of freedom is a formal bookkeeping device that holds irrespective of the numerical value of the prefactor. Thus the stabilisation of the electron is not an isolated electromagnetic phenomenon but a direct mechanical consequence of the full 7-dimensional phase space that the BCC lattice makes available to a charged soliton. The factor 8 reflects the eight primary propagation axes of the $Im\bar{3}m$ unit cell, ensuring that the nonlinear restoring force acts isotropically.

25.7 Geometric Estimate of the Degraded EMC Wall Radius

The nonlinear stabilisation described above depends on the extent of the EMC deficit region, which is bounded by the Degraded EMC Wall (Section 24.7). A natural length scale for the wall radius r_{wall} is provided by the electron Compton wavelength λ_C , whose geometric derivation is given in Section 23.3 (see Eq. 218). Numerically $\lambda_C \approx 2.426 \times 10^{-12}$ m. Since the classical

electron radius is $r_e = \lambda_C \cdot \alpha/2\pi$, the reduced Compton wavelength $\lambda_C/2\pi$ is approximately $r_e/\alpha \approx 137 r_e$. A plausible scale is

$$r_{\text{wall}} \sim \frac{\lambda_C}{2\pi} = \frac{r_e}{\alpha}. \quad (248)$$

Because α is itself derived from the BCC lattice geometry (Eq. 17), the relation $r_{\text{wall}} \sim r_e/\alpha$ does not introduce a new parameter; it is algebraically equivalent to the standard QED definition of the classical electron radius. The significance of this estimate lies not in its novelty but in its consistency: the same geometric framework that yields α and λ_C from r_ν and the $8\pi^7$ correction also predicts a natural length scale for the region in which the nonlinearity is active, providing a unified description of particle stability, forces, and annihilation. The considerable width of the wall — roughly $137 r_e$ — is not an arbitrary feature but a geometric consequence of the ratio $1/\alpha \approx 137$. Physically, this extended transition zone provides a broad buffer within which the nonlinearity can adiabatically relax the EMC density from its deep core deficit to the statutory background, suppressing the sharp gradients that would otherwise destabilise the standing wave.

Role of a Possible EMC Density Peak

The preceding estimate fixes the radial scale of the wall, but leaves open the question of its internal density profile. As noted in Section 11.11, two scenarios are compatible with the integral deficit $N_{\nu,\text{eff}}$: a monotonic approach to the statutory background, or a local density peak ($\rho_{\text{wall}} > \rho_0$) caused by the accumulation of EMCs expelled by the push-out mechanism. If such a peak exists, it would actively assist the nonlinear stabilisation in two ways. First, in Variant B the factor $(1 - \rho/\rho_0)$ would change sign within the peak, turning the self-focusing nonlinearity into a local defocusing barrier that prevents the standing wave from leaking outward. Second, even for the unmodulated Variants A and C, a density peak implies a locally higher elastic modulus, which acts as a partial reflector, increasing the Q -factor of the soliton resonator. Whether the peak is indispensable or merely auxiliary can only be decided once the full radial EMC profile is computed from the nonlinear wave equation, which remains a task for the OpenWave M4 simulation.

Internal Wave-Centre Dynamics and Spin

In the foundational Energy Wave Theory (Yee, 2019), an additional hypothesis was put forward that could further clarify the electron's stability: the electron's ten wave centres are not perfectly static but undergo continuous micro-motion around their nodal positions. According to this picture, a wave centre displaced from its node experiences a restoring force toward the local amplitude minimum; this perpetual shifting of centres is the origin of the 720° spin and, at the same time, a dynamic stabilisation mechanism. The tetrahedral 1-3-6 geometry ensures that only a fraction of the centres are off-node at any instant, while the remainder anchor the structure. Configurations such as $K_{WC} = 9$ or $K_{WC} = 11$ lack this balance and break apart. This hypothesis has not yet been proven mathematically and should be tested in the OpenWave M4 model once the nonlinear term and the Degraded EMC Wall are in place: comparing simulations with and without WC motion could reveal whether the internal dynamics is an essential ingredient or a secondary effect. In the language of the Dimensional Weight Operator (Section 24.4), this internal dynamics may be tentatively associated with the degree of freedom $\pi^{(6)}$ (soliton radius), which would thereby acquire a dynamic interpretation beyond its static geometric extent — a reinterpretation that remains speculative, is not used in deriving any numerical result in this paper, and applies to this one degree of freedom only.

25.8 Relation Between the Energetic and Topological Arguments

The present section advances two complementary conditions that together single out $K_{WC} = 10$: (i) an **energetic** condition based on the r^5/r^3 scaling disparity, which creates a deep potential well for intermediate wave-centre counts (Section 15.1); and (ii) a **topological** condition requiring an integer winding number Q on S^2 (Section 19.4). These are not independent post-hoc justifications of the same number; they are mutually reinforcing constraints. The r^5/r^3 argument explains why K_{WC} cannot be too small (shallow well) or too large (volumetric overload), while the topological argument explains why, within the allowed range, only discrete integer values are permitted. The specific value $K_{WC} = 10$ emerges as the unique integer that simultaneously satisfies both constraints under the octahedral symmetry of the $Im\bar{3}m$ BCC lattice. A rigorous proof that no other integer passes both filters remains a task for the OpenWave M4 simulation.

25.9 Summary

- The general soliton wave equation is $(\partial_t^2 - c^2 \nabla^2) \Psi + \mathcal{F}(\Psi) = 0$.
- The coupling coefficient is fixed by BCC geometry: $\gamma = 1/\epsilon_M = N_{\text{final}} \pi^3 \approx 2.41 \times 10^4$. Two variants are available: γ_{final} (anchored to α) and $\gamma_{\text{geo}} = 8\pi^7$ (ideal BCC limit); the 0.058% difference may be critical for K-selectivity.
- Three complementary forms for $\mathcal{F}$ are proposed:
 - Variant A: $\gamma \Psi^3$ (pure cubic, minimal implementation);
 - Variant B: $\gamma(1 - \rho/\rho_0) \Psi^3$ (density-modulated, links stability to the EMC push-out deficit);
 - Variant C: $\gamma \|\Psi\|^2 \Psi$ with an explicit 1-3-6 source operator $\mathbf{J}_{1-3-6}(\mathbf{r}, t)$ (directly encodes the discrete wave-centre architecture of the electron).
- The Degraded EMC Wall is proposed to provide the necessary boundary condition for the nonlinearity: its radius, estimated as $r_{\text{wall}} \sim r_e/\alpha$, is consistent with the Compton wavelength and may ensure a broad adiabatic transition zone ($\sim 137 r_e$) in which the density-modulated term can act without sharp gradients. A possible density peak within the wall would further assist stabilisation by creating a local defocusing barrier.
- The winding number $Q = 10$ on S^2 is proposed as the topological selection rule that would isolate the electron ground state under $Im\bar{3}m$ constraints; rigorous proof that $Q = 10$ is energetically preferred over other integers remains a task for the OpenWave M4 simulation.
- The 1-3-6 partition is the unique self-consistent candidate that simultaneously satisfies the topological winding, the cubic nonlinearity, and the octahedral symmetry of the BCC lattice; its stability is a postulate pending numerical verification.
- The energetic (r^5/r^3) and topological arguments are complementary: the former restricts the range of allowed K_{WC} , the latter restricts K_{WC} to integer values; together they single out $K_{WC} = 10$ as the only candidate satisfying both constraints.
- The coupling $\gamma \approx 8\pi^7$ (in its geometric limit) directly expresses the 7-dimensional charged-weak budget of the BCC lattice.
- The model is well-posed and falsifiable; the stability of the 1-3-6 configuration is a postulate whose numerical verification is the immediate objective of the OpenWave M4 simulation.

26 Predictive Power

The EWT model yields several precise, quantitative predictions that follow directly from the geometry of the Soliton and the BCC lattice. The results summarized in Table 20 demonstrate the internal consistency of the zero-calibration framework.

Table 20: Principal Geometric Predictions of the EWT Framework

Physical Quantity	Geometric Prediction	Target	Relative Error
G_{geom}	$6.6775199755 \times 10^{-11}$	6.674305×10^{-11}	0.048169%
$\alpha_{\text{geom}}^{-1}$	137.036262364	137.035999084	0.000192%
a_e^{EWT}	1159.916228 ppm	1159.652182 ppm	0.022769%
a_μ^{EWT}	1166.212608 ppm	1165.920610 ppm	0.025044%
a_τ^{EWT}	1176.838130 ppm	1177.210000 ppm	0.031589%

Note: Values for a_e , a_μ , and a_τ are full anomalous magnetic moments. All predictions are obtained from the geometric Python implementation described in the Appendix. Error for a_τ is compared with SM prediction.

26.1 Prediction of the Fundamental Value of G

The model predicts the value of G as a direct function of microscopic geometric parameters through the operator $\mathcal{U}$ (Section 11.5). For the zero-calibration inputs used in this work, the geometric identity gives

$$G_{\text{geom}} \approx 6.6775199755 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}. \quad (249)$$

Precise experimental measurements of G (e.g., torsion balance, atom interferometry) directly test this prediction.

26.2 Predictions for Lepton Properties: The First-Principles AMM Predictions

The full anomalous magnetic moments of the entire lepton family are computed from the same BCC lattice geometry that governs G and α_{geom} . The results form a compact monotonic sequence:

$$\boxed{a_e^{\text{EWT}} = 1.159916 \times 10^{-3}} \quad \boxed{a_\mu^{\text{EWT}} = 1.166213 \times 10^{-3}} \quad \boxed{a_\tau^{\text{EWT}} = 1.176838 \times 10^{-3}} \quad (250)$$

with relative errors 0.022769%, 0.025044%, and 0.031589%.

The projection rules $\mathcal{O}_e = \mathcal{O}_\tau = 1$ and $\mathcal{O}_\mu = 1/(4\pi^2)$ are fixed by the dimensional hierarchy and contain no generation-specific parameters. A fourth generation would require a second 2D projection and is geometrically forbidden by the saturation limit of the BCC lattice.

26.3 Magnetic Sensitivity $G(B)$ and Hypotheses Testing

The model predicts a dependence of the effective gravitational coupling on local magnetic coherence, as described in Section 18. In the zero-calibration formulation, the push-out operator

3105 is expressed through the geometric stiffness:

$$\mathbf{T}_{\text{II}} = \left(\frac{1}{A_{\pi} \cdot N_{\text{geom}}} \right)^3. \quad (251)$$

3106 The observable shift is expected to be small in ordinary environments:

$$\left(\frac{\Delta G}{G} \right)_{\text{obs}} = \xi(B) \cdot \mathbf{T}_{\text{II}}, \quad (252)$$

3107 where $\xi(B)$ is the coupling efficiency of the local magnetic field to the soliton's geometry. The
3108 sign of the effect distinguishes the hypotheses of Section 18.4.

3109 26.4 AMM as a Strong Test of Geometric Universality

3110 Because AMM measurements are exceptionally precise, they provide an ideal testing ground for
3111 the geometric mechanism. A single universal modulator ϵ_M , together with the recursive nodal
3112 structure, replaces the perturbative loop corrections of the Standard Model.

3113 26.5 Correlations with Neutrino Flux

3114 Since G emerges from the effective density $N_{\nu, \text{eff}}$, the model allows minute correlations between
3115 measured G and local neutrino flux densities. High neutrino flux may momentarily alter the
3116 local EMC background, leading to sub-ppm modulations of G .

3117 26.6 Spectral and Geometric Modal Tests

3118 The modal nature of the π^3 factor implies that internal geometric excitations may leave traces
3119 in the frequency spectrum of small fluctuations of G . The model predicts characteristic nodal
3120 frequencies of the soliton's standing modes.

3121 26.7 Tests using Gravitational Wave Observations

3122 If G is emergent from the BCC lattice, gravitational waves may exhibit small dispersion effects:

$$v_{\text{GW}} \simeq c(1 - \delta_{\text{dispersion}}), \quad (253)$$

3123 where $\delta_{\text{dispersion}}$ depends on the local nodal density. Arrival-time comparisons between GW and
3124 electromagnetic signals constrain this effect.

3125 26.8 Test on Bose-Einstein Condensates

3126 A BEC state synchronises the local magnetic coherence across a macroscopic volume. The model
3127 predicts a shift

$$G_{\text{eff}}^{(\text{BEC})} = G(1 + \delta_{\text{BEC}}), \quad \delta_{\text{BEC}} \propto \mathbf{T}_{\text{II}}. \quad (254)$$

3128 Consistent with the push-out mechanism, the primary prediction is $G_{\text{eff}} > G$. Atom interferom-
3129 etry on ultra-cold clouds before and after BEC formation can test this directly.

26.9 The Higgs Soliton vs. Fundamental Scalar Field

The EWT model identifies the Higgs as a resonant state ($K_{WC} = 117$) rather than a fundamental scalar field. This leads to specific predictions for Higgs-vector mixing:

$$\sin^2 \theta_{ZH} \approx 0.4686, \quad \sin^2 \theta_{WH} \approx 0.5906. \quad (255)$$

Angular asymmetries or interference patterns in Higgs decays at HL-LHC or FCC that match these geometric values would falsify the VEV mechanism of the Standard Model.

27 Falsifiability and Model Constraints

The following outcomes represent critical tests that can either **falsify the core tenets of the EWT model** or place stringent constraints on specific parameters and hypotheses. These tests are focused on the **existence or non-existence** of predicted effects, independent of the sign of the dynamic coupling (Hypotheses 1 and 2).

- **Falsification of Fundamental Constants (G):** Precise measurements of G consistently rejecting G_{model} (Eq. 26.1) beyond the level of uncertainty while maintaining the assumed values of microscopic parameters.
- **Falsification of Dynamic Coupling ($G(B)$):** The observation of a null effect, i.e., the absence of any observed $G(B)$ dependence above the detection threshold, assuming the technical feasibility of such a test. This outcome would support **Hypothesis 3 (No Influence)** and refute the core idea of magneto-geometric coupling.
- **Falsification of Correlations:** The absence of statistical correlations between fluctuations in G and external factors (neutrino flux, magnetic fields) within the range predicted by the model (ref. Section 26.5).
- **Falsification of Full Lepton AMM Predictions (a_e, a_μ, a_τ):** Future high-precision measurements of the electron, and muon, anomalous magnetic moments that deviate significantly from the full AMM predictions $a_e^{\text{EWT}} = 1.159916 \times 10^{-3}$, $a_\mu^{\text{EWT}} = 1.166213 \times 10^{-3}$, and $a_\tau^{\text{EWT}} = 1.176838 \times 10^{-3}$ (Table 9) would falsify the model in its present formulation. For the tau, the current reference value is the Standard Model prediction $a_\tau^{\text{SM}} = 1.177210 \times 10^{-3}$, not a direct measurement. Therefore the tau comparison is a consistency test against the SM prediction, not a direct experimental falsification. The projection rules underlying these predictions are completely determined by the dimensional hierarchy of the Geometric Ladder: $\mathcal{O}_e = 1$ (3D core), $\mathcal{O}_\mu = 1/(4\pi^2)$ (2D planar shell), and $\mathcal{O}_\tau = 1$ (3D volumetric shell). A significant deviation of any of the three measured anomalies from the predicted values would therefore directly challenge the geometric core of the model. The compact monotonic sequence of prediction errors— $0.022769\% \rightarrow 0.025044\% \rightarrow 0.031589\%$ —provides a stringent, correlated test: any future measurement that breaks this monotonicity would also indicate a structural tension with the BCC lattice framework.

27.1 Practical Considerations and Detection Limits

In practice, the expected magnitude of the relative change $\Delta G/G$ is governed by the robustness of the dynamic geometric equilibrium between energy scaling (r^5) and volumetric displacement (r^3). Due to this compensatory mechanism, the predicted deviations in near-equilibrium (low-field) environments are expected to be extremely subtle, likely requiring a detection sensitivity in the range of 10^{-9} to 10^{-12} .

While these requirements are stringent, they align precisely with the current state-of-the-art in quantum metrology and atom interferometry. Modern platforms, such as MAGIS-100 or advanced gravity gradiometers, are specifically designed to operate within these detection limits to probe dark matter and gravitational waves.

The model further posits that the detection threshold is fundamentally tied to the **magnetic flux density** or **quantum coherence level** of the source. In extreme high-field regimes or coherent macroscopic states (such as BEC), the compensatory mechanism is hypothesized to reach its non-linear limit, potentially shifting the magnitude of $\Delta G/G$ towards the 10^{-7} range. Consequently, even a null result within current experimental sensitivities would provide critical boundary values for the volumetric "stiffness" of the EMC medium governed by $\mathbf{T}_{II}$ and the saturation point of the geometric equilibrium. This ensures that the framework remains fully falsifiable, providing a clear mapping for the transition from the "low-field artefact" of invariant G to the dynamic G_{eff} regime predicted by EWT.

27.2 Mutual Falsification: The Ultimate Test

This creates a definitive scientific standoff:

- **Validation of SM:** If the Higgs boson continues to behave as a featureless, point-like scalar with no evidence of internal geometry or discrete mixing ratios, EWT's soliton model for heavy bosons will be challenged.
- **Validation of EWT:** If experiments reveal any discrete, geometric substructure in $H - W$ or $H - Z$ interactions, the Standard Model **automatically falls**, as its mathematical framework cannot accommodate a structured Higgs soliton without collapsing the VEV mechanism.

By providing these specific targets ($\sin^2 \theta_{WH,ZH}$), EWT transitions from a descriptive model to a **fully falsifiable predictive theory**, offering a clear roadmap for the next generation of experimental particle physics.

28 Robustness Analysis: Geometric Stability and Sensitivity

To demonstrate that the Enhanced EWT model is not a result of numerical coincidence, a series of sensitivity tests were performed. This analysis examines how small perturbations in the fundamental geometric inputs affect the physical outputs. By isolating each primary variable, the structural rigidity of the vacuum lattice is verified against experimental CODATA/PDG benchmarks.

28.1 Robustness Analysis of the Neutrino Soliton Radius and Geometric Identity

The structural integrity of the EWT framework is evaluated through the stability of the fundamental identity

$$\Delta N_\nu \sim \frac{1}{\epsilon_G}. \quad (256)$$

where ϵ_G is the gravitational deficit associated with the reference soliton. This equivalence suggests that the statutory constituent count, derived from the ratio of the neutrino radius to

the Planck scale, must inherently align with the geometric deficit required to scale gravity from its classical base G_{Base} to the observed value.

To verify the absence of numerical fine-tuning, a sensitivity analysis was conducted by applying relative perturbations of $\pm 20\%$ to the primary geometric inputs: the statutory neutrino radius r_ν and the fundamental EMC spacing λ_l . The resulting compliance curves are shown in Fig. 8. They demonstrate that even under significant variations in the input scales, the compliance ratios remain strictly within the $\pm 30\%$ tolerance boundaries.




Figure 8: **Structural Robustness Analysis of the Statutory Density $N_{\nu,\text{stat}}$** . The plot tracks the stability of the statutory background population $N_{\nu,\text{stat}}$ against relative fluctuations in the neutrino radius r_ν and the EMC spacing λ_l . Both curves remain within the established $\pm 30\%$ compliance boundaries (0.7 to 1.3 ratio). This stability ensures that the geometric vacuum parameters do not require fine-tuning.

This performance confirms that the EWT model is not a product of precise numerical coincidence, but is instead rooted in a robust geometric power-law relationship.

28.2 Gravitational Surface Transition and $N_{\nu,\text{eff}}$

The robustness test focuses on the relationship between the geometric volume deficit and the emergent gravitational constant G . As detailed in Section 11.10, gravity is interpreted as a surface effect resulting from a three-dimensional packing deficit within the EMC medium.

The results illustrated in Fig. 9 confirm the following physical constraints:

- **Transition trajectory:** The evolution of the curve follows the $G \propto N_\nu^{-1/2}$ scaling law. This confirms the gravitational interaction as a surface manifestation of the internal volumetric dilution.




Figure 9: Analysis of the G-constant surface transition. The green curve illustrates the evolution of G as a function of the effective volume deficit N_ν . The red intersection point marks the deterministic alignment with G_{CODATA} at the calculated $N_{\nu,\text{eff}}$.

- **Deterministic convergence:** The intersection with the G_{CODATA} threshold justifies the transition to the effective volume deficit

$$N_{\nu,\text{eff}} \approx 6.2472 \times 10^{48}, \quad (257)$$

obtained from the geometric dilution factor X_{eff} .

- **Model rigidity:** The steep slope of the transition curve indicates that the system is highly sensitive to the internal geometry. A deviation of even 1% in the value of N_ν results in a significant departure from G , proving that the EWT unification is not a result of arbitrary curve-fitting.

28.3 Sensitivity of α^{-1} to the Geometric Modulator N_{geom}

This test evaluates the stability of the fine-structure constant derivation by examining the influence of the dimensionless geometric stiffness N_{geom} . In the EWT framework, $\alpha_{\text{geom}}^{-1}$ is determined by the fixed vacuum geometry A_π modulated by the magnetic deficit $\epsilon_M = 1/(N_{\text{geom}}\pi^3)$.

The analysis of the results presented in Fig. 10 confirms:

- **Geometric necessity:** Precise alignment with $\alpha_{\text{CODATA}}^{-1}$ occurs only when the complete stability factor is applied. The “golden point” intersection at

$$N_{\text{geom}} \approx 778.8025 \quad (258)$$




Figure 10: Sensitivity analysis of the inverse fine-structure constant. The plot illustrates the convergence of the model toward the CODATA value as a function of the geometric stiffness coefficient N_{geom} .

proves that the fine-structure constant emerges from a rigid geometric base corrected by a finite magnetic deficit.

- **Force hierarchy indication:** While the modulator ϵ_M is shared by both G and α^{-1} , the gravitational constant operates on a significantly different scale through the cubic factor $(\epsilon_M/A_\pi)^3$ and the surface projection $N_{\nu,\text{eff}}^{-1/2}$.

28.4 Phase Space Analysis: EWT Trajectory vs Standard Model Interpretation

The phase-space mapping between the inverse fine-structure constant α^{-1} and the electron anomalous magnetic moment a_e reveals a fundamental geometric trajectory. By varying the vacuum stiffness parameter N , the model demonstrates that α^{-1} and a_e are not independent constants, but emergent properties of the same underlying geometric modulator.

The experimental CODATA point lies close to this EWT trajectory, as shown in Fig. 11, with a residual offset of approximately 0.022769%. In the EWT framework, this offset is not interpreted as model failure, but as a consequence of the structural difference between the geometric vacuum picture and the point-like renormalisation programme of the Standard Model.




Figure 11: Phase space correlation between the inverse fine-structure constant and the electron anomalous magnetic moment. The curve represents the EWT geometric trajectory, showing that α^{-1} and a_e emerge from the same vacuum stiffness.

28.5 Parametric Unification: The Stability Path of G and α

A parametric analysis of the relationship between G and α^{-1} was performed by varying the geometric modulator N . The resulting trajectory, derived from the geometric operator $\mathcal{U}$, shows that α^{-1} remains tightly constrained by the core geometry A_π , whereas G is significantly more sensitive to changes in the stiffness parameter.

The resulting trajectory is shown in Fig. 12. This sensitivity difference follows directly from the algebraic structure:

$$\alpha_{\text{geom}}^{-1} \sim A_\pi - \frac{1}{N\pi^3}, \quad G_{\text{geom}} \sim \frac{1}{N^3 \sqrt{N_{\nu, \text{eff}}}}. \quad (259)$$

Thus, small changes in N affect G much more strongly than α . This explains why the electromagnetic sector appears stable while the gravitational sector is more sensitive to microscopic geometric parameters.

28.6 EWT Unification: Convergence of G and a_e on the Vacuum Stiffness Isentrope

A fundamental consistency check is the interdependence between G and a_e at a single geometric stiffness value

$$N_{\text{geom}} \approx 778.8025. \quad (260)$$




Figure 12: The EWT Unification Path. The magenta line represents the geometric correlation between gravity and electromagnetism as a function of the vacuum stiffness parameter N .

At this point, G_{geom} and a_e^{EWT} are both determined from the same BCC lattice parameters, as illustrated in Fig. 13. This is not a fit: the value of N_{geom} is fixed by the packing geometry and the impedance ζ .

28.7 Robustness Analysis: Geometric Stability and Vacuum Elasticity

To evaluate the reliability of the derived gravitational constant, a sensitivity analysis of the coupling between the soliton volume and the vacuum stiffness was performed. The structural stability is governed by the relationship

$$N \propto N_{\nu, \text{eff}}^{-1/6}, \quad (261)$$

which follows from the volumetric scaling $N_{\nu} \propto r^3$ and the stiffness-radius relation $N \propto r^{-1/2}$.

The resulting stability relationship is illustrated in Fig. 14. This damping factor protects the geometric vacuum parameters from large fluctuations of the EMC density. In the current model, the relevant stiffness is N_{geom} , not an arbitrary fitted value.

28.8 Structural Robustness and Resonance Stability of the Lepton Hierarchy

The same geometric stiffness N_{geom} anchors the recursive lepton hierarchy. The sensitivity gradients of a_{μ} and a_{τ} are shown in Fig. 15. They differ because the muon is a 2D surface resonance,




Figure 13: EWT Unification: Direct convergence of the gravitational constant G and the electron anomaly a_e at the geometric stiffness N_{geom} .

while the tau is a 3D volumetric resonance. This is consistent with the dimensional hierarchy established in Section 3.

28.9 Electroweak and CKM Coincidence: Geometric Robustness of $\sin^2 \theta_W$ and $\sin \theta_C$

The electroweak and Cabibbo mixing sectors converge at the same geometric anchor N_{geom} , as shown in Fig. 16. The π^6 bosonic coupling is steeper than the π^5 fermionic coupling, reflecting the difference between volumetric and surface resonances.

The Cabibbo sector remains sensitive to the quality of the quark mass inputs. When PDG experimental quark masses are used, the geometric mixing operator C_{fermion} reproduces $\sin \theta_C$ to within 0.0055%. The remaining discrepancy with EWT-derived quark masses originates from the mass prediction sector, not from the mixing geometry itself.

28.10 Lock-In Test: Independent Determination of N_{geom}

The robustness analysis presented so far demonstrates that the model is stable under perturbations of its inputs. A complementary and more stringent test concerns whether the geometric stiffness N_{geom} is *over-determined* by independent physical sectors, or merely fitted to a single observable. The results of such a test are reported below.




Figure 14: Gravity Stability Equilibrium: The $r^{-1/2}$ trajectory of nodal stiffness N relative to the cubic scaling of soliton volume N_ν .

28.10.1 Method

For each of seven physical sectors — α^{-1} , G , a_e , a_μ , a_τ , $\sin^2 \theta_W$, and $\sin \theta_C$ — the following quantities are computed:

1. the value of the sector at $N_{\text{geom}} = 778.8025$;
2. the *drift* of that value when N is changed by ± 10 ;
3. the logarithmic derivative $\beta = d \ln O / d \ln N$ at N_{geom} , obtained by central difference in $\log N$ with step 10^{-6} ;
4. the *resolution*, defined as the relative error of the sector at N_{geom} divided by $|\beta|$, which measures how narrowly the sector can locate N ;
5. the *optimal* value N_{opt} for each sector, found by log-space scan followed by bisection.

The procedure is implemented in the OpenWave script `m4_11_lock_in_n_scan.py` [54]. The sectors are classified into three groups according to their resolution:

- **PRIMARY** — resolution $< 1\%$. These sectors can independently locate N at the sub-percent level.
- **WEAK** — resolution between 1% and 100% . These sectors are consistent with N_{geom} but cannot discriminate it from neighbouring values.




Figure 15: Stability Analysis of the Lepton Hierarchy: Normalized response of a_μ and a_τ to geometric perturbations.

- **NO CONSTRAINT** — resolution $> 100\%$. These sectors provide no information about N .

28.10.2 Results

The resolution analysis is summarised in Table 21. The grouping of sectors follows directly from the resolution column.

Table 21: Resolution of N_{geom} by independent physical sectors. The three groups are separated by horizontal rules.

Sector	$\beta = d \ln O / d \ln N$	Resolution	N_{opt}
<i>PRIMARY (resolves N, $< 1\%$)</i>			
G	-12.000002	4.0×10^{-5}	778.83
a_τ	+0.220297	1.4×10^{-3}	779.92
$\sin \theta_C$	-0.008921	6.2×10^{-3}	774.03
<i>WEAK (consistent, 1%–100%)</i>			
$\sin^2 \theta_W$	-0.045754	1.5×10^{-1}	910.04
a_e	+0.001285	1.8×10^{-1}	661.63
a_μ	+0.001415	1.8×10^{-1}	661.69
<i>NO CONSTRAINT ($> 100\%$)</i>			
α^{-1}	$+3.0 \times 10^{-7}$	6.4×10^0	105.85




Figure 16: Shift-Normalized Robustness Analysis: The intersection of the $\sin^2 \theta_W$ and $\sin \theta_C$ trajectories at the geometric lock-in point N_{geom} .

28.10.3 PRIMARY sectors: G , a_τ , and $\sin \theta_C$

Three sectors resolve N at the sub-percent level. The two strongest are the gravitational constant G (0.004%) and the tau anomalous magnetic moment a_τ (0.14%). Their separately optimal values are

$$N_{\text{opt}}(G) = 778.83, \quad N_{\text{opt}}(a_\tau) = 779.92, \quad (262)$$

which agree to within

$$\Delta = 0.14\% \quad \text{of } N_{\text{geom}}. \quad (263)$$

Crucially, the two sectors are constructed from completely different chains. G arises from the gravitational self-consistent derivation involving λ_l , $N_{\nu, \text{eff}}$, and the surface projection. a_τ arises from the recursive lepton shell damping governed by ϵ_M and the Fibonacci latching. Neither chain uses the other's inputs. The agreement between them is therefore not tautological.

The exponent $\beta_G = -12$ (rather than the naive -3) is a structural consequence of the self-consistent derivation of λ_l : the lattice spacing scales as $\lambda_l \propto N^{-6}$, and G inherits an additional N^{-9} dependence through $N_{\nu, \text{eff}}$, compounding with the explicit N^{-3} factor. This makes gravity by far the most sensitive probe of the lattice stiffness.

The Cabibbo angle $\sin \theta_C$ formally lands in the PRIMARY group with a resolution of 0.62%. This result is, however, conditioned on the central PDG values for the light quark masses. The experimental uncertainty on m_d and m_s spans approximately 14%, roughly fifteen times larger than the geometric correction being tested. At the low corner of that range no positive N

reproduces $\sin\theta_C$. The Cabibbo leg is therefore treated as *conditionally* primary, not as a fully independent constraint.

28.10.4 WEAK sectors: a_e , a_μ , and $\sin^2\theta_W$

Three sectors are consistent with N_{geom} but cannot discriminate it from neighbouring values. Their resolutions of approximately 15–18% reflect different suppression mechanisms.

For the electron anomalous magnetic moment a_e , the sensitivity to N is suppressed because the prediction is dominated by the nearly N -independent Schwinger term. The electron AMM therefore confirms the overall framework but does not constrain the lattice stiffness.

For the muon anomalous magnetic moment a_μ , the projection operator $\mathcal{O}_\mu = 1/(4\pi^2)$ suppresses the sensitivity of the muon shell. The muon leg is consistent with N_{geom} but, like the electron, cannot discriminate between neighbouring values.

For the Weinberg angle $\sin^2\theta_W$, the logarithmic derivative is small ($\beta \approx -0.046$), giving a resolution of 14.6%. The comparison between the model value $\sin^2\theta_W = 0.23276$ and the PDG reference uses the MS-bar scheme, while the on-shell value derived from the PDG masses is 0.22305. The geometric correction C_{gap} occupies the same slot as a known scheme-conversion factor of approximately 1.06%. Consequently, the Weinberg leg cannot yet carry independent weight.

28.10.5 NO CONSTRAINT: α^{-1}

The inverse fine-structure constant provides no constraint on N . Its logarithmic derivative is $\beta \approx 3 \times 10^{-7}$, giving a resolution of 635%. This is expected and consistent with the model structure: α^{-1} is dominated by the geometric core A_π , and the ϵ_M correction is tiny. α therefore confirms the model but does not constrain its central parameter. Its inclusion in the lock-in analysis serves as a consistency check: the absence of a constraint is itself a prediction of the model's structure.

28.10.6 Interpretation

The lock-in test demonstrates that N_{geom} is not a fitted parameter in the usual sense. It is a *node* at which two independent physical sectors — gravity and the tau anomalous magnetic moment — converge to within 0.14%, with completely different construction chains and different sensitivities to the lattice stiffness.

The PRIMARY sectors are those whose prediction chains contain N at high powers ($\beta_G = -12$, $\beta_{a_\tau} = 0.22$) and whose targets are known with sufficient precision. The WEAK sectors are those whose sensitivity is suppressed by projection operators or by dominant N -independent terms. The NO CONSTRAINT sector is dominated by the geometric core A_π and is essentially insensitive to N .

This hierarchy is a falsifiable structural feature of the model. If future measurements of a_τ (currently only bounded experimentally) or of G (currently known to 2.2×10^{-5}) were to shift either sector's optimal N beyond the common interval, the model would be challenged. If, on the other hand, the PRIMARY sectors were to remain in agreement while a WEAK sector moved into the PRIMARY group, the model would be strengthened by the addition of a third independent constraint.

29 Geometric Phase Transitions: From Neutrino to Black Hole with EMC Wall

In the EWT framework, the vacuum lattice is a physical medium with a finite redistribution capacity. The formation of a **Geometric Black Hole (GBH)** represents a phase transition where the vacuum is pushed to its absolute elastic and geometric limits.

29.1 Hierarchy of Vacuum States and the G-Operating Point

The hierarchy of states is governed by the depth of the density deficit relative to the statutory equilibrium:

- **Effective Point** ($N_{\nu,\text{eff}} \approx 6.25 \times 10^{48}$): The standard operating point of attractive gravity (approx. 99.98% deficit relative to background).
- **Statutory Limit** ($N_{\nu,\text{stat}} \approx 3.30 \times 10^{52}$): The saturation ceiling where gravity inverts into repulsion (**EMC Wall**).
- **Max Packing** ($N_{\nu,\text{max}} \approx 5.30 \times 10^{54}$): The asymptotic structural limit of the BCC lattice at the Planck scale.

The GBH Core: Finite Vacuum Exhaustion and Pure Wave Centers

Unlike General Relativity, EWT defines the interior of a GBH as a domain of **maximal individual energy concentration** (ρ_E) and **maximal medium deficit** (ρ). This state is realized by high-energy neutrino solitons acting as pure Wave Centers (WC).

Empirical Validation (Supernovae): The observation that **99% of supernova energy is released as neutrinos** constitutes empirical confirmation that the collapse of matter under extreme gravity leads to the conversion of the dominant mass into the **minimal geometric state** of the Neutrino (r_ν) at the point d_{crit} . This fact justifies the adoption of the Neutrino as the fundamental WC for the GBH model.

The EMC Wall and the Push-Out Mechanism

The EMC Wall is a physical density barrier in a vacuum created when the medium (BCC network) cannot keep up with the redistribution of displaced components (EMC). In ordinary solitons (such as leptons) this barrier exists in a **degraded** form – the *Degraded EMC Wall* discussed in Sec. 11.11 – where the local EMC density exceeds the statutory background $N_{\nu,\text{stat}}$ only moderately, and the wall acts as a passive geometric low-pass filter. In a Geometric Black Hole (GBH), however, the push-out mechanism is driven to its extreme: the core exhausts the vacuum so severely that the accumulated EMC density at the wall **significantly exceeds** the statutory background $N_{\nu,\text{stat}}$, turning the wall into an insurmountable barrier. The EMC wall then defines the geometric event horizon of the black hole. The relationship between the black hole core's energy and the wall's density is defined by the displacement flux:

$$\Delta\rho_{E(\text{core})} \xrightarrow{\text{push-out}} \Delta\rho_{(\text{wall})} \geq N_{\nu,\text{stat}} \quad (264)$$

Energy Dissipation and Degradation

Energy exchange between the Geometric Black Hole and the external environment remains possible; however, the emitted energy must adopt a **degraded form**. The EMC Wall acts as a

3414 mandatory transducer, where the extreme resistance of the saturated lattice forces all outgoing
3415 flux into states compatible with the medium's localized stiffness.

3416 **Astrophysical Jets and the Density Wall**

3417 The maximal vacuum exhaustion within the GBH core necessitates a **massive reciprocal ac-**
3418 **cumulation** of EMCs at the event horizon's boundary—the **EMC Density Wall**. This wall
3419 possesses extreme nodal stiffness, anchoring ultra-strong magnetic fields. Astrophysical jets act
3420 as **structural discharge mechanisms**, where the immense potential of the compressed lattice
3421 is released along the axes of least resistance (the poles).

3422 **The EMC Barrier and Orbital Stability**

3423 Beyond the **Statutory Limit** ($N_{\nu, \text{stat}}$), the density gradient undergoes a sign reversal ($\nabla \rho > 0$).
3424 This creates a localized **Repulsion Zone** (anti-gravity). The event horizon thus functions as a
3425 buffer; matter is either stabilized by this "gravitational conjunction" or **actively ejected** into
3426 jets upon encountering the super-statutory pressure of the wall.

3427 **The EMC Wall: Selective Frequency Response**

3428 The extreme nodal stiffness of the **EMC Wall** (approaching $N_{\nu, \text{max}}$) establishes a **high-pass**
3429 **filter**. Because the lattice bonds in the wall are pushed to their elastic limit, only waves with **ex-**
3430 **tremely high frequencies** (Gamma and X-rays) have the energy density required to overcome
3431 the medium's inertia.

3432 **The EMC Wall: Decomposition of Matter**

3433 As leptonic matter approaches the boundary, the transition on the wall's congestion creates an
3434 insurmountable **impedance mismatch**. The vacuum lattice becomes too rigid to maintain the
3435 wave structure of leptons, leading to the **structural breakdown** of the particle soliton.

3436 **Hypothesis: Gravitational Waves as Background Density Shifts**

3437 Within the EWT framework, gravitational waves can be interpreted as propagating pulses that
3438 momentarily increase the **statutory background density** ($N_{\nu, \text{stat}}$) of the BCC lattice. Cru-
3439 cially, the soliton's internal push-out mechanism is blocking EMCs; the matter does not absorb
3440 additional EMCs. Instead, the passing wave momentarily elevates the surrounding medium's
3441 density toward the EMC Wall threshold. As the background density $N_{\nu, \text{stat}}$ surges, the relative
3442 gradient between the soliton's core (the vacuum exhaustion zone) and the statutory environment
3443 is altered. This transient shift in the background reference point modulates the Ω_{Push} operator,
3444 manifesting as a temporary change in the "gravitational shadow". Once the peak of the back-
3445 ground EMC pulse passes, the lattice undergoes relaxation, restoring the statutory equilibrium.
3446 This interpretation suggests that gravitational waves are not a change in matter itself, but a
3447 moving "impedance spike" in the vacuum background that matter inhabits.

3448 In this scenario, the matter "does not notice" the wave because its internal structural iden-
3449 tity is preserved relative to the shifting background. The wave is only detectable via phase-
3450 sensitive measurements (interferometry), where the transient change in lattice impedance affects
3451 the propagation velocity of massless wave packets (photons) without disrupting the integrity of
3452 the leptonic solitons.

3453 Hypothesis: Stellar Stability and Local EMC Saturation

3454 In the context of stellar evolution, the EWT framework suggests that the equilibrium of a star
3455 is not solely a result of thermodynamic radiation pressure, but rather a consequence of the
3456 **structural impedance of the vacuum**. As stellar mass accumulates, the central region
3457 approaches a state of extreme vacuum exhaustion, while the surrounding layers may experience a
3458 density surge toward the **EMC Wall threshold** ($N_{\nu,\text{tmp}} \rightarrow N_{\nu,\text{stat}}$). This creates a "mechanical
3459 cushion" of high nodal density that effectively supports the matter against gravitational collapse.
3460 In this scenario, the stellar plasma "floats" on a layer of saturated lattice, where the push-out
3461 mechanism of the core solitons meets the resistance of a localized EMC Wall. This hypothesis
3462 could provide a more robust mechanical explanation for the stability of stars and the anomalous
3463 heating of the solar corona, interpreted here as a region of non-linear lattice relaxation and
3464 high-frequency nodal friction.

3465 Hypothesis: Variable Vacuum Impedance in Stellar Media

3466 It has been hypothesized that stars possess an EMC precursor wall. This local increase in $N_{\nu,\text{tmp}}$
3467 would manifest itself as a measurable change in vacuum impedance, potentially altering the local
3468 speed of light. Unlike general relativity (GR), which attributes such delays to metric curvature,
3469 EWT theory proposes a mechanical slowdown caused by increased BCC lattice stiffness near the
3470 saturation point. It remains an open question how far from the stellar core such a precursor
3471 could exist and what magnitude of deflection it assumes.

3472 Hypothesis: Residual EMC Walls and the Geometry of Weak Interactions

3473 It is hypothesized that every matter soliton is bounded by a **residual or "degraded" EMC**
3474 **wall**, defined by a local density gradient $N_{\nu,\text{stat}} > x > N_{\nu,\text{eff}}$. While the soliton core represents
3475 a zone of vacuum exhaustion, its boundary serves as a high-impedance interface where the push-
3476 out mechanism meets the statutory background of the BCC lattice. This boundary layer can
3477 provides a physical site for weak interactions.

3478 Hypothesis: The EMC Wall as a Dimensional Transformer ($\pi^6 \rightarrow \pi^7$)

3479 It is hypothesized that the residual EMC wall acts as a dimensional interface. Within this
3480 gradient, the interaction scales from the **volumetric bosonic coupling** (π^6) to the **full charge**
3481 **density resonance** (π^7). In this view, the electric charge is not an intrinsic "point-property"
3482 but the highest dimensional manifestation of the soliton's engagement with the BCC lattice.

3483 29.2 The Erasure of "Dark" Placeholders

3484 The success of the structural EWT framework renders several foundational "mysteries" of the
3485 Standard Model obsolete. By grounding physics in the properties of the BCC lattice, we eliminate
3486 the following "dark" placeholders:

- 3487 • **Dark Matter as a Geometric Push-Force:** Rather than invoking undiscovered WIMPs,
3488 EWT identifies galactic rotation anomalies as a consequence of the **massive EMC volume**
3489 **deficit**. The near-total geometric gap at the G -operating point creates a pervasive external
3490 pressure gradient that "pushes" matter toward regions of lower nodal density.
- 3491 • **Dark Energy as Vacuum Static Pressure:** Accelerated expansion is reinterpreted
3492 as the **restorative elastic response** of the BCC lattice. "Dark Energy" is simply the

3493 statutory static pressure of the EMC background acting upon the structural voids created
3494 by matter solitons—the vacuum's "desire" to return to its statutory equilibrium.

3495 • **Virtual Particles and Loop Complexity:** The thousands of Feynman diagrams are
3496 replaced by the **Nodal Stiffness** N . Vacuum fluctuations are revealed to be the deter-
3497 ministic, high-frequency oscillations of the lattice nodes.

3498 • **The Hierarchy Problem:** The discrepancy between gravity and electromagnetism is a
3499 simple ratio: gravity is a statistical effect of **missing nodes** (EMC deficit), while electro-
3500 magnetism is a dynamic effect of **node workload** (energetic load ρ_E).

3501 29.3 Resolution of Historically "Unsolvable" Paradoxes

3502 The mechanical properties of the BCC lattice, combined with the threshold-based dynamics of
3503 the **EMC Wall**, provide deterministic solutions to several long-standing paradoxes that have
3504 remained unresolved within the probabilistic framework of the Standard Model.

3505 The Horizon Problem: Nodal Saturation and Thermalization

3506 The large-scale uniformity of the universe is a consequence of the **Initial Saturation State**.
3507 In the earliest epoch, the vacuum medium existed at the **Max Packing limit** ($N_{\nu,\max} \approx 10^{54}$),
3508 where the BCC lattice is structurally incompressible.

3509 In this state of maximal density, nodal stiffness is absolute, and the velocity of structural equi-
3510 librium is effectively infinite. This "Super-Statutory" phase allowed the entire medium to act
3511 as a single, coupled resonator, achieving perfect thermal uniformity. The subsequent relaxation
3512 of the lattice toward the **Statutory Limit** ($N_{\nu,\text{stat}}$) and eventually the current **Effective Op-**
3513 **erating Point** ($N_{\nu,\text{eff}}$) preserved this initial homogeneity, rendering the speculative "Inflation"
3514 field unnecessary.

3515 The Information Paradox: Nodal Memory

3516 The "loss" of information in a black hole is prevented by the physical nature of the event horizon.
3517 The **EMC Wall** acts as a high-density storage medium. The decomposition of matter at the
3518 boundary is not an erasure, but a **transduction**: the soliton's wave frequency is encoded into the
3519 **Nodal Memory** of the saturated BCC lattice ($N_{\nu,\max}$). Information is preserved as a specific
3520 vibrational state of the wall's nodes.

3521 The GZK Limit: Lattice-Induced Drag

3522 The Greisen-Zatsepin-Kuzmin (GZK) limit, which restricts the energy of cosmic rays, is revealed
3523 to be a **structural speed limit**. As a particle soliton approaches extreme energies, it creates
3524 a "Micro-EMC Wall" (a local compression wave) directly ahead of its path. This results in a
3525 non-linear increase in vacuum resistance, effectively capping the maximum energy a soliton can
3526 maintain while moving through the BCC substrate.

3527 Matter-Antimatter Asymmetry: Asymmetric Phase-Lattice Transition

3528 In the EWT framework, following the geometric interpretation proposed by Yee and Gardi [26],
3529 antimatter is defined as a soliton state characterized by a 180° (π) phase-shift. The observable
3530 dominance of matter is reinterpreted as a result of **Non-linear Phase Conversion** occurring
3531 at the **trans-statutory boundary** ($N > N_{\nu,\text{stat}}$) of EMC Walls.

In this model, the transition through an EMC Wall is not phase-symmetrical. While the wall acts as a resonant processor that can theoretically shift phases in both directions, the current G -operating point of the global BCC lattice creates a bias that favors "locking" solitons into the matter-phase. Consequently, antimatter is a phase that less frequently exits the high-EMC density regions in its original form, being resonantly re-tuned to the dominant background. This suggests that the universe's matter-bias is a dynamic equilibrium maintained by the lattice's tuning, where high-EMC density regions act as filters that preferentially stabilize the primary soliton phase.

29.4 Density Duality: From Local Refraction to Cosmological Redshift

The Energy Density Profile $\rho_E(\mathbf{r})$ as the Origin of Local Refractive Index and EM Wave Slowing

While the **geometric packing density function** $\rho(\mathbf{r})$ describes the non-uniform distribution of the elastic medium and is fundamental to the emergence of the gravitational constant G , a profound and distinct consequence of the Soliton's internal structure is the role of its **localized energy concentration** in the propagation of electromagnetic (EM) waves.

The hypothesis is presented that the local **energy density** $\rho_E(\mathbf{r})$ (which defines the Soliton's mass $E = mc^2$) acts as a local, variable refractive index for EM waves. The speed of light c is modified locally to $v(r)$ based on the energy density profile:

$$v(r) = \frac{c}{n(r)}, \quad n(r) \approx 1 + C_E \cdot \rho_E(r), \quad (265)$$

where C_E is a constant of proportionality related to the electromagnetic field coupling with the Soliton's high energy density.

Compatibility with Density Duality (Clear Separation of Roles). This change establishes clear and distinct physical roles:

- **Refraction:** The **high energy density** (ρ_E) ensures $n(r) > 1$ and correctly predicts the **slowing of EM waves** inside the Soliton/Matter.
- **Gravity:** The **low geometric packing density** ($\rho(\mathbf{r})$) remains solely responsible for the geometric deficit ($N_{\nu, \text{effective}}$), which drives the **push force gravity**.

This framework suggests that the fundamental mechanism for the **slowing of EM waves in matter** (refraction) is the interaction with the combined, non-uniform $\rho_E(\mathbf{r})$ profiles of the atomic constituents. The geometric parameter that governs gravitational weakness ($N_{\nu, \text{effective}}$, which is derived from the **integral of $\rho(\mathbf{r})$**) is thus linked to a core electromagnetic phenomenon (refraction) controlled by a **separate density component** (ρ_E).

Refractive index and Nodal Delay. The slowing of EM waves inside the Soliton is not a function of the number of nodes (geometric density $\rho(r)$), but the "workload" of each node (energy density ρ_E). While the massive deficit of EMC components (the gap between statutory and effective N_{ν}) drives the gravitational push-force, the nodes are subject to extreme oscillatory stress. This increases the **nodal response time**, effectively lowering the propagation velocity $v(r) = c/n(r)$ as a function of $\rho_E(r)$.

Cosmological Implications and the Geometric Origin of Redshift

The geometric derivation of $\mathbf{G}$ and the structural formalization of the Soliton ($\rho(\mathbf{r})$) carry profound consequences for cosmology. The EWT model fundamentally shifts the interpretation

of observed phenomena away from dynamic spacetime expansion toward **wave propagation dynamics within the Elastic Medium Constituent (EMC)**.

The concept of gravity as a **push force** resulting from the volume deficit (N_ν) suggests an alternative framework for the accelerating expansion of the universe. This expansion, typically attributed to Dark Energy, can be re-interpreted as the **external pressure of the EMC background** acting on regions of matter deficit.

Furthermore, the dual role of the Soliton's energy density ($\rho_E(\mathbf{r})$) in governing both mass and local EM wave slowing (refraction) opens the possibility that **cosmological redshift (z)** may be partially or entirely a consequence of **"tired light" effects**—minimal energy loss of EM waves due to interaction with the bulk properties or temporal evolution of the EMC background over vast distances.

Synthesis of Geometric and Temporal Properties: While ϵ_M defines the fundamental geometric permeability (stiffness) of the vacuum in solitons, the **Nodal Delay** represents the temporal manifestation of this stiffness when under energetic load (ρ_E). This unifies the EMC deficit (ρ) with the dynamic slowing of light (refraction) and the cumulative energy loss over cosmological distances (redshift). This perspective effectively replaces the notion of "empty space" with a dynamic, elastic medium whose response time is intrinsically coupled to its energy density.

Although this model focuses on leptonic structure, the unification of geometric parameters suggests that the Nodal Delay mechanism may have a cumulative component at the cosmological scale. This allows for the hypothesis of a non-Doppler origin for a portion of the redshift, representing a promising direction for future research into the nature of the EMC background.

30 Outlook and Computational Tools

The formal establishment of the Magneto-Geometric Hypotheses (Section 18) provides clear, testable predictions for the effective gravitational constant G (Section 26.3). Beyond direct experimental verification, the computational modelling of the Soliton geometry and its dynamic coupling to the Elastic Medium Constituent (EMC) is being pursued on the open-source platform **OpenWave** (<https://github.com/openwave-labs/openwave>).

Progress in the Scalar M3 Model

In the scalar Wolff-LaFreniere model (M3), a spherical-phyllotaxis (golden-angle, $\sim 137.5^\circ$) configuration of wave centres was implemented and tested as a proof-of-concept. The results provided the first in-platform demonstration that a spherical, minimally interfering geometry can create K-selectivity: only $K_{WC} = 10$ formed a stable standing wave, while $K_{WC} = 9$ and $K_{WC} = 11$ decayed systematically. This validated the underlying principle that phyllotactic packing suppresses destructive interference, a principle that will be essential for the high-multiplicity recursive shells of the muon and tau. For the electron core itself, the native EWT topology (1-3-6 tetrahedron) remains the physically required configuration, as it alone guarantees the correct multipole structure for far-field forces. The golden-angle module, together with the logged simulation data, is available in the `m3_wolff_lafreniere` directory of the OpenWave repository.

Diagnostics in the Vector M4 Model and Implementation Plan

The same golden-angle arrangement does *not* stabilise the electron core in the vector model M4 (neither with nor without an initial spin), indicating that the missing ingredient in M4 is not geometric but nonlinear. The immediate goal is to stabilise the native 1-3-6 electron topology by

introducing the nonlinear self-trapping term derived in Section 25. Once the electron core is stable with its proper geometry, the golden-angle phyllotaxis will be applied to the high-multiplicity recursive shells of the muon and tau, where it is expected to be indispensable. Consequently, the following implementation roadmap has been adopted:

1. **Nonlinear term $\mathcal{F}(\Psi)$:** A cubic self-trapping term $\gamma \Psi^3$ will be added to the M4 wave kernel, with the coupling coefficient $\gamma = 1/\epsilon_M = N_{\text{final}}\pi^3 \approx 2.41 \times 10^4$ derived from the BCC geometry.
2. **Degraded EMC Wall:** A radial density gradient will be introduced to suppress shear instabilities and to provide a natural boundary where the phyllotaxis can later be introduced for the outer shells.
3. **Validation:** Once the nonlinearity is in place, the native EWT topology (1-3-6 tetrahedron) will be tested for K-selectivity.
4. **Shell application:** After the core is stabilised, the golden-angle arrangement will be deployed for the muon and tau shells.

Connection to the Nonlinear Stability Equation

A sketch of the required nonlinear wave equation was proposed in part by the OpenWave team during the collaborative development of the M3 and M4 engines. Building on that foundation, a formal nonlinear wave equation that guarantees the electron's stability is derived in Section 25, where the geometric constants ϵ_M and $\gamma = 1/\epsilon_M$ determine the self-trapping term. The full implementation of this equation in the OpenWave M4 kernel is the immediate next step.

This computational framework serves as a **theoretical complement to physical experiments**, allowing rapid iteration and falsification of the Soliton's micro-geometric constraints within the EWT Model.

31 Conclusion: The Geometric Paradigm, Correlated Lepton Anomalies, and Emergent Gravity

The formalization within the Energy Wave Theory establishes the **Minimal Wave Center** — the Neutrino Soliton — as a fundamental geometric structure. The central finding of this work is the derivation of the gravitational constant G as a geometric identity, together with the prediction of the lepton anomalous magnetic moments from the same underlying BCC lattice parameters. This necessitates a critical re-evaluation of the Soliton's geometric parameters.

This achievement realizes a principle that is increasingly recognized as a necessity for resolving the deepest tensions in modern physics. As highlighted by the CosmoVerse Network [48], addressing the observational crises in cosmology may ultimately require abandoning the notion of rigid, immutable fundamental constants in favour of dynamic, relational parameters. The Enhanced EWT model does not merely accommodate this principle; it provides its precise geometric mechanism. The observed “constants” — the gravitational coupling G , the electromagnetic coupling α , and the electron anomalous magnetic moment a_e — are not independent inputs but are scale-dependent projections of a single invariant: the structural impedance of the BCC vacuum lattice.

For the muon and tau generations, the model presently derives the anomalous magnetic moments as genuine predictions of the BCC lattice geometry, while the reference scale for comparison is obtained from the orbital mass relations. This constitutes a high-precision internal

3657 consistency test between the geometric and mass-generation sectors; the simultaneous, fully in-
 3658 dependent derivation of both mass and AMM for all generations remains an open objective.

3659 **Constants are not constant; only the relations between them are.**
 This relation, dictated by the fixed topology of the vacuum substrate, is the true
 fundamental law from which all measurable quantities emerge.

3660 31.1 Geometric Unification in a Nutshell: The ϵ_M Flowchart

3661 The Enhanced EWT Model operates on the principle of **Structural Monism**: all fundamental
 3662 coupling constants are derived from the same vacuum stiffness deficit. Rather than treating grav-
 3663 itation, electromagnetism, and lepton anomalies as independent phenomena, they are expressed
 3664 as direct geometric functions of the primary modulator ϵ_M .

$$\epsilon_M \Rightarrow \begin{cases} \mathbf{G} = \frac{\mathbf{G}_{\text{Base}}}{A_\pi} \cdot \left(\frac{\epsilon_M \cdot \pi^3}{A_\pi} \right)^3 \cdot \frac{1}{K_{WC} \cdot \sqrt{N_{\nu, \text{eff}}}} & \text{(Gravitational Scale)} \\ \alpha_{\text{geom}}^{-1} = A_\pi - \epsilon_M & \text{(Electromagnetic Scaling)} \\ a_e = \frac{\alpha_{\text{geom}}}{2\pi} \cdot (1 - \epsilon_M \cdot \pi^3) & \text{(Nodal Magnetic Deficit)} \\ C_{\text{local}} = \frac{\epsilon_M}{2\sqrt{2}} & \text{(Geometric Seed for Mixing Hierarchy)} \\ R_l = \text{recursive}(\epsilon_M, L_{\text{Fib}}) & \text{(Leptonic Shell Resonance)} \end{cases} \quad (266)$$

3665 In this zero-calibration formulation,

$$\epsilon_M = \frac{1}{N_{\text{geom}} \pi^3} = \frac{1}{8\pi^7(1 - \zeta)}, \quad (267)$$

3666 where $N_{\text{geom}} = 8\pi^4(1 - \zeta)$ is the effective geometric stiffness and ζ is the BCC packing impedance.

3667 31.2 The Unitary Geometric Path

3668 As illustrated in the schematic, ϵ_M acts as the **Universal Gear Ratio** of the vacuum lattice.
 3669 A single modulator simultaneously determines:

- 3670 • the gravitational coupling G ,
- 3671 • the electromagnetic coupling α_{geom} ,
- 3672 • the damping of the magnetic precession across generations, yielding a_e, a_μ, a_τ .

3673 This flowchart confirms that the Enhanced EWT Model is not a collection of independent
 3674 equations, but a **Unitary Geometric Circuit**. Any change in the value of ϵ_M would simulta-
 3675 neously shift all fundamental constants derived from it.

3676 31.3 The Geometry of the Constituent: From Cubic Grids to Spherical 3677 Units

3678 While the initial formalization of the EWT lattice utilizes the Body-Centered Cubic framework
 3679 for its nodal efficiency, the physical reality of the **Elastic Medium Constituent** is fundamen-

3680 tally spherical. This transition from a “cubic cell” to a “spherical grain” is a natural consequence
 3681 of the model’s convergence toward a minimum energy state and isotropic wave propagation.

3682 **Packing Fraction and the Vacuum Void**

3683 In a BCC lattice composed of hard spherical constituents, the maximum packing factor is

$$\eta_{\text{BCC}} = \frac{\sqrt{3}\pi}{8} \approx 0.680. \quad (268)$$

3684 This inherent geometric property implies that:

- 3685 • The vacuum is not a continuous “solid block” but a structured, granular medium with a
 3686 defined interstitial capacity.
- 3687 • The volume deficit N_ν identified in the gravitational derivation represents the displacement
 3688 of these spherical units.
- 3689 • The spherical nature of the EMC ensures isotropic nodal stiffness.

3690 **The Transition of Density Levels**

3691 The granular nature of the EMC directly dictates the three levels of vacuum density identified
 3692 in the EWT hierarchy:

- 3693 • **Maximum packing:** $N_{\nu,\text{max}} \approx 5.30 \times 10^{54}$.
- 3694 • **Statutory background:** $N_{\nu,\text{stat}} \approx 3.30 \times 10^{52}$.
- 3695 • **Effective gravitational density:** $N_{\nu,\text{eff}} \approx 6.25 \times 10^{48}$.

3696 **31.4 Pillar 1: The N_ν Deficit and Soliton Stability**

3697 A cornerstone of the EWT model is the derivation of the gravitational constant G from the
 3698 microscopic geometry of the neutrino soliton and the electron soliton. In the zero-calibration
 3699 formulation, the transition from the statutory background to the effective density is governed by
 3700 the geometric dilution factor X_{eff} :

$$N_{\nu,\text{eff}} = \frac{N_{\nu,\text{stat}}}{X_{\text{eff}}}, \quad (269)$$

3701 where X_{eff} is fixed by A_π , $K_{WC} = 10$, and the ideal BCC projection factor $L_p^{\text{geom}} = 2/\sqrt{3}$.
 3702 The resulting density hierarchy is:

$$\rho_{\text{eff}} < \rho_{\text{stat}} < \rho_{\text{max}}. \quad (270)$$

3703 Macroscopic gravitational attraction is the response of the external, higher-density vacuum
 3704 medium ($N_{\nu,\text{stat}}$) attempting to restore equilibrium by pressing toward the lower-density soliton
 3705 core.

31.5 Pillar 2: The Dynamic Scaling Operator Ω_{Push}

The dynamic push-out operator,

$$\Omega_{\text{Push}} \equiv G_{\text{Base}} \cdot A_{\pi}^{-1} \cdot (N_{\text{geom}} A_{\pi})^{-3}, \quad (271)$$

is the central mechanism in the G equation.

The final gravitational identity is:

$$G \equiv \underbrace{G_{\text{Base}} \cdot \frac{1}{A_{\pi}} \cdot \left(\frac{1}{N_{\text{geom}} A_{\pi}} \right)^3}_{\Omega_{\text{Push}}} \cdot \underbrace{\frac{1}{K_{WC} \sqrt{N_{\nu, \text{eff}}}}}_{T_{\text{Surface}}}. \quad (272)$$

The factor $\epsilon_M = (N_{\text{geom}} \pi^3)^{-1}$ is the master key of this operator. It dictates both the gravitational scale and the lepton anomalous magnetic moments.

31.6 Pillar 3: The Recursive Paradigm — Hierarchical Nodal Integration

The transition from the electron core to the muon and tau generations is viewed as a hierarchical integration of nested nodal shells, anchored by the Fibonacci invariants

$$L_{\mu} = 5, \quad L_{\tau} = 34. \quad (273)$$

The full AMM predictions are:

$$a_e^{\text{EWT}} = 1.159916 \times 10^{-3}, \quad a_{\mu}^{\text{EWT}} = 1.166213 \times 10^{-3}, \quad a_{\tau}^{\text{EWT}} = 1.176838 \times 10^{-3}. \quad (274)$$

The recursive architecture replaces the perturbative loop paradigm of the Standard Model with a single unified geometric identity.

31.7 The Soliton Radius and Neutrino Interaction Cross-Section

A common concern regarding geometric particle models is the apparent tension between a comparatively large geometric radius and the extremely small neutrino interaction cross-sections. This model resolves the issue by distinguishing between:

- the **geometric soliton radius** r_{geom} ,
- the **interaction cross-section** σ_{ν} .

The weakness of the neutrino interaction reflects the fact that only a minute fraction of the soliton's internal energy is available at its boundary for weak interactions. This dilution is quantified by the effective volume deficit:

$$\sigma_{\nu} \propto \frac{1}{N_{\nu, \text{eff}}}, \quad (275)$$

so that even a soliton with a substantial geometric radius can possess an exceedingly small interaction cross-section.

31.8 Paradigm Shift: The Planck Length as a Physical Boundary

In EWT, the Planck length is assigned a definitive physical and geometric role: it is the approximate magnitude of the base quantum distance λ_l , which defines the physical separation of the Elastic Medium Constituents.

In this framework, λ_l is not an external input; it is the self-consistent geometric Planck length derived in Section 11.6. All physics operates consistently at and above this scale, allowing the direct geometric calculation of macroscopic constants like G and α .

31.9 The ϵ_M Factor: Universal Geometric Modulator

The core tenet of EWT is that mass, charge, and spin are manifestations of the Soliton's wave geometry. The unified role of the magnetic deficit factor is governed by a fundamental geometric duality: while the Soliton's internal energy storage follows $E \propto r^5$, the volume of the displaced elastic medium follows $V \propto r^3$.

The factor ϵ_M acts as the universal modulator that reconciles these two scaling laws. It performs three distinct functions:

1. **Gravity:** It scales the volumetric deficit to the observed gravitational constant G .
2. **Electromagnetism:** It bridges the gap between α_{geom} and the soliton core.
3. **Spin/AMM:** It defines the recursive nodal integration of lepton anomalies.

The necessity of using the same factor ϵ_M across all these domains is the definitive proof of geometric unification.

31.10 Geometric Self-Consistency of the Planck–Gravity–Metric Triangle

A central result of the present work is the recognition that G_{geom} , $\hbar_{\text{geom}}$, λ_l , and c do not form a hierarchy of independent constants. They are linked by the closed geometric condition

$$\lambda_l = \sqrt{\frac{\hbar_{\text{geom}} G_{\text{geom}}}{c^3}}, \quad (276)$$

where

$$\hbar_{\text{geom}} = \frac{m_e c r_e}{\alpha_{\text{geom}}}, \quad G_{\text{geom}} = \frac{G_{\text{Base}}}{A_\pi} \left(\frac{1}{N_{\text{geom}} A_\pi} \right)^3 \frac{1}{K_{WC} \sqrt{N_{\nu, \text{eff}}}}. \quad (277)$$

The effective density $N_{\nu, \text{eff}}$ itself depends on λ_l through the statutory background,

$$N_{\nu, \text{eff}} = \frac{1}{X_{\text{eff}}} \left(\frac{r_\nu}{2e\lambda_l} \right)^3. \quad (278)$$

Thus Eq. (276) is not a definition imported from outside the model. It is an internal algebraic constraint that simultaneously fixes the gravitational scale, the quantum scale, and the fundamental lattice spacing from the same BCC geometry.

In natural units ($c = 1$), this constraint reduces to the purely geometric statement

$$\lambda_l^2 = \hbar_{\text{geom}} G_{\text{geom}}, \quad (279)$$

3759 showing that the square of the fundamental EMC length is the product of the geometric quantum
 3760 of action and the geometric gravitational coupling.

3761 This closure is a direct consequence of the BCC lattice structure. It is not an import from the
 3762 Standard Model: the Planck scale, the reduced Planck constant, and the gravitational constant
 3763 are reconstructed here as derived quantities, not assumed inputs.

3764 **31.11 Final Synthesis**

3765 The Enhanced EWT model demonstrates that the fundamental constants are integrated reso-
 3766 nances of a single BCC substrate. The geometric vacuum stiffness N_{geom} , the magnetic deficit
 3767 ϵ_M , and the geometric core A_π together determine the fine-structure constant, the gravitational
 3768 constant, the lepton anomalous magnetic moments, and the principal atomic scales.

3769 The model is not claimed to be absolutely parameter-free. It currently uses four experimental
 3770 anchors:

$$r_e, \quad m_e, \quad c, \quad e. \quad (280)$$

3771 These anchors are fixed by measurement and by the metric convention. They are not adjustable
 3772 calibration parameters: if any one of them were changed independently while keeping the BCC
 3773 geometry fixed, the entire set of geometric predictions would break. The model therefore contains
 3774 no free calibration constants; it is zero-free-parameter in the operational sense.

3775 Of these anchors, c is a metric conversion factor rather than a dynamical parameter. The
 3776 electron mass m_e may also be derivable in the future through the $E \propto r^5$ scaling law and the
 3777 $K_{WC} = 10$ stability condition. The true irreducible anchors would then reduce to r_e and e .
 3778 electromagnetism, and the lepton hierarchy.

3779

A Appendix

3780

A.1 List of Model Parameters and Physical Constants

3781

3782

3783

3784

The following table lists the fundamental physical constants and key dimensionless geometric factors used in the Geometric Soliton Model. The numerical consistency of the model relies on the exact values shown below, combining CODATA 2022 constants with the derived EWT geometric parameters.

Table 22: **EWT Parameters and CODATA 2022 Values Used in the Geometric Derivations**

Parameter	Symbol	Numerical Value	Source / Justification
I. Fundamental Constants (CODATA 2022)			
Speed of Light	c	299792458	Defined constant (m/s).
Electron Mass	m_e	$9.1093837015 \times 10^{-31}$	CODATA 2022 reference (kg).
Classical Radius	r_e	$2.8179403262 \times 10^{-15}$	CODATA 2022 reference (m).
II. Hierarchical Volume Deficit Parameters			
Absolute Maximum	$N_{\nu,\max}$	5.3004×10^{54}	Theoretical EMC packing limit.
Statutory Background	$N_{\nu,\text{stat}}$	3.2986×10^{52}	Eulerian vacuum capacity.
Effective Deficit	$N_{\nu,\text{eff}}$	6.2472×10^{48}	Integrated Soliton density.
Geometric Bridge Operator	X_{eff}	5275.7683	Transforms $N_{\nu,\text{stat}}$ into $N_{\nu,\text{eff}}$.
III. Geometric Operators			
Geometric Base	A_π	137.0363037759	$4\pi^3 + \pi^2 + \pi$. Static foundation.
Effective Stiffness	N_{geom}	778.8025179	$8\pi^4(1 - \zeta)$.
Lattice Impedance	ζ	0.0006033964	From BCC packing fraction.
Magnetic Deficit	ϵ_M	$4.141169769 \times 10^{-5}$	$1/(N_{\text{geom}}\pi^3)$.
Lattice Projection	L_p^{geom}	1.1547005384	$2/\sqrt{3}$, ideal BCC projection.
Unified Coupling	C_{unif}	1.1020116168	Using L_p^{geom} .

A.2 Computational Scripts (Python)

All numerical results in this work are produced by a set of Python scripts that implement the zero-calibration geometric model described in the main text. The scripts are self-contained and use only standard scientific libraries. The source code and console outputs are listed below. Persistent identifiers for each script will be added in the final published version.

A.2.1 Emergence Engine

`ewt_emergence_engine.py` is the core geometric engine. It derives the fine-structure constant, magnetic deficit, neutrino radius, Planck length, gravitational constant, lepton anomalous magnetic moments, and atomic scales from the BCC lattice geometry. The script also performs the rigidity tests described in Section 28.

```
# From manuscript version: 5.0.1
# =====
# SECTION 1: GLOBAL INPUTS & CONSTANTS
# -----
# This section contains:
#   - mathematical constants
#   - experimental reference values (CODATA 2022 / PDG 2022)
#   - pure BCC lattice geometry (independent of the EWT model)
#
# Every quantity is explicitly labelled:
#   INPUT      : experimental value used as an anchor or metric input
#   MATHEMATICAL : pure mathematical constant
#   BCC_GEOMETRY : crystallographic fact about the BCC lattice
#
# The model itself will not define any additional hidden numerical parameters
# in this section. All later parameters must be either DERIVED, POSTULATED,
# or explicitly marked as CALIBRATION (to be eliminated in future versions).
# =====

import math

# -----
# 1.1 MATHEMATICAL CONSTANTS
# -----

PI          = math.pi          # circle ratio
EULER       = math.e            # Euler's number
SQRT2       = math.sqrt(2.0)    # diagonal of unit square
SQRT3       = math.sqrt(3.0)    # diagonal of unit cube face

# -----
# 1.2 CODATA 2022 / PDG 2022 EXPERIMENTAL INPUTS
# -----

# Metric and electromagnetic scales
CO          = 299792458.0        # speed of light [m/s] (exact SI)
M_E         = 9.1093837015e-31   # electron mass [kg]
R_E         = 2.8179403262e-15   # classical electron radius [m]
ALPHA_INV_CODATA = 137.035999084 # inverse fine-structure constant
A_E_CODATA  = 0.0011596521816    # electron AMM (g-2)/2

# Gravitational scale
G_CODATA    = 6.674305e-11       # gravitational constant [m^3 kg^-1 s^-2]

# Electroweak and flavour scales
M_Z_CODATA  = 91.1876           # Z boson mass [GeV]
M_W_PDG     = 80.377            # W boson mass, PDG 2022 average [GeV]
M_W_CDFII   = 80.4335          # W boson mass, CDF II 2022 [GeV]
M_H_CODATA  = 125.25            # Higgs mass [GeV]
SIN2_THETA_W = 0.23122         # Weinberg angle sin^2(theta_W)
SIN_THETA_C_PDG = 0.2243       # Cabibbo angle sin(theta_C)
R_INF_CODATA = 10973731.568157  # Rydberg constant [m^-1]
AO_CODATA    = 5.29177210903e-11 # Bohr radius [m]
LAMBDA_C_CODATA = 2.42631023867e-12 # electron Compton wavelength [m]
```

```

3851 # Quark masses (MS-bar, PDG 2022) used in flavour mixing test
3852 M_D_PDG = 0.004692 # d-quark mass [GeV]
3853 M_S_PDG = 0.094954 # s-quark mass [GeV]
3854
3855 A_MU_EXP = 1.1659206100e-3 # muon AMM from Fermilab/Brookhaven
3856 A_TAU_EXP = 1.177210e-3 # # tau AMM: Standard Model prediction, not
3857     direct measurement
3858
3859 # Fundamental length scale (Planck charge interpreted as geometric amplitude)
3860 # This is currently treated as an INPUT, but will later be reclassified
3861 # as DERIVED once the geometric origin of q_P is explicitly shown.
3862 Q_P_INPUT = 1.87554603778e-18 # Planck charge as length [m]
3863 # Fundamental EMC spacing (Planck length) [m]
3864 LAMBDA_L = 1.6162e-35
3865 # Elementary charge (CODATA 2022) -- empirical anchor of the amplitude scale
3866 E_CHARGE_CODATA = 1.602176634e-19
3867
3868 # -----
3869 # 1.3 PURE BCC LATTICE GEOMETRY
3870 # -----
3871 # These are purely mathematical / crystallographic facts about the BCC lattice.
3872 # They are independent of the Enhanced EWT model and contain no physics inputs.
3873 # -----
3874
3875 BCC_NEIGHBOURS = 8 # nearest neighbours in BCC
3876 BCC_PACKING_FRACTION = (SQRT3 * PI) / 8.0 # ~0.6801748
3877 BCC_IDEAL_PROJECTION_LP = 2.0 / SQRT3 # inverse nearest-neighbour distance
3878
3879 # -----
3880 # 1.4 STATUS LABELS
3881 # -----
3882 # These labels will be used in all reporting functions to make the origin
3883 # of each quantity transparent.
3884
3885 STATUS = {
3886     "INPUT": "Experimental_input(CODATA/PDG)",
3887     "MATHEMATICAL": "Pure_mathematical_constant",
3888     "BCC_GEOMETRY": "Pure_BCC_lattice_geometry",
3889     "POSTULATE": "Geometric_postulate_of_the_model",
3890     "DERIVED": "Derived_from_model_postulates/constraints",
3891     "CALIBRATION": "Calibrated_parameter(to_be_eliminated)",
3892 }
3893
3894 # =====
3895 # SECTION 2: GEOMETRIC POSTULATES & ALPHA CORE EMERGENCE
3896 # -----
3897 # The alpha core emerges from three geometric postulates:
3898 # P1: charge = amplitude (x)
3899 # P2: emission surface = sphere + cone
3900 # P3: natural length ratios r = x, l = pi*x
3901 #
3902 # These postulates lead directly to the geometric ratio:
3903 # alpha_core^{-1} = S_total / x^2 = 4*pi^3 + pi^2 + pi
3904 # =====
3905
3906 def compute_emission_surface(x: float) -> dict:
3907     """
3908     Compute the total emission surface and the resulting alpha core
3909     from the geometric postulates.
3910
3911     Parameters
3912     -----
3913     x : float
3914         Wave amplitude (charge). The final ratio is independent of x.
3915
3916     Returns
3917     -----
3918     dict with:
3919         r : base radius of the cone (equals x)
3920         l : propagation distance (equals pi*x)
3921         S_sphere : surface area of the spherical part
3922         S_cone : total surface area of the conical part
3923         S_total : total emission surface

```

```

3924         A_pi      : alpha core inverse = S_total / x^2
3925     """
3926     r = x
3927     l = PI * x
3928
3929     S_sphere = 4.0 * PI * l**2
3930     S_cone = PI * r * l + PI * r**2
3931
3932     S_total = S_sphere + S_cone
3933     A_pi = S_total / x**2
3934
3935     return {
3936         "r": r,
3937         "l": l,
3938         "S_sphere": S_sphere,
3939         "S_cone": S_cone,
3940         "S_total": S_total,
3941         "A_pi": A_pi,
3942     }
3943
3944 def compute_alpha_core() -> float:
3945     """
3946     Compute the purely geometric core of the fine-structure constant.
3947     It is obtained from the emission surface ratio for a unit amplitude.
3948
3949     Returns
3950     -----
3951     float
3952         A_pi = 4*pi^3 + pi^2 + pi
3953     """
3954     return compute_emission_surface(1.0)["A_pi"]
3955
3956 # =====
3957 # SECTION 2B: GEOMETRIC LADDER EMERGENCE
3958 # -----
3959 # The geometric ladder is not an input. It is derived from the active
3960 # degrees of freedom of a soliton embedded in the BCC vacuum lattice.
3961 #
3962 # The central modulator eps_M is treated as an INPUT to this module.
3963 # In the current zero-calibration version it is obtained from pure BCC geometry:
3964 # - eps_M derived from packing fraction eta and lattice impedance zeta
3965 # - historical calibrated variant (N_final) is no longer used
3966 #
3967 # Later sections will compute eps_M and pass it here.
3968 # =====
3969
3970 def build_geometric_ladder(eps_M: float, source_label: str = "") -> dict:
3971     """
3972     Build the geometric ladder from the magnetic deficit eps_M.
3973
3974     Parameters
3975     -----
3976     eps_M : float
3977         Magnetic deficit parameter.
3978     source_label : str
3979         Optional label describing the source of eps_M.
3980
3981     Returns
3982     -----
3983     dict with:
3984         eps_M      : magnetic deficit (input)
3985         source     : source label
3986         C_local    : local coupling
3987         C_gap      : volumetric operator (pi^6)
3988         C_fermion  : surface operator (pi^5)
3989         ladder     : dictionary {3: pi^3, ..., 7: pi^7}
3990     """
3991     C_local = eps_M / (2.0 * SQRT2)
3992
3993     return {
3994         "eps_M": eps_M,
3995         "source": source_label,
3996         "C_local": C_local,

```

```

3997         "C_gap": 1.0 + PI**6 * C_local,
3998         "C_fermion": (1.0 + PI**5 * C_local) ** 2,
3999         "ladder": {n: PI**n for n in range(3, 8)},
4000     }
4001
4002
4003 def report_ladder(lad: dict) -> None:
4004     """Print the contents of a ladder dictionary in a readable form."""
4005     print(f"Source: {lad.get('source', 'unknown')}")
4006     print(f"eps_M: {lad['eps_M']:.15e}")
4007     print(f"C_local: {lad['C_local']:.15e}")
4008     print(f"C_gap: {lad['C_gap']:.15f}")
4009     print(f"C_fermion: {lad['C_fermion']:.15f}")
4010     print("Ladder:")
4011     for n in sorted(lad["ladder"]):
4012         print(f"pi^{n} = {lad['ladder'][n]:.15f}")
4013
4014 # =====
4015 # SECTION 3: BCC-DERIVED LATTICE IMPEDANCE
4016 # -----
4017 # Functions that derive eps_M from pure BCC geometry using the packing
4018 # fraction and lattice impedance.
4019 # =====
4020
4021 def lattice_impedance(N: float, eta: float = BCC_PACKING_FRACTION) -> float:
4022     """
4023     Compute the lattice impedance zeta for a given geometric stiffness N.
4024
4025     Formula:
4026         zeta = (1 - eta) / (eta * N)
4027
4028     Parameters
4029     -----
4030     N : float
4031         Geometric stiffness modulus (e.g. 8*pi^4, 8*pi^7).
4032     eta : float
4033         Packing fraction of the BCC lattice.
4034
4035     Returns
4036     -----
4037     float
4038         Lattice impedance.
4039     """
4040     return (1.0 - eta) / (eta * N)
4041
4042
4043 def derive_eps_M_from_BCC(N_input: float = 8.0 * PI**4) -> dict:
4044     """
4045     Derive eps_M from BCC geometry using the lattice impedance.
4046
4047     Parameters
4048     -----
4049     N_input : float
4050         Ideal geometric stiffness to use as base.
4051         Default is 8*pi^4 (BCC saturation stiffness).
4052
4053     Returns
4054     -----
4055     dict with:
4056         eta : BCC packing fraction
4057         N_ideal : input ideal stiffness
4058         zeta : lattice impedance
4059         N_geom : effective stiffness after impedance correction
4060         eps_M : magnetic deficit derived from BCC geometry
4061     """
4062     eta = BCC_PACKING_FRACTION
4063     zeta = lattice_impedance(N_input, eta)
4064     N_geom = N_input * (1.0 - zeta)
4065     eps_M = 1.0 / (N_geom * PI**3)
4066
4067     return {
4068         "eta": eta,
4069         "N_ideal": N_input,

```

```

4070         "zeta": zeta,
4071         "N_geom": N_geom,
4072         "eps_M": eps_M,
4073     }
4074
4075 # =====
4076 # SECTION 4: TEST FUNCTIONS (EXPERIMENTAL VALIDATION)
4077 # -----
4078 # These functions compare model predictions with experimental data.
4079 # They do NOT influence the model itself.
4080 # =====
4081
4082 def weinberg_sector(C_gap: float, M_Z: float, sin2_W: float) -> dict:
4083     """
4084     Compute the W boson mass from the Weinberg angle using the pi^6 operator.
4085
4086     Formula:
4087         M_W = M_Z * sqrt( (1 - sin^2(theta_W)) * C_gap )
4088
4089     Parameters
4090     -----
4091     C_gap : float
4092         Volumetric lattice operator from the pi^6 rung.
4093     M_Z : float
4094         Experimental Z boson mass [GeV].
4095     sin2_W : float
4096         Experimental Weinberg angle sin^2(theta_W).
4097
4098     Returns
4099     -----
4100     dict with:
4101         M_W_pred : predicted W mass [GeV]
4102         rel_err_PDG : relative error to PDG average
4103         rel_err_CDFII : relative error to CDF II measurement
4104     """
4105     M_W_pred = M_Z * math.sqrt((1.0 - sin2_W) * C_gap)
4106
4107     rel_err_PDG = abs(M_W_pred - M_W_PDG) / M_W_PDG * 100.0
4108     rel_err_CDFII = abs(M_W_pred - M_W_CDFII) / M_W_CDFII * 100.0
4109
4110     return {
4111         "M_W_pred": M_W_pred,
4112         "rel_err_PDG": rel_err_PDG,
4113         "rel_err_CDFII": rel_err_CDFII,
4114     }
4115
4116
4117 def cabibbo_sector(C_fermion: float, M_d: float, M_s: float) -> dict:
4118     """
4119     Compute the Cabibbo angle from quark masses using the pi^5 operator.
4120
4121     Formula:
4122         sin(theta_C) = sqrt( M_d / M_s ) * C_fermion
4123
4124     Parameters
4125     -----
4126     C_fermion : float
4127         Surface lattice operator from the pi^5 rung.
4128     M_d : float
4129         Down quark mass [GeV].
4130     M_s : float
4131         Strange quark mass [GeV].
4132
4133     Returns
4134     -----
4135     dict with:
4136         sin_C_pred : predicted sin(theta_C)
4137         rel_err : relative error to PDG target
4138     """
4139     sin_C_pred = math.sqrt(M_d / M_s) * C_fermion
4140     rel_err = abs(sin_C_pred - SIN_THETA_C_PDG) / SIN_THETA_C_PDG * 100.0
4141
4142     return {"sin_C_pred": sin_C_pred, "rel_err": rel_err}

```



```

4216
4217     Returns
4218     -----
4219     float
4220         Geometric inverse fine-structure constant.
4221     """
4222     A_pi = compute_alpha_core()
4223     return A_pi - eps_M
4224
4225 def test_alpha_sector(alpha_inv_pred: float) -> dict:
4226     """
4227     Compare the predicted inverse fine-structure constant with CODATA 2022.
4228
4229     Parameters
4230     -----
4231     alpha_inv_pred : float
4232         Predicted inverse fine-structure constant.
4233
4234     Returns
4235     -----
4236     dict with:
4237         alpha_inv_pred : predicted value
4238         abs_err        : absolute error vs CODATA
4239         rel_err        : relative error in percent
4240     """
4241     abs_err = abs(alpha_inv_pred - ALPHA_INV_CODATA)
4242     rel_err = abs_err / ALPHA_INV_CODATA * 100.0
4243
4244     return {
4245         "alpha_inv_pred": alpha_inv_pred,
4246         "abs_err": abs_err,
4247         "rel_err": rel_err,
4248     }
4249
4250 # =====
4251 # SECTION 6: NEUTRINO & ELECTRON ANCHORS
4252 # -----
4253 # This section derives hbar, Planck charge, uncorrected wavelength, and
4254 # the neutrino radius from geometric alpha and electron anchors.
4255 # =====
4256
4257 def derive_planck_charge_from_e(alpha_geom: float, e_charge: float) -> float:
4258     """
4259     Derive Planck charge (fundamental geometric amplitude) from
4260     geometric alpha and the elementary charge amplitude e.
4261
4262     Formula:
4263         q_P = e / sqrt(alpha_geom)
4264
4265     Note: This is NOT a tautology because alpha_geom is derived
4266     from pure BCC geometry (A_pi - eps_M), not from e or q_P.
4267     """
4268     return e_charge / math.sqrt(alpha_geom)
4269
4270
4271 def derive_hbar(alpha_geom: float, r_e: float, m_e: float, c0: float) -> float:
4272     """
4273     Derive the reduced Planck constant from geometric alpha and electron data.
4274
4275     Formula:
4276         hbar = m_e * c * r_e / alpha_geom
4277
4278     Parameters
4279     -----
4280     alpha_geom : float
4281         Geometric fine-structure constant (dimensionless).
4282     r_e : float
4283         Classical electron radius [m].
4284     m_e : float
4285         Electron mass [kg].
4286     c0 : float
4287         Speed of light [m/s].
4288

```

```

4289     Returns
4290     -----
4291     float
4292         Reduced Planck constant [kg m^2 / s].
4293     """
4294     return (m_e * c0 * r_e) / alpha_geom
4295
4296
4297 def derive_lambda_uncorr(q_P: float) -> float:
4298     """
4299     Compute the uncorrected neutrino wavelength from Planck charge.
4300
4301     Formula:
4302         lambda_uncorr = 2 * q_P * e^2
4303
4304     where e is Euler's number.
4305
4306     Parameters
4307     -----
4308     q_P : float
4309         Planck charge (in EWT interpreted as length).
4310
4311     Returns
4312     -----
4313     float
4314         Uncorrected wavelength [m].
4315     """
4316     return 2.0 * q_P * EULER**2
4317
4318
4319 def derive_neutrino_radius(alpha_geom: float, q_P: float) -> dict:
4320     """
4321     Derive the neutrino radius r_nu from the geometric fixed point of g_v.
4322
4323     The derivation uses the self-consistent equation:
4324         S_tot = alpha_geom_inv/(8+pi) + e + (1-g_v)*(sqrt2-1)
4325         S_tot = 2 e^2 / g_v
4326     which leads to a quadratic equation for g_v.
4327
4328     Parameters
4329     -----
4330     alpha_geom : float
4331         Geometric fine-structure constant.
4332     q_P : float
4333         Planck charge / geometric amplitude [m].
4334
4335     Returns
4336     -----
4337     dict with:
4338         alpha_inv_geom : geometric inverse alpha
4339         S_proj          : static lattice projection
4340         S_exp           : dynamic wave expansion (e)
4341         delta_imp       : lattice impedance term
4342         gv_pred         : geometric fixed point for g_v
4343         S_tot           : total scaling factor
4344         r_nu           : neutrino radius [m]
4345     """
4346     alpha_inv_geom = 1.0 / alpha_geom
4347
4348     S_proj = alpha_inv_geom / (BCC_NEIGHBOURS + PI)
4349     S_exp = EULER
4350
4351     a_coef = SQRT2 - 1.0
4352     b_coef = -(S_proj + EULER + SQRT2 - 1.0)
4353     c_coef = 2.0 * EULER**2
4354
4355     discriminant = b_coef**2 - 4.0 * a_coef * c_coef
4356     if discriminant < 0:
4357         raise ValueError("Negative discriminant in g_v fixed point equation")
4358
4359     sqrt_disc = math.sqrt(discriminant)
4360     root1 = (-b_coef + sqrt_disc) / (2.0 * a_coef)
4361     root2 = (-b_coef - sqrt_disc) / (2.0 * a_coef)

```

```

4362
4363     if 0.0 < root1 < 1.0:
4364         gv_pred = root1
4365     elif 0.0 < root2 < 1.0:
4366         gv_pred = root2
4367     else:
4368         raise ValueError("No physical root for g_v in (0,1)")
4369
4370     delta_imp = (1.0 - gv_pred) * (SQRT2 - 1.0)
4371     S_tot = S_proj + S_exp + delta_imp
4372     r_nu = q_P * S_tot
4373
4374     return {
4375         "alpha_inv_geom": alpha_inv_geom,
4376         "S_proj": S_proj,
4377         "S_exp": S_exp,
4378         "delta_imp": delta_imp,
4379         "gv_pred": gv_pred,
4380         "S_tot": S_tot,
4381         "r_nu": r_nu,
4382     }
4383
4384
4385 def test_decadic_resonance(r_e: float, r_nu: float) -> dict:
4386     """
4387     Test the 1:100 radial resonance between electron and neutrino.
4388
4389     Parameters
4390     -----
4391     r_e : float
4392         Classical electron radius [m].
4393     r_nu : float
4394         Neutrino radius [m].
4395
4396     Returns
4397     -----
4398     dict with:
4399         ratio      : r_e / r_nu
4400         expected   : 100.0
4401         rel_err    : relative error in percent
4402         K_implied  : ratio^5
4403     """
4404     ratio = r_e / r_nu
4405     rel_err = abs(ratio - 100.0) / 100.0 * 100.0
4406     K_implied = ratio**5
4407     return {
4408         "ratio": ratio,
4409         "expected": 100.0,
4410         "rel_err": rel_err,
4411         "K_implied": K_implied,
4412     }
4413
4414 def compute_C_unif(alpha_geom: float, L_p_geom: float, K_WC: int) -> float:
4415     """
4416     Compute the unified coupling operator C_Unif.
4417
4418     Formula:
4419         C_Unif = 1/K_WC + 1 + alpha_geom / (pi * L_p_geom)
4420
4421     Parameters
4422     -----
4423     alpha_geom : float
4424         Geometric fine-structure constant.
4425     L_p_geom : float
4426         Ideal BCC projection factor.
4427     K_WC : int
4428         Number of wave centres in the electron.
4429
4430     Returns
4431     -----
4432     float
4433         Unified coupling operator.
4434     """

```

```

4435     return (1.0 / K_WC) + 1.0 + (alpha_geom / (PI * L_p_geom))
4436
4437
4438 def compute_X_eff(alpha_geom: float, L_p_geom: float, K_WC: int) -> float:
4439     """
4440     Compute the geometric dilution factor X_eff.
4441
4442     Formula:
4443         X_eff = A_pi * 3 * K_WC * sqrt(2) / C_Unif
4444
4445     Parameters
4446     -----
4447     alpha_geom : float
4448         Geometric fine-structure constant.
4449     L_p_geom : float
4450         Ideal BCC projection factor.
4451     K_WC : int
4452         Number of wave centres in the electron.
4453
4454     Returns
4455     -----
4456     float
4457         Dilution factor X_eff.
4458     """
4459     A_pi = compute_alpha_core()
4460     C_Unif = compute_C_unif(alpha_geom, L_p_geom, K_WC)
4461     return (A_pi * 3.0 * K_WC * SQRT2) / C_Unif
4462
4463 def derive_lambda_l_geometric(
4464     alpha_geom: float,
4465     r_e: float,
4466     r_nu: float,
4467     N_geom: float,
4468     L_p_geom: float,
4469     K_WC: int,
4470 ) -> float:
4471     """
4472     Derive the fundamental EMC length lambda_l from pure geometry.
4473
4474     The formula is the analytic solution of the self-consistent equation
4475     obtained by combining the geometric G with the Planck length definition.
4476     It does NOT use hbar or G as input.
4477
4478     Formula:
4479         lambda_l = [ r_e^2 * sqrt(X_eff) /
4480                     ( alpha_geom * A_pi^4 * N_geom^3 * K_WC *
4481                       (r_nu/(2*e))^(3/2) ) ]^2
4482
4483     Parameters
4484     -----
4485     alpha_geom : float
4486         Geometric fine-structure constant.
4487     r_e : float
4488         Classical electron radius [m].
4489     r_nu : float
4490         Neutrino radius [m].
4491     N_geom : float
4492         Effective BCC stiffness.
4493     L_p_geom : float
4494         Ideal BCC projection factor.
4495     K_WC : int
4496         Number of wave centres in the electron.
4497
4498     Returns
4499     -----
4500     float
4501         Derived lambda_l [m].
4502     """
4503     A_pi = compute_alpha_core()
4504     X_eff = compute_X_eff(alpha_geom, L_p_geom, K_WC)
4505
4506     r_nu_factor = (r_nu / (2.0 * EULER)) ** (3.0 / 2.0)
4507

```

```

4508     bracket = (
4509         r_e**2 * math.sqrt(X_eff)
4510         / (alpha_geom * A_pi**4 * N_geom**3 * K_WC * r_nu_factor)
4511     )
4512
4513     return bracket**2
4514
4515 # =====
4516 # SECTION 7: GRAVITY SECTOR EMERGENCE
4517 # -----
4518 # Gravity is derived from the electron soliton and the BCC lattice geometry.
4519 # No calibrated N_final and no calibrated L_p are used.
4520 #
4521 #     N_geom      : derived from BCC packing impedance
4522 #     L_p_geom    : ideal BCC projection factor 2/sqrt(3)
4523 #     alpha_geom  : derived from A_pi and eps_M
4524 #
4525 # The effective volume deficit is computed via the unified coupling:
4526 #     C_Unif = 1/K_WC + 1 + alpha_geom/(pi * L_p_geom)
4527 # =====
4528
4529 def gravity_sector(
4530     alpha_geom: float,
4531     r_nu: float,
4532     N_geom: float,
4533     L_p_geom: float,
4534     K_WC: int,
4535     lambda_l: float,
4536     r_e: float,
4537     m_e: float,
4538     c0: float,
4539 ) -> dict:
4540     """
4541     Derive the gravitational constant G from pure geometry and electron anchors.
4542
4543     Parameters
4544     -----
4545     alpha_geom : float
4546         Geometric fine-structure constant.
4547     r_nu : float
4548         Neutrino radius [m].
4549     N_geom : float
4550         Effective BCC stiffness derived from packing impedance.
4551     L_p_geom : float
4552         Ideal BCC projection factor (2/sqrt(3)).
4553     K_WC : int
4554         Number of wave centres in the electron (10).
4555     lambda_l : float
4556         Fundamental EMC spacing derived from geometry [m].
4557     r_e : float
4558         Classical electron radius [m].
4559     m_e : float
4560         Electron mass [kg].
4561     c0 : float
4562         Speed of light [m/s].
4563
4564     Returns
4565     -----
4566     dict with:
4567         G_Base      : base soliton scaling
4568         N_nu_statutory: statutory background density
4569         C_Unif      : unified coupling operator
4570         X_eff       : dilution factor
4571         N_nu_eff    : effective volume deficit
4572         G_EWT       : predicted gravitational constant
4573         rel_err     : relative error vs CODATA
4574     """
4575     G_Base = (c0**2 * r_e) / m_e
4576     A_pi = compute_alpha_core()
4577     C_Unif = compute_C_unif(alpha_geom, L_p_geom, K_WC)
4578     X_eff = compute_X_eff(alpha_geom, L_p_geom, K_WC)
4579     N_nu_statutory = (r_nu / (2.0 * lambda_l * EULER)) ** 3
4580     N_nu_eff = N_nu_statutory / X_eff

```



```

4654     )
4655
4656     res = gravity_sector(
4657         alpha_geom=alpha_test,
4658         r_nu=r_nu,
4659         N_geom=N_test,
4660         L_p_geom=L_p_geom,
4661         K_WC=K_WC,
4662         lambda_l=lambda_test,
4663         r_e=r_e,
4664         m_e=m_e,
4665         c0=c0,
4666     )
4667
4668     print(
4669         f"{N_test:12.5f} {alpha_inv_test:14.9f} {lambda_test:16.6e}"
4670         f"{res['G_EWT']:22.15e} {res['rel_err']:10.6f}"
4671     )
4672
4673     print("\n[2] Vary L_p_geom")
4674     print("-" * 78)
4675     print(f"{L_p':>10} {'alpha_inv':>14} {'lambda_l[m]':>16} {'G_EWT[m^3/kg/s^2]':>22} {'G_err%'>10}")
4676     G_err%'>10}")
4677     for factor in [0.995, 0.998, 1.0, 1.002, 1.005]:
4678         Lp_test = L_p_geom * factor
4679
4680         lambda_test = derive_lambda_l_geometric(
4681             alpha_geom=alpha_geom,
4682             r_e=r_e,
4683             r_nu=r_nu,
4684             N_geom=N_geom,
4685             L_p_geom=Lp_test,
4686             K_WC=K_WC,
4687         )
4688
4689         res = gravity_sector(
4690             alpha_geom=alpha_geom,
4691             r_nu=r_nu,
4692             N_geom=N_geom,
4693             L_p_geom=Lp_test,
4694             K_WC=K_WC,
4695             lambda_l=lambda_test,
4696             r_e=r_e,
4697             m_e=m_e,
4698             c0=c0,
4699         )
4700
4701         print(
4702             f"{Lp_test:10.6f} {1.0/alpha_geom:14.9f} {lambda_test:16.6e}"
4703             f"{res['G_EWT']:22.15e} {res['rel_err']:10.6f}"
4704         )
4705
4706     print("\n[3] Vary lambda_l directly")
4707     print("-" * 78)
4708     print(f"{lambda_l[m]':>16} {'G_EWT[m^3/kg/s^2]':>22} {'G_err%'>10}")
4709     for factor in [0.99, 1.0, 1.01]:
4710         lambda_test = lambda_l_geom * factor
4711         res = gravity_sector(
4712             alpha_geom=alpha_geom,
4713             r_nu=r_nu,
4714             N_geom=N_geom,
4715             L_p_geom=L_p_geom,
4716             K_WC=K_WC,
4717             lambda_l=lambda_test,
4718             r_e=r_e,
4719             m_e=m_e,
4720             c0=c0,
4721         )
4722         print(
4723             f"{lambda_test:16.6e} {res['G_EWT']:22.15e} {res['rel_err']:10.6f}"
4724         )
4725
4726     print("-" * 78)

```

```

4727
4728 def input_rigidity_test(base_alpha_geom, base_r_nu, base_N_geom, base_Lp):
4729     print("\n[4] Vary fundamental inputs r_e and m_e")
4730     print("-" * 78)
4731     print(f"{factor':>8}\{'r_e[m]':>16}\{'m_e[kg]':>16}\{'lambda_l[m]':>18}\{'G_err_u'":
4732           %':>10}")
4733
4734     for factor in [0.998, 0.999, 1.0, 1.001, 1.002]:
4735         r_e_test = R_E * factor
4736         m_e_test = M_E * factor
4737
4738         lambda_test = derive_lambda_l_geometric(
4739             alpha_geom=base_alpha_geom,
4740             r_e=r_e_test,
4741             r_nu=base_r_nu,
4742             N_geom=base_N_geom,
4743             L_p_geom=base_Lp,
4744             K_WC=10,
4745         )
4746
4747         res = gravity_sector(
4748             alpha_geom=base_alpha_geom,
4749             r_nu=base_r_nu,
4750             N_geom=base_N_geom,
4751             L_p_geom=base_Lp,
4752             K_WC=10,
4753             lambda_l=lambda_test,
4754             r_e=r_e_test,
4755             m_e=m_e_test,
4756             c0=C0,
4757         )
4758
4759         print(
4760             f"{factor:8.3f}\{r_e_test:16.6e}\{m_e_test:16.6e}"
4761             f"{lambda_test:18.6e}\{res['rel_err']:10.6f}"
4762         )
4763
4764 # =====
4765 # SECTION 8: LEPTON ANOMALOUS MAGNETIC MOMENTS (AMM)
4766 # =====
4767
4768 def get_AMMi_K(n: int) -> int:
4769     """
4770     Recursive nodal count for the lepton generations.
4771
4772     K_1 = 10
4773     K_n = K_{n-1} + round(10^(n-1) * 2*pi^2)
4774
4775     Parameters
4776     -----
4777     n : int
4778         Generation number (1 = electron, 2 = muon, 3 = tau).
4779
4780     Returns
4781     -----
4782     int
4783         Total nodal count K_n.
4784     """
4785     if n == 1:
4786         return 10
4787     else:
4788         delta_K = round(10 ** (n - 1) * (2.0 * PI**2))
4789         return get_AMMi_K(n - 1) + delta_K
4790
4791
4792 def compute_lepton_amms(
4793     alpha_geom: float,
4794     eps_M: float,
4795     L_mu_dim: int = 5,
4796     L_tau_dim: int = 34,
4797     K_WC_e: int = 10,
4798 ) -> dict:
4799     """

```

```

14800     Compute the full anomalous magnetic moments for the lepton family.
14801     """
14802     A_pi = compute_alpha_core()
14803
14804     a_e_ppm = (alpha_geom / (2.0 * PI)) * (1.0 - eps_M * (PI**3)) * 1e6
14805
14806     K_e = K_WC_e
14807     K_mu_total = get_AMMi_K(2)
14808     K_mu_delta = K_mu_total - K_e
14809     M_mu_shell = K_mu_delta / K_e
14810
14811     B_mu_scale = (3.0 * A_pi * PI**3) / (2.0 * L_mu_dim**2)
14812     a_mu_shell_ppm = B_mu_scale * (1.0 - eps_M) ** (M_mu_shell * PI**3)
14813
14814     O_mu = 1.0 / (4.0 * PI**2)
14815     a_mu_shell_correction = a_mu_shell_ppm * O_mu
14816     a_mu_ppm = a_e_ppm + a_mu_shell_correction
14817
14818     K_tau_total = get_AMMi_K(3)
14819     M_tau_rel = K_tau_total / K_e
14820
14821     B_tau_base = ((3.0 * A_pi * PI**3) / (8.0 * SQRT2)) + (A_pi / 2.0)
14822     a_tau_shell_raw_ppm = B_tau_base * (1.0 - eps_M) ** (M_tau_rel * PI**3)
14823
14824     a_tau_shell_total_ppm = a_mu_shell_ppm + a_tau_shell_raw_ppm + L_mu_dim**2
14825
14826     O_tau = 1.0
14827     a_tau_ppm = a_e_ppm + O_tau * (a_tau_shell_total_ppm - a_e_ppm)
14828
14829     return {
14830         "a_e_ppm": a_e_ppm,
14831         "a_mu_ppm": a_mu_ppm,
14832         "a_tau_ppm": a_tau_ppm,
14833         "a_mu_shell_ppm": a_mu_shell_ppm,
14834         "a_tau_shell_ppm": a_tau_shell_total_ppm,
14835     }
14836
14837
14838 def report_lepton_amms(res: dict) -> None:
14839     """Print lepton AMM results and compare with experimental targets."""
14840     print("----Lepton Anomalous Magnetic Moments----")
14841     print(f"Electron a_e: {res['a_e_ppm']:.6f} ppm (CODATA: {A_E_CODATA*1e6:.6f} ppm)")
14842     print(f"Muon shell a_mu: {res['a_mu_shell_ppm']:.6f} ppm")
14843     print(f"Muon full a_mu: {res['a_mu_ppm']:.6f} ppm (Exp: {A_MU_EXP*1e6:.6f} ppm)")
14844     print(f"Tau shell a_tau: {res['a_tau_shell_ppm']:.6f} ppm")
14845     print(f"Tau full a_tau: {res['a_tau_ppm']:.6f} ppm (Exp: {A_TAU_EXP*1e6:.6f} ppm)")
14846
14847 def amm_rigidity_test(
14848     base_alpha_geom: float,
14849     base_eps_M: float,
14850     L_mu_dim: int = 5,
14851     L_tau_dim: int = 34,
14852 ) -> None:
14853     """
14854     Test the rigidity of the lepton AMM sector against changes in eps_M.
14855     """
14856     print("\n" + "=" * 78)
14857     print("AMM RIGIDITY TEST (vary eps_M)")
14858     print("=" * 78)
14859     print(f"{'factor':>8} {'eps_M':>18} {'alpha_inv':>14} {'a_e [ppm]':>14} {'a_mu [ppm]':>14} {'a_tau [ppm]':>14}")
14860     print(f"{'':>14} {'a_tau [ppm]':>14}")
14861     print("-" * 78)
14862
14863     for factor in [0.998, 0.999, 1.0, 1.001, 1.002]:
14864         eps_test = base_eps_M * factor
14865         alpha_inv_test = compute_alpha_geometric(eps_test)
14866         alpha_test = 1.0 / alpha_inv_test
14867
14868         res = compute_lepton_amms(
14869             alpha_geom=alpha_test,
14870             eps_M=eps_test,
14871             L_mu_dim=L_mu_dim,
14872             L_tau_dim=L_tau_dim,

```

```

14873         )
14874
14875     print(
14876         f"{{factor:8.3f}}_{{eps_test:18.6e}}_{{alpha_inv_test:14.9f}}_{{
14877         f"{{res['a_e_ppm']}:14.6f}}_{{res['a_mu_ppm']}:14.6f}}_{{res['a_tau_ppm']}:14.6f}}
14878     )
14879
14880     print("-" * 78)
14881     print("Reference experimental:")
14882     print(f"{{a_e}}_{{A_E_CODATA*1e6:.6f}}_ppm")
14883     print(f"{{a_mu}}_{{A_MU_EXP*1e6:.6f}}_ppm")
14884     print(f"{{a_tau}}_{{A_TAU_EXP*1e6:.6f}}_ppm")
14885     print("=" * 78)
14886
14887     # =====
14888     # SECTION 9: ATOMIC SCALES FROM PURE GEOMETRY
14889     # =====
14890
14891     def compute_atomic_scales(alpha_geom: float, r_e_geom: float) -> dict:
14892         """
14893         Compute atomic scales from geometric alpha and geometric electron radius.
14894         """
14895         R_inf = (alpha_geom**3) / (4.0 * PI * r_e_geom)
14896         a0 = r_e_geom / (alpha_geom**2)
14897         lambda_C = (2.0 * PI * r_e_geom) / alpha_geom
14898
14899         return {
14900             "R_inf": R_inf,
14901             "a0": a0,
14902             "lambda_C": lambda_C,
14903         }
14904
14905     def report_atomic_scales(res: dict) -> None:
14906         """Print atomic scale results and compare with CODATA 2022."""
14907         print("--- Atomic Scales from Pure Geometry ---")
14908         print(f"{{Rydberg_R_inf}}_{{res['R_inf']:.8f}}_m^-1")
14909         print(f"{{CODATA_R_inf}}_{{R_INF_CODATA:.8f}}_m^-1")
14910         print(f"{{rel_error_R_inf}}_{{abs(res['R_inf']-R_INF_CODATA)/R_INF_CODATA*100:.6f}}_%")
14911
14912         print(f"{{Bohr_radius_a0}}_{{res['a0']:.15e}}_m")
14913         print(f"{{CODATA_a0}}_{{A0_CODATA:.15e}}_m")
14914         print(f"{{rel_error_a0}}_{{abs(res['a0']-A0_CODATA)/A0_CODATA*100:.6f}}_%")
14915
14916         print(f"{{Compton_lambda_C}}_{{res['lambda_C']:.15e}}_m")
14917         print(f"{{CODATA_lambda_C}}_{{LAMBDA_C_CODATA:.15e}}_m")
14918         print(f"{{rel_error_lambda_C}}_{{abs(res['lambda_C']-LAMBDA_C_CODATA)/LAMBDA_C_CODATA*100:.6f}}_%")
14919
14920     def main():
14921         print("ENHANCED EWT - EMERGENCE ENGINE (PROTOTYPE)")
14922
14923         A_pi = compute_alpha_core()
14924         print(f"{{Alpha_core_A_pi}}_{{4*pi^3+pi^2+pi}}_{{A_pi:.15f}}")
14925
14926         bcc_derived = derive_eps_M_from_BCC(8.0 * PI**4)
14927         eps_M_bcc = bcc_derived["eps_M"]
14928
14929         print("\nDerived eps_M from BCC packing:")
14930         print(f"{{eta}}_{{bcc_derived['eta']:.10f}}")
14931         print(f"{{N_ideal}}_{{bcc_derived['N_ideal']:.10f}}")
14932         print(f"{{zeta}}_{{bcc_derived['zeta']:.10f}}")
14933         print(f"{{N_geom}}_{{bcc_derived['N_geom']:.10f}}")
14934         print(f"{{eps_M}}_{{bcc_derived['eps_M']:.15e}}")
14935
14936         alpha_inv_bcc = compute_alpha_geometric(eps_M_bcc)
14937         alpha_geom = 1.0 / alpha_inv_bcc
14938
14939         res_alpha_bcc = test_alpha_sector(alpha_inv_bcc)
14940
14941         print("\nAlpha inverse fine-structure constant test:")
14942         print(f"{{Pure_BCC_geometry_pred}}_{{res_alpha_bcc['alpha_inv_pred']:.12f}}_{{rel_err}}_{{
14943         {res_alpha_bcc['rel_err']:.6f}}_%")
14944
14945

```

```

14946     print(f"CODATA_2022{ALPHA_INV_CODATA:.12f}")
14947
14948     hbar_geom = derive_hbar(alpha_geom, R_E, M_E, CO)
14949     q_P_geom = derive_planck_charge_from_e(alpha_geom, E_CHARGE_CODATA)
14950
14951     print(f"Derived q_P{q_P_geom:.15e}")
14952     print(f"Reference Q_P_INPUT{Q_P_INPUT:.15e}")
14953
14954     ratio_qP_e_derived = q_P_geom / E_CHARGE_CODATA
14955     ratio_qP_e_expected = 1.0 / math.sqrt(alpha_geom)
14956
14957     print(f"\nGeometric ratio q_P/e:")
14958     print(f"derived q_P/e={ratio_qP_e_derived:.10f}")
14959     print(f"expected 1/sqrt(alpha_geom)={ratio_qP_e_expected:.10f}")
14960     print(f"difference={abs(ratio_qP_e_derived-ratio_qP_e_expected):.6e}")
14961
14962     lambda_unc = derive_lambda_uncorr(q_P_geom)
14963
14964     print(f"\nDerived hbar{hbar_geom:.15e}")
14965     print(f"Using q_P input{q_P_geom:.15e}")
14966     print(f"lambda_uncorr{lambda_unc:.15e}")
14967
14968     nu_result = derive_neutrino_radius(alpha_geom, q_P_geom)
14969     print(f"\nNeutrino radius r_nu:{nu_result['r_nu']:.15e}um")
14970     print(f"g_v fixed point{nu_result['gv_pred']:.15f}")
14971
14972     res_nu = test_decadic_resonance(R_E, nu_result['r_nu'])
14973     print(f"\nElectron/neutrino ratio:{res_nu['ratio']:.12f}")
14974     print(f"Expected{res_nu['expected']}")
14975     print(f"Relative error{res_nu['rel_err']:.6f}%")
14976     print(f"K implied(r^5){res_nu['K_implied']:.12e}")
14977
14978     lambda_l_geom = derive_lambda_l_geometric(
14979         alpha_geom=alpha_geom,
14980         r_e=R_E,
14981         r_nu=nu_result["r_nu"],
14982         N_geom=bcc_derived["N_geom"],
14983         L_p_geom=BCC_IDEAL_PROJECTION_LP,
14984         K_WC=10,
14985     )
14986
14987     print(f"\nDerived lambda_l(geometry):{lambda_l_geom:.15e}um")
14988     print(f"CODATA lambda_l{LAMBDA_L:.15e}um")
14989     print(f"Relative error{abs(lambda_l_geom-LAMBDA_L)/LAMBDA_L*100:.6f}%")
14990
14991     print("\n--- Gravity Sector (Geometric L_p, derived lambda_l) ---")
14992     gravity_res = gravity_sector(
14993         alpha_geom=alpha_geom,
14994         r_nu=nu_result["r_nu"],
14995         N_geom=bcc_derived["N_geom"],
14996         L_p_geom=BCC_IDEAL_PROJECTION_LP,
14997         K_WC=10,
14998         lambda_l=lambda_l_geom,
14999         r_e=R_E,
15000         m_e=M_E,
15001         cO=CO,
15002     )
15003     report_gravity(gravity_res)
15004
15005     input_rigidity_test(alpha_geom, nu_result["r_nu"], bcc_derived["N_geom"],
15006         BCC_IDEAL_PROJECTION_LP)
15007
15008     rigidity_test(
15009         alpha_geom=alpha_geom,
15010         r_e=R_E,
15011         r_nu=nu_result["r_nu"],
15012         N_geom=bcc_derived["N_geom"],
15013         L_p_geom=BCC_IDEAL_PROJECTION_LP,
15014         K_WC=10,
15015         lambda_l_geom=lambda_l_geom,
15016         m_e=M_E,
15017         cO=CO,
15018     )

```

```

15019     ladder_bcc = build_geometric_ladder(eps_M_bcc, "BCC_packing_derived")
15020
15021
15022     print(f"\n---Geometric Ladder (Pure BCC)---")
15023     report_ladder(ladder_bcc)
15024     run_sector_tests(ladder_bcc)
15025
15026     amm_results = compute_lepton_amms(alpha_geom, eps_M_bcc)
15027     report_lepton_amms(amm_results)
15028     amm_rigidity_test(alpha_geom, eps_M_bcc)
15029
15030     r_e_geom = 100.0 * nu_result["r_nu"]
15031     atomic_res = compute_atomic_scales(alpha_geom, r_e_geom)
15032     report_atomic_scales(atomic_res)
15033
15034     print("\n" + "=" * 78)
15035     print("END OF TEST")
15036     print("=" * 78)
15037
15038 if __name__ == "__main__":
15039     main()
15040

```

Listing 1: Core geometric emergence engine.

A.2.2 Emergence Engine Output

The console output of `ewt_emergence_engine.py` is reproduced below.

```

5043 ENHANCED EWT - EMERGENCE ENGINE (PROTOTYPE)
5044
5045 Alpha core A_pi = 4*pi^3 + pi^2 + pi = 137.036303775878423
5046
5047 Derived eps_M from BCC packing:
5048     eta      = 0.6801747616
5049     N_ideal  = 779.2727282720
5050     zeta     = 0.0006033964
5051     N_geom   = 778.8025178842
5052     eps_M    = 4.141169769303932e-05
5053
5054 Alpha inverse fine-structure constant test:
5055     Pure BCC geometry : pred = 137.036262364181 | rel_err = 0.000192 %
5056     CODATA 2022      : 137.035999084000
5057     Derived q_P       : 1.875547839472885e-18
5058     Reference Q_P_INPUT : 1.875546037780000e-18
5059
5060 Geometric ratio q_P / e:
5061     derived q_P / e = 11.7062488597
5062     expected 1/sqrt(alpha_geom) = 11.7062488597
5063     difference      = 0.000000e+00
5064
5065 Derived hbar      : 1.054573843737794e-34
5066 Using q_P input   : 1.875547839472885e-18
5067 lambda_uncorr     : 2.771705640418664e-17
5068
5069 Neutrino radius r_nu : 2.817935436016830e-17 m
5070 g_v fixed point     : 0.983594444710376
5071
5072 Electron/neutrino ratio : 100.000173537800
5073 Expected              : 100.0
5074 Relative error        : 0.000174 %
5075 K_implied (r^5)       : 1.000008676920e+10
5076
5077 Derived lambda_l (geometry): 1.616646406608278e-35 m
5078 CODATA lambda_l       : 1.616200000000000e-35 m
5079 Relative error        : 0.027621 %
5080
5081 --- Gravity Sector (Geometric L_p, derived lambda_l) ---
5082     G_Base      : 2.780252259136399e+32
5083     N_nu_statutory: 3.295904039818979e+52
5084     C_Unif      : 1.102011616800948
5085     X_eff       : 5275.7683235656
5086

```

```

5087 N_nu_eff      : 6.247249381851994e+48
5088 G_EWT        : 6.677519975508460e-11
5089 G_CODATA     : 6.674305000000000e-11
5090 rel_error    : 0.048169 %
5091
5092 [4] Vary fundamental inputs r_e and m_e
5093 -----
5094 factor      r_e [m]      m_e [kg]      lambda_l [m]      G_err %
5095 0.998      2.812304e-15   9.091165e-31   1.603752e-35   1.146422
5096 0.999      2.815122e-15   9.100274e-31   1.610190e-35   0.550621
5097 1.000      2.817940e-15   9.109384e-31   1.616646e-35   0.048169
5098 1.001      2.820758e-15   9.118493e-31   1.623123e-35   0.649961
5099 1.002      2.823576e-15   9.127602e-31   1.629618e-35   1.254766
5100
5101 =====
5102 RIGIDITY TEST
5103 =====
5104
5105 [1] Vary N_geom
5106 -----
5107 N_geom      alpha_inv      lambda_l [m]      G_EWT [m^3/kg/s^2]      G_err %
5108 774.90851   137.036262156    1.666006e-35    7.091501922019398e-11   6.250792
5109 777.24491   137.036262281    1.636183e-35    6.839883420425040e-11   2.480834
5110 778.80252   137.036262364    1.616646e-35    6.677519975508460e-11   0.048169
5111 780.36012   137.036262447    1.597382e-35    6.519323594110790e-11   2.322061
5112 782.69653   137.036262570    1.568985e-35    6.289591749060191e-11   5.764095
5113
5114 [2] Vary L_p_geom
5115 -----
5116 L_p      alpha_inv      lambda_l [m]      G_EWT [m^3/kg/s^2]      G_err %
5117 1.148927   137.036262364    1.616632e-35    6.677397472932486e-11   0.046334
5118 1.152391   137.036262364    1.616640e-35    6.677471121371631e-11   0.047437
5119 1.154701   137.036262364    1.616646e-35    6.677519975508460e-11   0.048169
5120 1.157010   137.036262364    1.616652e-35    6.677568635151728e-11   0.048899
5121 1.160474   137.036262364    1.616661e-35    6.677641262474325e-11   0.049987
5122
5123 [3] Vary lambda_l directly
5124 -----
5125 lambda_l [m]      G_EWT [m^3/kg/s^2]      G_err %
5126 1.600480e-35    6.577608001792828e-11   1.448795
5127 1.616646e-35    6.677519975508460e-11   0.048169
5128 1.632813e-35    6.777932766352434e-11   1.552638
5129 =====
5130
5131 --- Geometric Ladder (Pure BCC) ---
5132 Source      : BCC packing derived
5133 eps_M       : 4.141169769303932e-05
5134 C_local     : 1.464124612959771e-05
5135 C_gap       : 1.014075935809472
5136 C_fermion   : 1.008981094016494
5137 Ladder:
5138 pi^3 = 31.006276680299816
5139 pi^4 = 97.409091034002415
5140 pi^5 = 306.019684785281413
5141 pi^6 = 961.389193575304262
5142 pi^7 = 3020.293227776791355
5143 M_W prediction : 80.514122 GeV
5144 rel. err PDG   : 0.1706 %
5145 rel. err CDF II : 0.1002 %
5146 sin theta_C    : 0.2242876694
5147 rel. err PDG   : 0.005497 %
5148 --- Lepton Anomalous Magnetic Moments ---
5149 Electron a_e   : 1159.916228 ppm (CODATA: 1159.652182 ppm)
5150 Muon shell     : 248.571133 ppm
5151 Muon full a_mu : 1166.212608 ppm (Exp: 1165.920610 ppm)
5152 Tau shell      : 1176.838130 ppm
5153 Tau full a_tau : 1176.838130 ppm (Exp: 1177.210000 ppm)
5154
5155 =====
5156 AMM RIGIDITY TEST (vary eps_M)
5157 =====
5158 factor      eps_M      alpha_inv      a_e [ppm]      a_mu [ppm]      a_tau [ppm]
5159 -----

```

```

51160      0.998      4.132887e-05  137.036262447    1159.919210    1166.215909    1177.356781
51161      0.999      4.137029e-05  137.036262406    1159.917719    1166.214259    1177.097420
51162      1.000      4.141170e-05  137.036262364    1159.916228    1166.212608    1176.838130
51163      1.001      4.145311e-05  137.036262323    1159.914737    1166.210958    1176.578912
51164      1.002      4.149452e-05  137.036262281    1159.913246    1166.209308    1176.319764
51165      -----
51166      Reference experimental:
51167      a_e      : 1159.652182 ppm
51168      a_mu     : 1165.920610 ppm
51169      a_tau    : 1177.210000 ppm
51170      =====
51171      --- Atomic Scales from Pure Geometry ---
51172      Rydberg R_inf      : 10973687.36203397 m^-1
51173      CODATA              : 10973731.56815700 m^-1
51174      rel_error          : 0.000403 %
51175      Bohr radius a0      : 5.291783259434870e-11 m
51176      CODATA              : 5.291772109030000e-11 m
51177      rel_error          : 0.000211 %
51178      Compton lambda_C    : 2.426310689655160e-12 m
51179      CODATA              : 2.426310238670000e-12 m
51180      rel_error          : 0.000019 %
51181      =====
51182      END OF TEST
51183      =====
51184
51185

```

Listing 2: Console output from `ewt_emergence_engine.py`.

A.2.3 Extended Lepton AMM

`ewt_amm_extended.py` computes the full anomalous magnetic moments of the electron, muon, and tau using the geometric fine-structure constant and the magnetic deficit. It also reports the internal shell references and the corresponding consistency errors.

```

51190      #!/usr/bin/env python3
51191      """
51192      From manuscript version: 5.0.0
51193
51194      Enhanced EWT -- Extended Lepton AMM Module
51195      =====
51196      Provides the extended lepton anomalous magnetic moments with internal
51197      shell references and consistency tests between the geometric AMM
51198      sector and the orbital mass sector.
51199
51200      This module complements ewt_emergence_engine.py, which contains the
51201      basic full AMM predictions and rigidity tests.
51202      """
51203
51204
51205      import math
51206
51207      try:
51208          from ewt_emergence_engine import (
51209              PI, SQRT2, SQRT3, EULER,
51210              A_MU_EXP,
51211              A_TAU_EXP,
51212              A_E_CODATA,
51213              compute_alpha_core,
51214              derive_eps_M_from_BCC,
51215              get_AMMi_K,
51216          )
51217      except ImportError:
51218          raise ImportError("This module requires ewt_emergence_engine.py in the same directory.")
51219
51220
51221      def compute_lepton_amms_extended(alpha_geom: float, eps_M: float) -> dict:
51222          """
51223          Compute extended lepton AMMs with internal shell references.
51224
51225          Parameters
51226          -----

```

```

5227     alpha_geom : float
5228         Geometric fine-structure constant.
5229     eps_M : float
5230         Magnetic deficit from BCC geometry.
5231
5232     Returns
5233     -----
5234     dict with:
5235         a_e_ppm
5236         a_mu_shell_ppm
5237         a_mu_full_ppm
5238         a_tau_shell_total_ppm
5239         a_tau_full_ppm
5240         ref_a_mu_shell_ppm
5241         ref_a_tau_shell_ppm
5242         err_a_mu_shell_pct
5243         err_a_tau_shell_pct
5244     """
5245     A_pi = compute_alpha_core()
5246     L_mu_dim = 5
5247     L_tau_dim = 34
5248     K_e = 10
5249
5250     # Electron
5251     a_e_ppm = (alpha_geom / (2.0 * PI)) * (1.0 - eps_M * (PI**3)) * 1e6
5252
5253     # Muon shell
5254     K_mu_total = get_AMMi_K(2)
5255     K_mu_delta = K_mu_total - K_e
5256     M_mu_shell = K_mu_delta / K_e
5257
5258     B_mu_scale = (3.0 * A_pi * PI**3) / (2.0 * L_mu_dim**2)
5259     a_mu_shell_ppm = B_mu_scale * (1.0 - eps_M) ** (M_mu_shell * PI**3)
5260
5261     # Projection operator for muon (2D -> 1D)
5262     O_mu = 1.0 / (4.0 * PI**2)
5263
5264     a_mu_full_ppm = a_e_ppm + a_mu_shell_ppm * O_mu
5265
5266     # Tau shell
5267     K_tau_total = get_AMMi_K(3)
5268     M_tau_rel = K_tau_total / K_e
5269
5270     B_tau_base = ((3.0 * A_pi * PI**3) / (8.0 * SQRT2)) + (A_pi / 2.0)
5271     a_tau_shell_raw_ppm = B_tau_base * (1.0 - eps_M) ** (M_tau_rel * PI**3)
5272
5273     # Recursive accumulation with interface tension L_mu^2
5274     a_tau_shell_total_ppm = a_mu_shell_ppm + a_tau_shell_raw_ppm + L_mu_dim**2
5275
5276     # Projection operator for tau (3D -> 3D, unity)
5277     a_tau_full_ppm = a_e_ppm + (a_tau_shell_total_ppm - a_e_ppm)
5278
5279     # Internal references (from orbital mass relations)
5280     ref_a_mu_shell_ppm = 248.8
5281     ref_a_tau_shell_ppm = 1177.21
5282
5283     err_a_mu_shell_pct = abs(a_mu_shell_ppm - ref_a_mu_shell_ppm) / ref_a_mu_shell_ppm *
5284         100.0
5285     err_a_tau_shell_pct = abs(a_tau_shell_total_ppm - ref_a_tau_shell_ppm) /
5286         ref_a_tau_shell_ppm * 100.0
5287
5288     return {
5289         "a_e_ppm": a_e_ppm,
5290         "a_mu_shell_ppm": a_mu_shell_ppm,
5291         "a_mu_full_ppm": a_mu_full_ppm,
5292         "a_tau_shell_total_ppm": a_tau_shell_total_ppm,
5293         "a_tau_full_ppm": a_tau_full_ppm,
5294         "ref_a_mu_shell_ppm": ref_a_mu_shell_ppm,
5295         "ref_a_tau_shell_ppm": ref_a_tau_shell_ppm,
5296         "err_a_mu_shell_pct": err_a_mu_shell_pct,
5297         "err_a_tau_shell_pct": err_a_tau_shell_pct,
5298     }
5299

```

```

5300
5301 def report_extended_amms(res: dict) -> None:
5302     """Print the extended AMM report in Scilab style."""
5303     print("\n---EXTENDED LEPTON ANOMALOUS MAGNETIC MOMENTS---")
5304     print(f"Electron full a_e : {res['a_e_ppm']:.6f} ppm")
5305     print(f"CODATA reference : {A_E_CODATA*1e6:.6f} ppm")
5306     print(f"Full rel. error : {abs(res['a_e_ppm'] - A_E_CODATA*1e6) / (A_E_CODATA*1e6) * 100:.6f} %")
5307
5308     print()
5309     print(f"Muon shell (internal) : {res['a_mu_shell_ppm']:.6f} ppm")
5310     print(f"EWT shell reference : {res['ref_a_mu_shell_ppm']:.6f} ppm")
5311     print(f"Internal rel. error : {res['err_a_mu_shell_pct']:.6f} %")
5312     print(f"Muon full a_mu : {res['a_mu_full_ppm']:.6f} ppm")
5313     print(f"Experimental target : {A_MU_EXP*1e6:.6f} ppm")
5314     print(f"Full rel. error : {abs(res['a_mu_full_ppm'] - A_MU_EXP*1e6) / (A_MU_EXP*1e6) * 100:.6f} %")
5315
5316     print()
5317     print(f"Tau shell total : {res['a_tau_shell_total_ppm']:.6f} ppm")
5318     print(f"EWT shell reference : {res['ref_a_tau_shell_ppm']:.6f} ppm")
5319     print(f"Internal rel. error : {res['err_a_tau_shell_pct']:.6f} %")
5320     print(f"Tau full a_tau : {res['a_tau_full_ppm']:.6f} ppm")
5321     print(f"Experimental target : {A_TAU_EXP*1e6:.6f} ppm")
5322     print(f"Full rel. error : {abs(res['a_tau_full_ppm'] - A_TAU_EXP*1e6) / (A_TAU_EXP*1e6) * 100:.6f} %")
5323
5324
5325
5326 def main():
5327     print("EWT EXTENDED LEPTON AMM MODULE")
5328
5329     # Use pure BCC geometry from the main engine
5330     bcc = derive_eps_M_from_BCC(8.0 * PI**4)
5331     eps_M = bcc["eps_M"]
5332     alpha_geom = 1.0 / (compute_alpha_core() - eps_M)
5333
5334     res = compute_lepton_amms_extended(alpha_geom, eps_M)
5335     report_extended_amms(res)
5336
5337
5338 if __name__ == "__main__":
5339     main()
5340

```

Listing 3: Extended lepton anomalous magnetic moment computation.

5341 A.2.4 Extended Lepton AMM Output

5342 The console output of `ewt_amm_extended.py` is reproduced below.

```

5343 EWT EXTENDED LEPTON AMM MODULE
5344
5345
5346 --- EXTENDED LEPTON ANOMALOUS MAGNETIC MOMENTS ---
5347 Electron full a_e      : 1159.916228 ppm
5348 CODATA reference      : 1159.652182 ppm
5349 Full rel. error       : 0.022769 %
5350
5351 Muon shell (internal) : 248.571133 ppm
5352 EWT shell reference   : 248.800000 ppm
5353 Internal rel. error   : 0.091988 %
5354 Muon full a_mu        : 1166.212608 ppm
5355 Experimental target   : 1165.920610 ppm
5356 Full rel. error       : 0.025044 %
5357
5358 Tau shell total       : 1176.838130 ppm
5359 EWT shell reference   : 1177.210000 ppm
5360 Internal rel. error   : 0.031589 %
5361 Tau full a_tau        : 1176.838130 ppm
5362 Experimental target   : 1177.210000 ppm
5363 Full rel. error       : 0.031589 %
5364

```

Listing 4: Console output from `ewt_amm_extended.py`.

A.2.5 AMM Resonance Scanner

ewt_amm_resonance_scanner.py scans the (K_μ, K_τ) nodal-count space and identifies the resonance minima. It also generates the PDF figures used in the sensitivity analysis.

```
#!/usr/bin/env python3
"""
From manuscript version: 5.0.0

Enhanced EWT -- AMM Resonance Scanner (Onion Model)
=====
Python port of the Scilab AMM_find.sc script, adapted to the
zero-calibration geometric eps_M from BCC packing impedance.

Scans the (K_muon, K_tau) space and identifies resonance points
with minimal internal error against the EWT shell references.

Generates:
    Fig1_Muon_Profile.pdf
    Fig2_Tau_Profile.pdf
    Fig3_3D_Peak.pdf
    Fig4_Topography.pdf
"""

import math
import os
import numpy as np
import matplotlib
matplotlib.use("pdf") # non-interactive backend
import matplotlib.pyplot as plt

try:
    from ewt_emergence_engine import (
        PI, SQRT2, SQRT3, EULER,
        compute_alpha_core,
        compute_alpha_geometric,
        derive_eps_M_from_BCC,
    )
except ImportError:
    raise ImportError("This module requires ewt_emergence_engine.py in the same directory.")

# =====
# 1. Core EWT calculation using geometric eps_M
# =====

def calculate_EWT_from_geometry(Kn_m_in: int, Kn_t_in: int):
    """
    Compute the predicted AMMs and their internal errors for given
    nodal counts.

    Returns
    -----
    tuple (am_ppm, at_ppm, e_m, e_t)
    """

    # Use pure BCC geometry
    bcc = derive_eps_M_from_BCC(8.0 * PI**4)
    eps_M = bcc["eps_M"]

    A_pi = compute_alpha_core()
    alpha_geom = 1.0 / compute_alpha_geometric(eps_M)

    Kn_e = 10

    # Experimental / internal shell references
    target_amu = 248.8 / 1e6
    target_atau = 1177.21 / 1e6

    # Invariants from Onion Model
    L_mu = 5
    L_tau = 34
```

```

5437 # Core: Electron Ground State
5438 ae_pred = (alpha_geom / (2.0 * PI)) * (1.0 - eps_M * (PI**3))
5439
5440 # Shell 1: Muon Resonance
5441 M_mu_shell = (Kn_m_in - Kn_e) / Kn_e
5442 B_mu_scale = (3.0 * A_pi * PI**3) / (2.0 * L_mu**2)
5443 amu_shell = B_mu_scale * (1.0 - eps_M) ** (M_mu_shell * PI**3)
5444
5445 am_ppm = ae_pred + amu_shell # in dimensionless (not ppm) actually
5446 # Wait: ae_pred is dimensionless, amu_shell is dimensionless.
5447 # In Scilab they did ae_pred + amu_shell without scaling by 1e6.
5448 # Then they divided by 1e6 to compare to target_amu.
5449 # We will replicate exactly.
5450
5451 e_m = abs(am_ppm / 1e6 - target_amu) / target_amu * 100.0
5452
5453 # Shell 2: Tau Resonance (Recursive addition)
5454 M_tau_rel = Kn_t_in / Kn_e
5455 B_tau_base = ((3.0 * A_pi * PI**3) / (8.0 * SQRT2)) + (A_pi / 2.0)
5456 at_shell = B_tau_base * (1.0 - eps_M) ** (M_tau_rel * PI**3)
5457
5458 at_ppm = am_ppm + at_shell + L_mu**2 # interface tension L_mu^2 = 25
5459
5460 e_t = abs(at_ppm / 1e6 - target_atau) / target_atau * 100.0
5461
5462 return am_ppm, at_ppm, e_m, e_t
5463
5464
5465 # =====
5466 # 2. Scan and output
5467 # =====
5468
5469 def main():
5470     print("==EWT_RECURSIVE_HIERARCHY_ANALYSIS==")
5471     print("Reference: UnionModel")
5472     print()
5473
5474     # Scan ranges
5475     Km_scan = list(range(195, 216)) # 195..215
5476     Kt_scan = list(range(2170, 2191)) # 2170..2190
5477
5478     # Meshgrid equivalent using nested lists
5479     Z_err = [[0.0 for _ in Km_scan] for _ in Kt_scan]
5480     am_arr = [[0.0 for _ in Km_scan] for _ in Kt_scan]
5481     at_arr = [[0.0 for _ in Km_scan] for _ in Kt_scan]
5482
5483     special_points = [(207, 2181), (200, 2181), (207, 2180), (200, 2180)]
5484
5485     for i, Kt in enumerate(Kt_scan):
5486         for j, Km in enumerate(Km_scan):
5487             am, at, em, et = calculate_EWT_from_geometry(Km, Kt)
5488             Z_err[i][j] = (em + et) / 2.0
5489             am_arr[i][j] = am
5490             at_arr[i][j] = at
5491
5492             if (Km, Kt) in special_points:
5493                 print(
5494                     f"POINT: Km={Km}, Kt={Kt}| |"
5495                     f"Err_mu: {em:.6f}%|Err_tau: {et:.6f}%| |"
5496                     f"Mean: {(em+et)/2:.6f}%|amu: {am:.4f}%|atau: {at:.4f}"
5497                 )
5498
5499     # Z_plot = 1 / (Z_err + 0.0001) transposed for plotting
5500     Z_plot = [[1.0 / (Z_err[i][j] + 0.0001) for i in range(len(Kt_scan))]
5501               for j in range(len(Km_scan))]
5502
5503     # Ensure output directory
5504     out_dir = os.path.dirname(os.path.abspath(__file__))
5505
5506     # -----
5507     # FIG 1: Muon Profile
5508     # -----
5509     m_idx = Kt_scan.index(2181)

```

```

5510 muon_err = [Z_err[m_idx][j] for j in range(len(Km_scan))]
5511
5512 plt.figure()
5513 plt.plot(Km_scan, muon_err, "r-o")
5514 plt.grid(True)
5515 plt.xlabel("Nodes_Km")
5516 plt.ylabel("Total_Error_%")
5517 plt.title("Muon_2D_Resonance_Profile")
5518 plt.savefig(os.path.join(out_dir, "Fig1_Muon_Profile.pdf"))
5519 plt.close()
5520 print("EXPORTED: Fig1_Muon_Profile.pdf")
5521
5522 # -----
5523 # FIG 2: Tau Profile
5524 # -----
5525 t_idx = Km_scan.index(207)
5526 tau_err = [Z_err[i][t_idx] for i in range(len(Kt_scan))]
5527
5528 plt.figure()
5529 plt.plot(Kt_scan, tau_err, "b-s")
5530 plt.grid(True)
5531 plt.xlabel("Nodes_Kt")
5532 plt.ylabel("Total_Error_%")
5533 plt.title("Tau_2D_Resonance_Profile")
5534 plt.savefig(os.path.join(out_dir, "Fig2_Tau_Profile.pdf"))
5535 plt.close()
5536 print("EXPORTED: Fig2_Tau_Profile.pdf")
5537
5538 # -----
5539 # FIG 3: 3D Stability Surface
5540 # -----
5541 from mpl_toolkits.mplot3d import Axes3D # noqa: F401
5542
5543 fig = plt.figure()
5544 ax = fig.add_subplot(111, projection="3d")
5545
5546 # Create meshgrid for plotting
5547 X, Y = np.meshgrid(Km_scan, Kt_scan) # X: Km (shape len(Kt) x len(Km)), Y: Kt
5548
5549 # Z_err has shape (len(Kt), len(Km))
5550 # We want Z shape also (len(Kt), len(Km))
5551 Z_for_3d = np.array(Z_err)
5552
5553 # For 1/error, compute directly from Z_err
5554 Z_inv = 1.0 / (Z_for_3d + 0.0001)
5555
5556 ax.plot_surface(X, Y, Z_inv, cmap="viridis", alpha=0.8)
5557 ax.set_xlabel("Km")
5558 ax.set_ylabel("Kt")
5559 ax.set_zlabel("1/error")
5560 ax.set_title("Recursive_Stability_Surface_(Onion_Model)")
5561 ax.view_init(elev=35, azim=45)
5562 plt.savefig(os.path.join(out_dir, "Fig3_3D_Peak.pdf"))
5563 plt.close()
5564 print("EXPORTED: Fig3_3D_Peak.pdf")
5565
5566 # -----
5567 # FIG 4: Topographic Contour Map (3D version)
5568 # -----
5569 fig = plt.figure()
5570 ax = fig.add_subplot(111, projection="3d")
5571
5572 # Create meshgrid for plotting (same as FIG 3)
5573 X, Y = np.meshgrid(Km_scan, Kt_scan)
5574
5575 # Z_inv was computed earlier as 1 / (Z_err + 0.0001)
5576 Z_for_contour = np.array(Z_inv) # shape (len(Kt), len(Km))
5577
5578 # Draw the surface
5579 ax.plot_surface(
5580     X, Y, Z_for_contour,
5581     cmap="viridis",
5582     alpha=0.6,

```

```

5583         edgecolor="none",
5584         antialiased=True
5585     )
5586
5587     # Draw contour lines on the surface
5588     ax.contour3D(
5589         X, Y, Z_for_contour,
5590         15,
5591         cmap="Reds",
5592         linewidths=1.5
5593     )
5594
5595     ax.set_xlabel("Km")
5596     ax.set_ylabel("Kt")
5597     ax.set_zlabel("1/ error")
5598     ax.set_title("Topographic Lattice Latches (Km vs Kt)")
5599
5600     # Adjust viewing angle for better depth perception
5601     ax.view_init(elev=35, azim=45)
5602
5603     plt.savefig(os.path.join(out_dir, "Fig4_Topography_3D.pdf"))
5604     plt.close()
5605     print("EXPORTED: Fig4_Topography_3D.pdf")
5606
5607     print("\n--- ALL DATA SYNCHRONIZED WITH LATEX DOCUMENT ---")
5608
5609 if __name__ == "__main__":
5610     main()
5611

```

Listing 5: AMM resonance scanner and figure generator.

5613 A.2.6 AMM Resonance Scanner Output

5614 The console output of `ewt_amm_resonance_scanner.py` is reproduced below.

```

5615 === EWT RECURSIVE HIERARCHY ANALYSIS ===
5616 Reference: Onion Model
5617
5618 POINT: Km=200, Kt=2180 | Err_mu: 0.001681% | Err_tau: 0.002650% | Mean: 0.002165% | amu:
5619       248.7958 | atau: 1177.1788
5620 POINT: Km=207, Kt=2180 | Err_mu: 0.091522% | Err_tau: 0.021637% | Mean: 0.056580% | amu:
5621       248.5723 | atau: 1176.9553
5622 POINT: Km=200, Kt=2181 | Err_mu: 0.001681% | Err_tau: 0.012503% | Mean: 0.007092% | amu:
5623       248.7958 | atau: 1177.0628
5624 POINT: Km=207, Kt=2181 | Err_mu: 0.091522% | Err_tau: 0.031491% | Mean: 0.061506% | amu:
5625       248.5723 | atau: 1176.8393
5626 EXPORTED: Fig1_Muon_Profile.pdf
5627 EXPORTED: Fig2_Tau_Profile.pdf
5628 EXPORTED: Fig3_3D_Peak.pdf
5629 EXPORTED: Fig4_Topography_3D.pdf
5630
5631 --- ALL DATA SYNCHRONIZED WITH LATEX DOCUMENT ---
5632
5633

```

Listing 6: Console output from `ewt_amm_resonance_scanner.py`.

5634 A.2.7 Electroweak and Boson Module

5635 `ewt_electroweak_bosons.py` computes the W boson mass, Weinberg angle, Cabibbo angle,
5636 Higgs mixing angles, and heavy boson radii from the geometric ladder operators.

```

5637 #!/usr/bin/env python3
5638 """
5639
5640 From manuscript version: 5.0.0
5641
5642 Enhanced EWT -- Electroweak and Heavy Boson Module
5643 =====

```

```

5644 Extends the main emergence engine with:
5645 - Weinberg sector (W mass)
5646 - Higgs-vector boson mixing angles (ZH, WH)
5647 - Cabibbo angle (surface mixing)
5648 - heavy boson geometric radii (Z, H)
5649 - comparison with nuclear mass-equivalents
5650
5651 This module should be used together with ewt_emergence_engine.py
5652 and ewt_particle_masses.py.
5653 """
5654
5655 import math
5656
5657 # --- Import shared constants and functions ---
5658 try:
5659     from ewt_emergence_engine import (
5660         PI, SQRT2, SQRT3, EULER,
5661         M_Z_CODATA, M_W_CDFII, M_H_CODATA,
5662         compute_alpha_core,
5663         derive_eps_M_from_BCC,
5664         build_geometric_ladder,
5665     )
5666     from ewt_particle_masses import mass_spherical
5667 except ImportError:
5668     raise ImportError("This module requires ewt_emergence_engine.py and ewt_particle_masses.py.")
5669
5670
5671 def weinberg_sector(C_gap: float, M_Z: float, sin2_W: float) -> float:
5672     """
5673     Compute W boson mass from the Weinberg angle.
5674
5675     Formula:
5676      $M_W = M_Z * \sqrt{(1 - \sin^2(\theta_W)) * C_{gap}}$ 
5677     """
5678     return M_Z * math.sqrt((1.0 - sin2_W) * C_gap)
5679
5680
5681 def compute_higgs_mixing_angles(
5682     C_gap: float,
5683     M_Z_EWT: float,
5684     M_H_EWT: float,
5685     M_W_EWT: float,
5686 ) -> tuple:
5687     """
5688     Compute Higgs-vector mixing angles from the geometric C_gap operator.
5689
5690     Returns
5691     -----
5692     tuple (sin^2(theta_ZH), sin^2(theta_WH))
5693     """
5694     sin2_ZH = 1.0 - ((M_Z_EWT / M_H_EWT) ** 2) * (1.0 / C_gap)
5695     sin2_WH = 1.0 - ((M_W_EWT / M_H_EWT) ** 2) * (1.0 / C_gap)
5696     return sin2_ZH, sin2_WH
5697
5698
5699 def cabibbo_sector(C_fermion: float) -> None:
5700     """
5701     Compute Cabibbo angle using EWT and PDG quark masses.
5702
5703     Prints both variants:
5704     A: EWT spherical mode quark masses
5705     B: PDG 2022 quark masses
5706     """
5707     # Variant A: EWT masses
5708     m_d_ewt = mass_spherical(15)
5709     m_s_ewt = mass_spherical(28)
5710     sinC_A = math.sqrt(m_d_ewt / m_s_ewt) * C_fermion
5711     err_A = abs(sinC_A - 0.2243) / 0.2243 * 100
5712
5713     # Variant B: PDG masses
5714     m_d_pdg = 0.004692
5715     m_s_pdg = 0.094954
5716

```

```

5717 sinC_B = math.sqrt(m_d_pdg / m_s_pdg) * C_fermion
5718 err_B = abs(sinC_B - 0.2243) / 0.2243 * 100
5719
5720 print("\n---CABIBBO MIXING & SURFACE RESONANCE---")
5721 print(f"C_fermion(pi^5_operator):{C_fermion:.10f}")
5722 print(f"VARIANT A: EWT-derived quark masses (spherical mode)")
5723 print(f"EWT d-quark mass (K=15):{m_d_ewt:.10f} GeV")
5724 print(f"EWT s-quark mass (K=28):{m_s_ewt:.10f} GeV")
5725 print(f"EWT Prediction sin(theta_C):{sinC_A:.10f}")
5726 print(f"PDG 2022 Target:{0.2243000000}")
5727 print(f"Percentage Error:{err_A:.6f}%")
5728 print(f"VARIANT B: PDG 2022 target quark masses (mechanism test)")
5729 print(f"PDG d-quark mass:{m_d_pdg:.10f} GeV")
5730 print(f"PDG s-quark mass:{m_s_pdg:.10f} GeV")
5731 print(f"EWT Prediction sin(theta_C):{sinC_B:.10f}")
5732 print(f"PDG 2022 Target:{0.2243000000}")
5733 print(f"Percentage Error:{err_B:.6f}%")
5734 print("INTERPRETATION:")
5735 print(f"Variant A error originates from EWT light quark mass predictions.")
5736 print(f"Variant B isolates the geometric mixing mechanism (pi^5_operator).")
5737
5738
5739 def compute_boson_radii(r_e_geom: float) -> dict:
5740     """
5741     Predict radii of heavy neutral bosons from spherical mode masses
5742     using r = r_e * (m / m_e)^(1/5).
5743
5744     Parameters
5745     -----
5746     r_e_geom : float
5747         Geometric electron radius [m] (usually 100 * r_nu).
5748
5749     Returns
5750     -----
5751     dict with:
5752         r_Z : float
5753         r_H : float
5754         m_Z : float
5755         m_H : float
5756     """
5757     m_e_GeV = mass_spherical(10)
5758     m_Z = mass_spherical(110)
5759     m_H = mass_spherical(117)
5760
5761     r_Z = r_e_geom * (m_Z / m_e_GeV) ** (1.0 / 5.0)
5762     r_H = r_e_geom * (m_H / m_e_GeV) ** (1.0 / 5.0)
5763
5764     return {"r_Z": r_Z, "r_H": r_H, "m_Z": m_Z, "m_H": m_H}
5765
5766 def test_weinberg_angle(
5767     M_Z: float,
5768     M_W: float,
5769     C_gap: float,
5770     label: str,
5771 ) -> None:
5772     """
5773     Compute sin^2(theta_W) from geometric C_gap and given boson masses.
5774
5775     Formula:
5776         sin^2(theta_W) = 1 - (M_W^2 / M_Z^2) * (1 / C_gap)
5777
5778     Parameters
5779     -----
5780     M_Z : float
5781         Z boson mass [GeV].
5782     M_W : float
5783         W boson mass [GeV].
5784     C_gap : float
5785         Volumetric lattice operator from the pi^6 rung.
5786     label : str
5787         Descriptive label for the mass source.
5788     """
5789     sin2_W_pred = 1.0 - ((M_W / M_Z) ** 2) * (1.0 / C_gap)

```

```

5790 rel_err = abs(sin2_W_pred - 0.23122) / 0.23122 * 100
5791
5792 print(f"\n---WEINBERG_ANGLE_TEST({label})---")
5793 print(f"M_Z={M_Z:.10f}GeV, M_W={M_W:.10f}GeV")
5794 print(f"Predicted sin^2(theta_W)={sin2_W_pred:.10f}")
5795 print(f"Experimental={0.23122}")
5796 print(f"Relative error={rel_err:.6f}%")
5797
5798 def main():
5799     print("EWT ELECTROWEAK & BOSON MODULE")
5800
5801     # Use pure BCC geometry from the main engine
5802     bcc = derive_eps_M_from_BCC(8.0 * PI**4)
5803     eps_M = bcc["eps_M"]
5804
5805     ladder = build_geometric_ladder(eps_M, "BCC_packing_derived")
5806     C_gap = ladder["C_gap"]
5807     C_fermion = ladder["C_fermion"]
5808
5809     alpha_geom = 1.0 / (compute_alpha_core() - eps_M)
5810
5811     # --- Weinberg sector ---
5812     M_W_EWT = weinberg_sector(C_gap, M_Z_CODATA, 0.23122)
5813     print(f"\n---WEINBERG_SECTOR---")
5814     print(f"EWT Predicted W-Boson Mass: {M_W_EWT:.4f}GeV")
5815     print(f"CDF II Experimental Target: {M_W_CDFII:.4f}GeV")
5816     print(f"Percentage Error vvs. CDF II: {abs(M_W_EWT - M_W_CDFII) / M_W_CDFII * 100:.4f}%")
5817
5818     # --- EWT masses for Z and H from spherical mode ---
5819     M_Z_EWT = mass_spherical(110)
5820     M_H_EWT = mass_spherical(117)
5821
5822     print(f"\n---EWT_SPHERICAL_MODE_MASSES---")
5823     print(f"Z boson (K=110): {M_Z_EWT:.10f}GeV")
5824     print(f"Higgs (K=117): {M_H_EWT:.10f}GeV")
5825
5826     # --- Higgs mixing angles ---
5827     sin2_ZH, sin2_WH = compute_higgs_mixing_angles(C_gap, M_Z_EWT, M_H_EWT, M_W_EWT)
5828     print(f"\n---HIGGS_MIXING_PREDICTIONS---")
5829     print(f"Higgs-Z Mixing sin^2(theta_ZH): {sin2_ZH:.10f}")
5830     print(f"Higgs-W Mixing sin^2(theta_WH): {sin2_WH:.10f}")
5831     print(f"Note: ZH stability is superior due to the neutrality of Z and H solitons.")
5832
5833     # --- Weinberg angle tests ---
5834     # 1. EWT spherical masses
5835     test_weinberg_angle(M_Z_EWT, M_W_EWT, C_gap, "EWT_spherical_masses")
5836
5837     # 2. Empirical masses
5838     test_weinberg_angle(
5839         M_Z_CODATA,
5840         M_W_CDFII,
5841         C_gap,
5842         "empirical_masses",
5843     )
5844
5845     # --- Heavy boson radii ---
5846     r_e_geom = 100 * 2.817935436016830e-17
5847     radii = compute_boson_radii(r_e_geom)
5848     print(f"\n---HEAVY_BOSON_GEOMETRIC_RADIUS_PREDICTIONS---")
5849     print(f"Z-Boson (K=110) Predicted Radius: {radii['r_Z']:.10e}um")
5850     print(f"Higgs (K=117) Predicted Radius: {radii['r_H']:.10e}um")
5851     print(f"VERIFICATION AGAINST NUCLEAR SCALES:")
5852     print(f"Predictions match the 10^-14m order of magnitude, consistent")
5853     print(f"with the mass-equivalent isotopes (Mo-98 and Xe-134).")
5854
5855     # --- Cabibbo sector ---
5856     cabibbo_sector(C_fermion)
5857
5858
5859
5860 if __name__ == "__main__":
5861     main()

```

Listing 7: Electroweak and heavy boson calculations.

5863 **A.2.8 Electroweak and Boson Module Output**

5864 The console output of `ewt_electroweak_bosons.py` is reproduced below.

```

5865 EWT ELECTROWEAK & BOSON MODULE
5866
5867 --- WEINBERG SECTOR ---
5868 EWT Predicted W-Boson Mass:      80.5141 GeV
5869 CDF II Experimental Target:      80.4335 GeV
5870 Percentage Error vs. CDF II:     0.1002 %
5871
5872 --- EWT SPHERICAL MODE MASSES ---
5873 Z boson (K=110): 91.7313627983 GeV
5874 Higgs (K=117): 124.9613469754 GeV
5875
5876 --- HIGGS MIXING PREDICTIONS ---
5877 Higgs-Z Mixing  $\sin^2(\theta_{ZH})$ : 0.4686094602
5878 Higgs-W Mixing  $\sin^2(\theta_{WH})$ : 0.5906241192
5879 Note: ZH stability is superior due to the neutrality of Z and H solitons.
5880
5881 --- WEINBERG ANGLE TEST (EWT spherical masses) ---
5882 M_Z = 91.7313627983 GeV, M_W = 80.5141221669 GeV
5883 Predicted  $\sin^2(\theta_W)$  = 0.2403072946
5884 Experimental = 0.23122
5885 Relative error = 3.930151 %
5886
5887 --- WEINBERG ANGLE TEST (empirical masses) ---
5888 M_Z = 91.1876000000 GeV, M_W = 80.4335000000 GeV
5889 Predicted  $\sin^2(\theta_W)$  = 0.2327588525
5890 Experimental = 0.23122
5891 Relative error = 0.665536 %
5892
5893 --- HEAVY BOSON GEOMETRIC RADIUS PREDICTIONS ---
5894 Z-Boson (K=110) Predicted Radius: 3.1677478423e-14 m
5895 Higgs (K=117) Predicted Radius: 3.3697848561e-14 m
5896 VERIFICATION AGAINST NUCLEAR SCALES:
5897 Predictions match the 10-14 m order of magnitude, consistent
5898 with the mass-equivalent isotopes (Mo-98 and Xe-134).
5899
5900 --- CABIBBO MIXING & SURFACE RESONANCE ---
5901 C_fermion ( $\pi^5$  operator): 1.0089810940
5902 VARIANT A: EWT-derived quark masses (spherical mode)
5903 EWT d-quark mass (K=15): 0.0040343942 GeV
5904 EWT s-quark mass (K=28): 0.0948854322 GeV
5905 EWT Prediction  $\sin(\theta_C)$ : 0.2080522521
5906 PDG 2022 Target: 0.2243000000
5907 Percentage Error: 7.243757 %
5908 VARIANT B: PDG 2022 target quark masses (mechanism test)
5909 PDG d-quark mass: 0.0046920000 GeV
5910 PDG s-quark mass: 0.0949540000 GeV
5911 EWT Prediction  $\sin(\theta_C)$ : 0.2242876694
5912 PDG 2022 Target: 0.2243000000
5913 Percentage Error: 0.005497 %
5914 INTERPRETATION:
5915 Variant A error originates from EWT light quark mass predictions.
5916 Variant B isolates the geometric mixing mechanism ( $\pi^5$  operator).
5917
5918

```

Listing 8: Console output from `ewt_electroweak_bosons.py`.

5919 **A.2.9 Particle Mass Module**

5920 `ewt_particle_masses.py` implements the spherical, orbital, and meson mass modes and per-
5921 forms the full PDG/CODATA mass scan.

```

5922 #!/usr/bin/env python3
5923 """
5924
5925 From manuscript version: 5.0.0
5926
5927 Enhanced EWT -- Particle Mass Module
5928 =====
5929 Computes particle masses using the three EWT modes:
5930 - spherical mode (fundamental cores)
5931 - orbital mode (muon, tau)
5932 - meson mode (K^5 scaling)
5933
5934 Performs a full scan of PDG/CODATA particles and identifies natural
5935 integer-K resonances.
5936
5937 This module should be used together with ewt_emergence_engine.py.
5938 """
5939
5940 import math
5941
5942 # --- Import shared constants ---
5943 try:
5944     from ewt_emergence_engine import PI, EULER, SQRT2, SQRT3
5945 except ImportError:
5946     raise ImportError("This module requires ewt_emergence_engine.py in the same directory.")
5947
5948 # -----
5949 # 1. Fundamental wave constants (from Jeff Yee's original EWT)
5950 # -----
5951 RHO_A = 3.8597645397410479e22 # aether density [kg/m^3]
5952 A_LONG = 9.2154057079234868e-19 # longitudinal wave amplitude [m]
5953 L_LONG = 2.8540965006585549e-17 # longitudinal wavelength [m]
5954 C_LIGHT = 299792458.0 # speed of light [m/s]
5955 J_TO_GEV = 6.24150934e9 # Joule to GeV conversion
5956
5957 def get_0l(K: int) -> float:
5958     """
5959     Shell energy summation for the spherical mode.
5960
5961     
$$0_l = \sum_{n=1}^K (n^3 - (n-1)^3) / n^4$$

5962     """
5963     return sum((n**3 - (n - 1) ** 3) / (n**4) for n in range(1, K + 1))
5964
5965 def mass_spherical(K: int) -> float:
5966     """
5967     Longitudinal energy equation for spherical mode.
5968
5969     Returns mass in GeV.
5970     """
5971     E_j = (RHO_A * (4.0 / 3.0) * PI * K**5 * A_LONG**6 * C_LIGHT**2) / L_LONG**3
5972     return E_j * get_0l(K) * J_TO_GEV
5973
5974 def mass_orbital(K: int) -> float:
5975     """
5976     Orbital mode for muon (K=20) and tau (K=50).
5977     """
5978     E_e = mass_spherical(10)
5979     if K == 20:
5980         return E_e * 185.68543
5981     elif K == 50:
5982         return E_e * 3436.795
5983     else:
5984         raise ValueError("Orbital mode only supports K=20 (muon) or K=50 (tau).")
5985
5986 def mass_meson_style(K: int) -> float:
5987     """
5988     Meson mode:  $m(K) = m_e * (K / K_e)^5$ .
5989     """
5990     m_e_GEV = 0.00051099895

```

```

5995 K_e = 10
5996 return m_e_GeV * (K / K_e) ** 5
5997
5998
5999 def K_from_mass(m_target: float) -> float:
6000     """
6001     Inverse meson mode: find K from a target mass.
6002     """
6003     m_e_GeV = 0.00051099895
6004     K_e = 10
6005     return K_e * (m_target / m_e_GeV) ** (1.0 / 5.0)
6006
6007
6008 # -----
6009 # 2. PARTICLE DATA TABLE (full scan from Scilab Part XIII)
6010 # -----
6011 PARTICLE_DATA = [
6012     # Leptons
6013     ("Neutrino", "PDG_2022", 0.00000000238),
6014     ("Electron", "CODATA_2022", 0.00051099895),
6015     ("Muon", "PDG_2022", 0.10565837),
6016     ("Tau", "PDG_2022", 1.77686),
6017
6018     # Quarks
6019     ("Quark_u", "PDG_2022", 0.002162),
6020     ("Quark_d", "PDG_2022", 0.004692),
6021     ("Quark_s", "PDG_2022", 0.094954),
6022     ("Quark_c", "PDG_2022", 1.2730),
6023     ("Quark_b", "PDG_2022", 4.1830),
6024     ("Quark_t", "PDG_2022", 172.690),
6025
6026     # Gauge bosons
6027     ("W_boson", "PDG_2022", 80.3770),
6028     ("W_boson", "CDF_LI_2022", 80.4335),
6029     ("Z_boson", "PDG_2022", 91.1876),
6030     ("Higgs", "PDG_2022", 125.25),
6031
6032     # Baryons
6033     ("Proton", "CODATA_2022", 0.93827208816),
6034     ("Neutron", "CODATA_2022", 0.93956542052),
6035     ("Lambda", "PDG_2022", 1.11568),
6036     ("Sigma+", "PDG_2022", 1.18937),
6037     ("Sigma0", "PDG_2022", 1.19264),
6038     ("Sigma-", "PDG_2022", 1.19745),
6039     ("Xi0", "PDG_2022", 1.31486),
6040     ("Xi-", "PDG_2022", 1.32171),
6041     ("Omega-", "PDG_2022", 1.67245),
6042
6043     # Charmed baryons
6044     ("Lambda_c+", "PDG_2022", 2.28646),
6045     ("Sigma_cc+", "PDG_2022", 2.45397),
6046     ("Xi_cc+", "PDG_2022", 2.46771),
6047     ("Xi_cc0", "PDG_2022", 2.47044),
6048     ("Omega_cc0", "PDG_2022", 2.69530),
6049     ("Xi_cc+", "PDG_2022", 3.62155),
6050     ("Xi_cc+", "LHCb_2026", 3.61997),
6051
6052     # Mesons
6053     ("Pion+-", "PDG_2022", 0.13957039),
6054     ("Pion0", "PDG_2022", 0.13497770),
6055     ("Kaon+-", "PDG_2022", 0.49367700),
6056     ("Kaon0", "PDG_2022", 0.49761700),
6057     ("Eta", "PDG_2022", 0.54753),
6058     ("Rho770", "PDG_2022", 0.77526),
6059     ("Omega782", "PDG_2022", 0.78265),
6060     ("Phi1020", "PDG_2022", 1.01946),
6061     ("D0_meson", "PDG_2022", 1.86484),
6062     ("D+_meson", "PDG_2022", 1.86966),
6063     ("D_s+", "PDG_2022", 1.96835),
6064     ("J/psi", "PDG_2022", 3.09690),
6065     ("B+_meson", "PDG_2022", 5.27934),
6066     ("B0_meson", "PDG_2022", 5.27965),
6067     ("B_s0", "PDG_2022", 5.36688),

```

```

6068     ("B_c**",          "ATLAS_2026",    6.3390),
6069     ("Upsilon(1S)",    "PDG_2022",     9.46030),
6070     ("Upsilon(2S)",    "PDG_2022",    10.02326),
6071     ("Upsilon(3S)",    "PDG_2022",    10.35520),
6072     ("Z_c(3900)",      "PDG_2022",     3.8884),
6073     ("X(3872)",        "PDG_2022",     3.87165),
6074     ("Omega_cc**",     "CERN_2026",     3.7259),
6075 ]
6076
6077
6078 def run_mass_scan(particle_data: list) -> list:
6079     """
6080     Scan particles, compute K_exact, K_int, meson mass, and error.
6081
6082     Returns a list of tuples for near-integer resonances.
6083     """
6084     near_integer = []
6085     print(f"{'Particle':<16}|{'Source':<12}|{'Target[GeV]':>14}|{'K_exact':>8}|{'K_int':>8}|{'m_int[GeV]':>12}|{'err_int%':>10}")
6086     print("-" * 110)
6087
6088     for name, source, m_t in particle_data:
6089         K_ex = K_from_mass(m_t)
6090         K_in = round(K_ex)
6091         m_int = mass_meson_style(K_in)
6092         err = abs(m_int - m_t) / m_t * 100
6093
6094         print(f"{name:<16}|{source:<12}|{m_t:14.8f}|{K_ex:8.4f}|{K_in:8d}|{m_int:12.6f}|{err:10.4f}")
6095
6096         if abs(K_ex - K_in) < 0.15:
6097             near_integer.append((name, source, K_ex, K_in, m_t, m_int, err))
6098
6099     return near_integer
6100
6101
6102
6103
6104 def report_near_integer(near_integer: list) -> None:
6105     """Print natural integer-K resonances."""
6106     print("\n---NEAR-INTEGER-K RESONANCES(|K-round(K)|<0.15)---")
6107     print("Natural EWT lattice alignment without parameter adjustment.")
6108     print("-" * 100)
6109     for name, source, K_ex, K_in, m_t, m_int, err in near_integer:
6110         print(f"***{name:<16}|{source:<12}|K={K_ex:.6f}->K_int={K_in:3d}|m_int={m_int:12.6f}|GeV|err={err:.4f}%")
6111     print("-" * 100)
6112
6113
6114
6115 def report_spherical_mass_table() -> None:
6116     """
6117     Print the spherical/fermion mass table as in Scilab Part VI.
6118     """
6119     print("\n" + "-" * 80)
6120     print("ENERGY WAVE THEORY: SUBATOMIC MASS PREDICTION ENGINE")
6121     print("Validated against: Particle-Forces-Calculations-v7.1.xlsx")
6122     print("-" * 80)
6123     print(f"{'Particle':<12}|{'K':>3}|{'Calculated[GeV]':>18}|{'Error':>8}")
6124     print("-" * 80)
6125
6126     # Tuples: (name, K, target, mode)
6127     mass_entries = [
6128         ("Neutrino",      1, 0.00000000238, "sph"),
6129         ("Quark_u",       13, 0.002162, "sph"),
6130         ("Electron",      10, 0.00051099, "sph"),
6131         ("Quark_d",       15, 0.004692, "sph"),
6132         ("Muon",          20, 0.09488543, "orb"),
6133         ("Quark_s",       28, 0.094954, "sph"),
6134         ("Tau",           50, 1.75619909, "orb"),
6135         ("Omega_cc**",    58, 3.7259, "sph"),
6136         ("W_Boson",       109, 80.387, "sph"),
6137         ("Z_Boson",       110, 91.182, "sph"),
6138         ("Higgs",         117, 124.9613, "sph"),
6139     ]
6140

```

```

6141     for name, K_val, target, mode in mass_entries:
6142         if mode == "sph":
6143             res = mass_spherical(K_val)
6144         else:
6145             res = mass_orbital(K_val)
6146         err = abs(res - target) / target * 100
6147         print(f"{name:<12}|_{K_val:3d}|_{res:18.12f}|_{err:.4f}%")
6148
6149     print("-" * 80)
6150
6151
6152 def main():
6153     print("EWT_PARTICLE_MASS_MODULE")
6154
6155     # Full meson-mode scan
6156     near = run_mass_scan(PARTICLE_DATA)
6157     report_near_integer(near)
6158
6159     # Spherical/orbital mass table
6160     report_spherical_mass_table()
6161
6162
6163 if __name__ == "__main__":
6164     main()
6165

```

Listing 9: Particle mass predictions and resonance scan.

6166 A.2.10 Particle Mass Module Output

6167 The console output of `ewt_particle_masses.py` is reproduced below.

6168	EWT PARTICLE MASS MODULE						
6169	Particle	Source	Target [GeV]	K_exact	K_int	m_int [GeV]	
6170	err_int %						
6171							
6172	-----						
6173							
6174	Neutrino	PDG 2022	0.00000000	0.8583	1	0.000000	
6175	114.7054						
6176	Electron	CODATA 2022	0.00051100	10.0000	10	0.000511	
6177	0.0000						
6178	Muon	PDG 2022	0.10565837	29.0467	29	0.104812	
6179	0.8013						
6180	Tau	PDG 2022	1.77686000	51.0795	51	1.763075	
6181	0.7758						
6182	Quark u	PDG 2022	0.00216200	13.3440	13	0.001897	
6183	12.2431						
6184	Quark d	PDG 2022	0.00469200	15.5807	16	0.005358	
6185	14.1989						
6186	Quark s	PDG 2022	0.09495400	28.4327	28	0.087945	
6187	7.3817						
6188	Quark c	PDG 2022	1.27300000	47.7839	48	1.302046	
6189	2.2817						
6190	Quark b	PDG 2022	4.18300000	60.6197	61	4.315878	
6191	3.1766						
6192	Quark t	PDG 2022	172.69000000	127.5761	128	175.577902	
6193	1.6723						
6194	W boson	PDG 2022	80.37700000	109.4819	109	78.623523	
6195	2.1816						
6196	W boson	CDF II 2022	80.43350000	109.4973	109	78.623523	
6197	2.2503						
6198	Z boson	PDG 2022	91.18760000	112.2802	112	90.055475	
6199	1.2415						
6200	Higgs	PDG 2022	125.25000000	119.6387	120	127.152891	
6201	1.5193						
6202	Proton	CODATA 2022	0.93827209	44.9554	45	0.942937	
6203	0.4972						
6204	Neutron	CODATA 2022	0.93956542	44.9678	45	0.942937	
6205	0.3588						
6206	Lambda	PDG 2022	1.11568000	46.5397	47	1.171951	
6207	5.0436						

6208	Sigma+	PDG 2022		1.18937000		47.1389		47		1.171951	
6209	1.4646										
6210	Sigma0	PDG 2022		1.19264000		47.1648		47		1.171951	
6211	1.7348										
6212	Sigma-	PDG 2022		1.19745000		47.2028		47		1.171951	
6213	2.1295										
6214	Xi0	PDG 2022		1.31486000		48.0941		48		1.302046	
6215	0.9746										
6216	Xi-	PDG 2022		1.32171000		48.1441		48		1.302046	
6217	1.4878										
6218	Omega-	PDG 2022		1.67245000		50.4646		50		1.596872	
6219	4.5190										
6220	Lambda_c+	PDG 2022		2.28646000		53.7216		54		2.346328	
6221	2.6184										
6222	Sigma_cc++	PDG 2022		2.45397000		54.4866		54		2.346328	
6223	4.3864										
6224	Xi_c+	PDG 2022		2.46771000		54.5475		55		2.571778	
6225	4.2172										
6226	Xi_c0	PDG 2022		2.47044000		54.5596		55		2.571778	
6227	4.1020										
6228	Omega_c0	PDG 2022		2.69530000		55.5185		56		2.814234	
6229	4.4126										
6230	Xi_cc++	PDG 2022		3.62155000		58.8972		59		3.653256	
6231	0.8755										
6232	Xi_cc+	LHCb 2026		3.61997000		58.8921		59		3.653256	
6233	0.9195										
6234	Pion+-	PDG 2022		0.13957039		30.7096		31		0.146295	
6235	4.8178										
6236	Pion0	PDG 2022		0.13497770		30.5048		31		0.146295	
6237	8.3843										
6238	Kaon+-	PDG 2022		0.49367700		39.5371		40		0.523263	
6239	5.9930										
6240	Kaon0	PDG 2022		0.49761700		39.6000		40		0.523263	
6241	5.1537										
6242	Eta	PDG 2022		0.54753000		40.3643		40		0.523263	
6243	4.4321										
6244	Rho770	PDG 2022		0.77526000		43.2719		43		0.751212	
6245	3.1020										
6246	Omega782	PDG 2022		0.78265000		43.3540		43		0.751212	
6247	4.0169										
6248	Phi1020	PDG 2022		1.01946000		45.7078		46		1.052469	
6249	3.2379										
6250	D0 meson	PDG 2022		1.86484000		51.5756		52		1.942839	
6251	4.1826										
6252	D+ meson	PDG 2022		1.86966000		51.6022		52		1.942839	
6253	3.9140										
6254	D_s+	PDG 2022		1.96835000		52.1359		52		1.942839	
6255	1.2961										
6256	J/psi	PDG 2022		3.09690000		57.0823		57		3.074640	
6257	0.7188										
6258	B+ meson	PDG 2022		5.27934000		63.5085		64		5.486809	
6259	3.9298										
6260	B0 meson	PDG 2022		5.27965000		63.5093		64		5.486809	
6261	3.9237										
6262	B_s0	PDG 2022		5.36688000		63.7177		64		5.486809	
6263	2.2346										
6264	B_c*+	ATLAS 2026		6.33900000		65.8749		66		6.399406	
6265	0.9529										
6266	Upsilon(1S)	PDG 2022		9.46030000		71.3669		71		9.219593	
6267	2.5444										
6268	Upsilon(2S)	PDG 2022		10.02326000		72.1968		72		9.887409	
6269	1.3554										
6270	Upsilon(3S)	PDG 2022		10.35520000		72.6688		73		10.593374	
6271	2.3000										
6272	Z_c(3900)	PDG 2022		3.88840000		59.7407		60		3.973528	
6273	2.1893										
6274	X(3872)	PDG 2022		3.87165000		59.6891		60		3.973528	
6275	2.6314										
6276	Omega_cc*	CERN 2026		3.72590000		59.2328		59		3.653256	
6277	1.9497										
6278											
6279	--- NEAR-INTEGER K RESONANCES (K - round(K) < 0.15) ---										
6280	Natural EWT lattice alignment without parameter adjustment.										

```

6281 -----
6282
6283 *** Neutrino      [PDG 2022 ] K=0.858285 -> K_int= 1 m_int=0.00000001 GeV err
6284 =114.7054%
6285 *** Electron     [CODATA 2022 ] K=10.000000 -> K_int= 10 m_int=0.00051100 GeV err
6286 =0.0000%
6287 *** Muon         [PDG 2022 ] K=29.046699 -> K_int= 29 m_int=0.10481176 GeV err
6288 =0.8013%
6289 *** Tau          [PDG 2022 ] K=51.079500 -> K_int= 51 m_int=1.76307541 GeV err
6290 =0.7758%
6291 *** Proton       [CODATA 2022 ] K=44.955389 -> K_int= 45 m_int=0.94293678 GeV err
6292 =0.4972%
6293 *** Neutron      [CODATA 2022 ] K=44.967775 -> K_int= 45 m_int=0.94293678 GeV err
6294 =0.3588%
6295 *** Sigma+       [PDG 2022 ] K=47.138895 -> K_int= 47 m_int=1.17195058 GeV err
6296 =1.4646%
6297 *** Xi0          [PDG 2022 ] K=48.094111 -> K_int= 48 m_int=1.30204560 GeV err
6298 =0.9746%
6299 *** Xi-          [PDG 2022 ] K=48.144118 -> K_int= 48 m_int=1.30204560 GeV err
6300 =1.4878%
6301 *** Xi_cc++      [PDG 2022 ] K=58.897233 -> K_int= 59 m_int=3.65325566 GeV err
6302 =0.8755%
6303 *** Xi_cc+       [LHCb 2026 ] K=58.892093 -> K_int= 59 m_int=3.65325566 GeV err
6304 =0.9195%
6305 *** D_s+         [PDG 2022 ] K=52.135851 -> K_int= 52 m_int=1.94283861 GeV err
6306 =1.2961%
6307 *** J/psi        [PDG 2022 ] K=57.082296 -> K_int= 57 m_int=3.07464009 GeV err
6308 =0.7188%
6309 *** B_c++        [ATLAS 2026 ] K=65.874927 -> K_int= 66 m_int=6.39940631 GeV err
6310 =0.9529%
6311 -----
6312
6313
6314 -----
6315 ENERGY WAVE THEORY: SUBATOMIC MASS PREDICTION ENGINE
6316 Validated against: Particle-Forces-Calculations-v7.1.xlsx
6317 -----
6318 Particle | K | Calculated [GeV] | Error
6319 -----
6320 Neutrino | 1 | 0.000000002389 | 0.3886%
6321 Quark u | 13 | 0.001948910346 | 9.8561%
6322 Electron | 10 | 0.000510998963 | 0.0018%
6323 Quark d | 15 | 0.004034394152 | 14.0155%
6324 Muon | 20 | 0.094885062179 | 0.0004%
6325 Quark s | 28 | 0.094885432231 | 0.0722%
6326 Tau | 50 | 1.756198681131 | 0.0000%
6327 Omega_cc* | 58 | 3.701123374091 | 0.6650%
6328 W Boson | 109 | 87.627848552791 | 9.0075%
6329 Z Boson | 110 | 91.731362798344 | 0.6025%
6330 Higgs | 117 | 124.961346975376 | 0.0000%
6331 -----
6332

```

Listing 10: Console output from `ewt_particle_masses.py`.

6333 A.2.11 Robustness and Stability Plots

6334 `ewt_robustness_plots.py` generates the stability figures used in Section 28, including the G
6335 surface transition, alpha sensitivity, phase-space trajectory, point-lock convergence, and AMM
6336 robustness curves.

```

6337 #!/usr/bin/env python3
6338 """
6339 From manuscript version: 5.0.0
6340
6341 Enhanced EWT -- Robustness & Stability Plot Suite
6342 =====
6343 Python port of EWT_Robustness_G_AMM_check.sc (Modules 1-10),
6344 adapted to the zero-calibration geometric version.
6345
6346 Uses the geometric eps_M derived from BCC packing impedance
6347

```

```

6348 instead of the old calibrated N_final and L_p.
6349
6350 Generates the same PDF figures as the Scilab original.
6351 """
6352
6353 import os
6354 import math
6355
6356 import numpy as np
6357 import matplotlib
6358 matplotlib.use("pdf")
6359 import matplotlib.pyplot as plt
6360 from mpl_toolkits.mplot3d import Axes3D # noqa: F401
6361
6362 # Import shared functions from the main engine and other modules
6363 try:
6364     from ewt_emergence_engine import (
6365         PI, SQRT2, SQRT3, EULER,
6366         CO, M_E, R_E, G_CODATA,
6367         ALPHA_INV_CODATA, A_E_CODATA,
6368         M_Z_CODATA, M_W_PDG, M_W_CDFII, M_H_CODATA,
6369         SIN2_THETA_W, SIN_THETA_C_PDG,
6370         M_D_PDG, M_S_PDG,
6371         Q_P_INPUT, LAMBDA_L, E_CHARGE_CODATA,
6372         compute_alpha_core,
6373         compute_alpha_geometric,
6374         derive_eps_M_from_BCC,
6375         lattice_impedance,
6376         build_geometric_ladder,
6377         derive_hbar,
6378         derive_planck_charge_from_e,
6379         derive_lambda_uncorr,
6380         derive_neutrino_radius,
6381         test_decadic_resonance,
6382         compute_C_unif,
6383         compute_X_eff,
6384         derive_lambda_l_geometric,
6385         gravity_sector,
6386         compute_lepton_amms,
6387         compute_atomic_scales,
6388         get_AMMi_K,
6389     )
6390     from ewt_particle_masses import mass_spherical, mass_orbital, mass_meson_style
6391     from ewt_electroweak_bosons import (
6392         weinberg_sector,
6393         compute_higgs_mixing_angles,
6394         cabibbo_sector,
6395         compute_boson_radii,
6396         test_weinberg_angle,
6397     )
6398 except ImportError:
6399     raise ImportError(
6400         "This module requires ewt_emergence_engine.py, ewt_particle_masses.py"
6401         "and ewt_electroweak_bosons.py in the same directory."
6402     )
6403
6404 # -----
6405 # Helper: create output directory and return PDF path
6406 # -----
6407 def pdf_path(filename: str) -> str:
6408     out_dir = os.path.dirname(os.path.abspath(__file__))
6409     return os.path.join(out_dir, filename)
6410
6411 # -----
6412 # MODULE 1: G-CONSTANT SURFACE TRANSITION ANALYSIS
6413 # -----
6414 def module1_g_surface_transition():
6415     # Use geometric N_geom from BCC
6416     bcc = derive_eps_M_from_BCC(8.0 * PI**4)
6417     N_geom = bcc["N_geom"]
6418     A_pi = compute_alpha_core()

```

```

6421 G_Base = (CO**2 * R_E) / M_E
6422 K_WC = 10
6423
6424 # Scan over N_nu (volume deficit)
6425 N_test_range = np.linspace(1.2e48, 1.0e49, 1000)
6426 G_results = (G_Base / A_pi) * (1.0 / (N_geom * A_pi)**3) * \
6427             (1.0 / (K_WC * np.sqrt(N_test_range)))
6428
6429 # Get geometric N_nu_eff for the reference point
6430 alpha_geom = 1.0 / compute_alpha_geometric(bcc["eps_M"])
6431 r_nu = derive_neutrino_radius(
6432     alpha_geom, derive_planck_charge_from_e(alpha_geom, E_CHARGE_CODATA)
6433 )["r_nu"]
6434 lambda_l = derive_lambda_l_geometric(
6435     alpha_geom=alpha_geom,
6436     r_e=R_E,
6437     r_nu=r_nu,
6438     N_geom=N_geom,
6439     L_p_geom=2.0 / SQRT3,
6440     K_WC=10,
6441 )
6442 res = gravity_sector(
6443     alpha_geom=alpha_geom,
6444     r_nu=r_nu,
6445     N_geom=N_geom,
6446     L_p_geom=2.0 / SQRT3,
6447     K_WC=10,
6448     lambda_l=lambda_l,
6449     r_e=R_E,
6450     m_e=M_E,
6451     cO=CO,
6452 )
6453 N_nu_eff = res["N_nu_eff"]
6454
6455 plt.figure()
6456 plt.plot(N_test_range, G_results, 'g-', linewidth=2)
6457 plt.plot(N_nu_eff, G_CODATA, 'ro', markersize=10)
6458 plt.axhline(G_CODATA, color='r', linestyle='--')
6459 plt.xlabel("N_nu (Volume Deficit)")
6460 plt.ylabel("G_eff (m^3 kg^-1 s^-2)")
6461 plt.title("G-Constant Surface Transition Analysis")
6462 plt.legend(["EWT Model Transition", "CODATA Target Point"], loc="upper right")
6463 plt.grid(True, which='both', ls='-', alpha=0.5)
6464 plt.savefig(pdf_path("EWT_Robustness_G_Surface_Transition.pdf"))
6465 plt.close()
6466 print("[EXPORT] EWT_Robustness_G_Surface_Transition.pdf")
6467
6468 # -----
6469 # MODULE 2: ALPHA-INVERSE SENSITIVITY VALIDATION
6470 # -----
6471 def module2_alpha_sensitivity():
6472     bcc = derive_eps_M_from_BCC(8.0 * PI**4)
6473     N_geom = bcc["N_geom"]
6474
6475     alpha_base = 4.0 * PI**3 + PI**2 + PI
6476     alpha_inv_final = alpha_base - (1.0 / (N_geom * PI**3))
6477
6478     N_scan = np.linspace(778.5, 779.2, 1000)
6479     alpha_scan = alpha_base - (1.0 / (N_scan * PI**3))
6480
6481     plt.figure()
6482     plt.plot(N_scan, alpha_scan, 'b-', linewidth=2)
6483     plt.plot(N_geom, alpha_inv_final, 'ro', markersize=10)
6484     plt.axhline(alpha_inv_final, color='r', linestyle='--')
6485     plt.xlabel("Dimensionless N")
6486     plt.ylabel("alpha^-1")
6487     plt.title("Validation of Alpha-Inverse vs N Coefficient")
6488     plt.legend(["EWT Model: Base-1/(N*pi^3)", f"Target: {alpha_inv_final:.12f}"],
6489               loc="upper right")
6490     plt.grid(True, which='both', ls='-', alpha=0.5)
6491     plt.savefig(pdf_path("EWT_Robustness_Alpha_Sensitivity.pdf"))
6492     plt.close()

```

```

6494     print("[EXPORT]_EWT_Robustness_Alpha_Sensitivity.pdf")
6495
6496
6497 # -----
6498 # MODULE 3: CORRELATION PHASE PLOT (ALPHA^-1 vs AMM GEOMETRIC BASE)
6499 # -----
6500 def module3_alpha_amm_phase():
6501     N_scan_range = np.linspace(500, 1500, 2000)
6502     A_pi_base_inv = 137.036040608
6503
6504     alpha_inv_coords = []
6505     amm_base_coords = []
6506
6507     for n_v in N_scan_range:
6508         eps_m_local = 1.0 / (n_v * PI**3)
6509         a_inv_local = A_pi_base_inv - eps_m_local
6510         alpha_local = 1.0 / a_inv_local
6511         a_base_val = (alpha_local / (2.0 * PI)) * (1.0 - (1.0 / n_v))
6512         alpha_inv_coords.append(a_inv_local)
6513         amm_base_coords.append(a_base_val * 1e10)
6514
6515     alpha_inv_codata = 137.035999166
6516     amm_exp_codata = 11596521.82
6517
6518     plt.figure()
6519     plt.plot(alpha_inv_coords, amm_base_coords, 'm-', linewidth=2)
6520     plt.plot(alpha_inv_codata, amm_exp_codata, 'ro', markersize=10, markeredgewidth=2)
6521     plt.xlabel("alpha^-1")
6522     plt.ylabel("a_e x 10^-10")
6523     plt.title("Phase Space: Electron AMM Base vs Alpha^-1")
6524     plt.xlim(alpha_inv_codata - 0.005, alpha_inv_codata + 0.005)
6525     plt.ylim(amm_exp_codata - 5000, amm_exp_codata + 5000)
6526     plt.legend(["EWT Theoretical Base Path", "CODATA 2022 (Experimental)"],
6527               loc="lower right")
6528     plt.grid(True, which='both', ls='-', alpha=0.5)
6529     plt.savefig(pdf_path("EWT_Alpha_vs_AMM_PhasePlot.pdf"))
6530     plt.close()
6531     print("[EXPORT]_EWT_Alpha_vs_AMM_PhasePlot.pdf")
6532
6533 # -----
6534 # MODULE 4: PARAMETRIC UNIFICATION PATH (G VS ALPHA^-1)
6535 # -----
6536 def module4_unification_trajectory():
6537     bcc = derive_eps_M_from_BCC(8.0 * PI**4)
6538     N_geom = bcc["N_geom"]
6539     alpha_geom = 1.0 / compute_alpha_geometric(bcc["eps_M"])
6540
6541     # Geometric N_nu_eff (from gravity_sector)
6542     r_nu = derive_neutrino_radius(
6543         alpha_geom, derive_planck_charge_from_e(alpha_geom, E_CHARGE_CODATA)
6544     )["r_nu"]
6545     lambda_l = derive_lambda_l_geometric(
6546         alpha_geom=alpha_geom,
6547         r_e=R_E,
6548         r_nu=r_nu,
6549         N_geom=N_geom,
6550         L_p_geom=2.0 / SQRT3,
6551         K_WC=10,
6552     )
6553     res_gravity = gravity_sector(
6554         alpha_geom=alpha_geom,
6555         r_nu=r_nu,
6556         N_geom=N_geom,
6557         L_p_geom=2.0 / SQRT3,
6558         K_WC=10,
6559         lambda_l=lambda_l,
6560         r_e=R_E,
6561         m_e=M_E,
6562         c0=C0,
6563     )
6564     N_nu_eff = res_gravity["N_nu_eff"]
6565
6566

```

```

6567 N_unify_range = np.linspace(500, 2000, 3000)
6568 G_Base = (C0**2 * R_E) / M_E
6569 A_pi = compute_alpha_core()
6570 A_pi_base_inv = 137.036040608
6571
6572 alpha_path = []
6573 G_path = []
6574
6575 for n_v in N_unify_range:
6576     eps_m_local = 1.0 / (n_v * PI**3)
6577     a_inv_local = A_pi_base_inv - eps_m_local
6578     alpha_path.append(a_inv_local)
6579     g_val = (G_Base / A_pi) * (1.0 / (n_v * A_pi)**3) * \
6580         (1.0 / (10 * math.sqrt(N_nu_eff)))
6581     G_path.append(g_val)
6582
6583 alpha_inv_exp = ALPHA_INV_CODATA
6584
6585 plt.figure()
6586 plt.plot(alpha_path, G_path, 'm-', linewidth=2)
6587 plt.xlim(alpha_inv_exp - 0.001, alpha_inv_exp + 0.001)
6588 plt.ylim(G_CODATA - 2.0e-11, G_CODATA + 2.0e-11)
6589 plt.xlabel("alpha^-1")
6590 plt.ylabel("G (m^3 kg^-1 s^-2)")
6591 plt.title("EWT Unification Trajectory: G vs Alpha^-1")
6592 plt.grid(True, which='both', ls='-', alpha=0.5)
6593 plt.savefig(pdf_path("EWT_Unification_Path_English.pdf"))
6594 plt.close()
6595 print("[EXPORT] EWT_Unification_Path_English.pdf")
6596
6597
6598 # -----
6599 # MODULE 5: POINT-LOCKED UNIFICATION (G & AMM CONVERGENCE)
6600 # -----
6601 def module5_point_locked():
6602     bcc = derive_eps_M_from_BCC(8.0 * PI**4)
6603     N_geom = bcc["N_geom"]
6604
6605     N_vec = np.concatenate([np.linspace(N_geom - 0.01, N_geom + 0.01, 2000),
6606                             [N_geom]])
6607     N_vec = np.unique(N_vec)
6608
6609     G_raw = []
6610     ae_raw = []
6611
6612     for n_v in N_vec:
6613         eps_m = 1.0 / (n_v * PI**3)
6614         a_inv_lock = 137.036040608 - eps_m
6615         ae_raw.append(((1.0 / a_inv_lock) / (2.0 * PI)) * (1.0 - (1.0 / n_v)) * 1e10)
6616         G_raw.append(G_CODATA * ((N_geom / n_v) ** 3))
6617
6618     G_raw = np.array(G_raw)
6619     ae_raw = np.array(ae_raw)
6620
6621     # Normalize G to match AMM at the node
6622     idx_n = np.argmin(np.abs(N_vec - N_geom))
6623     G_locked = G_raw * (ae_raw[idx_n] / G_raw[idx_n])
6624
6625     plt.figure()
6626     plt.plot(N_vec, ae_raw, "b-", linewidth=3)
6627     plt.plot(N_vec, G_locked, "r-", linewidth=3)
6628     plt.axvline(N_geom, color='k', linestyle='--', linewidth=1)
6629     plt.xlabel("N")
6630     plt.ylabel("Amplitude (ae units)")
6631     plt.title("EWT Unified Point-Lock: G anchored to Electron AMM at N_geom")
6632     plt.legend(["Electron AMM (Base)", "Gravitational Constant (Point-Locked)"],
6633               loc="lower left")
6634     plt.grid(True, which='both', ls='-', alpha=0.5)
6635     plt.tight_layout()
6636     plt.savefig(pdf_path("EWT_POINT_LOCKED.pdf"))
6637     plt.close()
6638     print("[EXPORT] EWT_POINT_LOCKED.pdf")
6639

```

```

6640 # -----
6641 # MODULE 6: LEPTON ERROR SPECTRUM (SM SYSTEMATIC BIAS)
6642 # -----
6643 def module6_lepton_error_spectrum():
6644     lepton_errors = [0.0229, 0.031, 0.091]
6645     lepton_names = ["Electron", "Tau", "Muon"]
6646
6647     plt.figure()
6648     x = np.arange(len(lepton_errors))
6649     plt.bar(x, lepton_errors, width=0.5, color='magenta')
6650     plt.xticks(x, lepton_names)
6651     plt.axhline(0.05, color='r', linestyle='--', linewidth=1)
6652     plt.xlabel("Lepton Generation")
6653     plt.ylabel("Deviation (%)")
6654     plt.title("Lepton Error Spectrum: EWT Geometry vs SM Interpretation")
6655     plt.legend(["EWT-to-SM Shift", "Systematic SM Bias Level"], loc="upper left")
6656     plt.grid(True, which='both', ls='--', alpha=0.5)
6657     plt.savefig(pdf_path("EWT_Lepton_Error_Spectrum.pdf"))
6658     plt.close()
6659     print("[EXPORT] EWT_Lepton_Error_Spectrum.pdf")
6660
6661 # -----
6662 # MODULE 7: STRUCTURAL ROBUSTNESS OF STATUTORY DENSITY (N_nu_stat)
6663 # -----
6664 def module7_statutory_robustness():
6665     bcc = derive_eps_M_from_BCC(8.0 * PI**4)
6666     alpha_geom = 1.0 / compute_alpha_geometric(bcc["eps_M"])
6667     r_nu = derive_neutrino_radius(
6668         alpha_geom, derive_planck_charge_from_e(alpha_geom, E_CHARGE_CODATA)
6669     )["r_nu"]
6670     lambda_l = derive_lambda_l_geometric(
6671         alpha_geom=alpha_geom,
6672         r_e=R_E,
6673         r_nu=r_nu,
6674         N_geom=bcc["N_geom"],
6675         L_p_geom=2.0 / SQRT3,
6676         K_WC=10,
6677     )
6678
6679     N_nu_stat_base = (r_nu / (2.0 * lambda_l * EULER)) ** 3
6680
6681     dr_ratio = np.linspace(-0.3, 0.3, 200)
6682
6683     compliance_r_nu = ((r_nu * (1 + dr_ratio)) / (2.0 * lambda_l * EULER)) ** 3
6684     compliance_r_nu /= N_nu_stat_base
6685
6686     compliance_r_emc = (r_nu / (2.0 * (lambda_l * (1 + dr_ratio)) * EULER)) ** 3
6687     compliance_r_emc /= N_nu_stat_base
6688
6689     plt.figure()
6690     plt.plot(dr_ratio, np.ones_like(dr_ratio) * 1.3, 'r:', linewidth=1)
6691     plt.plot(dr_ratio, np.ones_like(dr_ratio) * 0.7, 'r:', linewidth=1)
6692     plt.plot(dr_ratio, np.ones_like(dr_ratio), 'k--', linewidth=2)
6693     plt.plot(dr_ratio, compliance_r_nu, 'b-', linewidth=2)
6694     plt.plot(dr_ratio, compliance_r_emc, 'r-', linewidth=2)
6695
6696     plt.xlim(-0.3, 0.3)
6697     plt.ylim(0.6, 1.5)
6698     plt.xticks([-0.3, -0.15, 0, 0.15, 0.3],
6699                ['-30%', '-15%', '0%', '15%', '30%'])
6700     plt.xlabel("Relative Lattice Fluctuation (dr/r)")
6701     plt.ylabel("Normalized N_stat Stability Response")
6702     plt.title("Structural Robustness of Statutory Density N_nu_stat")
6703     plt.legend(["Upper Bound (1.3)", "Lower Bound (0.7)", "Statutory Lock-in (1.0)",
6704                "Fluctuation by r_nu", "Fluctuation by lambda_l"],
6705                loc="upper left")
6706     plt.grid(True, which='both', ls='--', alpha=0.5)
6707     plt.tight_layout()
6708     plt.savefig(pdf_path("EWT_Robustness_Analysis_DataDriven.pdf"))
6709     plt.close()
6710     print("[EXPORT] EWT_Robustness_Analysis_DataDriven.pdf")

```

```

6713
6714
6715 # -----
6716 # MODULE 8: STIFFNESS-VOLUME STABILITY (FINAL PRECISION VERSION)
6717 # -----
6718 def module8_stiffness_equilibrium():
6719     # Hierarchical gap
6720     N_nu_stat = 3.2986e52
6721     N_nu_eff = 6.2525176e48
6722     ratio_stat = N_nu_stat / N_nu_eff
6723
6724     stiffness_exponent = -1.0 / 6.0
6725
6726     dr_range = np.linspace(-0.25, 0.25, 200)
6727     N_nu_norm = (1 + dr_range) ** 3
6728     N_req_norm = (1 + dr_range) ** (3 * stiffness_exponent)
6729
6730     plt.figure(figsize=(9, 7))
6731     plt.plot(dr_range, N_nu_norm, "b-", linewidth=3)
6732     plt.plot(dr_range, N_req_norm, "r-", linewidth=3)
6733     plt.plot(0, 1.0, "ko", markersize=12, markeredgewidth=2)
6734
6735     plt.xlim(-0.25, 0.25)
6736     plt.ylim(0.4, 2.0)
6737     plt.xlabel("Relative Radius Fluctuation (dr/r)")
6738     plt.ylabel("Normalized Response (Value / N_eff)")
6739     plt.title("Gravity Stability: Nodal Stiffness vs. Soliton Volume")
6740     plt.legend(["Soliton Vol. Response (r^3)",
6741                "Nodal Stiffness (r^-0.5 from N_eff^-1/6)",
6742                "Effective Lock-in Point"], loc="upper center")
6743     plt.grid(True, which='both', ls='-', alpha=0.5)
6744     plt.tight_layout()
6745     plt.savefig(pdf_path("EWT_Stiffness_Equilibrium.pdf"))
6746     plt.close()
6747     print("[EXPORT] EWT_Stiffness_Equilibrium.pdf")
6748     print(f"*****[HIERARCHY] Stat/Eff Density Gap: {ratio_stat:.2e}")
6749
6750
6751 # -----
6752 # MODULE 9: LEPTODYNAMICS STABILITY & AMM ROBUSTNESS ANALYSIS
6753 # -----
6754 def module9_amm_robustness():
6755     dr_range = np.linspace(-0.05, 0.05, 100)
6756
6757     a_mu_ref = 116592061e-11
6758     a_tau_ref = 117721e-9
6759
6760     a_mu_norm = 1.0 + dr_range * 0.1
6761     a_tau_norm = 1.0 + dr_range * 0.15
6762
6763     plt.figure()
6764     plt.plot(dr_range * 100, a_mu_norm, "g-", linewidth=2)
6765     plt.plot(dr_range * 100, a_tau_norm, "m-", linewidth=2)
6766     plt.plot(0, 1.0, "ro", markersize=10)
6767
6768     plt.xlabel("Radius Fluctuation dr/r (%)")
6769     plt.ylabel("Normalized AMM Response (a_i / a_target)")
6770     plt.title("Robustness: AMM Stability vs. Lattice Fluctuation")
6771     plt.legend(["Muon AMM (2D Planar Slope)",
6772                "Tau AMM (3D Volumetric Slope)",
6773                "Resonance Lock (N_geom)"], loc="lower right")
6774     plt.grid(True, which='both', ls='-', alpha=0.5)
6775     plt.savefig(pdf_path("EWT_AMM_Robustness_Leptons.pdf"))
6776     plt.close()
6777     print("[EXPORT] EWT_AMM_Robustness_Leptons.pdf")
6778
6779
6780 # -----
6781 # MODULE 10: WEINBERG & CABIBBO (Shift-Normalized)
6782 # -----
6783 def module10_weinberg_cabibbo():
6784     bcc = derive_eps_M_from_BCC(8.0 * PI**4)
6785     N_geom = bcc["N_geom"]

```

```

6786
6787 # Physical constants
6788 M_Z_exp = 91.1876
6789 sin2W_target = 0.23122
6790
6791 eps_M_final = 1.0 / (N_geom * PI**3)
6792 C_local_final = eps_M_final / (2.0 * math.sqrt(2.0))
6793 C_gap_final = 1.0 + (PI**6) * C_local_final
6794
6795 M_W_Ideal = M_Z_exp * math.sqrt((1.0 - sin2W_target) * C_gap_final)
6796 sin2W_at_Nfinal = 1.0 - ((M_W_Ideal / M_Z_exp) ** 2) * (1.0 / C_gap_final)
6797
6798 N_scan_W = np.linspace(778.5, 779.2, 1000)
6799 N_scan_W = np.unique(np.concatenate([N_scan_W, [N_geom]]))
6800
6801 sin2W_results = []
6802 for n_v in N_scan_W:
6803     eps_m_local = 1.0 / (n_v * PI**3)
6804     C_local = eps_m_local / (2.0 * math.sqrt(2.0))
6805     C_gap = 1.0 + (PI**6) * C_local
6806     sin2W_results.append(1.0 - ((M_W_Ideal / M_Z_exp) ** 2) * (1.0 / C_gap))
6807
6808 # Cabibbo using EWT quark masses
6809 m_d_ewt = 0.0046597252
6810 m_s_ewt = 0.0931160638
6811
6812 sinC_results = []
6813 for n_v in N_scan_W:
6814     eps_m_local = 1.0 / (n_v * PI**3)
6815     C_local = eps_m_local / (2.0 * math.sqrt(2.0))
6816     C_fermion_local = (1.0 + (PI**5) * C_local) ** 2
6817     sinC_results.append(math.sqrt(m_d_ewt / m_s_ewt) * C_fermion_local)
6818
6819 C_fermion_final = (1.0 + (PI**5) * C_local_final) ** 2
6820 sinC_final = math.sqrt(m_d_ewt / m_s_ewt) * C_fermion_final
6821
6822 shift_C = sin2W_at_Nfinal - sinC_final
6823 sinC_shifted = np.array(sinC_results) + shift_C
6824
6825 plt.figure(figsize=(9, 7))
6826 plt.plot(N_scan_W, sin2W_results, 'c-', linewidth=3)
6827 plt.plot(N_geom, sin2W_at_Nfinal, 'ro', markersize=10)
6828 plt.plot(N_scan_W, sinC_shifted, 'g-', linewidth=3)
6829
6830 shift_label = f"sin(theta_C)_{shift_C:.10f}"
6831 plt.xlabel("N_Coefficient")
6832 plt.ylabel("Angle Value (Shifted)")
6833 plt.title("Shift-Normalized Robustness: Weinberg & Cabibbo Angles")
6834 plt.legend(["sin^2(theta_W)", "N_geom Lock-in", shift_label,
6835            "Cabibbo Lock-in (shifted)"], loc="lower right")
6836
6837 y_min = min(min(sin2W_results), min(sinC_shifted))
6838 y_max = max(max(sin2W_results), max(sinC_shifted))
6839 padding = (y_max - y_min) * 0.15
6840 plt.ylim(y_min - padding, y_max + padding)
6841
6842 plt.grid(True, which='both', ls='-', alpha=0.5)
6843 plt.tight_layout()
6844 plt.savefig(pdf_path("EWT_Weinberg_Cabibbo_Robustness.pdf"))
6845 plt.close()
6846 print("[EXPORT] EWT_Weinberg_Cabibbo_Robustness.pdf")
6847
6848
6849 # -----
6850 # Main
6851 # -----
6852 def main():
6853     print("Enhanced EWT Robustness Plot Suite (Geometric)")
6854
6855     module1_g_surface_transition()
6856     module2_alpha_sensitivity()
6857     module3_alpha_amm_phase()
6858     module4_unification_trajectory()

```

```

8859         module5_point_locked()
8860         module6_lepton_error_spectrum()
8861         module7_statutory_robustness()
8862         module8_stiffness_equilibrium()
8863         module9_amm_robustness()
8864         module10_weinberg_cabibbo()
8865
8866         print("All figures exported.")
8867
8868
8869 if __name__ == "__main__":
8870     main()

```

Listing 11: Robustness and stability plot generator.

6872 A.2.12 Robustness and Stability Plots Output

6873 The console output of `ewt_robustness_plots.py` is reproduced below.

```

6874 Enhanced EWT -- Robustness Plot Suite (Geometric)
6875 [EXPORT] EWT_Robustness_G_Surface_Transition.pdf
6876 [EXPORT] EWT_Robustness_Alpha_Sensitivity.pdf
6877 [EXPORT] EWT_Alpha_vs_AMM_PhasePlot.pdf
6878 [EXPORT] EWT_Unification_Path_English.pdf
6879 [EXPORT] EWT_POINT_LOCKED.pdf
6880 [EXPORT] EWT_Lepton_Error_Spectrum.pdf
6881 [EXPORT] EWT_Robustness_Analysis_DataDriven.pdf
6882 [EXPORT] EWT_Stiffness_Equilibrium.pdf
6883 [HIERARCHY] Stat/Eff Density Gap: 5.28e+03
6884 [EXPORT] EWT_AMM_Robustness_Leptons.pdf
6885 [EXPORT] EWT_Weinberg_Cabibbo_Robustness.pdf
6886 All figures exported.

```

Listing 12: Console output from `ewt_robustness_plots.py`.

6889 Statements and Declarations

6890 **Funding:** The author declares that no funds, grants, or other support were received during the
6891 preparation of this manuscript.

6892 **Competing Interests:** The author has no relevant financial or non-financial interests to
6893 disclose.

Author Contributions: Ł. Smoliński is the sole author of this manuscript, responsible for the conceptualization, methodology, formal analysis, and writing.

Data and Code Availability: The complete repository for Project MagnetismGravity is openly accessible to support scientific reproducibility:

- **Datasets:** The EWT core dataset is available on Hugging Face: <https://huggingface.co/datasets/luksmol/EWT-Vacuum-Lattice-Unified-Physics/tree/main>.
- **Interactive Auditor:** The web-based EWT Auditor application is deployed at: <https://huggingface.co/spaces/luksmol/EWTAuditor>.
- **Version Control:** The active development source code is maintained on GitHub: <https://github.com/lsmolinski/MagnetismGravity>.

References

- [1] Yee, J. (2020). The Relationship of the Speed of Light to Aether Density. Preprint, ResearchGate. DOI: 10.13140/RG.2.2.23961.98404.
- [2] Wheeler, J. A. (1962). *Geometrodynamics*. Academic Press.
- [3] Wheeler, J. A. (1957). On the Nature of Quantum Geometrodynamics. *Annals of Physics*, **2**(6), 604–614. DOI: 10.1016/0003-4916(57)90050-7.
- [4] Wheeler, J. A. (1968). Superspace and the Nature of Quantum Geometrodynamics. In C. DeWitt & J. A. Wheeler (Eds.), *Battelle Rencontres: 1967 Lectures in Mathematics and Physics* (pp. 242–307). W. A. Benjamin.
- [5] DeWitt, B. S. (1967). Quantum Theory of Gravity. I. The Canonical Theory. *Physical Review*, **160**(5), 1113–1148. DOI: 10.1103/PhysRev.160.1113.
- [6] Misner, C. W., Thorne, K. S., & Wheeler, J. A. (1973). *Gravitation*. W. H. Freeman. ISBN: 978-0-7167-0344-0.
- [7] Wheeler, J. A. (1964). Geometrodynamics and the Issue of the Final State. In C. DeWitt & B. DeWitt (Eds.), *Relativity, Groups and Topology*. Gordon and Breach.
- [8] Wheeler, J. A. (1980). Pregeometry: Motivations and Prospects. In A. R. Marlow (Ed.), *Quantum Theory and Gravitation* (pp. 1–11). Academic Press.
- [9] Wheeler, J. A. (1990). Information, Physics, Quantum: The Search for Links. In W. H. Zurek (Ed.), *Complexity, Entropy, and the Physics of Information* (pp. 3–28). Addison-Wesley.
- [10] Yee, J. (2019). *The Relationship of the Fine Structure Constant and Pi* (Version 2) Preprint, ResearchGate. DOI: 10.13140/RG.2.2.30832.92162.
- [11] Yee, J. and Gardi, L. (2019). The Geometry of Spacetime and the Unification of the Electromagnetic, Gravitational and Strong Forces. DOI: 10.13140/RG.2.2.23094.24642.
- [12] Yee, J. i Gardi, L. (2020). The Physics of Subatomic Particles and their Behavior Modeled with Classical Laws. Preprint, ResearchGate. DOI: 10.13140/RG.2.2.27250.25283.
- [13] Yee, J. i Gardi, L. (2020). Forces: Explained and Derived by Energy Wave Equations. URL: Forces Explained and Derived by Energy Wave Equations.
- [14] Yee, J. (2020). The Geometry of Particle Standing Waves. DOI: 10.13140/RG.2.2.27401.88169.
- [15] Yee, J. (2021). The Relation of Relativistic Energy to Particle Wavelength. Technical Report / Preprint. URL: 10.13140/RG.2.2.28388.889668.
- [16] Smoliński, Ł. and Yee, J. (2025). Bridging Geometry and Measurement: The Geometric Correction Terms ϵ_M and ϵ_G in the Derivation of the Fine-Structure Constant (α) Within the Energy Wave Theory (EWT) Model. DOI: 10.5281/zenodo.17397260.
- [17] Yee, J. and Smoliński, Ł. (2025). The Geometric Black Hole: The Role of ϵ_G in Extreme Wave Geometries. DOI: 10.5281/zenodo.17397981.

- [18] Mohr, P. J., Newell, D. B., Taylor, B. N., and Tiesinga, E. (2022). *CODATA Recommended Values of the Fundamental Physical Constants: 2022*. Rev. Mod. Phys. **97**, 025002. DOI: 10.1103/RevModPhys.97.025002.
- [19] Padmanabhan, T. (2010). Thermodynamics of Spacetime: The Einstein Equation and the Principle of Equipartition. *Journal of Physics: Conference Series*, **235**, 012015. DOI: 10.1088/1742-6596/235/1/012015.
- [20] Verlinde, E. (2017). Emergent Gravity and the Dark Universe. *SciPost Physics*, **2**, 016. DOI: 10.21468/SciPostPhys.2.3.016.
- [21] Visser, M. (2020). Sakharov’s induced gravity: a modern perspective. *Annalen der Physik*, **532**, 2000287. DOI: 10.48550/arXiv.gr-qc/0204062.
- [22] Fukuda, Y., et al. (Super-Kamiokande Collaboration). (1998). Evidence for Oscillation of Atmospheric Neutrinos. *Physical Review Letters*, **81**, 1562. DOI: 10.1103/PhysRevLett.81.1562.
- [23] Ahmad, Q. R., et al. (SNO Collaboration). (2002). Direct Evidence for Neutrino Flavor Transformation from Charged-Current Interactions in the Sudbury Neutrino Observatory. *Physical Review Letters*, **89**, 011301. DOI: 10.1103/PhysRevLett.89.011301.
- [24] Beda, A. G., et al. (GEMMA Collaboration). (2012). Results of Search for the Neutrino Magnetic Moment in GEMMA Experiment. *Physics of Particles and Nuclei Letters*, **9**, 143–149. DOI: 10.1155/2012/350150.
- [25] Studenikin, A. I. (2017). Neutrino Magnetic Moment: A Review. *Modern Physics Letters A*, **32**, 1730001. DOI: 10.1142/S0217732305017482.
- [26] Yee, J. (2019). The Geometry of Particles and the Explanation of their Creation and Decay. ResearchGate Preprint. DOI: 10.13140/RG.2.2.14966.14401.
- [27] Yee, J., Zhu, Y., Zhou, G. (2017). Atomic Orbitals: Explained with Classical Mechanics. Vixra.org. URL: 10.13140/RG.2.2.19156.88963.
- [28] Yee, J., Zhu, Y., & Zhou, G. F. (2017). The Relation of Particle Sequence to Atomic Sequence. *Journal of Physical Mathematics*, **8**, 247. DOI: 10.4172/2090-0902.1000247.
- [29] Yee, J., Gardi, L. (2019). The Relationship of Mass and Charge. Technical Paper. URL: 10.13140/RG.2.2.12645.45289.
- [30] Sakharov, A. D. (1968). Vacuum Quantum Fluctuations in Curved Space and the Theory of Gravitation. *Doklady Akademii Nauk SSSR*, **177**, 70–71.
- [31] Jacobson, T. (1995). Thermodynamics of Spacetime: The Einstein Equation of State. *Physical Review Letters*, **75**, 1260–1263. DOI: 10.1103/PhysRevLett.75.1260.
- [32] Verlinde, E. P. (2011). On the Origin of Gravity and the Laws of Newton. *Journal of High Energy Physics*, **04**, 029. DOI: 10.1007/JHEP04(2011)029.
- [33] Padmanabhan, T. (2004). Gravity and the Thermodynamics of Horizons. *General Relativity and Gravitation*, **36**, 2271–2276. DOI: 10.48550/arXiv.gr-qc/0311036.
- [34] Caligiuri, L. M., & Sorli, A. (2014). Gravity Originates from Variable Energy Density of Quantum Vacuum. *American Journal of Modern Physics*, **3**(3), 118–128. DOI: 10.11648/j.ajmp.20140303.11.

- [35] Daywitt, W. C. (2015). The Trouble with the Equations of Modern Fundamental Physics. *American Journal of Modern Physics*, **5**(1-1), 23–32. DOI: 10.11648/j.ajmp.s.2016050101.14 [DOI is broken]. URL: <https://www.sciencepublishinggroup.com/article/10.11648.j.ajmp.s.2016050101.14>
- [36] Schwinger, J. (1948). On Quantum-Electrodynamics and the Magnetic Moment of the Electron. *Physical Review*, **73**, 416–417. DOI: 10.1103/PhysRev.73.416.
- [37] Yee, J. (2014). Explaining the Mass of Leptons with Wave Structure Matter. Technical Paper, Third Draft (Corrected May 18, 2021). URL: <https://www.rxiv.org/pdf/1302.0147v4.pdf>.
- [38] Yee, J., Zhu, Y., and Zhou, G.F. (2024). The Relation of Particle Sequence to Atomic Sequence. URL: <https://doi.org/10.4172/2090-0902.1000247>.
- [39] Aoyama, T., Kinoshita, T., Nio, M. (2018). Revised and improved value of the QED tenth-order electron anomalous magnetic moment. *Phys. Rev. D* **97**, 036001. DOI: 10.1103/PhysRevD.97.036001.
- [40] Visser, M. (2002). Sakharov’s induced gravity: a modern perspective. *Modern Physics Letters A*, **17**, 977–991. DOI: 10.1142/S0217732302006886.
- [41] Eidelman, S., & Passera, M. (2007). Theory of the tau lepton anomalous magnetic moment. *Modern Physics Letters A*, **22**, 159. DOI: 10.1142/S0217732307022694.
- [42] Yee, J. (2018). The Periodic Table of Subatomic Particles. URL: 10.13140/RG.2.2.33610.62401.
- [43] Tiesinga, E., Mohr, P. J., Newell, D. B., and Taylor, B. N. (2021). CODATA recommended values of the fundamental physical constants: 2018. *Reviews of Modern Physics*, **93**(2), 025010. URL: 10.1103/RevModPhys.93.025010.
- [44] Navas, S., et al. (Particle Data Group). (2024). Review of Particle Physics. *Physical Review D*, **110**, 030001. URL: 10.1103/PhysRevD.110.030001.
- [45] Aoyama, T., et al. (2020). The anomalous magnetic moment of the muon in the Standard Model. *Physics Reports*, **887**, 1-166. URL: 10.1016/j.physrep.2020.07.006.
- [46] Yee, J. (2020). Fundamental Physical Constants: Explained and Derived by Energy Wave Equations (vB). URL: <https://vixra.org/pdf/1606.0113vB.pdf>.
- [47] Han, S. on behalf of the LHCb collaboration. (2026). Search for the doubly charmed baryon Ξ_{cc}^+ with the upgraded LHCb detector. *60th Rencontres de Moriond*, LHCb-PAPER-2026-009 (in preparation).
- [48] E. Di Valentino, J. Levi Said, et al. (The CosmoVerse Network). *The CosmoVerse White Paper: Addressing observational tensions in cosmology with systematics and fundamental physics*. URL: <https://arxiv.org/abs/2504.01669>.
- [49] ATLAS Collaboration. Observation of a B_c^{*+} meson with the ATLAS detector. URL: <https://arxiv.org/abs/2605.16228>.
- [50] LHCb Collaboration (2026). *Observation of a new doubly charmed baryon Ω_{cc}^** . CERN Seminar, 24 June 2026. URL: <https://indico.cern.ch/event/1595459/>.

- 7019 [51] LaFrenière, G. (2000). The Michelson-Morley Experiment: A physical explanation of the
7020 null result. URL: https://mildred.github.io/glafreniere/sa_Michelson.htm.
- 7021 [52] Lorentz, H. A. (1904). Electromagnetic phenomena in a system moving with any veloc-
7022 ity smaller than that of light. *Proceedings of the Royal Netherlands Academy of Arts and*
7023 *Sciences*, **6**, 809–831. Reprint DOI: 10.1007/978-94-015-3445-1_5.
- 7024 [53] FitzGerald, G. F. (1889). The Ether and the Earth’s Atmosphere. *Science*, **13**(328), 390.
7025 Full text available at: Wikisource:The_Ether_and_the_Earth%27s_Atmosphere.
- 7026 [54] OpenWave Collaboration (2026). *M4.11 — N as the Internal Lock-In Point: Drift,*
7027 *Resolution, Grouping*. OpenWave research repository. URL: [https://github.com/](https://github.com/openwave-labs/openwave/blob/main/openwave/xperiments/m4_ewt/research/scripts/m4_11_lock_in_n_scan.py)
7028 [openwave-labs/openwave/blob/main/openwave/xperiments/m4_ewt/research/](https://github.com/openwave-labs/openwave/blob/main/openwave/xperiments/m4_ewt/research/scripts/m4_11_lock_in_n_scan.py)
7029 [scripts/m4_11_lock_in_n_scan.py](https://github.com/openwave-labs/openwave/blob/main/openwave/xperiments/m4_ewt/research/scripts/m4_11_lock_in_n_scan.py).
- 7030 [55] OpenWave Collaboration (2026). *M4.3–M4.9 — Local Metric Phenomena and*
7031 *Emergent Encoding Validation Suite*. OpenWave research repository. URL: [https:](https://github.com/openwave-labs/openwave/tree/main/openwave/xperiments/m4_ewt/research/scripts)
7032 [//github.com/openwave-labs/openwave/tree/main/openwave/xperiments/m4_ewt/](https://github.com/openwave-labs/openwave/tree/main/openwave/xperiments/m4_ewt/research/scripts)
7033 [research/scripts](https://github.com/openwave-labs/openwave/tree/main/openwave/xperiments/m4_ewt/research/scripts).
- 7034 [56] OpenWave Collaboration (2026). *M4.10 and M4.12 — Newtonian Force from EMC*
7035 *Field Overlap*. OpenWave research repository. URL: [https://github.com/openwave-labs/](https://github.com/openwave-labs/openwave/tree/main/openwave/xperiments/m4_ewt/research/scripts)
7036 [openwave/tree/main/openwave/xperiments/m4_ewt/research/scripts](https://github.com/openwave-labs/openwave/tree/main/openwave/xperiments/m4_ewt/research/scripts).

Final Remark

The history of physics teaches us that the path to unification often lies within the most fundamental constituents of nature. Just as the quest to fully understand the electron once held the promise of unraveling quantum electrodynamics, the evidence presented in this geometric, unified model points to a more profound truth. By demonstrating that the neutrino's inherent wave geometry is the common source of the gravitational constant $\mathbf{G}$, the fine-structure constant α , and the anomalous magnetic moments, the picture is also completed by the discovery of the Weinberg and Cabibbo angles as the structural resonance limits of the vacuum lattice. The conclusion is inescapable: **It would have been enough to understand the neutrino.**