

- ▶ I looked into the distortion from the metabolic image of 2019-12-12 based on the BW per pixel and frequency offset between metabolites.
- ▶ I tested the execution time of the 2DFT sequence on Supershop.
- ▶ I implemented diffusion for the ems.
- ▶ I wrote a script to generate a DWI pulse sequence.
- ▶ I simulated em diffusion + a DWI sequence.

## Distortion from metabolic image

# Summary



- ▶ I found the BW per pixel to be 125 Hz.
- ▶ I found the frequency offset to be -1277 Hz.



$$\text{int} \left( \frac{-1277 \text{ Hz}}{125 \text{ Hz}} \right) = -10$$

- ▶ Take a look at the image.

# Image from 2019-12-12



# Comments

- ▶ The bottommost em is shifted 10 pixels to the left (if you wrap around to the other side of the image). Checks out!

Execution time of 2DFT sequence on Supershop

# Summary

- ▶ I ran the same script as 2019-12-12 for testing the execution time on Supershop.
- ▶ The total execution time was 129.8 seconds.
- ▶ This means 129.8 microseconds per em per time step.

# Comments

- ▶ Why is the execution time so much slower on Supershop than on my Macbook? The execution time on my Macbook was 5.41 microseconds per em per time step.



Em diffusion

# Summary

- ▶ I looked into diffusion simulation methods.
- ▶ I implemented diffusion in the Em class.
- ▶ I instantiated one em and took  $10^4$  diffusion steps and plotted the trajectory.

# Simulating diffusion


- ▶ Assume space is isotropic.
- ▶ Let  $D(x, y, z)$  be the diffusion coefficient at position  $(x, y, z)$ .
- ▶ Let  $\Delta t$  be the time step of the simulation.
- ▶ Update the position of each em as

$$\begin{aligned}x &\leftarrow x + Q_1 \\y &\leftarrow y + Q_2 \\z &\leftarrow z + Q_3,\end{aligned}$$

where  $Q_i \sim \mathcal{N}(0, 2D(x, y, z)\Delta t)$ .

- ▶ I am uncertain whether  $Q_i \sim \mathcal{N}(0, 2D(x, y, z)\Delta t)$  is correct or if it should be  $Q_i \sim \mathcal{N}(0, D(x, y, z)\Delta t)$ .
- ▶ **This** reference says the former, **this** reference says the latter.
- ▶ I am sticking with  $Q_i \sim \mathcal{N}(0, 2D(x, y, z)\Delta t)$  for now.

## Example trajectory of diffusing em



# Comments

- ▶ Trajectory looks reasonable.
- ▶ The execution time for  $10^4$  diffusion steps for a single em is 0.0783 seconds  $\Rightarrow$  7.828 microseconds per em per time step.
- ▶ Compare with the 5.41 microseconds per em per time step of the 2DFT sequence simulation with stationary ems.
- ▶ Generating the normally distributed random numbers required to model diffusion increases the computation time substantially.


## DWI pulse sequence

# Summary

- ▶ I modified the 2DFT sequence to produce a DWI sequence.
- ▶ The user specifies the amplitude of the diffusion gradient in each of the 3 spatial directions, the duration of these diffusion gradients, and the time between the positive and negative lobes.
- ▶ See the following figure for an example DWI sequence.



# DWI pulse sequence example



# Comments



- ▶ Does that look right?

## DWI sequence simulation

# Summary

- ▶ I defined the diffusion coefficient as  
 $(D_x, D_y, D_z) = (5 \times 10^{-6}, \times 10^{-6}, 0) \text{ m}^2/\text{s}$  for  $x > 0$  and  
 $(D_x, D_y, D_z) = (0, 0, 0)$  for  $x \leq 0$ .
- ▶ I simulated a DWI sequence with diffusion gradients of  
 $(5 \times 10^{-3}, 5 \times 10^{-3}, 5 \times 10^{-3}) \text{ uT/cm}$ , 1 ms diffusion gradient pulses, positive and negative lobes separated by 4 ms.

# Results



# Comments

- ▶ The right side of the image (with a nonzero diffusion coefficient) is less intense than the left side (with zero diffusion coefficient).
- ▶ It looks like the diffusion-weighting is working (?).