

- tions [M]. Cambridge: Cambridge Univ Press.
- Franz M, Goller F. 2003. Respiratory patterns and oxygen consumption in singing zebra finches [J]. The journal of Experimental biology, 206 (6): 967-978.
- Hunter ML, Krebs JR. 1979. Geographical variation in the song of the great tit (*Parus major*) in relation to ecological factors [J]. J of Animal Ecology, 48(3): 759-785.
- Konishi M. 1985. Birdsong: from behavior to neuron [J]. Annual review of Neuroscience, 8(1): 125-170.
- Krebs J, Ashcroft R, Webber M. 1978. Song repertoires and territory defence in the great tit [J]. Nature, 271(5645): 539-542.
- Kroodsma DE, Miller EH (eds). 1982. Acoustic communication in bird, 1 & 2 [M]. New York: Academic Press.
- Lambrechts MM, Dhondt AA. 1987. Differences in singing performance between male great tits [J]. Ardea, 75: 43-52.
- Lambrechts MM, Dhondt AA. 1988. The anti-exhaustion hypothesis: a new hypothesis to explain song performance and song switching in the great tit [J]. Animal Behaviour, 36(2): 327-334.
- Lambrechts MM. 1997. Song frequency plasticity and composition of phrase versions in great tits *Parus major* [J]. Ardea, 85: 99-109.
- McGregor PK, Krebs JR. 1982. Song types in a population of Great Tits (*Parus major*): their distribution, abundance, and acquisition by individuals [J]. Behaviour, 79(2): 126-152.
- Päckert M, Martens J, ECK S, et al. 2005. The great tit (*Parus major*) – a misclassified ring species [J]. Biological Journal of the Linnean Society, 86(2): 153-174.
- Rivera-Gutierrez HF, Matthysen E, Adriaensen F, et al. 2010. Repertoire Sharing and Song Similarity between Great Tit Males Decline with Distance between Forest Fragments [J]. Ethology, 116(10): 951-960.
- Slabbekoorn H, den Boer-Visser A. 2006. Cities Change the Songs of Birds [J]. Current Biology, 16(23): 2326-2331.

西藏自治区鸟类分布新纪录——蒙古沙雀

2012 年 7 月 8 日,笔者在西藏自治区西南部札达县(北纬 31°42′33.5″,东经 79°42′31.2″,海拔 4251 m)进行野外考察期间,拍摄到一只雀形目 Passeriformes 鸟类,经鉴定为燕雀科 Fringillidae 的蒙古沙雀 *Rhodopechys mongolicus* 雄鸟。此鸟体羽以沙褐色为主,腹部色淡;眉纹、喉部和胁部沾粉红色;初级飞羽黑褐色染橙红色外缘,次级飞羽具大块白斑,覆羽黑褐色具橙红色斑;尾羽黑褐色具白色边缘;嘴短粗,上嘴较下嘴色深;脚肉粉色,爪灰黑色。发现地为荒漠草原生境,发现时此鸟正与角百灵 *Eremophila alpestris*、黄嘴朱顶雀 *Carduelis flavirostris* 等雀形目鸟类聚集在水坑边饮水。

蒙古沙雀在我国以往记录分布于内蒙古、宁夏、甘肃、青海、新疆等省区(傅桐生等,1998;郑光美等,2011),此次发现为西藏自治区鸟类分布新纪录。蒙古沙雀为典型的荒漠鸟类,从海拔 500 m 的丘陵到海拔 4200 m 的高山都有分布(傅桐生等,1998;MacKinnon & Phillipps, 2000)。在国外,蒙古沙雀分布于中亚、西亚以及南亚的克什米尔地区(Vaurie, 1959),札达县所在的我国阿里地区与克什米尔地区接壤,在以往有关阿里地区乃至整个西藏自治区的鸟类调查报告中均未记录到蒙古沙雀(李德浩,王祖祥,1979;中国科学院青藏高原综合科学考察队,1983)。因此,蒙古沙雀在西藏的分布范围以及是否在西藏境内繁殖或居留,还需要进一步调查。



王宁*, 邓涛

(中国科学院古脊椎动物与古人类研究所,北京 100044)

* 通讯作者 Corresponding author, E-mail: wangning@ivpp.ac.cn