- 1.1先用y=exp(a1*x+a2)+a3作为拟合函数,得到化石燃料排放的二氧化碳随时间变化的曲线,画出真实值和拟合后的图,得到拟合函数里的三个参数,然后在不考虑buffer effect的情况下设置碳循环函数,画出没有buffer effect的二氧化碳大气浓度。
 1.2重新定义有buffer effect的函数,画图。
 1.3利用二氧化碳年平均数据,再把上面两个图结合起来,得到类似文献中的图2的二氧化程度以上观测值的变化数据图
- 碳模拟与观测值的变化趋势图。

Bonus 先选取需要使用的数据,计算各种k值和初始值,然后积分计算box每部分的变化,最后画图。

```
In [1]: # Import SciPy
    import scipy
    import pandas as pd
    import numpy as np
    from scipy import constants
    from scipy.interpolate import UnivariateSpline
    from numpy import exp
    from scipy import integrate
    from scipy import optimize
    import numpy as np
    import matplotlib.pyplot as plt
    import numpy as np
    import xlrd
    %matplotlib inline
    from scipy.integrate import odeint
```

1.1build a two-box model to compute the atmospheric CO2 level in ppm (parts per million) from 1987 to 2004(without the buffer effect)


```
In [2]: df1=pd. read_csv('global. 1751_2014. csv') df1
```

Out[2]:

	Year	Total carbon emissions from fossil fuel consumption and cement production (million metric tons of C)	Carbon emissions from gas fuel consumption	Carbon emissions from liquid fuel consumption	Carbon emissions from solid fuel consumption	Carbon emissions from cement production	Carbon emissions from gas flaring	Pe em
0	1751	3	0	0	3	0	0	
1	1752	3	0	0	3	0	0	
2	1753	3	0	0	3	0	0	
3	1754	3	0	0	3	0	0	
4	1755	3	0	0	3	0	0	
259	2010	9128	1696	3107	3812	446	67	
260	2011	9503	1756	3134	4055	494	64	
261	2012	9673	1783	3200	4106	519	65	
262	2013	9773	1806	3220	4126	554	68	
263	2014	9855	1823	3280	4117	568	68	

264 rows × 8 columns


```
In [3]: |# got help from my senior fellow apprentice Zhao Wangchao
        # 用y=exp(a1*x+a2)+a3作为拟合函数,得到化石燃料排放的二氧化碳随时间变化的曲线
        def fossil emiss(x, a1, a2, a3):
            return np. \exp(a1*x+a2)+a3
        # 设置拟合的初始值
        a1=0.1
        a2=0.1
        a3=0
        p0=[a1, a2, a3]
        # 调用拟合函数
        df1['Year'].astype(int)
        df1['Total carbon emissions from fossil fuel consumption and cement production (milli-
        para, cov = optimize.curve_fit(fossil_emiss, df1['Year'], df1['Total carbon emissions
        # 画真实值和拟合后的图
        plt.plot(df1['Year'], df1['Total carbon emissions from fossil fuel consumption and cem
        plt.plot(df1['Year'], fossil_emiss(df1['Year'], *para), '-', label='fit')
        plt. show()
        print (para)
```



[2.39596800e-02 -3.90147674e+01 -1.22229282e+02]

```
In [4]: #https://zhuanlan.zhihu.com/p/484167038 # 定义没有buffer效应的碳循环函数 def model1(y, t, k12, k21): N1, N2, a = y #a是 γ dydt=[-k12*N1+k21*N2+a, k12*N1-k21*N2, 2. 39596800e-02*a+2. 39596800e-02*1. 222292826 return dydt
```

```
In [5]: # 设置初始值
         t1=np. linspace (1987, 2005)
         a=fossil emiss(t1,*para)
         k12=105/740
         k21=102/900
         N1 = 740 \times 1000
         N2=900*1000
         y0=[N1, N2, a[0]]
         #转化单位
         sol = odeint(modell, y0, t1, args=(k12, k21))/1000/740*347
In [6]: # 画图
         from matplotlib.ticker import MaxNLocator
         plt.plot(t1, sol[:, 0], 'k', label='CO2 in atmosphere(ppm)')
         plt.legend(loc='best')
         plt. xlabel ('year')
         # 设置横坐标轴刻度为整数
         plt.gca().xaxis.set major locator(MaxNLocator(integer=True))
         plt.title('1987-2004 CO2 concentration without the buffer effect')
```



1.2 build a two-box model to compute the atmospheric CO2 level in ppm from 1987 to 2004(with the buffer effect)

```
In [7]: # 定义有buffer效应的函数
         def model2(y, t, k12, k21, N0):
             N1, N2, a2 = y
             bf=3.69+1.86e-2*(N1/740/1000*347)-1.8e-6*((N1/740/1000*347)**2) #buffer factor
             dydt = [-k12*N1+k21*(N0+bf*(N2-N0))+a2,
                   k12*N1-k21*(N0+bf*(N2-N0)),
                   2. 39596800e-02*a2+2. 39596800e-02*1. 22229282e+02
             return dydt
In [8]: # 设置初始值
         t2=np. linspace (0, 253, 253)+1751
         a2=fossil emiss(t2,*para)
         k12=105/740
         k21=102/900
         N0=821*1000
         N1=618*1000
         N2=821*1000
         y0=[N1, N2, a2[0]]
         sol2 = odeint(model2, y0, t2, args=(k12, k21, N0))/740/1000*347 #计算微分方程,并且转化
In [9]: so12
Out[9]: array([[ 2.89791892e+02,
                                   3. 84982432e+02, -4. 84577482e-02],
                [ 2.91238045e+02,
                                   3.83487737e+02, -4.82420907e-02],
                2.91660746e+02,
                                   3. 83016713e+02, -4. 80211828e-02,
                2.91763603e+02,
                                   3.82865757e+02, -4.77948966e-02],
                                   3.82814589e+02, -4.75631011e-02],
                2.91766902e+02,
                                   3.82794521e+02, -4.73256623e-02],
                 2.91739337e+02,
                                   3. 82784156e+02, -4.70824426e-02],
                2.91702311e+02,
                [ 2.91662489e+02,
                                   3.82776834e+02, -4.68333014e-02],
                                   3.82770480e+02, -4.65780946e-02],
                2.91621951e+02,
                2.91581350e+02,
                                   3. 82764449e+02, -4. 63166743e-02],
                2.91540891e+02,
                                   3.82758540e+02, -4.60488894e-02,
                                   3.82752693e+02, -4.57745849e-02],
                  2. 91500644e+02,
                [ 2.91460633e+02,
                                   3.82746889e+02, -4.54936021e-02],
                [ 2.91420870e+02,
                                   3.82741121e+02, -4.52057783e-02],
                                   3.82735390e+02, -4.49109471e-02,
                 [ 2.91381363e+02,
                                   3.82729695e+02, -4.46089378e-02],
                 2.91342120e+02,
                                   3.82724038e+02, -4.42995755e-02],
                 [ 2.91303146e+02,
                2.91264449e+02,
                                   3.82718418e+02, -4.39826815e-02],
                                   3.82712838e+02, -4.36580724e-02],
```


9 00707900 - 109

4 220EECO2 A01

2.91226034e+02, F 0 01107000 - 100

```
In [10]: # 画图 plt.plot(t2[235:253], sol2[235:253,0], 'k', label='CO2 in atmosphere(ppm)') plt.legend(loc='best') plt.xlabel('year') plt.gca().xaxis.set_major_locator(MaxNLocator(integer=True)) plt.title('1987-2004 CO2 concentration with the buffer effect') plt.show()
```





1.3 reproduce Figure 2 in Tomizuka (2009)


```
In [11]: # 导入观测值数据 df2=pd. read_csv('co2_annmean_mlo.csv') df2
```

Out[11]:

	year	mean	unc
0	1959	315.98	0.12
1	1960	316.91	0.12
2	1961	317.64	0.12
3	1962	318.45	0.12
4	1963	318.99	0.12
59	2018	408.72	0.12
60	2019	411.65	0.12
61	2020	414.21	0.12
62	2021	416.41	0.12
63	2022	418.53	0.12

64 rows × 3 columns

```
In [12]: # 画图
         plt.plot(t1, sol[:, 0], linewidth=2, color='grey') # without buffer effect
         plt.plot(t2[235:253], sol2[235:253,0], linewidth=4, color='black') # with buffer effe
         plt. scatter(df2['year'][30:47], df2['mean'][30:47], color='gray') # observed data
         # 设置图例
         plt.legend(['without buffer effect', 'with buffer effect', 'observations'], loc='best')
         plt. xlabel ('Year')
         # 设置横纵坐标
         plt.gca().xaxis.set_major_locator(MaxNLocator(integer=True))
         plt.xticks([1985, 1990, 1995, 2000], fontsize=12)
         plt.yticks([360, 380, 400, 420], fontsize=12)
         # 隐藏右边和上边的边框
         plt.gca().spines['right'].set_visible(False)
         plt.gca().spines['top'].set visible(False)
         plt. title('1987-2004 CO2 trend')
         plt.show()
         # 不考虑buffer effect的曲线与文献中图2略有区别,可能是函数拟合不够好
```



[Bonus] Following equation 5-13, compute the atmospheric CO2 level in ppm and reproduce Figure 4 in Tomizuka (2009).

```
In [13]: # got help from my classmate Xu Jiwen
    co2_observations = pd.read_csv('1750-2000C02.csv')
    land_use_data = pd.read_excel('Global_land-use_flux-1750_2005.xls')
    ff_emissions = pd.read_csv('global.1751_2014.csv')

land_use_data = land_use_data[['Year', 'Global']]
    land_use_data['LandUseChange'] = land_use_data['Global'] / (1000 * 2.13)

ff_emissions = ff_emissions[['Year', 'Total carbon emissions from fossil fuel consump
    ff_emissions['FossilFuelEmissions'] = ff_emissions.iloc[:, 1] - ff_emissions.iloc[:, ff_emissions['EmissionFactor'] = ff_emissions['FossilFuelEmissions'] / (1000 * 2.13)
```

```
In [14]: k12, k21, k23, k24, k32, k34, k43, k45, k51, k67, k71 = [60 / 615, 60 / 842, 9 / 842,
          N2 \ 0 = 842 / 2.13
          initial conditions = [615 / 2.13, 842 / 2.13, 9744 / 2.13, 26280 / 2.13, 90000000 /
          f0 = 62 / 2.13
          P0 = 615 / 2.13
          # Explore the Beta value
          beta values = [0.38, 0.5]
          results = []
          for beta in beta values:
              N1, N2, N3, N4, N5, N6, N7 = initial_conditions.copy()
              atmosphere = [N1]
              for year in range (1751, 2001):
                  gamma = ff emissions[ff emissions['Year'] == year]['EmissionFactor'].values[0
                  delta = land_use_data[land_use_data['Year'] == year]['LandUseChange'].values[
                  xi = 3.69 + 0.0186 * N1 - 0.0000018 * N1**2
                  f = f0 * (1 + beta * np. log(N1 / P0))
                  # Calculate the rate of change for each part
                  dN1_dt = -k12 * N1 + k21 * (N2_0 + xi * (N2 - N2_0)) + gamma - f + delta + l
                  dN2_dt = k12 * N1 - k21 * (N2_0 + xi * (N2 - N2_0)) - k23 * N2 + k32 * N3 - N2_0
                  dN3 dt = k23 * N2 - k32 * N3 - k34 * N3 + k43 * N4
                  dN4 dt = k34 * N3 - k43 * N4 + k24 * N2 - k45 * N4
                  dN5 dt = k45 * N4 - k51 * N5
                  dN6 dt = f - k67 * N6 - 2 * delta
                  dN7 dt = k67 * N6 - k71 * N7 + delta
                  # Update the values of each section
                  N1 += dN1 dt
                  N2 += dN2_dt
                  N3 += dN3 dt
                  N4 += dN4_dt
                  N5 += dN5 dt
                  N6 += dN6 dt
                  N7 += dN7 dt
                  atmosphere.append(N1)
              results. append (atmosphere)
```

```
In [15]: plt.figure(figsize=(12, 8))

plt.scatter(co2_observations['year'], co2_observations['mean'], label='Observations',

plt.text(1850, 300, 'Calculations', fontsize=12)

plt.text(1900, 290, 'Observations', fontsize=12)

plt.text(1950, 345, 'β=0.38', fontsize=12, color='red')

plt.text(1980, 320, 'β=0.50', fontsize=12, color='blue')

# Draw the results of each beta value directly, with the colors red and blue

plt.plot(range(1750, 2001), results[0], color='red', label='β=0.38')

plt.plot(range(1750, 2001), results[1], color='blue', label='β=0.50')

plt.xlabel('Year', fontsize=14)

plt.ylabel('CO2 Concentration (ppm)', fontsize=14)

plt.legend()

plt.show()
```

