



# Arctic Mixed-Phase Cloud Dissipation and its Relationship to Low CCN Concentrations

Lucas Sterzinger, Adele L. Igel

Atmospheric Science Graduate Group  
Department of Land, Air, and Water Resources - University of California, Davis

**UCDAVIS**  
**ATMOSPHERIC SCIENCE**

## Overview

*Can a lack of environmental CCN/aerosol be a primary factor for Arctic cloud dissipation?*

- Persistent mixed-phase boundary layer clouds are important regulators for Arctic (and global) climate.
- Accurately modeling Arctic clouds are important to properly simulate the global climate system.
- Unlike in lower latitudes, Arctic aerosol concentrations have been hypothesized to be low enough to inhibit cloud formation
  - Mauritsen et al. (2011) coined the term "tenuous clouds" in which cloud structure was limited by aerosol concentration

## Cases and Simulation Setup

Data from the ASCOS field campaign and from the DOE ARM site at Oliktok Point, Alaska, analysed to find cases of potential aerosol-limited cloud dissipation.

Two potential cases have been identified:

- Oliktok Point - May 12th, 2017
  - Northern slope of Alaska - ocean/land boundary
- ASCOS - August 31st, 2008
  - Arctic ocean ice floe

The plots below show radar reflectivity (a) and aerosol concentration (b).