

BAYESIAN BEST PRACTICES

BAYESIAN STATISTICS FOR ECOLOGISTS

IGB 12. TO 19. NOVEMBER 2018

FOUR STEPS TO AN ANALYSIS

1. Specify the joint posterior distribution of outcomes (i.e., response variables) and all unknowns/parameters
2. Draw from posterior distribution using MCMC
3. Evaluate model and revise if necessary (return to step 1)
4. Use posterior predictive distribution for inference

1. JOINT POSTERIOR—LIKELIHOOD

1. JOINT POSTERIOR—LIKELIHOOD

- ▶ Specify a **generative model** (the distribution from which the observations are generated) as the likelihood

1. JOINT POSTERIOR—LIKELIHOOD

- ▶ Specify a **generative model** (the distribution from which the observations are generated) as the likelihood
 - ▶ whenever possible use knowledge of the appropriate processes

1. JOINT POSTERIOR—LIKELIHOOD

- ▶ Specify a **generative model** (the distribution from which the observations are generated) as the likelihood
 - ▶ whenever possible use knowledge of the appropriate processes
 - ▶ transform parameters and sample in transformed space to improve behaviour

1. JOINT POSTERIOR—LIKELIHOOD

- ▶ Specify a **generative model** (the distribution from which the observations are generated) as the likelihood
 - ▶ whenever possible use knowledge of the appropriate processes
 - ▶ transform parameters and sample in transformed space to improve behaviour
 - ▶ use $\log(\sigma)$ instead of σ to avoid impossible negative variance in sampler (Stan does this automagically)

1. JOINT POSTERIOR—LIKELIHOOD

- ▶ Specify a **generative model** (the distribution from which the observations are generated) as the likelihood
 - ▶ whenever possible use knowledge of the appropriate processes
 - ▶ transform parameters and sample in transformed space to improve behaviour
 - ▶ use $\log(\sigma)$ instead of σ to avoid impossible negative variance in sampler (Stan does this automagically)
- ▶ Models should be specified to be **scale independent**

1. JOINT POSTERIOR—LIKELIHOOD

- ▶ Specify a **generative model** (the distribution from which the observations are generated) as the likelihood
 - ▶ whenever possible use knowledge of the appropriate processes
 - ▶ transform parameters and sample in transformed space to improve behaviour
 - ▶ use $\log(\sigma)$ instead of σ to avoid impossible negative variance in sampler (Stan does this automagically)
- ▶ Models should be specified to be **scale independent**
 - ▶ This is most easily accomplished by scaling your predictors to have mean=0, sd=1

1. JOINT POSTERIOR—LIKELIHOOD

- ▶ Specify a **generative model** (the distribution from which the observations are generated) as the likelihood
 - ▶ whenever possible use knowledge of the appropriate processes
 - ▶ transform parameters and sample in transformed space to improve behaviour
 - ▶ use $\log(\sigma)$ instead of σ to avoid impossible negative variance in sampler (Stan does this automagically)
- ▶ Models should be specified to be **scale independent**
 - ▶ This is most easily accomplished by scaling your predictors to have mean=0, sd=1
 - ▶ Be careful scaling outcomes (y variables)—it can affect your generative model.

1. JOINT POSTERIOR—LIKELIHOOD

- ▶ Specify a **generative model** (the distribution from which the observations are generated) as the likelihood
 - ▶ whenever possible use knowledge of the appropriate processes
 - ▶ transform parameters and sample in transformed space to improve behaviour
 - ▶ use $\log(\sigma)$ instead of σ to avoid impossible negative variance in sampler (Stan does this automagically)
- ▶ Models should be specified to be **scale independent**
 - ▶ This is most easily accomplished by scaling your predictors to have mean=0, sd=1
 - ▶ Be careful scaling outcomes (y variables)—it can affect your generative model.
 - ▶ Prefer to “scale” outcomes via a **link function**

1. JOINT POSTERIOR—PRIOR

1. JOINT POSTERIOR—PRIOR

- ▶ All remaining unknowns specified with a **prior** (or possibly a hierarchical generative model and hyperpriors)

1. JOINT POSTERIOR—PRIOR

- ▶ All remaining unknowns specified with a **prior** (or possibly a hierarchical generative model and hyperpriors)
- ▶ Prefer regularising priors to vague priors

1. JOINT POSTERIOR—PRIOR

- ▶ All remaining unknowns specified with a **prior** (or possibly a hierarchical generative model and hyperpriors)
- ▶ Prefer regularising priors to vague priors
 - ▶ Normal(0, 5) instead of Normal(0,500)

1. JOINT POSTERIOR—PRIOR

- ▶ All remaining unknowns specified with a **prior** (or possibly a hierarchical generative model and hyperpriors)
- ▶ Prefer regularising priors to vague priors
 - ▶ Normal(0, 5) instead of Normal(0,500)
- ▶ Avoid improper priors: Uniform(-Inf, Inf)

1. JOINT POSTERIOR—PRIOR

- ▶ All remaining unknowns specified with a **prior** (or possibly a hierarchical generative model and hyperpriors)
- ▶ Prefer regularising priors to vague priors
 - ▶ Normal(0, 5) instead of Normal(0,500)
- ▶ Avoid improper priors: Uniform(-Inf, Inf)
- ▶ Forget conjugacy unless you know what you are doing and why

1. JOINT POSTERIOR—PRIOR

1. JOINT POSTERIOR—PRIOR

- ▶ Avoid hard boundaries

1. JOINT POSTERIOR—PRIOR

- ▶ Avoid hard boundaries
 - ▶ ~~Uniform(0, 1000)~~

1. JOINT POSTERIOR—PRIOR

- ▶ Avoid hard boundaries
 - ▶ ~~Uniform(0, 1000)~~
 - ▶ Exponential(0.1)

1. JOINT POSTERIOR—PRIOR

- ▶ Avoid hard boundaries
 - ▶ ~~Uniform(0, 1000)~~
 - ▶ Exponential(0.1)
- ▶ With informative priors, be sure to specify reasonable initial values

1. JOINT POSTERIOR—PRIOR

- ▶ Avoid hard boundaries
 - ▶ ~~Uniform(0, 1000)~~
 - ▶ Exponential(0.1)
- ▶ With informative priors, be sure to specify reasonable initial values
- ▶ Begin inference with weaker priors, gradually strengthen once you know the model is working

1. JOINT POSTERIOR—PRIOR

- ▶ Avoid hard boundaries
 - ▶ ~~Uniform(0, 1000)~~
 - ▶ Exponential(0.1)
- ▶ With informative priors, be sure to specify reasonable initial values
- ▶ Begin inference with weaker priors, gradually strengthen once you know the model is working
- ▶ Specify priors for everything—avoid defaults

1. JOINT POSTERIOR—PRIOR

- ▶ Avoid hard boundaries
 - ▶ ~~Uniform(0, 1000)~~
 - ▶ Exponential(0.1)
- ▶ With informative priors, be sure to specify reasonable initial values
- ▶ Begin inference with weaker priors, gradually strengthen once you know the model is working
- ▶ Specify priors for everything—avoid defaults
- ▶ Can be useful to draw out your model as a **digraph** to make sure you don't miss anything

2. SAMPLE FROM POSTERIOR

2. SAMPLE FROM POSTERIOR

- ▶ Start with a small run (10s-100s of iterations) to catch errors, make sure nothing surprising happens

2. SAMPLE FROM POSTERIOR

- ▶ Start with a small run (10s-100s of iterations) to catch errors, make sure nothing surprising happens
- ▶ Increase to a few thousand to view convergence

2. SAMPLE FROM POSTERIOR

- ▶ Start with a small run (10s-100s of iterations) to catch errors, make sure nothing surprising happens
- ▶ Increase to a few thousand to view convergence
- ▶ Select starting values and run until convergence (1.000s for Stan, 10.000s–100.000s or more for Metropolis)

3. EXAMINE MODEL—EFFICIENCY & CONVERGENCE

3. EXAMINE MODEL—EFFICIENCY & CONVERGENCE

- ▶ Positive-recurrent Markov chains have an asymptotic stationary distribution

3. EXAMINE MODEL—EFFICIENCY & CONVERGENCE

- ▶ Positive-recurrent Markov chains have an asymptotic stationary distribution
- ▶ Once a chain reaches this distribution, it will stay there

3. EXAMINE MODEL—EFFICIENCY & CONVERGENCE


- ▶ Positive-recurrent Markov chains have an asymptotic stationary distribution
- ▶ Once a chain reaches this distribution, it will stay there
- ▶ For Bayesian MCMCs, this distribution approximates the **joint posterior**

3. EXAMINE MODEL—EFFICIENCY & CONVERGENCE

- ▶ Positive-recurrent Markov chains have an asymptotic stationary distribution
- ▶ Once a chain reaches this distribution, it will stay there
- ▶ For Bayesian MCMCs, this distribution approximates the **joint posterior**
- ▶ **Trace plots** are one way to visually examine this property


3. TRACE PLOTS

- ▶ Low autocorrelation
- ▶ Thorough coverage of range of parameter values




3. TRACE PLOTS

- ▶ High autocorrelation; can increase Metropolis step size
- ▶ Run longer
- ▶ Use thinning (not recommended)




3. TRACE PLOTS

- ▶ Bad starting value
- ▶ Select new start
- ▶ Use burn-in (not the same as warm up!)




3. TRACE PLOTS

- ▶ A “burn-in” period selects a number of iterations to discard
- ▶ The idea is that the algorithm hasn’t “forgotten” it’s starting value
- ▶ Everything after burn-in approximates the stationary distribution



3. CONVERGENCE DIAGNOSTICS

- ▶ Using multiple chains allows us to compare within- and among-chain variance
- ▶ This is the **Gelman-Rubin statistic**; provided by Stan as r-hat
- ▶ Target value of 1.0. Less than 1.1 for all parameters is probably ok



3.5 EVALUATE AND COMPARE MODELS

3.5 EVALUATE AND COMPARE MODELS

- ▶ Before moving to inference, it is necessary to:

3.5 EVALUATE AND COMPARE MODELS

- ▶ Before moving to inference, it is necessary to:
 - ▶ select a model (or, to assign probabilities to competing models)

3.5 EVALUATE AND COMPARE MODELS

- ▶ Before moving to inference, it is necessary to:
 - ▶ select a model (or, to assign probabilities to competing models)
 - ▶ evaluate the fit of the model to data

3.5 EVALUATE AND COMPARE MODELS

- ▶ Before moving to inference, it is necessary to:
 - ▶ select a model (or, to assign probabilities to competing models)
 - ▶ evaluate the fit of the model to data
- ▶ Bayesian methods for model comparison range from fairly simple (DIC) to difficult (model averaging) to bewildering (RJ-MCMC) – will be covered Friday

3.5 EVALUATE AND COMPARE MODELS

- ▶ Before moving to inference, it is necessary to:
 - ▶ select a model (or, to assign probabilities to competing models)
 - ▶ evaluate the fit of the model to data
- ▶ Bayesian methods for model comparison range from fairly simple (DIC) to difficult (model averaging) to bewildering (RJ-MCMC) – will be covered Friday
- ▶ Model evaluation is a more complex topic – briefly on Friday, in more detail if time allows

4. INFERENCE

4. INFERENCE

- ▶ There are multiple levels of inference

4. INFERENCE

- ▶ There are multiple levels of inference
- ▶ Examine posterior distributions of **parameters** (most informative for simple models)

4. INFERENCE

- ▶ There are multiple levels of inference
- ▶ Examine posterior distributions of **parameters** (most informative for simple models)
- ▶ **Posterior predictive distributions** – mean and standard error of outcomes

4. INFERENCE

- ▶ There are multiple levels of inference
- ▶ Examine posterior distributions of **parameters** (most informative for simple models)
- ▶ **Posterior predictive distributions** – mean and standard error of outcomes
- ▶ **Posterior predictive simulations** – generate new outcomes from model (incorporates all uncertainty, including process error)