Annex 4 – A. Policy adoption, 2024

WHO region Country/area	Testing	Treatment				Chemoprevention		
	Malaria diagnosis with microscopy or RDT is free in public sector	ACT for treatment of malaria is free in public sector	ACT is delivered in the community	Single low dose of PQ with ACT to reduce transmissibility of *P. falciparum*[1]	Pre-referral treatment with rectal artesunate suppositories at community level	IPTp is used	c-IPTp is used (aligned with WHO recommendation)	IPTsc is used
AFRICAN								
Angola	🟢	🟢	🟢	🟣	⚫	🟢	⚫	⚫
Benin	🟢	🟢	⚫	⚫	⚫	🟢	⚫	⚫
Botswana	🟢	🟢	⚫	🟢	⚫	NA	NA	NA
Burkina Faso	🟢	🟢	🟢	⚫	🟢	🟢	🟢	⚫
Burundi	🟢	🟢	🟢	🟣	🟢	🟢	🟢	⚫
Cameroon	⚫	🟢	🟢	⚫	🟢	🟢	⚫	⚫
Central African Republic	🟢	🟢	🟢	⚫	🟢	🟢	🟢	⚫
Chad	🟢	🟢	🟢	⚫	⚫	🟢	🟢	⚫
Comoros	🟢	🟢	🟢	🟣	⚫	⚫	⚫	⚫
Congo	🟢	🟢	🟢	⚫	⚫	🟢	⚫	⚫
Côte d'Ivoire	🟢	–	–	⚫	🟣	🟢	⚫	⚫
Democratic Republic of the Congo	🟢	🟢	🟢	⚫	🟢	🟢	🟢	⚫
Equatorial Guinea	🟣	🟣	⚫	⚫	⚫	🟢	⚫	⚫
Eritrea	🟢	🟢	⚫	🟣	🟢	⚫	⚫	⚫
Eswatini	🟢	🟢	⚫	🟢	🟢	NA	NA	NA
Ethiopia	🟢	🟢	🟢	🟣	🟢	⚫	⚫	⚫
Gabon	🟢	🟢	⚫	⚫	🟢	🟢	🟢	⚫
Gambia	🟢	🟢	🟢	🟢	⚫	🟢	⚫	⚫
Ghana	🟢	🟢	⚫	⚫	🟣	🟢	⚫	⚫
Guinea	🟢	🟢	🟢	⚫	🟢	🟢	🟢	🟢
Guinea-Bissau	🟢	🟢	🟢	🟣	🟢	🟢	⚫	⚫
Kenya	🟢	🟢	🟢	⚫	⚫	🟢	⚫	⚫
Liberia	🟢	🟢	🟢	⚫	🟢	🟢	⚫	⚫
Madagascar	🟢	🟢	🟢	🟣	🟢	🟢	🟢	⚫
Malawi	🟢	🟢	🟢	⚫	🟢	🟢	⚫	⚫
Mali	🟢	🟢	🟢	⚫	🟢	🟢	🟢	⚫
Mauritania	🟢	–	–	⚫	🟣	🟢	⚫	⚫
Mozambique	🟢	🟢	🟢	⚫	🟢	🟢	🟢	🟢
Namibia	🟢	🟢	🟢	🟣	⚫	NA	NA	NA
Niger	🟢	–	–	⚫	🟣	🟢	⚫	⚫
Nigeria[3]	⚫	⚫	🟢	⚫	🟢	🟢	⚫	⚫
Rwanda	🟢	–	–	⚫	🟣	NA	NA	NA
Sao Tome and Principe	🟢	🟢	🟢	🟣	⚫	🟣	⚫	⚫
Senegal	🟢	🟢	🟢	🟢	🟢	🟢	🟢	⚫
Sierra Leone	🟢	🟢	🟢	⚫	🟣	🟢	🟣	⚫
South Africa	🟢	🟢	🟢	🟢	⚫	NA	NA	NA
South Sudan[4]	🟢	🟣	🟣	⚫	🟣	🟢	⚫	⚫
Togo	🟢	🟢	🟢	⚫	⚫	🟢	⚫	⚫
Uganda	🟢	🟢	🟢	⚫	🟢	🟢	🟣	🟣
United Republic of Tanzania[5]								
Mainland	🟢	🟢	🟢	🟣	⚫	🟢	⚫	🟢
Zanzibar	🟢	🟢	🟢	🟢	⚫	NA	NA	NA
Zambia[3]	🟢	🟢	🟢	🟣	🟢	🟢	⚫	⚫
Zimbabwe	🟢	🟢	🟢	🟣	🟣	🟢	⚫	⚫
AMERICAS								
Bolivia (Plurinational State of)	🟣	⚫	⚫	⚫	⚫	NA	NA	NA
Brazil	🟢	🟢	🟢	🟢	⚫	NA	NA	NA
Colombia	🟢	🟢	🟢	🟢	⚫	NA	NA	NA
Costa Rica	🟢	🟢	⚫	⚫	⚫	NA	NA	NA
Dominican Republic	🟢	🟢	🟢	🟣	⚫	NA	NA	NA
Ecuador	🟢	🟢	🟢	🟢	⚫	NA	NA	NA
French Guiana	🟢	🟢	⚫	🟣	⚫	NA	NA	NA
Guatemala	🟢	🟢	–	NA	NA	NA	NA	NA
Guyana	🟢	–	–	🟣	🟣	NA	NA	NA
Haiti	🟢	⚫	⚫	⚫	⚫	NA	NA	NA
Honduras	🟢	🟢	⚫	⚫	⚫	NA	NA	NA
Mexico	🟢	🟣	🟣	🟢	⚫	NA	NA	NA
Nicaragua	🟢	🟢	⚫	⚫	⚫	NA	NA	NA
Panama	🟢	🟣	⚫	🟣	⚫	NA	NA	NA
Peru	🟢	🟣	🟣	🟣	⚫	NA	NA	NA
Venezuela (Bolivarian Republic of)	🟢	–	–	🟢	⚫	NA	NA	NA

Chemoprevention		Surveillance		Vector control					
SMC is used	PMC is used	Malaria is a notifiable disease	Case reporting from the private sector is mandatory	WHO-recommended ITNs are distributed	ITNs are distributed free of charge to all age groups through mass campaigns[2]	ITNs are distributed free of charge to all age groups through routine channels	Durability of ITNs is monitored	IRS is used by the NMP	Insecticide resistance is monitored and reported to WHO
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
NA	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	–	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
NA	●	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	–	–	●	–	●	–	●	–
●	●	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	–	●	●
●	●	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	–	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	–
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
NA	●	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	–	●	–
NA	NA	●	●	●	●	●	–	●	–
NA	NA	●	●	●	●	●	–	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	–	●	–

Annex 4 – A. Policy adoption, 2024

WHO region Country/area	Testing	Treatment				Chemoprevention		
	Malaria diagnosis with microscopy or RDT is free in public sector	ACT for treatment of malaria is free in public sector	ACT is delivered in the community	Single low dose of PQ with ACT to reduce transmissibility of *P. falciparum*[1]	Pre-referral treatment with rectal artesunate suppositories at community level	IPTp is used	c-IPTp is used (aligned with WHO recommendation)	IPTsc is used
EASTERN MEDITERRANEAN								
Afghanistan	🟢	🟢	🟢	🟢	⚫	NA	NA	NA
Djibouti	🟢	🟢	🟣	🟢	⚫	NA	NA	NA
Iran (Islamic Republic of)	🟢	🟢	🟢	🟢	🟢	NA	NA	NA
Pakistan	🟢	🟢	🟢	🟢	⚫	NA	NA	NA
Somalia	🟢	🟢	🟢	🟢	🟢	🟢	⚫	⚫
Sudan	🟢	🟢	🟢	⚫	🟣	🟢	⚫	⚫
Yemen	🟢	🟢	🟢	🟢	⚫	NA	NA	NA
SOUTH-EAST ASIA								
Bangladesh	🟢	🟢	🟢	🟢	🟢	NA	NA	NA
Democratic People's Republic of Korea	🟢	NA	NA	NA	NA	NA	NA	NA
India	🟢	🟢	🟢	🟣	⚫	NA	NA	NA
Myanmar	🟢	🟢	🟢	🟢	🟣	NA	NA	NA
Nepal	🟢	🟢	⚫	🟢	⚫	NA	NA	NA
Thailand	🟢	🟢	🟢	🟢	⚫	NA	NA	NA
WESTERN PACIFIC								
Cambodia	🟢	🟢	🟢	🟢	🟣	NA	NA	NA
Indonesia[6]	🟢	🟢	🟢	🟢	⚫	NA	NA	NA
Lao People's Democratic Republic	🟢	🟢	🟢	🟢	⚫	NA	NA	NA
Papua New Guinea	🟢	🟢	🟢	🟣	🟢	🟣	🟣	⚫
Philippines	🟢	🟢	🟢	🟢	⚫	NA	NA	NA
Republic of Korea	🟢	NA	NA	NA	NA	NA	NA	NA
Solomon Islands	🟢	–	–	🟣	🟣	🟣	⚫	⚫
Vanuatu	🟢	🟢	🟢	🟣	🟣	NA	NA	NA
Viet Nam	🟢	🟢	🟢	🟢	⚫	NA	NA	NA

ACT: artemisinin-based combination therapy; c-IPTp: community-based delivery of IPTp; IPTp: intermittent preventive treatment of malaria in pregnancy; IPTsc: intermittent preventive treatment of malaria in school-aged children; IRS: indoor residual spraying; ITN: insecticide-treated mosquito net; NMP: national malaria programme; *P.: Plasmodium*; PMC: perennial malaria chemoprevention; PQ: primaquine; RDT: rapid diagnostic test; SMC: seasonal malaria chemoprevention; WHO: World Health Organization.

[1] Single dose of PQ (0.75 mg base/kg) for countries in the WHO Region of the Americas.

[2] Mass campaigns do not include targeted campaigns directed towards specific sub-groups in the population.

[3] Data reported to the Alliance for Malaria Prevention were used where data reported to WHO were missing or incomplete.

[4] In May 2013, South Sudan was reassigned to the WHO African Region (resolution WHA66.21, https://apps.who.int/gb/ebwha/pdf_files/WHA66/A66_R21-en.pdf).

[5] Where national data for the United Republic of Tanzania are unavailable, refer to Mainland and Zanzibar.

[6] As of 27 May 2025, Indonesia has been reassigned to the WHO Western Pacific Region (resolution WHA78.25, https://apps.who.int/gb/ebwha/pdf_files/WHA78/A78_R25-en.pdf).

Method: Information on existing policies and whether they were implemented in 2024 was reported by NMPs. Policy implementation in 2024 was adjusted for the following variables, based on whether supporting data were available and reported by NMPs to the world malaria report database: distribution of ITNs; IPTp; IPTsc; SMC; PMC; RDTs used at community level and ACTs used for the treatment of *Plasmodium falciparum* infection. A setting of "not applicable" was automatically assigned to countries where the following interventions did not apply; IPTp, IPTpsc, SMC and PMC.

Chemoprevention		Surveillance		Vector control					
SMC is used	PMC is used	Malaria is a notifiable disease	Case reporting from the private sector is mandatory	WHO-recommended ITNs are distributed	ITNs are distributed free of charge to all age groups through mass campaigns[2]	ITNs are distributed free of charge to all age groups through routine channels	Durability of ITNs is monitored	IRS is used by the NMP	Insecticide resistance is monitored and reported to WHO
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
●	●	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	NA	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	●	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	–
NA	●	●	–	●	●	●	–	●	–
NA	NA	●	●	●	●	●	●	●	●
NA	NA	●	●	●	●	●	●	●	●

Data as of 7 November 2025

● = Policy exists and has been implemented this year.

● = Policy exists but was not implemented this year or no data exist to support implementation.

● = Policy does not exist or policy has been discontinued.

NA = Policy does not apply.

– = Question not answered, and there is no information from previous years.

Annex 4 – B. Antimalarial drug policy in malaria endemic countries and areas, 2024

WHO region Country/area	*P. falciparum*				*P. vivax*
	Uncomplicated unconfirmed	Uncomplicated confirmed	Severe	Prevention during pregnancy	Treatment
AFRICAN					
Angola	AL; DHA-PPQ; AS+AQ	AL; DHA-PPQ; AS+AQ	AS	SP(IPT)	AL; DHA-PPQ; AS+AQ
Benin	NA	AL	AS	SP(IPT)	NA
Botswana	NA	AL-PQ	AS	NA	AL-PQ
Burkina Faso	AL; AS+AQ	AL; AS-PY; AS+AQ; DHA-PPQ	AS	SP(IPT)	NA
Burundi	AL	AL	ART; AS	SP(IPT)	NA
Cameroon	NA	AL; AS-PY; AS+AQ; DHA-PPQ	AS; AM	SP(IPT)	NA
Central African Republic	AL	AL	AS	SP(IPT)	NA
Chad	AL; AS+AQ	AL; AS+AQ	ART; AS; QN	SP(IPT)	NA
Comoros	AL	AL; AL+PQ	AS	SP(IPT)	NA
Congo	AL	AL	AS	SP(IPT)	NA
Côte d'Ivoire	AL; AS+AQ	AL; AS+AQ	QN	SP(IPT)	NA
Democratic Republic of the Congo	AL; AS+AQ	AL; AS-PY; AS+AQ	AS	SP(IPT)	NA
Equatorial Guinea	AL	AL	AS	SP(IPT)	NA
Eritrea	AS+AQ	AS+AQ	AS	NA	AS+AQ
Eswatini	NA	AL-PQ	AS	NA	PQ
Ethiopia	AL+PQ	AL-PQ	AS; AL+PQ	NA	CQ+PQ
Gabon	AL; DHA-PPQ; AS+MQ; AS+AQ	AL; DHA-PPQ; AS+MQ; AS+AQ	AS	SP(IPT)	NA
Gambia	AL	AL+PQ	AS	SP(IPT)	NA
Ghana	AL; AS+AQ; DHA-PPQ	AL; AS+AQ	AM; AS; QN	SP(IPT)	AL+PQ; DHA-PPQ+PQ
Guinea	AL	AL	ART+AL	SP(IPT)	NA
Guinea-Bissau	NA	AL	AS	SP(IPT)	NA
Kenya	AL	AL	AS	SP(IPT)	AL-PQ
Liberia	AS+AQ	AL; AS+AQ	AM; AS; QN	SP(IPT)	NA
Madagascar	AS+AQ	AS+AQ	AS	SP(IPT)	AS+AQ
Malawi	AL	AL	AS	SP(IPT)	NA
Mali	AL	AL	AS	SP(IPT)	NA
Mauritania	AS+AQ	AS+AQ	AS	SP(IPT)	NA
Mozambique	AL	AS+AQ	AS	SP(IPT)	NA
Namibia	NA	AL+PQ	AS	NA	AL+PQ
Niger	AL	AL	AS	SP(IPT)	AL
Nigeria	AL; AS+AQ	AL; AS-PY; AS+AQ; DHA-PPQ	AS	SP(IPT)	NA
Rwanda	AL	AL	AS; QN	NA	NA
Sao Tome and Principe	NA	AS+MQ+PQ	AS	SP(IPT)	NA
Senegal	NA	AL; AS+AQ+PQ	AS	SP(IPT)	NA
Sierra Leone	AL	AL	AS	SP(IPT)	NA
South Africa	AL	AL-PQ	AS; QN	NA	AL
South Sudan[1]	AS+AQ	AS+AQ	AS	SP(IPT)	AS-PY
Togo	NA	AL	AS; AM; QN	SP(IPT)	NA
Uganda	NA	AL	AS	SP(IPT)	AL
United Republic of Tanzania					
Mainland	AL; AS+AQ; DHA-PPQ	AL; AS+AQ; DHA-PPQ	AS; AL	SP(IPT)	NA
Zanzibar	NA	AS+AQ+PQ	AS	NA	PQ
Zambia	AL	AL	AS	SP(IPT)	NA
Zimbabwe	NA	AL	AS	SP(IPT)	NA
AMERICAS					
Bolivia (Plurinational State of)	NA	AL+PQ	AS	NA	CQ+PQ
Brazil	NA	AL+PQ; AS+MQ+PQ	AS	CQ	CQ+PQ; CQ+TQ
Colombia	AL+PQ	AL; AL-PQ	AS	CQ	CQ+PQ
Costa Rica	CQ+PQ	CQ+PQ	AS	NA	CQ+PQ

WHO region Country/area	*P. falciparum*				*P. vivax*
	Uncomplicated unconfirmed	Uncomplicated confirmed	Severe	Prevention during pregnancy	Treatment
AMERICAS					
Dominican Republic	NA	CQ+PQ	AS	NA	CQ+PQ
Ecuador	NA	AL+AM+PQ	AS	NA	CQ+PQ
French Guiana	NA	AL; DHA-PPQ	AS	NA	CQ+PQ
Guatemala	CQ+PQ	AL	AS	NA	CQ+PQ
Guyana	NA	AL+PQ	AM; AS-QN; QN+CL	NA	CQ+PQ
Haiti	NA	CQ+PQ	AS	NA	CQ+PQ
Honduras	NA	CQ+PQ	AS	NA	CQ+PQ
Mexico	NA	AL; AL-PQ	AL; AL+PQ; AS	NA	CQ+PQ
Nicaragua	CQ+PQ	CQ+PQ	AS	CQ	CQ+PQ
Panama	AL-PQ	AL+PQ	AS	NA	CQ+PQ
Peru	AS+MQ+PQ	AS+MQ+PQ	AS+PQ+CL	NA	CQ+PQ
Venezuela (Bolivarian Republic of)	NA	AL+PQ	AS	NA	CQ+PQ
EASTERN MEDITERRANEAN					
Afghanistan	CQ	AL+PQ	AM; AS; QN	NA	CQ+PQ
Djibouti	NA	AL+PQ	AS	NA	AL+PQ
Iran (Islamic Republic of)	NA	AL-PQ	AS	NA	CQ+PQ
Pakistan	CQ	AL-PQ	AS	NA	CQ+PQ
Somalia	AL	AL-PQ	AS	SP(IPT)	AL+PQ
Sudan	NA	AL	AS; QN	SP(IPT)	AL+PQ
Yemen	AL	AL+PQ	AS	NA	AL-PQ
SOUTH-EAST ASIA					
Bangladesh	NA	AL+PQ	AS; AL+PQ	NA	CQ+PQ
Democratic People's Republic of Korea	NA	NA	NA	NA	CQ+PQ
India	NA	AL+PQ; AS+SP+PQ	AS	NA	CQ+PQ
Myanmar	NA	AL+PQ	ART+AL	NA	CQ+PQ
Nepal	CQ+PQ; AL+PQ	AL-PQ	AS	NA	CQ+PQ
Thailand	NA	DHA-PPQ+PQ; AS-PY	CQ+PQ	NA	CQ+PQ
WESTERN PACIFIC					
Cambodia	AS+MQ	AS+MQ+PQ	AS	NA	AS+MQ+PQ
Indonesia[2]	NA	DHA-PPQ+PQ	AS	NA	DHA-PPQ+PQ
Lao People's Democratic Republic	NA	AL+PQ	AS	NA	AL-PQ
Papua New Guinea	AL+PQ	AL+PQ	DHA-PPQ	SP(IPT)	AL-PQ
Philippines	NA	AL+PQ	AS	NA	AL+PQ
Republic of Korea	AT-PG; PY-AS; MQ	PY; MQ	AS; MQ; AT-PG	NA	CQ+PQ
Solomon Islands	AL+PQ	AL+PQ	AS	CQ	AL+PQ
Vanuatu	AL+PQ	AL+PQ	AS	CQ	AL+PQ
Viet Nam	AS-PY; DHA-PPQ+PQ	DHA-PPQ+PQ	AS	AL	CQ+PQ

Data as of 29 September 2025

AL: artemether–lumefantrine; AM: artemether; AQ: amodiaquine; ART: artemisinin; AS: artesunate; AT-PG: atovaquone–proguanil; CL: clindamycin; CQ: chloroquine; DHA-PPQ: dihydroartemisinin–piperaquine; IPT: intermittent preventive treatment of malaria; MQ: mefloquine; NA: not applicable; *P*: *Plasmodium*; PQ: primaquine; PY: pyronaridine; QN: quinine; SP: sulfadoxine–pyrimethamine; TQ: tafenoquine; WHO: World Health Organization.

[1] In May 2013, South Sudan was reassigned to the WHO African Region (resolution WHA66.21, https://apps.who.int/gb/ebwha/pdf_files/WHA66/A66_R21-en.pdf).

[2] As of 27 May 2025, Indonesia has been reassigned to the WHO Western Pacific Region (resolution WHA78.25, https://apps.who.int/gb/ebwha/pdf_files/WHA78/A78_R25-en.pdf).

Notes:
Co-blistered products are denoted by "+" (e.g. artesunate + amodiaquine).
Co-formulated products are denoted by "–" (e.g. artemether–lumefantrine).