Empirical measures in dynamical systems

Lua Viana Reis

December 3, 2025

"Everyone else has an attractor — why can't we have one of our own? Stranger than all the rest. Look at it: it's the most beautiful thing on the map."

— Maria, "Unstable Orbits in the Space of Lies" (Greg Egan).

Contents

Contents ii

1 Ergodic theory and dynamics **1**

1.1 A categorical zoo . . . 1

Examples . . . 2

Invariant states and observables . . . 3

1.2 Ergodic theory and ergodicity . . . 4

Ergodic theorems . . . 4

1.3 Extensions . . . 6

Invertible extensions . . . 7

The solenoid . . . 9

1.4 Cocycles and skew products . . . 9

Lyapunov exponents . . . 9

1.5 Empirical measures . . . 9

1.6 Partitions . . . 10

Lattice structure . . . 11

An interlude of zero measure . . . 11

1.7 Disintegrations . . . 11

Uniqueness . . . 12

Measurability . . . 12

Conditional measures . . . 13

1.8 Entropy . . . 13

Topological entropy . . . 13

Measure-theoretic entropy . . . 13

Pesin entropy formula . . . 13

2 Schwarzian derivatives and cylinder maps **14**

2.1 Cross ratios and the Schwarzian derivative . . . 14

2.2 Skew products . . . 18

2.3 Transversal Lyapunov exponents . . . 18

2.4 Intermingled basins . . . 20

2.5 Negative Schwarzian . . . 22

Hausdorff dimension . . . 24
2.6 Positive Schwarzian . . . 24

3 Historical behavior **26**
3.1 History . . . 26
Genericity of historical orbits . . . 27
3.2 Martingale-coboundary decomposition . . . 27
3.3 Historical behavior for skew products . . . 29
3.4 Recent developments . . . 31

4 Partially hyperbolic dynamics **32**
4.1 Partial hyperbolicity . . . 32
4.2 Invariance Principle . . . 32

A Functional Analysis **33**

B Measure and probability **34**
B.1 Probability spaces . . . 34
Products . . . 34
Borel Spaces . . . 34
The Kolmogorov Extension Theorem . . . 34
B.2 Random variables . . . 34
B.3 Weak convergence . . . 35
B.4 Brownian motion . . . 35
The space of càdlàg functions . . . 36
The arcsine law . . . 37
Weak invariance principles . . . 38
Reverse martingale difference sequences . . . 40

C A categorical lens **42**
C.1 A category . . . 42
C.2 Functors . . . 42
C.3 Natural transformations . . . 42
C.4 Limits and colimits . . . 42

Bibliography **45**

Chapter 1

Ergodic theory and dynamics

1.1 A categorical zoo

The starting point of the field of dynamical systems is a space X, endowed with some structure, along with some notion of "time evolution", which could be a map $X \to X$ preserving the said structure. For instance, in topological dynamics, X is usually a topological space and the time evolution is given by a continuous map, while in ergodic theory the space is endowed with the structure of a measure and the maps are measure-preserving. But it is also of equal interest to the field to look at more versatile settings, like the ones in which there are multiple maps — or perhaps, the ones in which the maps do not send send each given state to a single next state, but to a set, or distribution, of future possibilities.

Of course, we could have a separate definition for each (and that is usually what we do). But it is also fun to see how general we can make a reasonable definition that still conveys what most such systems are. One interesting way to do that, which nicely bundles in a single notion the totality of the examples above, is the one of a *monoid object* in a category.

Following [1], we can define a monoid object in a category C as a functor $M \to C$, where M is *monoid*, that is, a category with a single object. An usual monoid is just a group without the inverse operation, namely, a set M endowed with a special identity element 1_M and a multiplication $(a, b) \in X \times X \mapsto a \cdot b \in X$ satisfying

i. $1_M \cdot a = a = a \cdot 1_M$ for all $a \in M$;

ii. $(a \cdot b) \cdot c = a \cdot (b \cdot c)$ for all $a, b, c \in M$.

Given such a monoid M, the category composed of a single object $\bigstar$, set of morphisms $\mathrm{Hom}(\bigstar, \bigstar) = M$ and composition given by the monoid multiplication will satisfy the definition of a category. Conversely, any (small) category with a single object $\bigstar$ can be seen as a monoid, by taking the carrier set to be the set $M = \mathrm{Hom}(\bigstar, \bigstar)$ and the product, to be the composition. Let's see how many familiar examples of dynamics fit neatly in this definition.

Examples

One map

The simplest example is that of a set X and a map $f\colon X \to X$. This fits the categorical definition by taking $C = \mathbf{Set}$ and $M = \mathbb{N}$ (the natural numbers under addition, which forms a monoid). The functor $F\colon M \to \mathbf{Set}$ sends the single object $\bigstar$ of M to X, and each morphism $n \in \mathbb{N} = \mathrm{Hom}(\bigstar, \bigstar)$ to the n-fold composition $f^n = f \circ f \circ \cdots \circ f$ (with $f^0 = \mathrm{id}_X$). The functor axioms ensure that $F(0) = \mathrm{id}_X$ and $F(n+m) = F(n) \circ F(m) = f^n \circ f^m = f^{n+m}$, which is precisely the monoid homomorphism property.

Two maps

To illustrate the definition, if $f_a, f_b\colon X \to X$ are two measure-preserving maps, then the associated monoid is the one-object category for the free monoid with two generators $\{a, b\}$, and the functor sends $\bigstar$ to the object X and words to the compositions of f_a and f_b in the category of measure-preserving maps. For instance, the word *abba* is sent to the map $f_a \circ f_b \circ f_b \circ f_a\colon X \to X$.

Markov kernels

Another example is that of Markov kernels, or Markov chains. Often, it is useful to model stochastic dynamics by having the maps send each point $x \in X$ not to a point $f(x) \in X$, but to a *probability measure* $f(x) \in \mathcal{M}_1(X)$. The probability measure $f(x)$ then encodes the distribution of expected outcomes given that we started in the state x.

How we can compose iterates of f and regard it as a dynamic on X? Semantically, we want the iterates to encode the distribution of expected outcomes after n steps given that the system started at the point x. To work it out, we can start with the case where X is finite, so any $f(x) \in \mathcal{M}_1(X)$ is completely determined by some transition probabilities $\mathbb{P}(f(x) = y) := f(x)(\{y\})$. Then, supposing that the probability of first making the transition $x \mapsto y$ and then $y \mapsto z$ is $\mathbb{P}(f(x) = y) \cdot \mathbb{P}(f(y) = z)$, we should have

$$\mathbb{P}(f^2(x) = z) = \sum_{y \in X} \mathbb{P}(f(x) = y) \cdot \mathbb{P}(f(y) = z).$$

If we rewrite the above with a measure-theoretic language, we see that we ought to define

$$f^2(x)(E) = \int_X f(y)(E)\ d(f(x))\{y\}.$$

More generally, given $f\colon X \to \mathcal{M}_1(Y)$, $g\colon Y \to \mathcal{M}_1(Z)$, we can define a kernel composition by

$$(g \circ f)(x)(E) = \int_X g(y)(E)\ d(f(x))\{y\}$$

so that $g \circ f\colon X \to \mathcal{M}_1(Z)$.

A map with type $X \to \mathcal{M}_1(Y)$ is called a *Markov kernel* between the measurable spaces X and Y. The category whose objects are measurable spaces and morphisms are Markov kernels,

with compositions given by the rule above, will be called **Stoch**. We remark that the identity morphisms in this category are the Dirac masses $\delta(x) = \delta_x \in \mathcal{M}_1(X)$.

An interesting aspect of this category is that deterministic measurable maps still live inside it — given $g\colon X \to Y$ a measurable function, $f_g(x) = \delta_y$ is a Markov kernel. That is to say, this correspondence defines a (faithful) functor from the category **Meas** of measurable functions to **Stoch**. This means that dynamics in **Stoch** can be seen a generalization of deterministic measurable systems, where we now also allow the maps to be stochastic.

Another observation is that a probability measure μ in a measurable space X can be represented in this category by a Markov kernel $m\colon \{\star\} \to \mathcal{M}_1(X)$, where $\{\star\}$ is a measurable space with one element. Namely, we set $m(\star) = \mu$.

Invariant states and observables

Two important concepts in the study of dynamical systems are the notions of invariant states and invariant observables. In [1], a systematic treatment of those objects is suggested using the categorical language which we have been developing here.

A *cone* over a functor $F\colon \mathcal{D} \to \mathcal{C}$ is defined as an object $C \in \mathcal{C}$ and a natural transformation $T\colon \Delta_C \Rightarrow F$, where $\Delta_C\colon \mathcal{D} \to \mathcal{C}$ is the constant functor at the identity of C. In the case we are interested, $\mathcal{D} = M$ is a monoid and F is given by the dynamics. This means a cone is given as an object $C \in \mathcal{C}$ and a morphism $c\colon C \to X$ with the property that for every $m \in M$,

$$\begin{tikzcd} & C \arrow[dl, "c"'] \arrow[dr, "c"] & \\ X \arrow[rr, "m"] & & X \end{tikzcd}$$

commutes.

A cone (C, c) is *universal* for $M \to \mathcal{C}$ if it is a limit of the above diagram, that is, for every other cone (D, d), there is a unique arrow $D \to C$ so that

$$\begin{tikzcd} & D \arrow[d, dashed] \arrow[ddl, "d"'] \arrow[ddr, "d"] & \\ & C \arrow[dl] \arrow[dr] & \\ X \arrow[rr, "m"] & & X \end{tikzcd}$$

commutes. If a cone is to be seen as a collection of invariant states, then the universal cone is the biggest of them. As we shall see in appendix C.4, a universal cone is a special case of a *limit*. Since categorical limits are unique, if a universal cone exists, it is unique up to isomorphism as well.

Example 1.1. Let $f\colon X \to X$ be any function from a set to itself. Like in previous example, this will define a dynamical system in **Set** over the monoid $\mathbb{Z}_{\geq 0}$, mapping $n \in \mathbb{Z}_{\geq 0}$ to the iterate f^n. If $x \in X$ is a fixed point of f, then we get a cone by defining $C = \{x\}$ and picking the morphism $c\colon C \to X$ to be the inclusion. Since x is a fixed point, it's clear that the associated diagram will commute. Conversely, if (C, c) is a cone and $x \in C$, then the commutativity of the triangle means $f(c(x)) = c(x)$, and therefore $c(x) \in X$ is a fixed point for f.

In this case, the *universal* cone for f, then, will correspond to the set $\mathrm{Fix}(f)$ of all fixed points (which may be empty) paired with the set inclusion $\mathrm{Fix}(f) \hookrightarrow X$. Any other cone will map uniquely to this set, by matching each of its points with the corresponding fixed point in $\mathrm{Fix}(f)$.

Example 1.2. Let $f\colon X \to \mathcal{M}_1(X)$ be a *Markov kernel* from a measurable set to itself. Examples of cones for the associated system in **Stoch** over $\mathbb{Z}_{\geq 0}$ are *stationary measures* $m\colon \{\star\} \to \mathcal{M}_1(X)$, that is, measures with the property that

$$(f \circ m)(\star) = \int_X f(x)\, d\mu\{x\} = \mu,$$

where $\mu = m(\star)$. In other words, the following diagram commutes in the category **Stoch**:

$$\begin{tikzcd} & \{\star\} \arrow[dl, "m"'] \arrow[dr, "m"] & \\ X \arrow[rr, "f"] & & X \end{tikzcd}$$

More generally, a kernel $g\colon Y \to \mathcal{M}_1(X)$ will define a cone if for every $y \in Y$, $g(y) \in \mathcal{M}_1(X)$ is invariant. As we shall see later, if X is a Borel space and f comes from a deterministic map (that is, every $f(x)$ is a Dirac mass), then the universal cone exists and is given by what is called the *ergodic decomposition*.

Example 1.3. For an iterated function system of contractions, the universal cone is the Hutchinson attractor. It is the map sending $\star$ to the compact set $A = \lim_{n\to\infty} H^i(x)$, which is independent of our starting point x.

1.2 Ergodic theory and ergodicity

Ergodic theorems

The fundamental tools in ergodic theory.

Below we reproduce a succinct proof of the pointwise ergodic theorem, described for example in [2, p. 136][1].

Theorem 1.4 (Birkhoff Ergodic Theorem)**.** *Let μ be an invariant probability measure and $\phi\colon X \to \mathbb{R}$ an integrable function. Then, for μ-almost every $x \in X$,*

$$\frac{1}{n}\sum_{i=0}^{n-1} \phi(f^i(x)) \to \mathbb{E}_\mu(\phi \mid \mathcal{I})(x),$$

where $\mathcal{I}$ is the sub-σ-algebra of (strictly) invariant sets.

Proof. Let

$$F_n(x) = \max_{1\leq k\leq n} \sum_{i=0}^{k} \phi(f^i(x)).$$

[1] This is not exactly relevant to this dissertation, but the author has formally verified this proof in the Lean 4 ITP. The code is currently available at `https://github.com/lua-vr/pointwise-birkhoff`. It is in the process of being upstreamed to the Mathlib library.

Claim 1. $F_{n+1} = \phi + \max\{0, F_n \circ f\}$.

To see this, note that $\sum_{i=0}^{k} \phi \circ f^i = \phi + \sum_{i=0}^{k-1} \phi \circ f^{i+1}$, so

$$F_{n+1} = \phi + \max_{1 \le k \le n+1} \sum_{i=0}^{k-1} \phi \circ f^{i+1} = \phi + \max\left\{0, \max_{1 \le k \le n} \sum_{i=0}^{k} \phi \circ f^{i+1}\right\} = \phi + \max\{0, F_n \circ f\}.$$

Claim 2. $F_{n+1} - F_n \circ f = \phi - \min\{0, F_n \circ f\}$.

This claim follows from the previous one by rewriting

$$F_{n+1} + \min\{0, F_n \circ f\} = \phi + \max\{0, F_n \circ f\} + \min\{0, F_n \circ f\} = \phi + F_n \circ f.$$

Two very useful corollaries of the claim are that for each $x \in X$, $F_{n+1}(x) - F_n(f(x))$ is a non-increasing sequence (because $F_n(f(x))$ is non-decreasing by definition), and that if $F_n(f(x)) \to \infty$, then the sequence will converge monotonically to $\phi(x)$.

Now, we define the set

$$S = \{x \in X \mid F_n(x) \to +\infty\}.$$

The first observation is that $S \in \mathcal{I}$, because it's measurable and strictly invariant. Since μ is invariant by f, this implies $\int_S F_n \circ f \, d\mu = \int_S F_n \, d\mu$. Also, the observations from the last claim show that $F_{n+1}(x) - F_n(f(x)) \to \phi(x)$ for $x \in S$. Thus, by the dominated convergence theorem,

$$0 \le \int_S F_{n+1} - F_n \, d\mu = \int_S F_{n+1} - F_n \circ f \, d\mu \to \int_S \phi \, d\mu \tag{1.1}$$

which means $\int_S \phi \, d\mu \ge 0$.

The second observation is that if $x \notin S$, then $(F_n(x))_n$ is bounded above, so

$$\limsup_{n\to\infty} \frac{1}{n} \sum_{i=0}^{n-1} \phi(f^i(x)) \le \limsup_{n\to\infty} \frac{1}{n} F_n(x) \le 0. \tag{1.2}$$

Putting the two together, we conclude that if ϕ is such that $\mathbb{E}(\phi \mid \mathcal{I}) < 0$ almost everywhere, then by eq. (1.1) and the fact $S \in \mathcal{I}$, the set S has measure zero, so eq. (1.2) is valid almost everywhere.

The theorem then follows by noticing that for any $\varepsilon > 0$, if we replace ϕ by $\tilde{\phi} = \phi - \mathbb{E}(\phi \mid \mathcal{I}) - \varepsilon$, then $\mathbb{E}(\tilde{\phi} \mid \mathcal{I}) < 0$, so

$$\limsup_{n\to\infty} \frac{1}{n} \sum_{i=0}^{n-1} \tilde{\phi}(f^i(x)) = \limsup_{n\to\infty} \frac{1}{n} \sum_{i=0}^{n-1} \phi(f^i(x)) - \mathbb{E}(\phi \mid \mathcal{I}) - \varepsilon \le 0.$$

If we combine this with the analogous inequality for $-\phi$, we get

$$\mathbb{E}(\phi \mid \mathcal{I}) - \varepsilon \le \liminf_{n\to\infty} \frac{1}{n} \sum_{i=0}^{n-1} \phi(f^i(x)) \le \limsup_{n\to\infty} \frac{1}{n} \sum_{i=0}^{n-1} \phi(f^i(x)) \le \mathbb{E}(\phi \mid \mathcal{I}) + \varepsilon$$

and the result follows by letting $\varepsilon \to 0$. □

1.3 Extensions

We can relate different dynamical systems by the concept of *factors*, or *extensions*, expressing a sense in which one system is a simplification of another. In the categorical language we have been developing, an extension is a natural transformation $F \Rightarrow G$ between two dynamical systems F and G. In this situation, we then say that G is a *factor* of F. The single morphism $F(\star) \to G(\star)$ specified by the natural transformation is also called the *factor map*.

Definition 1.5. Let $F, G\colon M \to \mathcal{C}$ be two dynamical systems on the category $\mathcal{C}$. An extension $\eta\colon F \Rightarrow G$ consists of a morphism $\eta\colon F(\star) \to G(\star)$ of $\mathcal{C}$ (the factor map), such that for all $m \in M$, the diagram

$$\begin{tikzcd} F(\star) \arrow[r, "F(m)"] \arrow[d, "\eta"] & F(\star) \arrow[d, "\eta"] \\ G(\star) \arrow[r, "G(m)"] & G(\star) \end{tikzcd}$$

commutes.

Now, we conclude our tour around the zoo and focus on our beloved species, measure-preserving systems.

Definition 1.6. Let $(X, \mathcal{B}_X, \mu, f)$ and $(Y, \mathcal{B}_Y, \nu, g)$ be measure-preserving transformations. $(Y, \mathcal{B}_Y, \nu, g)$ is a **factor** of $(X, \mathcal{B}_X, \mu, f)$ if there exists a measure-preserving map $\phi : X \to Y$ such that

$$\begin{tikzcd} X \arrow[r, "f"] \arrow[d, "\phi"] & X \arrow[d, "\phi"] \\ Y \arrow[r, "g"] & Y \end{tikzcd}$$

commutes.

Example 1.7. Every system admits a trivial factor:

$$\begin{tikzcd} X \arrow[r, "f"] \arrow[d, "\phi"] & X \arrow[d, "\phi"] \\ \{\star\} \arrow[r, "\mathrm{id}"] & \{\star\} \end{tikzcd}$$

"For Borel standard systems, extensions are classified by invariant sub-σ-algebras"

Lemma 1.8. *Let $(X, \mathcal{B}_X, \mu, f)$ and $(Y, \mathcal{B}_Y, \nu, g)$ be invertible systems, and let $\phi : X \to Y$ be a factor map. Then $\mathcal{A} = \phi^{-1}(\mathcal{B}_Y) \subset \mathcal{B}_X$ is an invariant sub-σ-algebra.*

Theorem 1.9. *Let $(X, \mathcal{B}, \mu, f)$ be a measure-preserving Borel probability system, and let $\mathcal{A} \subseteq \mathcal{B}$ be an f-invariant sub-σ-algebra. Then, there is a measure-preserving system $(Y, \mathcal{B}_Y, \nu, g)$ on a Borel probability space and a factor map $\phi : X \to Y$ with $\mathcal{A} = \phi^{-1}(\mathcal{B}_Y) \pmod{\mu}$.*

Proof.

- $Y = \mathcal{M}(X)$

- $S : Y \longrightarrow S$ defined by $S(\eta) = T_*\eta$
- $\phi : X \longrightarrow Y$ defined by $\phi(x) = \mu_x^{\mathcal{A}}$
- $\nu = \phi_*\mu$

□

Corollary 1.10. *A system $(X, \mathcal{B}, \mu, f)$ is ergodic if, and only if, its only factors are the trivial factor and the identity factor, up to isomorphism.*

Lemma 1.11. *Suppose $p\colon (X, f) \longrightarrow (Y, g)$ is a factor map. Then, if $\mu \in \mathcal{M}_f(X)$ is ergodic for f then $p_*\mu \in \mathcal{M}_g(Y)$ is ergodic for g.*

Proof. Suppose $A \subseteq Y$ is measurable and invariant for g. Then, $p^{-1}(A)$ is an invariant set for f, because

$$f^{-1}(p^{-1}(A)) = p^{-1}(g^{-1}(A)) = p^{-1}(A).$$

Therefore, $p_*\mu(A) = \mu(p^{-1}(A)) = 0$. □

Invertible extensions

$$\begin{tikzcd}
(\widetilde{X}, \tilde{f}) \arrow[dd, "\pi"'] & \\
& (Y, g) \arrow[ul, dashed, "\exists! h"'] \arrow[dl, "p"] \\
(X, f) &
\end{tikzcd} \tag{1.3}$$

Theorem 1.12. *Let $(X, \mathcal{B}, \mu, f)$ be a measure-preserving system on a Borel space. Then, the natural invertible extension exists.*

Proof. Apply Kolmogorov extension theorem.

Let $\widetilde{X} \subseteq X^{\mathbb{Z}}$ be the subset defined by

$$\widetilde{X} = \bigcap_{i \in \mathbb{Z}} \{x \in X^{\mathbb{Z}} \mid x_{i+1} = f(x_i)\},$$

which is closed in the product topology by definition.

, with $\pi\colon \widetilde{X} \longrightarrow X$ being the projection on the zeroth coordinate, that is, $\pi(\dots, x_{-1}, x_0, x_1, \dots) = x_0$.

First, define $\tilde{f}$ to be □

Lemma 1.13. *Let $\pi\colon (\widetilde{X}, \tilde{f}, \tilde{\mu}) \longrightarrow (X, f, \mu)$ be a natural invertible extension. Then, we have a converse for lemma 1.11: if μ is ergodic for f, then $\tilde{\mu}$ is ergodic for $\tilde{f}$.*

A failure of existence

[3]

Let $X = \mathbb{S}^1$ be the circle with the Borel σ-algebra $\mathcal{B}$ and $f\colon \mathbb{S}^1 \to \mathbb{S}^1$ be an irrational rotation. With the Axiom of Choice, we can form a set E with exactly one point from each orbit of f. Since f is a Lebesgue measure-preserving bijection, $E_n = f^n(E)$ are disjoint, have the same outer measure and $\bigcup_{n\in\mathbb{Z}} E_n = \mathbb{S}^1$. In particular, the E_n cannot be measurable, thus $m^*(E) > 0$.

If K is a subset of positive outer measure and $\mathcal{E} = \sigma(\{K\} \cup \mathcal{B})$, and if we define $\mu^*\colon 2^X \to [0,1]$ as

$$\mu^*(A) = \frac{m^*(A \cap K)}{m^*(K)},$$

then μ^* is an probability outer measure that restricts itself to a probability measure μ on $\mathcal{E}$. It is also the case that $\mathcal{E} = \mathcal{B} \cup \{A \cap K : A \in \mathcal{B}\} \cup \{A \cap K^c : A \in \mathcal{B}\}$. Let C_K be the completion by μ^* of the σ-algebra $\mathcal{E}$.

Let's apply the above construction for $K = \bigcup_{n=1}^{\infty} f^n(E)$. Note that $E \cap K = \varnothing$, therefore $\mu^*(E) = 0$ and $E \in C_K$. We conclude that $f^{-1}(K) = E \cup K$ is measurable as well, so for every $A \in \mathcal{B}$, $f^{-1}(A \cap K) = f^{-1}(A) \cap f^{-1}(K)$ is measurable, thus $f^{-1}(\mathcal{E}) \subseteq C_K$.

Now, for any $A \in C_K$, we can write $A = B \cup Z$ with $B \in \mathcal{E}$ and $\mu^*(Z) = 0$; thus, $f^{-1}(A) = f^{-1}(B) \cup f^{-1}(Z)$. Note that

$$\mu^*(f^{-1}(Z)) = m^*(Z \cap f(K)) \le m^*(Z \cap K) = \mu^*(Z) = 0,$$

therefore $f^{-1}(C_K) \subset C_K$.

The same calculation shows that $\mu^*(f^{-1}(A)) \le \mu^*(A)$ for any $A \subseteq X$. Thus, if A is measurable,

$$1 - \mu(f^{-1}(A)) = \mu(f^{-1}(A^c)) \le \mu(A^c) = 1 - \mu(A),$$

so $f_*\mu = \mu$ on C_K.

Consider $\Sigma = \{(x_i)_{i\le 0} : f(x_i) = x_{i+1}\} \subseteq X^{\mathbb{Z}_{\le 0}}$ as the inverse limit of (X, f, C_K), endowed with the restriction of the product σ-algebra. Note that $\tilde{f}\colon \Sigma \to \Sigma$ given by

$$\tilde{f}((x_i)_{i\le 0}) = (f(x_i))_{i\le 0} = (\ldots, x_{-1}, x_0, f(x_0))$$

is invertible and measurable, since

$$\tilde{f}([A_{-k}, \ldots, A_0] \cap \Sigma) = [A_{-k}, \ldots, A_0, X] \cap \Sigma.$$

For each $n \in \mathbb{Z}_{\le 0}$, let $K_n = [n \mid K] = \{(x_i)_{i\le 0} : x_n \in K\}$. Suppose there exists $\tilde{\mu}$, a measure on Σ such that $\pi_*\tilde{\mu} = \mu$, where $\pi((x_i)_{i\le 0}) = x_0$. Then

$$\tilde{\mu}(K_n) = \tilde{\mu}(\tilde{f}^n(\pi^{-1}(K))) = \mu(K) = 1$$

for every n. On the other hand, since f is invertible,

$$\begin{aligned}\bigcap_{n\le 0} K_n &= \{(x_i)_{i\le 0} : \forall i, x_i \in K\} \\ &= \{(x_i)_{i\le 0} : \forall n \ge 0, f^{-n}(x_0) \in K\}.\end{aligned}$$

But this set is empty because

$$\{(x_i)_{i\leq 0} : \forall n \geq 0, f^{-n}(x_0) \in K\} = \bigcap_{n=0}^{\infty} f^n(K) = \limsup_n E_n = \varnothing.$$

The solenoid

[4] [5]

1.4 Cocycles and skew products

$$\begin{tikzcd} X \times Y \arrow[r] \arrow[d, "\pi_X"] & X \times Y \arrow[d, "\pi_X"] \\ X \arrow[r, "f"] & X \end{tikzcd}$$

Example 1.14 (Iterated function system)**.** IFSs are examples of skew products.

Lyapunov exponents

Define for linear cocycles. Oseledets theorem.

1.5 Empirical measures

Improve this, this is the main concept of the dissertation, so it deserves more love.

Given a map $f\colon X \to X$ on a measurable space and a point $x \in X$, we can define a sequence of *empirical measures*

$$e_n(x) = \frac{1}{n}\sum_{i=0}^{n-1} \delta_{f^i(x)},$$

encoding the averages along the orbit up to a time n. These measures are, indeed, uniquely defined by the property that for any observable $\phi\colon X \to \mathbb{R}$, its integral with respect to $e_n(x)$ equals the Birkhoff average up to time n:

$$\int \phi \; d(e_n(x)) = \frac{1}{n}\sum_{i=0}^{n-1} \phi(f^i(x)),$$

and it is this quality of being defined by the values observed on a finite number of steps along an orbit that should justify the nomenclature empirical. The direct relation to Birkhoff averages also highlights these objects as having a central role in ergodic theory.

It's not always the case that this sequence converges (weakly) to some measure, but when it does, the limit is invariant and represents the asymptotic distribution of the orbit. Reciprocally, For each invariant measure $\mu \in \mathcal{M}_f(X)$, we can ask whether the empirical measures of a point will converge weakly to it. The set $\mathcal{B}(\mu) \subseteq X$ of points with this property is called the *basin* of the measure μ.

Describe the overall problem of generically finding finite set of limiting measures, whose basins cover almost all points of the manifold. Describe the robust counterexample to finiteness by Newhouse. Describe that the problem of historical behavior is open.

It may happen that the set of points whose empirical measures do not converge is large, in the sense that it has positive volume. Systems with this property are said to exhibit *historical behavior*, and the existence of diffeomorphisms exhibiting historical behavior in a robust way is an important open problem in dynamics.

Proposition 1.15. *Let X be a compact metric space with the Borel σ-algebra. For every $x \in X$, the limit points in the weak* topology of the sequence of empirical measures*

$$\mathcal{M}(x) = \{e_n(x)\}'_{n\geq 0} = \bigcap_{i=0}^{\infty} \overline{(\{e_n(x)\}_{n\geq i})}$$

form a compact connected subset of the invariant measures.

Proof. The set is closed by definition, so by Banach-Alaoglu it's compact; for connectedness, suppose $\mathcal{M}(x) = K_1 \cup K_2$ where K_1, K_2 are closed and disjoint. Let $\{\phi_i\}_{i\in\mathbb{N}} \subseteq B_{C(X)}$ be a dense subset and define

$$d(\mu, \nu) = \sum_{i=1}^{\infty} \frac{|\mu(\phi_i) - \nu(\phi_i)|}{2^i},$$

which is a metric inducing the weak* topology on the set of probability measures (see A.3). Since it is a metric, there are two open sets $U_1 \supseteq K_1, U_2 \supseteq K_2$ with positive distance $s > 0$ from each other. We claim $\lim_{n\to\infty} d(e_n(x), e_{n+1}(x)) = 0$, which would contradict $s > 0$ because the definition of $\mathcal{M}(x)$ implies $e_n(x) \in U_1$ and $e_{n+1}(x) \in U_2$ for infinitely many n. Indeed,

$$\begin{aligned} d(e_n(x), e_{n+1}(x)) &= \sum_{i=1}^{\infty} \frac{|\frac{1}{n}\sum_{i=0}^{n-1} \phi_i \circ f^i(x) - \frac{1}{n+1}\sum_{i=0}^{n} \phi_i \circ f^i(x)|}{2^i} \\ &\leq \sum_{i=1}^{\infty} \frac{2}{n}\frac{\|\phi_i\|_\infty}{2^i} \leq \frac{2}{n} \to 0. \end{aligned}$$

□

Lemma 1.16. *If μ is invariant for f, then $\mu(\mathcal{B}(\mu)) \in \{0, 1\}$. Moreover, $\mu(\mathcal{B}(\mu)) = 1$ if, and only if, μ is ergodic for f.*

1.6 Partitions

A *partition* of a set S is a collection of disjoint subsets $\mathcal{P} \subseteq 2^S$ whose union is S. It's important to notice that $\mathcal{P}$ can be seen either as a set of subsets or (with some abuse of notation) as a function $\mathcal{P}\colon S \to 2^S$ assigning to each point $p \in S$ the unique $\mathcal{P}(p) \in \mathcal{P}$ such that $p \in \mathcal{P}(p)$. To make matters clearer, henceforward we shall denote by $\overline{\mathcal{P}} \subseteq 2^S$ the image of $\mathcal{P}\colon S \to 2^S$ whenever we need to emphasize that the set of subsets is the object of our attentions.

Lattice structure

The partitions $\mathbf{P}(S)$ of a set S form a lattice under the refinement order. Namely, we endow $\mathbf{P}(S)$ with the partial order where $\mathcal{P} \preceq \mathcal{Q}$ if and only if for all $Q \in \overline{\mathcal{Q}}$, there is some $P \in \overline{\mathcal{P}}$ such that $Q \subseteq P$, and whenever this happens we say that $\mathcal{Q}$ is *finer* and $\mathcal{P}$ is *coarser* relative to each other. Given two partitions $\mathcal{P}, \mathcal{Q} \in \mathbf{P}(S)$, there is a *join* operation defined by

$$(\mathcal{P} \vee \mathcal{Q})(x) = \mathcal{P}(x) \cap \mathcal{Q}(x)$$

that is the least upper bound of $\mathcal{P}$ and $\mathcal{Q}$ (the coarsest partition finer than both $\mathcal{P}$ and $\mathcal{Q}$). There is also a *meet* operation $\mathcal{P} \wedge \mathcal{Q}$, giving the greatest lower bound, which is less straightforward to construct: on the union $\overline{\mathcal{P}} \cup \overline{\mathcal{Q}}$, we say P and Q are related if they overlap, that is, $P \cap Q \neq \varnothing$, and by taking the transitive closure of this relation we get an equivalence relation on the set $\overline{\mathcal{P}} \cup \overline{\mathcal{Q}}$. The elements of $\mathcal{P} \wedge \mathcal{Q}$, then, are precisely the unions of sets in the same equivalence class:

$$\mathcal{P} \wedge \mathcal{Q} = \left\{ \bigcup_{S \in [P]} S \,\middle|\, [P] \in \overline{\mathcal{P}} \cup \overline{\mathcal{Q}}/\sim \right\}.$$

The partitions $\mathbf{P}(S)$ form, in fact, a *complete* lattice, meaning that all subsets $\mathsf{P} \subseteq \mathbf{P}(S)$ have both a least upper bound $\bigvee_{\mathcal{P} \in \mathsf{P}} \mathcal{P} \in \mathbf{P}(S)$ and a greatest lower bound $\bigwedge_{\mathcal{P} \in \mathsf{P}} \mathcal{P} \in \mathbf{P}(S)$. Their construction is very similar to their binary versions: the join can be defined as $\left(\bigvee_{\mathcal{P} \in \mathsf{P}} \mathcal{P}\right)(x) = \bigcap_{\mathcal{P} \in \mathsf{P}} \mathcal{P}(x)$, and we can use the join to define the meet as

$$\bigwedge_{\mathcal{P} \in \mathsf{P}} \mathcal{P} = \bigvee_{\substack{\mathcal{Q} \text{ is an upper} \\ \text{bound for } \mathsf{P}}} \mathcal{Q}.$$

In the case of $\mathsf{P} = \{\mathcal{P}, \mathcal{Q}\}$, they reduce to the binary operations of the paragraph above.

An interlude of zero measure

In a measure space, it is often useful to relax the notions of equality and order on the partitions so that zero measure sets are disregarded. This is done in the obvious way, namely by saying $\mathcal{P} \preceq \mathcal{Q}$ if for all $Q \in \overline{\mathcal{Q}}$, there is a $P \in \overline{\mathcal{P}}$ such that $Q \setminus P$ has zero measure. Going a bit further, the partitions themselves may be allowed to cover only a total measure subset of the space, and their elements could overlap on zero measure sets and still be regarded as disjoint. Formally, the idea here is to quotient the subsets by the almost-surely equality relation, and notice that the usual binary operators on the sets lift to well-defined operators on the quotient.

1.7 Disintegrations

In the Calculus course, we learn that it's most often easier to find the area of a region in the plane if we reduce the problem back to a single dimension, by taking integrals of the length along parallel lines. This is the essence of Fubini's theorem, which states that the integral of a function in a higher dimensional region can be written as nested integrals of the function along one-dimensional subsets. Then we go to the probability course, and the lecturer talks about

conditional expectations and the whole philosophy of Bayes of taking probabilities conditioned to some event. It may not seem at first that this has anything to do with the aforementioned theorem in the Calculus course, at least not until we take the Ergodic theory course and learn about measures and disintegrations.

While Fubini remains true for product measures, in general the problem of decomposing an arbitrary measure into a family of measures defined in smaller subsets is not as straightforward. Neither is straightforward to see how could one talk about the probability of an event conditioned to another event of *zero probability*, even though in practice this happens very often. Both situations are instances of the general problem the disintegration of measures solves: given a measure μ and a partition $\mathcal{P}$ of the space X, we want to find a family of measures $\{\mu_x\}_{x\in X}$, each μ_x giving full measure to $\mathcal{P}(x)$, with the property that $\mu(E) = \int \mu_x(E)\, d\mu(x)$.

Definition 1.17 (Disintegration)**.** A *disintegration* of a measure $\mu \in \mathcal{M}(X)$ with respect to a partition $\mathcal{P}$ is a family $\{\mu_x\}_{x\in X} \subseteq \mathcal{M}(X)$ such that for every measurable set $E \subseteq X$,

1. $\mu_x(\mathcal{P}(x)) = 1$ for μ-almost every x;
2. the map $x \mapsto \mu_x(E)$ is measurable;
3. $\mu(E) = \int \mu_x(E)\, d\mu(x)$.

Uniqueness

Measurability

Not all partitions are created equal. Some may be pathological enough that there may be no hope in disintegrating over them. In fact, such partitions appear *very commonly* in dynamics. A prime example is the partition consisting of orbits of a map without too many periodic points.

Example 1.18. Suppose $f\colon X \to X$ is invertible and μ is an invariant measure for f such that $\mu(X \setminus \mathrm{Per}(f)) > 0$. Then, the partition $\mathcal{O}_f\colon X \to 2^X$ of two-sided orbits of f is such that μ does not admit a disintegration with respect to $\mathcal{O}_f$.

Proof. Suppose that such disintegration $\{\mu_x\}_{x\in X}$ exists. We claim that $\{f_*\mu_x\}_{x\in X}$ is also a disintegration. Indeed, all $\mathcal{O}_f(x)$ are invariant, so we have

$$(f_*\mu_x)(\mathcal{O}_f(x)) = \mu_x(f^{-1}(\mathcal{O}_f(x))) = \mu_x(\mathcal{O}_f(x)) = 1$$

for a.e. x, and similarly, we have that $x \mapsto (f_*\mu_x)(E) = \mu_x(f^{-1}(E))$ is measurable, and $\mu(E) = \mu(f^{-1}(E)) = \int f_*\mu_x(E)\, d\mu(x)$. By uniqueness, this implies that $f_*\mu_x = \mu_x$ almost everywhere.

By item 1, the hypothesis and the paragraph above, we can find some $x \notin \mathrm{Per}(f)$ such that $f_*\mu_x = \mu_x$ and $\mu_x(\mathcal{O}_f(x)) = 1$. But this is impossible: since $\mathcal{O}_f(x)$ is infinite and $\mu_x(\{f^k(x)\}) = \mu_x(\{x\})$ for every k,

$$1 = \mu_x(\mathcal{O}_f(x)) = \sum_{k\in\mathbb{Z}} \mu_x(\{f^k(x)\}) = \sum_{k\in\mathbb{Z}} \mu_x(\{x\}) \in \{0, \infty\}.$$

□

Remark 1. A very basic instance of example 1.18 would be with $X = S^1$, $f\colon S^1 \to S^1$ an irrational rotation, and μ the unique ergodic measure, because f has no periodic points.

Remark 2. The example 1.18 also shows that it is not sufficient for the *elements* of the partition to be measurable sets. In the remark above, the orbits are countable sets of points in the circle, and thus are Lebesgue (or even Borel) measurable, and yet the partition does not admit a disintegration.

The right notion of a "well-behaved" partition with respect to disintegrations will turn out to be the notion of *measurable partitions*.

Definition 1.19. A partition $\mathcal{P}$ is said to be *measurable* if there is a sequence $\mathcal{P}_1, \mathcal{P}_2, \dots$ of partitions, all of which composed of a countable number of measurable sets, such that $\mathcal{P} = \bigvee_{n\in\mathbb{N}} \mathcal{P}_n$.

Example 1.20. The partition $\perp$ of all points of a second-countable T_1 topological space X is measurable in the Borel σ-algebra.

Proof. Let $\{U_n\}_{n\in\mathbb{N}}$ be a countable basis for the topology of X, and define $\mathcal{P}_n = \{U_n, U_n^c\}$. Let $P \in \mathcal{P} = \bigvee_{n\in\mathbb{N}} \mathcal{P}_n$, which is measurable by definition, and $p \in P$. If $q \in P$, then we must have $\mathcal{P}_n(p) = \mathcal{P}_n(q)$ for all $n \in \mathbb{N}$, or in other words, $p \in U_n \iff q \in U_n$, for all n. The T_1 property then implies $p = q$, so $P = \{p\}$. This shows $\mathcal{P} = \perp$. □

Example 1.21. Any partition $\mathcal{P}$ of a compact metric space X whose elements are closed is measurable in the Borel σ-algebra.

Proof. Since they are closed, the elements of $\mathcal{P}$ are measurable in the Borel algebra. To show the partition is countably generated, we can define a metric on $\overline{\mathcal{P}}$ by letting $d(P_1, P_2)$ be the distance between the compact sets $P_1, P_2 \in \overline{\mathcal{P}}$ (this is the Hausdorff metric, which we detail in **??**). With this metric, the surjection $\mathcal{P}\colon X \to \overline{\mathcal{P}}$ is a contraction, so $\overline{\mathcal{P}}$ is a compact metric space as well. By the example 1.20, the partition $\perp$ of points of $\overline{\mathcal{P}}$ is countably generated, thus so is its preimage by $\mathcal{P}$. □

Conditional measures

1.8 Entropy

Topological entropy

Define.

Measure-theoretic entropy

Define.

Pesin entropy formula

State the theorem, to be used in the invariance principle section.

Chapter 2

Schwarzian derivatives and cylinder maps

In this chapter, we follow the exposition by Bonifant and Milnor [6] on a family of endomorphisms of a cylinder.

2.1 Cross ratios and the Schwarzian derivative

Suppose we start with a field $\mathbb{K}$ and, not being happy with the fact that we cannot divide by zero, decide to add a point at infinity; this gives us the *projective line* $P^1(\mathbb{K}) = \mathbb{K} \sqcup \{\infty\}$, whose arithmetic operations are the same as those of $\mathbb{K}$ except for the additional rules $\frac{1}{0} = \infty$, $\frac{1}{\infty} = 0$, $x \cdot \infty = \infty$ if $x \neq 0$, and $x + \infty = \infty$ if $x \neq \infty$. It's not possible to define such operations everywhere without introducing inconsistent rules, so the operations are partial — we cannot multiply $\infty \cdot 0$ or add $\infty + \infty$.

Our new space $P^1(\mathbb{K})$ has an interesting property: given $z \in P^1(\mathbb{K})$, we have three types of operations that we can apply to z: translations $z \mapsto z + x$, where $x \in \mathbb{K}$, homotheties $z \mapsto xz$ for $x \in \mathbb{K} \setminus \{0\}$, and the inverse $z \mapsto 1/z$, which was not defined before if $z = 0$, but now makes sense for every z. These three families of actions, which are invertible, generate the *group of homographies* with coefficients in $\mathbb{K}$.

A second way to construct $P^1(\mathbb{K})$, which makes the role of the group of homographies as a symmetry group of this space even clearer, is to think of the space of "formal fractions", that is, pairs of numbers (x, y), where $x, y \in \mathbb{K}$ are not both zero, where we identify "equivalent fractions", in the sense that $(\lambda x, \lambda y) \sim (x, y)$ for every $\lambda \in \mathbb{K} \setminus \{0\}$. The quotient by this equivalence relation gives us $P^1(\mathbb{K})$ as the space of lines passing through the origin of the vector space $\mathbb{K}^2$, and the operations are like fraction operations: $(x, y) + (z, w) = (wx + yz, yw)$, $(x, y) \cdot (z, w) = (xz, yw)$, and $(x, y)^{-1} = (y, x)$. Despite being seemingly more complicated, the interesting fact about this construction is that it emphasizes how the symmetry group of the space $\mathbb{K}^2$, the linear group $\mathrm{GL}_2(\mathbb{K})$, naturally induces a symmetry group on the quotient, the projective group $\mathrm{PGL}_2(\mathbb{K})$, which is precisely the group of homographies.

This group is a quotient of $\mathrm{GL}_2(\mathbb{K})$ by a subgroup of dimension 1, so it has dimension 3. Furthermore, it is 3-transitive: given two sets of three distinct points in $P^1(\mathbb{K})$, there exists an element $g \in \mathrm{PGL}_2(\mathbb{K})$ that takes the first set to the second. But not only that, $P^1(\mathbb{K})$ has three special points: 0, 1, and ∞. Thus, given $x_1, x_2, x_3 \in P^1(\mathbb{K})$ distinct, we can ask which $g \in \mathrm{PGL}_2(\mathbb{K})$ such that $g(x_1) = \infty$, $g(x_2) = 0$, and $g(x_3) = 1$. With some calculations, we arrive at

$$g(z) = \frac{(x_2 - z)(x_3 - x_1)}{(x_1 - z)(x_3 - x_2)}$$

The *cross ratio* of four points x_0, x_1, x_2, x_3 is the value of $g(x_0)$ for the element g of the homography group that sends $x_1 \mapsto \infty$, $x_2 \mapsto 0$, and $x_3 \mapsto 1$.

Definition 2.1 (Cross ratio)**.**

$$\rho(x_0, x_1, x_2, x_3) = \frac{(x_2 - x_0)(x_3 - x_1)}{(x_1 - x_0)(x_3 - x_2)}$$

Definition 2.2. Let $f\colon I \to I'$ be a C^3 diffeomorphism between two intervals of the real line. The *Schwarzian derivative* of f is the function

$$\mathcal{S}f = \frac{f'''}{f'} - \frac{3}{2}\left(\frac{f''}{f'}\right)^2.$$

We shall write $\mathcal{S}f \prec 0$ if there exists a dense open set $U \subseteq I$ such that $\mathcal{S}f(y) < 0$ for every $y \in U$. Similarly, we write $\mathcal{S}f \succ 0$ if there exists $U \subseteq I$ open and dense such that $\mathcal{S}f(y) > 0$ for every $y \in U$.

The Schwarzian derivative has some important properties that we highlight in the Proposition below, as described in [6, Proposition B.1].

Proposition 2.3. *The Schwarzian derivative has the following properties:*

1. *The Schwarzian is a second-order derivation [7]*
   If $\mathcal{S}f \succ 0$ and $\mathcal{S}g \succ 0$, then $\mathcal{S}(f \circ g) \succ 0$, and similarly for $\prec 0$. Moreover, if $\mathcal{S}f \equiv 0$, then $\mathcal{S}(f \circ g) = \mathcal{S}g$, and if $\mathcal{S}g \equiv 0$, then $\mathcal{S}(f \circ g) = (\mathcal{S}f) \circ g$.
2. *If $\mathcal{S}f \prec 0$, then $\mathcal{S}f^{-1} \succ 0$.*
3. *$\mathcal{S}f \prec 0$ if, and only if, $\varphi(y) = 1/\sqrt{|f'(y)|}$ is strictly convex.*
4. *$\mathcal{S}f \equiv 0$ if, and only if, f is a fractional linear transformation.*

Proof.

1. We have

$$\mathcal{S}(f \circ g) = \frac{(f \circ g)'''}{(f \circ g)'} - \frac{3}{2}\left(\frac{(f \circ g)''}{(f \circ g)'}\right)^2.$$

By the Faà di Bruno's (beautiful) formula [8, p. 479] in its combinatorial form,

$$(f \circ g)^{(n)}(x) = \sum_{\pi \in \Pi} f^{(|\pi|)}(g(x)) \cdot \prod_{B \in \pi} g^{(|B|)}(x)$$

where the sum is taken over the set Π of all partitions of $\{1, \dots, n\}$, and $B \in \pi$ means B is an element of the partition π (therefore, a subset of $\{1, \dots, n\}$).
Specializing Faà di Bruno's formula to $n = 1, 2, 3$, we have

$$\begin{aligned}(f \circ g)' &= (f' \circ g) \cdot g' \\ (f \circ g)'' &= (f'' \circ g) \cdot (g')^2 + (f' \circ g) \cdot g'' \\ (f \circ g)''' &= (f''' \circ g) \cdot (g')^3 + 3(f'' \circ g) \cdot g'' \cdot g' + (f' \circ g) \cdot g'''\end{aligned}$$

and thus

$$\begin{aligned}\mathcal{S}(f \circ g) &= \frac{(f''' \circ g) \cdot (g')^3 + 3(f'' \circ g) \cdot g'' \cdot g' + (f' \circ g) \cdot g'''}{(f' \circ g) \cdot g'} \\ &\qquad - \frac{3}{2}\left(\frac{(f'' \circ g) \cdot (g')^2 + (f' \circ g) \cdot g''}{(f' \circ g) \cdot g'}\right)^2 \\ &= \frac{(f''' \circ g) \cdot (g')^2}{f' \circ g} + 3\frac{(f'' \circ g) \cdot g''}{f' \circ g} + \frac{g'''}{g'} - \frac{3}{2}\left(\frac{(f'' \circ g) \cdot g'}{f' \circ g} + \frac{g''}{g'}\right)^2 \\ &= \left(\frac{f''' \circ g}{f' \circ g} - \frac{3}{2}\left(\frac{f'' \circ g}{f' \circ g}\right)^2\right) \cdot (g')^2 + \frac{g'''}{g'} - \frac{3}{2}\left(\frac{g''}{g'}\right)^2 \\ &= (\mathcal{S}f) \circ g \cdot (g')^2 + \mathcal{S}g. \qquad (2.1)\end{aligned}$$

Since g is a diffeomorphism, $g'(x) \neq 0$ for all $x \in I$, and therefore $(g')^2 > 0$. In particular, g is a homeomorphism, so that if $\mathcal{S}f(x) > 0$ for all x the open dense set U, and $\mathcal{S}g(x) > 0$ in the open dense set V, then $\mathcal{S}(f \circ g) > 0$ in the open dense set $g^{-1}(U) \cap V$. Similarly, from eq. (2.1) we see that if $\mathcal{S}f \equiv 0$, then the first term is zero, therefore $\mathcal{S}(f \circ g) = \mathcal{S}g$.

2. Applying eq. (2.1) to $\mathrm{id} = f^{-1} \circ f$, we see that

$$0 = \mathcal{S}(\mathrm{id}) = (\mathcal{S}f^{-1}) \circ f \cdot (f')^2 + \mathcal{S}f,$$

and by the same argument of the paragraph above, we conclude $\mathcal{S}f \prec 0 \iff \mathcal{S}f^{-1} \succ 0$.

3. Since f is a diffeomorphism between intervals, either $f' > 0$ or $f' < 0$. As a first step, assume $f' > 0$. Taking the second derivative of $\varphi(y) = 1/\sqrt{f'(y)} = f'(y)^{-1/2}$,

$$\varphi'' = \left(-\frac{1}{2}(f')^{-3/2} \cdot f''\right)' = \frac{3}{4}(f')^{-5/2} \cdot (f'')^2 - \frac{1}{2}(f')^{-3/2} \cdot f'''.$$

Thus

$$2\varphi''/\varphi = \frac{3}{2}(f')^{-4/2} \cdot (f'')^2 - (f')^{-2/2} \cdot f''' = \frac{3}{2}\left(\frac{f''}{f'}\right)^2 - \frac{f'''}{f'} = -\mathcal{S}f. \qquad (2.2)$$

From this, and noticing that $\varphi > 0$, we see that if $\mathcal{S}f(x) < 0$ if and only if $\varphi''(x) > 0$. A C^2 function $\varphi : I \to \mathbb{R}$ will be strictly convex if and only if $\varphi'' \geq 0$ and φ'' does not vanish on any open interval, so $\mathcal{S}f \prec 0$ implies that ϕ'' is positive in an open dense set, thus strictly convex. Reciprocally, if φ is strictly convex, then the set $\{x \mid \varphi''(x) > 0\} = \{x \mid \mathcal{S}f(x) < 0\}$ is open and its complement does not contain any open interval, and therefore is dense.
If $f' < 0$, we define $g \colon I \to -I'$, $g(y) = -f(y)$, so that $g' > 0$ and we fall back to the previous case. The argument is finished by noticing that $\mathcal{S}g = \mathcal{S}f$ and $1/\sqrt{|f'|} = 1/\sqrt{|g'|}$.

4. From eq. (2.2), we see that $\mathcal{S}f \equiv 0 \iff \varphi'' \equiv 0$. This, in turn, is equivalent to φ being linear, say $\varphi(y) = cy + d$. We thus have
$$f'(y) = \frac{1}{(cy+d)^2}.$$
Integrating both sides, by the Fundamental Theorem of Calculus we have
$$f(y) - f(a) = \int_a^y \frac{1}{(ct+d)^2}\,dt = \frac{1}{c}\left(\frac{1}{ca+d} - \frac{1}{cy+d}\right) = \frac{y-a}{c(ca+d)y + d(ca+d)}$$
which is a fractional linear transformation, as desired.

□

Lemma 2.4. *If $f\colon I \to I'$ is C^3 and $\mathcal{S}f \prec 0$, then f has at most three fixed points. When it has three fixed points, the middle one is strictly repelling and the other two are strictly attracting.*

Proof. Suppose f had four fixed points $y_1 < y_2 < y_3 < y_4$. By the Mean Value Theorem, in each of the three open intervals (y_i, y_{i+1}), for $i = 1, 2, 3$, we would have a point $c_i \in (y_i, y_{i+1})$ where $f'(c_i) = 1$. But this is impossible, since by item 3 $\varphi(y) = 1/\sqrt{|f'(y)|}$ is strictly convex, which would imply $\varphi(c_2) < \varphi(c_1) = \varphi(c_3) = 1$.

Now, suppose there are exactly three fixed points $y_1 < y_2 < y_3$. By the same reasoning, there are two points $c_1 \in (y_1, y_2)$ and $c_2 \in (y_2, y_3)$ where $\varphi(c_1) = \varphi(c_2) = 1$. Knowing that φ is strictly convex, this implies $\varphi(y_2) > 1$ and $\varphi(y_1), \varphi(y_3) < 1$, and therefore $|f'(y_2)| < 1$ and $|f'(y_1)|, |f'(y_3)| > 1$. □

Lemma 2.5. *If $f\colon I \to I'$ is C^3, then $\mathcal{S}f \prec 0$ if, and only if, f increases cross ratios, in the sense that $\rho(f(y_1), f(y_2), f(y_3), f(y_4)) > \rho(y_1, y_2, y_3, y_4)$ for all $y_1 < y_2 < y_3 < y_4$.*

Proof. Given $y_1 < y_2 < y_3 < y_4$, let φ be the linear fractional transformation that sends $f(y_1)$ to y_1, $f(y_2)$ to y_2 and $f(y_4)$ to y_4. Since from proposition 2.3, we have that $\mathcal{S}(\varphi \circ f) = \mathcal{S}f$, and we know fractional linear transformations also preserve cross-ratios, without loss of generality we may assume that f itself fixes y_1, y_2 and y_4. Now, by lemma 2.4, $f(y) > y$ on the interval (y_2, y_4), and therefore $f(y_3) > y_3$. It thus follows easily by the definition of cross-ratios that $\rho(f(y_1), f(y_2), f(y_3), f(y_4)) > \rho(y_1, y_2, y_3, y_4)$. □

Lemma 2.6. *If $f\colon I \to I$ is a diffeomorphism that preserves orientation and $\mathcal{S}f \prec 0$, where $I = [a, b]$, then $f'(a)f'(b) < 1$.*

Proof. Let's consider the case where $f'(a) = f'(b)$ first. If we had $f'(a)^2 = f'(b)^2 = f'(a)f'(b) \geq 1$, then $\varphi(a) = \varphi(b) \leq 1$ which by strict convexity of φ implies $\varphi|_{(a,b)} < 1$. This, in turn, means that $f'|_{(a,b)} < 1$, which (by the mean value theorem) is not possible for a diffeomorphism that sends an interval to itself.

The general case is shown by noticing that if we define the function $g = r \circ f \circ r\colon I \to I$, where $r(x) = a + b - x$ (and thus g is the reflection of f along the anti-diagonal), then $g'(a) = f'(b)$ and $g'(b) = f'(a)$. By the chain rule,
$$(f \circ g)'(a) = (f \circ g)'(b) = f'(a)f'(b),$$
and by item 1, $\mathcal{S}(f \circ g) \prec 0$ because $\mathcal{S}(r) \equiv 0$ and thus $\mathcal{S}(g) = \mathcal{S}(f) \circ r \prec 0$. □

Note that by proposition 2.3, analogous results to the last two lemmas hold if $\mathcal{S}f > 0$. For example, if $\mathcal{S}f > 0$ then f decreases cross ratios and $f'(0)f'(1) > 1$.

2.2 Skew products

move things to the skew product section under extensions

We call a map $F\colon X \times Y \to X \times Y$ a *skew product* if it is of the form $F(x, y) = (g(x), f_x(y))$ for some $g\colon X \to X$ and a family of functions $\{f_x\}_{x\in X}$, $f_x\colon Y \to Y$. We are interested in the case where X is a compact metric space with the Borel σ-algebra, $g\colon X \to X$ is a continuous map preserving an ergodic probability measure $\mu \in \mathcal{M}(X)$, $Y = I$ is the interval $[0, 1]$, and $f_x\colon I \to I$ are C^3 diffeomorphisms preserving orientation. We will see how, depending on the Schwarzian derivative of the maps f_x, we observe different behaviors of the dynamics with respect to the invariant measures $\nu_0 = \mu \otimes \delta_0$ and $\nu_1 = \mu \otimes \delta_1$. Note that for $i \in \{0, 1\}$, $\operatorname{supp} \nu_i = \mathcal{A}_i$, where $\mathcal{A}_i = X \times \{i\}$ are the upper and lower "boundaries" of $X \times I$. Let $\mathcal{B}_i = \mathcal{B}(\nu_i)$ be the basins of the respective measures.

To simplify notation, let's denote by $f_x^n(y)$ the second coordinate of $F^n(x, y)$, that is, $f_x^n = f_{g^{n-1}(x)} \circ \cdots \circ f_x$.

Lemma 2.7. *If $(x, y_1) \in \mathcal{B}_0$, then $(x, y_2) \in \mathcal{B}_0$ for all $y_2 \le y_1$. Similarly, if $(x, y_1) \in \mathcal{B}_1$, then $(x, y_2) \in \mathcal{B}_0$ for all $y_2 \ge y_1$.*

Proof. We will prove the case $i = 0$, as the situation is symmetric. Let $(x, y_1) \in \mathcal{B}_0$ and $y_2 \le y_1$. If $\varphi \in C(X \times I)$, and $\varepsilon > 0$, by the uniform continuity of φ, there exists $\delta > 0$ such that $y < \delta$ implies $|\varphi(x, 0) - \varphi(x, y)| < \varepsilon$ for all $x \in X$. Since $U = X \times [0, \delta)$ is a continuity set for ν_0 and $e_n(x, y_2) \to \nu_0$, by Portmanteau

$$\lim_{n\to\infty} e_n(x, y_2)(U) = \nu_0(U) = 1.$$

Define $\tau_n = e_n(x, y_2)(U) = \frac{1}{n}\#\{0 \le i < n \mid f_x^n(y_2) < \delta\}$. Note that if $f_x^n(y_2) < \delta$, then $f_x^n(y_1) < \delta$ since the diffeomorphisms preserve orientation. Thus,

$$\begin{aligned}|e_n(x, y_1)(\varphi) - e_n(x, y_2)(\varphi)| &\le \frac{1}{n}\sum_{i=0}^{n-1}\left|\varphi(g^n(x), f_x^n(y_1)) - \varphi(g^n(x), f_x^n(y_2))\right| \\ &\le \varepsilon\tau_n + 2\|\varphi\|_\infty(1 - \tau_n).\end{aligned}$$

Since $\tau_n \to 1$, there exists N such that $n \ge N$ implies that the right side is less than 2ε, hence both integrals on the left converge to $\nu_0(\varphi)$. □

2.3 Transversal Lyapunov exponents

To each point (x, y) of either boundary $\mathcal{A}_i$, for $i \in \{0, 1\}$, we can associate a number that expresses the average expansion rate of tangent vectors along the direction normal to the skew product boundary. This is a specific case of the general definition of Lyapunov exponents, whose existence is given by the Oseledets theorem. make the link to section 1.4

Definition 2.8 (Transversal Lyapunov exponent)**.**

$$\mathrm{Lyap}_{\mathcal{A}_i}(x) = \lim_{n\to\infty} \frac{1}{n} \log\left|\partial_y F^n(x,i)\right|.$$

By the product rule, and noting that $\{0,1\}$ are fixed points of the fiber diffeomorphisms, the right side of the equation above can be rewritten as

$$\lim_{n\to\infty} \frac{1}{n} \sum_{k=0}^{n-1} \log\left|\partial_y F(g^k(x),i)\right| = \lim_{n\to\infty} \frac{1}{n} \sum_{k=0}^{n-1} \log f'_{g^k(x)}(i).$$

Thus, by the ergodicity of μ for g and the Birkhoff ergodic theorem, for μ-almost every $x \in X$ it holds that

$$\mathrm{Lyap}_{\mathcal{A}_i}(x) = \int \log f'_x(i)\ d\mu(x), \tag{2.3}$$

and we shall denote this common value by $\mathrm{Lyap}(\mathcal{A}_i)$.

The version of the statement below is slightly more general than the corresponding one in [6, lema A.1], that asks for F to be C^2. In our context, $F(x,y) = (g(x), f_x(y))$ where g is an ergodic map for μ and $f_x \in C^1[0,1]$ for all x. In particular, if $(x,y) \mapsto f'_x(y)$ is continuous, the following equicontinuity hypothesis is satisfied.

Lemma 2.9. *If the family $\{f'_x\}_{x\in X}$ is equicontinuous at $i \in \{0,1\}$ and $\mathrm{Lyap}(\mathcal{A}_i) < 0$, then the basin of $\mu \otimes \delta_i$ has positive $\mu \otimes \mathrm{Leb}$ measure. Conversely, if $\mathrm{Lyap}(\mathcal{A}_i) > 0$, then the corresponding basin has null measure.*

Proof. We will assume $i = 0$. Let $\varepsilon > 0$ be such that

$$\int \log(f'_x(0) + \varepsilon)\ d\mu(x) < 0.$$

Let $a(x) = \log(f'_x(0) + \varepsilon)$. By the Birkhoff ergodic theorem, for μ-almost every $x \in X$

$$\lim_{n\to\infty} \frac{1}{n} \sum_{k=0}^{n-1} a(g^k(x)) = \int a(x)\ d\mu(x) < 0,$$

which implies $\lim_{n\to\infty} \sum_{k=0}^{n-1} a(g^k(x)) = -\infty$. This means that the function

$$A(x) = \max_{n\geq 1} \sum_{k=0}^{n-1} a(g^k(x)) \geq 0$$

is well-defined in almost every x.

By equicontinuity, there is $\delta > 0$ such that $0 \leq y \leq \delta$ implies $f'_x(x) \leq f'_x(0) + \varepsilon$ for all $x \in X$. Thus, if $y < \delta e^{-A(x)}$, we claim that

$$f^n_x(y) \leq \exp\left(\sum_{k=0}^{n-1} a(g^k(x)) - A(x)\right)\delta \leq \delta. \tag{2.4}$$

is true by induction. Indeed, the $n = 0$ case follows from the hypothesis on y, and if (2.4) is true for n, then by the mean value theorem,

$$\begin{aligned} f_x^{n+1}(y) &\leq \left(\max f'_{g^n(x)}([0, f_x^n(y)])\right) f_x^n(y) \\ &\leq (f'_{g^n(x)}(0) + \varepsilon) f_x^n(y) \\ &\leq \exp\left(a(g^n(x)) + \sum_{k=0}^{n-1} a(g^k(x)) - A(x)\right)\delta. \end{aligned}$$

Since $\lim_{n\to\infty} \sum_{k=0}^{n-1} a(g^k(x)) = -\infty$, we have $f_x^{n+1}(y) \to 0$. Thus, if $x \in \mathcal{B}(\mu)$ (which has full measure), then $e_n(x, y) \xrightarrow{w} \nu_0$ whenever $y \leq \delta e^{-A(x)}$.

□

Corollary 2.10. *If $(x, y) \mapsto \mathcal{S}f_x(y)$ is continuous and $\mathcal{S}f_x < 0$ for μ-almost every $x \in X$, then* $\mathrm{Lyap}(\mathcal{A}_0) + \mathrm{Lyap}(\mathcal{A}_1) < 0$. *In particular, at least one of the basins has positive measure.*

Proof. If suffices to integrate the logarithm of the inequality $f'_x(0)f'_x(1) < 1$ from lemma 2.6, and then apply lemma 2.9. □

2.4 Intermingled basins

The basins of the examples we saw before are continuity sets for the volume measure — that is, they are open sets modulo a zero volume set — and the union of all basins have full Lebesgue measure. But should basins of attractors constitute well-behaved sets in general? In some cases, such as for uniformly hyperbolic diffeomorphisms, this indeed the case, and there is a finite number of physical measures whose basins are continuity sets, and their union has full volume [9]. Outside the realm uniform hyperbolicity, though, this may not be the case — Kan [10] gave the first example of a partially hyperbolic endomorphism on a surface with two physical measures whose basins are intermingled, in the sense that for every open set U, the intersection of this set with either basin has positive volume [9]. This implies both basins are dense in the surface, and therefore not continuity sets.

Refer to discussion in [11]

In this section we shall describe the construction of that flavor of example in a more general setting of skew products that also encompasses the original example. As before, let X be a compact metric space, $g\colon X \to X$ be ergodic for $\mu \in \mathcal{M}_1(X)$ and $F\colon X \times I \to X \times I$ a continuous skew-product endomorphism of the form $F(x, y) = (g(x), f_x(y))$. We shall understand by "volume" the reference measure $m = \mu \otimes \mathrm{Leb}$. As mentioned in lemma 2.9, under some mild hypothesis for F, if the transverse Lyapunov exponents at the invariant boundaries $\mathcal{A}_i = X \times \{i\}$, given by

$$\mathrm{Lyap}(\mathcal{A}_i) = \int \log f'_x(i)\, d\mu(x)$$

for $i = 0, 1$ are both negative, then both basins have positive volume. Assuming this, the main property that will produce the intermingled phenomenon is the following hypothesis.

Hypothesis 1. There exists two periodic points x^+ and x^- in X, of periods n^+, n^-, such that for every $y \in (0,1)$, $f_{x^+}^{n^+}(y) > y$ and $f_{x^-}^{n^-}(y) < y$.

Note that if we define the measures $\mu_i(A) = \mathrm{Leb}(A \cap \mathcal{B}_i)$, then the basins are intermingled if, and only if, $\operatorname{supp} \mu_i = X \times I$ for both $i = 0, 1$. We will show this is the case for $\operatorname{supp} \mu_0$, as the proof for $i = 1$ follows by symmetry. First, we shall collect some lemmas.

Lemma 2.11. *Given $x \in X$, $\operatorname{supp} \mu_0 \cap \{x\} \times I$ is a segment $\{x\} \times [0, a_x]$.*

Proof. By lemma 2.7, $\mathcal{B}_0$ is of the above form. Since we know that the support of a measure is closed, we can take $y < 1$ such that $(x, y) \in \operatorname{supp} \mu_0$ and show that if $z < y$ then $(x, z) \in \operatorname{supp} \mu_0$. Let $U = B_\varepsilon(x, z)$ be a ball centered at (x, z) of radius smaller than $1 - y$, and $V = B_\varepsilon(x, y)$. Note that

$$V \cap \mathcal{B}_0 \subseteq \{(u, v + y - z) \mid (u, v) \in U \cap \mathcal{B}_0\},$$

so since $(x, y) \in \operatorname{supp} \mu_0$, and Leb is invariant by vertical translations, $\mathrm{Leb}(U \cap \mathcal{B}_0) \geq \mathrm{Leb}(V \cap \mathcal{B}_0) > 0$. □

Lemma 2.12. *The support $\operatorname{supp} \mu_0$ is an invariant set, that is, $F^{-1}(\operatorname{supp} \mu_0) = \operatorname{supp} \mu_0$.*

Proof. If $p \in F^{-1}(\operatorname{supp} \mu_0)$ and U is a neighborhood of p, then since F is an open map, $F(U)$ is a neighborhood of $F(p) \in \operatorname{supp} \mu_0$, so $\mu_0(F(U)) > 0$. Since F is Lipschitz, if $\mu_0(U)$ were 0, then $\mu_0(F(U)) = \mathrm{Leb}(F(U) \cap \mathcal{B}_0) = \mathrm{Leb}(F(U \cap \mathcal{B}_0)) = 0$, which is not the case, so $\mu_0(U) > 0$. □

Corollary 2.13. $\operatorname{supp} \mu_0 = X \times I$.

WRITE this

Proof. By symmetry, it is enough to show $\mathcal{A}_1 \subseteq K$. Let $(x, y) \in K$ be the point in the statement. Inductively, we can construct a backward orbit $\cdots \mapsto x_{-2} \mapsto x_{-1} \mapsto x_0$ of g such that $x_{-n-1} \in g^{-1}(\{x_{-n}\})$ is the closest preimage of x_-. Since x_- is a repelling fixed point, the construction implies that $x_{-n} \to x_-$, so by hypothesis 1, for every $y \in (0,1)$ there exists n_0 such that $n \geq n_0$ implies $f_{x_{-n}}^{-1}(y_0) > y$. In particular,

$$y_{-n} = f_{x_{-n}}^{-1} \circ \cdots \circ f_{x_{-1}}^{-1}(y_0)$$

is such that $y_{-n} \to 1$. Since $\operatorname{supp} \mu_0$ is closed and $(x_{-n}, y_{-n}) \in F^{-n}(\operatorname{supp} \mu_0)$, we conclude that $(x_-, 1) \in \operatorname{supp} \mu_0$. Now, notice that $E = \bigcup_{n=0}^{\infty} g^{-1}(\{x_-\})$ is dense in X and 1 is a fixed point of the f_x. Thus, $E \times \{1\} \subseteq \operatorname{supp} \mu_0$ and consequently $\mathcal{A}_1 \subseteq \operatorname{supp} \mu_0$. □

Corollary 2.14. *Suppose hypothesis 1 holds and both basins have positive measure. Then the basins $\mathcal{B}_0$ and $\mathcal{B}_1$ are intermingled for the Lebesgue measure on the cylinder.*

Proof. By definition, the basins are strictly invariant sets. If we define the measures $\mu_i(A) = \mathrm{Leb}(A \cap \mathcal{B}_i)$, then the basins are intermingled if, and only if, $\operatorname{supp} \mu_i = X \times I$ for both $i = 0, 1$. We will show this is the case for $\operatorname{supp} \mu_0$, as the proof for $i = 1$ is analogous.

Claim 1. The support $\operatorname{supp} \mu_0$ is backwards-invariant.

Indeed, if $p \in F^{-1}(\operatorname{supp} \mu_0)$ and U is a neighborhood of p, then since F is an open map, $F(U)$ is a neighborhood of $F(p) \in \operatorname{supp} \mu_0$, so $\mu_0(F(U)) > 0$. Since F is Lipschitz, if $\mu_0(U)$ were 0, then $\mu_0(F(U)) = \operatorname{Leb}(F(U) \cap \mathcal{B}_0) = \operatorname{Leb}(F(U \cap \mathcal{B}_0)) = 0$, which is not the case, so $\mu_0(U) > 0$.

Claim 2. There exists a point $(x_0, y_0) \in \operatorname{supp} \mu_0 \setminus X \times \{0\}$.

Indeed, $\mu_0(\operatorname{supp} \mu_0) \geq \operatorname{Leb}(\mathcal{B}_0) > 0$, so $\mu_0(\operatorname{supp} \mu_0 \setminus X \times \{0\}) > 0$ and the set cannot be empty.

Claim 3. Given $x \in X$, $\operatorname{supp} \mu_0 \cap \{x\} \times I$ is a segment $\{x\} \times [0, a_x]$.

By lemma 2.7, $\mathcal{B}_0$ is of the above form. To show that the support also has this property, we can take $y < 1$ such that $(x, y) \in \operatorname{supp} \mu_0$ and show that if $z < y$ then $(x, z) \in \operatorname{supp} \mu_0$. Let $U = B_\varepsilon(x, z)$ be a ball centered at (x, z) of radius smaller than $1 - y$, and $V = B_\varepsilon(x, y)$. Note that $V \cap \mathcal{B}_0 \subseteq (U \cap \mathcal{B}_0) + (0, y - z)$, and since $(x, y) \in \operatorname{supp} \mu_0$, $\operatorname{Leb}(U \cap \mathcal{B}_0) \geq \operatorname{Leb}(V \cap \mathcal{B}_0) > 0$. This shows the claim.

Now, by the first two claims and **??**, $\operatorname{supp} \mu_0$ contains both boundaries. By the last claim, it is composed of vertical segments, so it must be equal to $X \times I$. □



Figure 2.1: Picture of the basins from example 2.15.

Example 2.15. Define $q_a(y) = y + ay(1 - y)$ and $p(x) = \varepsilon \cos(2\pi x)$ for some $0 < \varepsilon < 1$. If $|a| < 1$, then q_a is a diffeomorphism of the interval I. If $F(x) = (3x, q_{p(x)}(y))$, then $x^+ = 0$ and $x^- = 1/2$ are fixed points of the map $g(x) = 3x$ (where we identify S^1 with $[0, 1]$) that satisfy hypothesis 1, because $q_a(y) > y$ if $a > 0$ and $q_a(y) < y$ if $a < 0$ for all $y \in (0, 1)$. Moreover, since q_a is a degree two polynomial, it has negative Schwarzian whenever $a \neq 0$, and from the symmetry of p we see both basins of $F(x, y) = (kx, q_{p(x)}(y))$ must have positive measure, because at least one of them has. So F displays intermingled basins.

2.5 Negative Schwarzian

The statement of the theorem below corresponds to [6, Theorem 3.2], but without the hypothesis that both basins have positive measure. The question of whether this hypothesis could be removed was left open in the article, and our proof below confirms that it is unnecessary.

Theorem 2.16. *If $(x, y) \mapsto \mathcal{S}f_x(y)$ is continuous and $\mathcal{S}f_x \prec 0$ almost everywhere in $x \in X$, then there exists a measurable function $\sigma\colon X \to I$ defined almost everywhere such that*

$$(x, y) \in \mathcal{B}_0 \text{ whenever } y < \sigma(x) \quad \text{and} \quad (x, y) \in \mathcal{B}_1 \text{ whenever } y > \sigma(x).$$

Consequently, the union $\mathcal{B}_0 \cup \mathcal{B}_1$ has total $\mu \otimes \mathrm{Leb}$ measure.

Proof. We start by defining functions $\xi_0, \xi_1 : X \to I$ that separate the basins in the following sense:

$$\xi_0(x) = \sup\{y \in I : (x, y) \in \mathcal{B}_0\}, \qquad \xi_1(x) = \inf\{y \in I : (x, y) \in \mathcal{B}_1\}.$$

By lemma 2.7, if $y < \xi_0(x)$ or $y > \xi_1(x)$, the empirical measures $e_n(x, y)$ converge to $\nu_0 = \mu \otimes \delta_0$ or to $\nu_1 = \mu \otimes \delta_1$, and if $\xi_0(x) < y < \xi_1(x)$, they do not converge to either ν_0 or ν_1.

It is sufficient to show that $A = \{x \in X : \xi_0(x) \neq \xi_1(x)\}$ has measure zero, since in this case we can define σ as the common value of the two functions. Suppose by contradiction that this set does not have measure zero. Since the f_x are bijective, A is an invariant set under g, so the ergodicity of μ implies that A has full measure.

Since we are under the hypotheses of corollary 2.10, we can assume without loss of generality that $\mathrm{Leb}(\mathcal{A}_1) > 0$. By ergodicity, we also have $\xi_1(x) < 1$ for μ-almost every $x \in X$. Thus, given $k \in \mathbb{N}$, there exists $\delta_k < 1$ such that $E_k = \{x \in X : \xi_1(x) < \delta_k\}$ has measure $\mu(E_k) > 1 - 1/k$. Applying the Birkhoff ergodic theorem to the functions 1_{E_k}, we obtain $B \subseteq X$ of full measure such that for all $x \in B$ and k,

$$\lim_{n\to\infty} \frac{1}{n} \#\{0 \le i < n \mid \xi_1(g^i(x)) < \delta_k\} = \mu(E_k) > 1 - 1/k. \tag{2.5}$$

By the ergodicity of μ, there exists another set $C \subseteq X$ of full measure whose empirical measures converge to μ, and since $\mathcal{S}f_x \prec 0$ almost everywhere in $x \in X$, there exists a set D of full measure such that $\mathcal{S}f_{g^n(x)} \prec 0$ for all $n \ge 0, x \in D$. Let $x_0 \in A \cap B \cap C \cap D$ and $y_0 \in (\xi_0(x), \xi_1(x))$, and let $\nu \in \mathcal{M}(x, y) \setminus \{\nu_0\}$ be a limit point of the empirical measures different from ν_0. Note that since $x_0 \in B$, necessarily $(\pi_X)_*\nu = \mu$, but since $\nu \neq \nu_0$, it follows that $\nu(X \times \{0\}) < 1$. Thus, $\operatorname{supp} \nu$ is not contained in $X \times \{0\}$, and there exists $\varepsilon > 0$ such that $U = X \times (\varepsilon, 1]$ satisfies $\nu(U) > 0$. Let k be such that $\nu(U) > 1/k$, and $n_j \to \infty$ such that $\liminf_{j\to\infty} e_{n_j}(x_0, y_0)(U) > 1/k$. In other words, we have

$$\liminf_{j\to\infty} \frac{1}{n_j} \#\{0 \le i < n_j \mid f^i_{x_0}(y_0) > \varepsilon\} > 1/k. \tag{2.6}$$

Now, consider the sequence $r_n = \rho(0, f^n_{x_0}(y_0), f^n_{x_0}(\xi_1(x_0)), 1)$, which is well-defined and positive since $0 < f^n_{x_0}(y_0) < f^n_{x_0}(\xi_1(x_0)) < 1$ by the definition of the points. Note also that the graph of ξ_1 is strictly invariant under F, so $f^n_{x_0}(\xi_1(x_0)) = \xi_1(g^n(x_0))$.

Since $x \in D$, it follows by lemma 2.5 that r_n is an increasing sequence. On one hand, it is not possible for $r_n \to \infty$, because from eqs. (2.5) and (2.6) we know that there exist $n \in \mathbb{N}$ arbitrarily large such that $0 < \varepsilon < f^n_{x_0}(y_0) < f^n_{x_0}(\xi_1(x_0)) < \delta_k < 1$. On the other hand, if $r_n \to r < \infty$, then for each $x \in X$ there exists a unique $\xi(x)$ such that $\rho(0, \xi(x), \xi_1(x), 1) = r$. In fact, a calculation shows that

$$\xi(x) = \frac{\xi_1(x)}{r(1 - \xi_1(x)) + \xi_1(x)}.$$

Note that $r_n \to r$ implies $\lim_{n\to\infty}|\xi(g^n(x_0)) - f^n_{x_0}(y_0)| = 0$. Since we know that x_0 is in the basin of μ, necessarily $e_n(x_0, y_0) \xrightarrow{w} \eta$, where $\eta = (\text{graph}\,\xi)_*\mu$ and $(\text{graph}\,\xi)(x) = (x, \xi(x))$. Thus, η is an invariant measure. But then,

$$\varphi(x, y) = \frac{\rho(0, y, \xi_1(x), 1)}{1 + \rho(0, y, \xi_1(x), 1)}$$

is an integrable function defined at η-almost every point, with the property that $\varphi(F(x, y)) > \varphi(x, y)$ at η-almost every point. This is impossible, as it would imply $\int \varphi \, d\eta < \int \varphi \circ F \, d\eta = \int \varphi \, d\eta$. □

Example 2.17. In example 2.15, we have $q_a'(y) = 1 + a(1 - 2y)$, $q_a''(y) = -2a$ and $q_a''' \equiv 0$, so $\mathcal{S}q_a(y) < 0$ whenever $a \neq 0$. Since $q(x) = 0$ only at two points, by theorem 2.16 there exists $\sigma\colon S^1 \to I$ that separates the two basins.

Hausdorff dimension

An interesting question to consider is what is the Hausdorff dimension of the set graph ξ in the situation of intermingled basins. Since this set is a graph, it is clearly at least 1 and at most 2.

In this situation of intermingled basins, what is the Hausdorff dimension of the graph separating the basins, or of the basins themselves?

It should be strictly greater than 1: this is a corollary of regularity.

2.6 Positive Schwarzian

the paragraph below conflicts, or should be moved to, the section which defines the empirical measures, as it's not clear whether it concerns an asymptotic measure for each point, or a single one for almost every point. If necessary, adjust the theorem statement.

If $f\colon M \to M$ is a diffeomorphism on a manifold, we say that an invariant measure μ is *asymptotic* with respect to a reference measure if the empirical measures of almost every point of the reference measure converges to μ. Intuitively, μ describes the entire statistical behavior of the system, up to a set of zero volume.

Since theorem 2.16 does not require that both basins have positive measure, we also obtain a more general statement than the version in [6, Theorem 5.1], which requires both Lyapunov exponents to be positive. The proof below is also somewhat different, more similar to the proof of theorem 2.16.

Theorem 2.18. *If $\mathcal{S}f_x > 0$ almost everywhere, then F has a unique asymptotic measure ν for $\mu \otimes \text{Leb}$-almost every point. Furthermore, for each $i \in \{0, 1\}$, $\text{Lyap}(\mathcal{A}_i) > 0$ implies $\nu(\mathcal{A}_i) = 0$.*

The idea of the proof is to pass to the invertible extension of the system and observe that the inverses of the fiber maps have negative Schwarzian. By applying theorem 2.16, we obtain an invariant graph that separates the two basins of the inverse transformation. Since the orbits diverge from the graph for the inverse of the extension, they converge to the graph for the

extension. Thus, the graph describes an asymptotic measure for the extension. The pushforward of this measure is the asymptotic measure for the original map.

Proof. Let $(\tilde{X}, \tilde{g}, \tilde{\mu})$ be the invertible extension of (X, g, μ), $\pi\colon (\tilde{X}, \tilde{g}, \tilde{\mu}) \to (X, g, \mu)$ the associated conjugation, and $\tilde{F}\colon \tilde{X} \times I \to \tilde{X} \times I$ given by

$$\tilde{F}(\tilde{x}, y) = (\tilde{g}(\tilde{x}), f_{\pi(\tilde{x})}(y)).$$

Note that $\tilde{F}^{-1}(x, y) = (\tilde{g}^{-1}(x), f^{-1}_{\pi(\tilde{g}^{-1}(x))}(y))$. Since $\pi_*\tilde{\mu} = \mu$ and $\tilde{g}_*\tilde{\mu} = \tilde{\mu}$, by proposition 2.3 we have $f^{-1}_{\pi(\tilde{g}^{-1}(\tilde{x}))} < 0$ for almost every $\tilde{x} \in \tilde{X}$. Thus, we can apply theorem 2.16 to F^{-1} and obtain a function $\sigma\colon \tilde{X} \to I$ that separates the basins $\tilde{\mathcal{B}}_1$ and $\tilde{\mathcal{B}}_0$.

The hyperbolic metric (which coincides with the Hilbert metric) on $(0, 1)$ is given by

$$d_H(r, s) = \log|\rho(0, r, s, 1)|.$$

This metric induces the usual topology of the interval. Since maps with positive Schwarzian decrease cross-ratios, the condition $\mathcal{S}f_x > 0$ means that f_x is a contraction (not necessarily uniform) for this metric.

Let $A \subseteq \tilde{X}$ be a set of full $\tilde{\mu}$ measure where $\mathcal{S}f_{\pi(\tilde{g}^n(\tilde{x}))} > 0$ for all $n \geq 0, \tilde{x} \in A$, and $B \subseteq \tilde{X}$ be another set of full measure where the empirical measures of $\tilde{x} \in B$ converge to the ergodic measure $\tilde{\mu}$. It follows that for all $\tilde{x} \in A \cap B$ and $y \in (0, 1)$, the limit $r = \lim_{n\to\infty} d_H(f^n_{\pi(\tilde{x})}(y), \sigma(\tilde{g}^n(\tilde{x})))$ exists and is zero if and only if the orbit of $(\tilde{x}, y)$ converges in distance to the graph of σ under $\tilde{F}$, which in turn implies that $e_n(\tilde{x}, y) \xrightarrow{w} \tilde{\nu} = (\operatorname{graph}\sigma)_*\tilde{\mu}$.

Suppose $r > 0$. Then, as in the previous proof, there would exist another graph ξ, different from σ almost everywhere, such that $e_n(\tilde{x}, y) \xrightarrow{w} \eta = (\operatorname{graph}\xi)_*\tilde{\mu}$. Since $\varphi(\tilde{x}, y) = d_H(y, \sigma(\tilde{x}))$ is such that $\varphi(F(\tilde{x}, y)) < \varphi(\tilde{x}, y)$ whenever $(\tilde{x}, y) \in \operatorname{graph}\xi$ and η is an invariant measure for F with $\eta(\operatorname{graph}\xi) = 1$, we reach a contradiction.

This shows that $E = (A \cap B) \times (0, 1)$ is a set of full $\tilde{\mu} \otimes \mathrm{Leb}$ measure such that $(\tilde{x}, y) \in E$ implies $e_n(\tilde{x}, y) \xrightarrow{w} \tilde{\nu}$. Let $\nu = \pi_*\tilde{\nu} \in \mathcal{M}(X \times I)$. The measure ν is invariant, since

$$F_*\nu = (F \circ \pi)_*\tilde{\nu} = (\pi \circ \tilde{F})_*\tilde{\nu} = \pi_*\tilde{\nu} = \nu.$$

Furthermore, the basin of ν has full Lebesgue measure. In fact, if $\varphi \in C(X \times I)$, then $\varphi \circ \pi \in C(\tilde{X} \times I)$ and $e_n(\pi(\tilde{x}), y)(\varphi) = e_n(\tilde{x}, y)(\varphi \circ \pi)$. Therefore, $e_n(x, y) \to \nu$ if and only if $e_n(\tilde{x}, y) \to \tilde{\nu}$ for any $\tilde{x} \in \pi^{-1}(x)$, that is, $(\pi \times \mathrm{id})^{-1}(\mathcal{B}(\nu)) = \mathcal{B}(\tilde{\nu})$. It follows that $(\mu \otimes \mathrm{Leb})(\mathcal{B}(\nu)) = (\tilde{\mu} \otimes \mathrm{Leb})(\mathcal{B}(\tilde{\nu})) = 1$. □

Perhaps a section should be dedicated to the hyperbolic metric, or the Hilbert metric. Perhaps put it next to the Hausdorff metric.

Chapter 3

Historical behavior

Ergodic theory tries to find order in chaos by showing that even if a dynamical system looks complicated, the statistical properties of its orbits are often well behaved. But what if the statistics don't even exist for most orbits? When this happens for all orbits starting in a set of positive reference measure, the phenomenon is known as *non-statistical*, or *historical* behavior.

More precisely, by existence of statistics, we mean the weak convergence of the empirical measures $e_n(x) \xrightarrow{w} \mu$ to an asymptotic measure μ. To say that $(e_n(x))_n$ does not converge, therefore, is equivalent (in the compact setting) to the existence of a continuous observable $\varphi\colon X \to \mathbb{R}$ such that

$$\lim_{n\to\infty} \frac{1}{n} \sum_{i=0}^{n-1} \varphi(f^i(x)) \tag{3.1}$$

does not exist.[1] We then say that a diffeomorphism f of a compact manifold presents historical behavior if $(e_n(x))_n$ does not converge for x in a set of positive Lebesgue measure.

3.1 History

The "historical" terminology was introduced by Ruelle in [12], suggesting that for eq. (3.1) not to converge, the sequence must keep "changing its mind" as n goes to infinity, and therefore has some sort of history. For instance, in the absence of external forces like human activity, we could expect that general patterns in climate would follow some predictable statistics, while with climate change we see that such statistics are not consistent along history. In the article, Ruelle recognized the problem of *nonrecurrent* behavior of orbits as going against the expectation that recurrent behavior is generic in the space of diffeomorphisms, and presented the open problem of finding whether historical behavior can exist in a persistent manner, making some physical justifications for this possibility. He cites the theory of spin glasses, where the time evolution of the system is akin to a random walk in a random potential: a particle stays arbitrarily large amounts of time exploring some valley in the potential before eventually crossing the barrier to another valley, in such a way that time averages like eq. (3.1) do not exist. As we shall see in

[1] The fact we don't need to quantify over measures that could be limits here is a consequence of corollary A.2.

this chapter, we can indeed produce historical behavior by constructing a diffeomorphism whose orbits behave like random walks, albeit not necessarily persistently.

Later, Floris Takens exposed the problem again in [13], giving some promising examples that were known at the time and also making some remarks about generiticity of points without asymptotic measures. For this reason and his work in related matters, the open question also became known as *Taken's Last Problem*.

Example 3.1 (Logistic family)**.** The family of interval maps $f_a(x) = ax(1-x)$, for $0 < a \leq 4$, is known as the logistic family, and is one of the earliest examples where historical behavior is present abundantly: in [14], the authors showed that for an uncountable number of parameters a, Lebesgue almost every orbit in $[0,1]$ has no asymptotic measure.

figure

Example 3.2 (Bowen's eye)**.** Another example was given by Bowen, although it's also not persistent.

figure

Genericity of historical orbits

Under relatively few hypothesis, historical orbits form a residual subset of the manifold.

[15]

3.2 Martingale-coboundary decomposition

The key idea for applying theorem B.20 in a dynamical system $(X, \mathcal{F}, \mu, f)$, where μ is a f-invariant probability, is finding a sub-σ-algebra $\mathcal{G} \subseteq \mathcal{F}$ for which $\varphi \in L^2(\mu)$ can be decomposed as a sum

$$\varphi = \varphi_0 + \phi \circ f - \phi \tag{3.2}$$

such that $\phi \in L^2(\mu)$, φ_0 is $\mathcal{G}$-measurable and $\eta = (\varphi_0 \circ f^{i-1})_{i \geq 1}$ is a reversed martingale difference sequence. Indeed, this allows the transference of the weak invariant principle for the original φ: let $\xi = (\varphi \circ f^{i-1})_{i \geq 1}$, and notice that

$$S_n^\xi = S_n^\eta + \phi \circ f^n - \phi,$$

so

$$D_n^\xi(x)(t) = \frac{S_{\lfloor nt \rfloor}^\xi(x)}{\sqrt{n}} = D_n^\eta(x)(t) + \frac{1}{\sqrt{n}}\big(\phi \circ f^{\lfloor nt \rfloor} - \phi\big)(x).$$

It suffices to show that for μ-almost every x, the second term in the sum converges uniformly to zero (as a function of t). Given $\varepsilon > 0$, we look at the sets $B_n = \{x : (\phi^2 \circ f^n)(x) \geq n\varepsilon\}$. Since

$$\sum_{n=1}^{\infty} \mu(B_n) = \sum_{n=1}^{\infty} \mu(\{\phi^2 \geq n\varepsilon\}) = \int \lfloor \phi^2/\varepsilon \rfloor \, d\mu \leq \frac{1}{\varepsilon}\|\phi\|_2^2,$$

by Borel-Cantelli, $\mu(\limsup_{n\to\infty} B_n) = 0$. If we take the intersection over k of the sets $\liminf_{n\to\infty} B_n^c$ for $\varepsilon = 1/k$, we get a full-measure set G with the property that for every $x \in G$ and $\varepsilon > 0$, there exists N such that $n \geq N$ implies $\frac{1}{\sqrt{n}}|\phi(f^n(x))| \leq \varepsilon$. Now, let $M \geq N$ be big enough so that $n \geq M$ implies $\frac{1}{\sqrt{n}}|\phi(f^i(x))| \leq \varepsilon$ for all $0 \leq i \leq N$. So for $n \geq M$,

$$\sup_{t\in[0,1]} \frac{1}{\sqrt{n}}\left|\phi \circ f^{\lfloor nt \rfloor} - \phi\right|(x) \leq \max_{0\leq i\leq N} \frac{|\phi \circ f^i(x)|}{\sqrt{n}} + \max_{N\leq i\leq n} \frac{|\phi \circ f^i(x)|}{\sqrt{i}} \leq 2\varepsilon$$

as desired.

Existence of martingale-coboundary decomposition

From now on, we assume that μ is an ergodic measure for f.

Theorem 3.3. *Suppose $\varphi \in L^2(\mu)$ and $\mathcal{F}_j = f^{-j}(\mathcal{F})$ are such that*

$$\sum_{n=0}^{\infty} \|\mathbb{E}_\mu(\varphi \mid \mathcal{F}_j)\|_2 < \infty.$$

Then a decomposition as in eq. (3.2) exists. Moreover, this implies that

$$\int \varphi \, d\mu = \int \varphi_0 \, d\mu = 0$$

and that the series in

$$\sigma^2 = \|\varphi_0\|_2^2 = \|\varphi\|_2^2 + 2\sum_{n=1}^{\infty} \int \varphi \cdot \varphi \circ f^n \, d\mu$$

converges absolutely. By the preceding arguments, if $\varphi_0 \neq 0$ then $(D_n^\xi)_*\mu \xrightarrow{w} \sigma W$.

Proof. We begin by writing $\phi = \phi^- + \phi^+$, where $\phi^+ = \mathbb{E}_\mu(\phi \mid \mathcal{G}_0)$.

write □

Remark 3. If φ was a coboundary $\varphi = \phi \circ f - \phi$ to begin with, then it is clear that $S_n^\xi = \phi \circ f^n - \phi$, so the iterates of the skew product $F(x, y) = (f(x), y + \varphi(x))$ satisfy

$$F^n(x_0, y_0) = (f^n(x_0), \phi(f^n(x_0)) - \phi(x_0) + y_0) \subseteq \mathrm{graph}(\psi)$$

and are thus contained in the graph of ψ, where $\psi(x) = \phi(x) - \phi(x_0) + y_0$. In particular, F is not ergodic for $\mu \otimes \mathrm{Leb}$.

Corollary 3.4. *Suppose φ is such that there exists a summable sequence $c_n > 0$ satisfying*

$$\int (\psi \circ f^n)\varphi \, d\mu \leq \|\psi\|_2 c_n$$

for all $\psi \in L^2(\mathcal{G})$, where $\mathcal{G}$ is as in theorem 3.3. Then

$$\sum_{n=0}^{\infty} \|\mathbb{E}(\varphi \mid f^{-n}(\mathcal{G}))\|_2 < \infty.$$

Remark 4. The book [16] proves this condition is satisfied for smooth expanding maps, Hölder-continuous φ and $\mathcal{G} = \mathcal{F}$, so this includes the $f(x) = kx$ maps. The second condition is trivial when $\mathcal{G} = \mathcal{F}$.

3.3 Historical behavior for skew products

Let (X, μ) be a Borel probability space, where X is compact and metrizable, and $F\colon X \times I \to X \times I$ a skew product of the form

$$F(x, y) = (g(x), f_x(y))$$

where $g\colon X \to X$ is ergodic and measure-preserving. Suppose there is a homeomorphism $\gamma\colon I \to \overline{\mathbb{R}}$ and a measurable function $p\colon X \to \mathbb{R}$ such that, if we set $t_x(y) = y + p(x)$, we have

$$t_x \circ \gamma = \gamma \circ f_x$$

for all $x \in X$, where t_x is to be interpreted as the translation by $p(x)$.

$$\begin{tikzcd} I \arrow[r, "f_x"] \arrow[d, "\gamma"] & I \arrow[d, "\gamma"] \\ \overline{\mathbb{R}} \arrow[r, "t_x"] & \overline{\mathbb{R}} \end{tikzcd}$$

Since the t_x are translations, there is no loss of generality in assuming $\gamma(0) = -\infty$, $\gamma(1) = +\infty$ and $\gamma\left(\frac{1}{2}\right) = 0$, so we shall adopt this convention for simplicity.

Theorem 3.5. *If the weak invariance principle is valid for the pair (g, p), in the sense $(D_n^\xi)_*\mu \xrightarrow{w} \sigma W$ for $\sigma > 0$ and $\xi = (p \circ g^{i-1})_{i \geq 1}$, then F has historical behavior. More specifically,*

$$\{\delta_-, \delta_+\} \subseteq \{e_n(x, y)\}'_{n \in \mathbb{N}} \tag{3.3}$$

where $\delta_- = \mu \otimes \delta_0$ and $\delta_+ = \mu \otimes \delta_1$ and the limit points are taken in the weak topology.*

Proof. Let $\delta > 0$ and $s_\delta\colon [0, 1] \to \mathbb{R}$ be any continuous function such that $s_\delta(0) = 0$ and $s_\delta(t) > 1 + \delta$ for $t > \delta$. Let $U_\delta = B_\delta(s_\delta)$ be an open ball of radius δ around s_δ in the uniform topology. By theorem B.11, we know $W(U_\delta) > 0$, so by proposition B.3

$$\liminf_{n \to \infty} ((D_n^\xi)_*\mu)(U_\delta) \geq \sigma W(U_\delta) > 0.$$

Therefore,

$$S_\delta = \limsup_{n \to \infty} (D_n^\xi)^{-1}(U_\delta) \subseteq X$$

has positive measure. This set is also invariant for g because

$$\begin{aligned} g^{-1}(S_\delta) &= \limsup_{n \to \infty} \{x \in X \mid D_n^\xi(g(x)) \in U_\delta\} \\ &= \left\{ x \in X : \liminf_{n \to \infty} \sup_t \left| \frac{S^\xi_{\lfloor nt \rfloor + 1}(x) - p(x)}{\sqrt{n}} - s_\delta(t) \right| < \delta \right\} \\ &= \left\{ x \in X : \liminf_{n \to \infty} \sup_t \left| \frac{S^\xi_{\lfloor nt' \rfloor}(x)}{\sqrt{n}} - s_\delta(t) \right| < \delta \right\} \text{ where } t' = t + 1/n \\ &= \left\{ x \in X : \liminf_{n \to \infty} \sup_t \left| \frac{S^\xi_{\lfloor nt \rfloor}(x)}{\sqrt{n}} - s_\delta(t) \right| < \delta \right\} = S_\delta. \end{aligned}$$

Therefore $\mu(S_\delta) = 1$. Now, notice that if $x \in S_\delta$, then there is $n_k^+ \to \infty$ for which for every k, $D_{n_k}^\xi(x) \in U_\delta$. This in turn implies that

$$\#\left\{0 \le i < n_k^+ \,\middle|\, S_i^\xi(x) > \sqrt{n_k^+}\right\} \ge (1-\delta)n_k^+.$$

Analogously, by repeating the argument with $-s_\delta$, we find $n_k^- \to \infty$ such that for every k,

$$\#\left\{0 \le i < n_k^- \,\middle|\, S_i^\xi(x) < -\sqrt{n_k^-}\right\} \ge (1-\delta)n_k^-.$$

By considering $E = \bigcap_{i=1}^\infty U_{1/i} \subseteq X$, a set of full measure, and applying a diagonalization argument to the previous two observations, for each $x \in E$ we find $n_k^+, n_k^- \to \infty$ for which

$$\frac{1}{n_k^+}\#\left\{0 \le i < n_k^+ \,\middle|\, S_i^\xi(x) > \sqrt{n_k^+}\right\} \to 1$$

and

$$\frac{1}{n_k^-}\#\left\{0 \le i < n_k^- \,\middle|\, S_i^\xi(x) < -\sqrt{n_k^-}\right\} \to 1. \tag{3.4}$$

Now, by ergodicity of g and lemma 1.16, let $S = E \cap \mathcal{B}(\mu)$ be a full measure set where

$$\frac{1}{n}\sum_{i=0}^{n-1} \varphi(g^i(x)) \to \int_X \varphi(z)\, d\mu(z)$$

for all $\varphi \in C(X)$ and $x \in S$. It suffices to show that for all $x \in S$,

$$e_{n_k^-}(x,y) \xrightarrow{w} \delta_- \quad \text{and} \quad e_{n_k^+}(x,y) \xrightarrow{w} \delta_+.$$

We only show the first convergence, since the second is analogous.

Let $\phi \in C(X \times I)$ and $\varepsilon > 0$. Since ϕ is uniformly continuous, there is some k_1 for which

$$\sup\left\{|\phi(x,0) - \phi(x,s)| : x \in X, s \in [0, 1/k_1]\right\} < \varepsilon.$$

Let $D = \{i \mid f_x^{(i)}(y) < 1/k_1\}$, and let N be big enough so that for $k \ge N$,

$$\left|\frac{1}{n_k^-}\sum_{i=0}^{n_k^- - 1} \phi(F^i(x,0)) - \int_X \phi(z,0)\, d\mu(z)\right| < \varepsilon,$$

which is possible since $x \in S$ and $\varphi(x) = \phi(x,0)$ is continuous, and also big enough so that

$$\frac{1}{n_k^-}\#\{0 \le i < n_k^- \mid i \notin D\} < \varepsilon/\|\phi\|_\infty,$$

which is due to eq. (3.4). Therefore, we can estimate

$$
\begin{aligned}
&|e_{n_k^-}(x,y)(\phi) - \delta_-(\phi)| \\
&\quad = \left| \frac{1}{n_k^-} \sum_{i=0}^{n_k^- - 1} \phi(F^i(x,y)) - \int_X \phi(z,0)\, d\mu(z) \right| \\
&\quad \le \left| \frac{1}{n_k^-} \sum_{i=0}^{n_k^- - 1} \phi(F^i(x,y)) - \frac{1}{n_k^-} \sum_{i=0}^{n_k^- - 1} \phi(F^i(x,0)) \right| + \varepsilon \\
&\quad \le \frac{1}{n_k^-} \sum_{\substack{0 \le i < n_k^- \\ i \in D}} \left| \phi(F^i(x,y)) - \phi(F^i(x,0)) \right| + \frac{2}{n_k^-} \sum_{\substack{0 \le i < n_k^- \\ i \notin D}} \|\phi\|_\infty + \varepsilon \\
&\quad \le \frac{1}{n_k^-} \sum_{\substack{0 \le i < n_k^- \\ i \in D}} \left| \phi(F^i(x,y)) - \phi(F^i(x,0)) \right| + 3\varepsilon \\
&\quad \le \varepsilon + 3\varepsilon = 4\varepsilon.
\end{aligned}
$$

This completes the proof. □

Corollary 3.6. *If for any compact $K \subseteq \mathbb{R}$, the vising time of the processes S_n^ξ to K tends to zero almost surely as $n \to \infty$, that is,*

$$\int 1_K \circ S_n^\xi(x)(t)\, d\,\mathrm{Leb} \to 0$$

then

$$\{e_n(x,y)\}'_{n\in\mathbb{N}} = [\delta_-, \delta_+] \tag{3.5}$$

for μ-almost every x and all $y \in (0,1)$.

Proof.

Claim 1. If $\eta \in \mathcal{M}(x,y)$, then $\operatorname{supp}\eta \subseteq X \times \{0,1\}$ and $\eta \in [\delta_-, \delta_+]$.

We use **??** again for this. Let $e_{n_k}(x,y) \xrightarrow{w} \eta$ be a limit point, $x_0 \in X$ and $y_0 \in (0,1)$, so there is a neighborhood U of (x_0, y_0) such that $\overline{U} \subseteq X \times (0,1)$. By the lemma, $e_{n_k}(x,y)(U) \to 0$, so by Portmanteau

$$\eta(U) \le \liminf_{k\to\infty} e_{n_k}(x,y)(U) = 0.$$

This shows $(x_0, y_0) \notin \operatorname{supp}\eta$.

We know η is a limiting measure for x, so it is invariant by F. The fact $x \in S$ implies by the definition of S that $(\pi_1)_*\eta = \mu$. This means that $\eta \ll \delta_- + \delta_+$, and since δ_- and δ_+ are ergodic measures, η must be a convex combination of the two.

We have shown $\mathcal{M}(x,y) \subseteq [\delta_-, \delta_+]$. To conclude eq. (3.3), it suffices to notice that by proposition 1.15, $\mathcal{M}(x,y)$ is connected, and therefore, since it contains both endpoints δ_- and δ_+, it must be the whole segment. □

3.4 Recent developments

Conclude with an overview of the recent work of Kiriri, Soma, Vargas et al.

Chapter 4

Partially hyperbolic dynamics

4.1 Partial hyperbolicity

4.2 Invariance Principle

Definition 4.1. A partially hyperbolic diffeomorphism is *accessible* if between any two points in the manifold, there is a path composed of segments tangent to either E^s or E^u.

Definition 4.2. A partially hyperbolic diffeomorphism is *dynamically coherent* if there exists invariant foliations $\mathcal{F}^{cs}$ and $\mathcal{F}^{cu}$ tangent to $E^c \oplus E^s$ and $E^c \oplus E^u$, respectively.

Theorem 4.3. *Let $f\colon M \to M$, where $M = X \times Y$, be a skew product diffeomorphism with an ergodic invariant measure μ. Then, $\lambda^c(\mu) \leq 0$ implies that $\{\mu_x^c\}_{x\in M}$ is invariant by unstable holonomies, meaning that $(h^u_{x,y})_*\mu_x^c = \mu_y^c$.*

Sketch the proof as in the notes taken on ICTP.
What does it say about the Bonifant-Milnor example.

Appendix A

Functional Analysis

Perhaps it's not worth to split this appendix into sections, and instead have just a collection of miscellanea.

Theorem A.1 (Riesz–Markov–Kakutani)**.** *Let K be a compact space. Then, for every continuous linear functional $L\colon C(K) \to \mathbb{R}$, there exists a unique (possibly signed) Radon measure $\mu \in \mathcal{M}(K)$ such that*

$$L(\varphi) = \int \varphi \, d\mu, \quad \forall \varphi \in C(K).$$

Proof. See [17, Theorem 7.10.4], or most textbooks on functional analysis. □

Corollary A.2. *If X is a compact space, $(\nu_n)_n \in \mathcal{M}(X)^{\mathbb{N}}$ is a sequence of measures and for all $\varphi \in C(X)$, the limit $L(\varphi) = \lim_{n\to\infty} \int \varphi \, d\nu_n$ exists, then $(\nu_n)_n$ converges in the weak* topology. Moreover, if the ν_n are probabilities, then the weak limit $\nu = \lim_{n\to\infty} \nu_n$ is a probability.*

Proof. By theorem A.1 and definition of weak* convergence, it suffices to show that L is a continuous linear functional. This, in turn, is a direct application of Banach-Steinhaus to the sequence of linear operators $L_n(\varphi) \mapsto \int \varphi \, d\nu_n$, since the hypotheses imply $C(X)$ is a Banach space. The last observation is simply the fact $\nu(X) = \lim_{n\to\infty} L_n(1) = 1$. □

Proposition A.3. *If a normed vector space E is separable, then the weak* topology on its dual E' is metrizable. For instance, for a choice $\{\phi_i\}_{i\in\mathbb{N}} \subseteq E$ of a countable dense set,*

$$d(\mu, \nu) = \sum_{i=1}^{\infty} \frac{|\mu(\phi_i) - \nu(\phi_i)|}{2^i},$$

defines a metric.

Proof. See [18, proposition 6.5.1]. □

Appendix B

Measure and probability

B.1 Probability spaces

Products

We can take arbitrary products of probability measures. See [17]

Borel Spaces

The Kolmogorov Extension Theorem

What if we want to construct a measure on a product space that is not the product measure? Kolmogorov to the rescue.

The following theorem is known as the *Kolmogorov Extension Theorem*, as quoted from [17, theorem 7.7.1].

Theorem B.1. *Suppose that for every finite set $\Lambda \subset T$, we are given a probability measure μ_Λ on $(\Omega_\Lambda, \mathcal{B}_\Lambda)$ such that the following consistency condition is fulfilled: if $\Lambda_1 \subset \Lambda_2$, then the image of the measure μ_{Λ_2} under the natural projection from Ω_{Λ_2} to Ω_{Λ_1} coincides with μ_{Λ_1}. Suppose that for every $t \in T$, the measure μ_t on $\mathcal{B}_t$ possesses an approximating compact class $\mathcal{K}_t \subset \mathcal{B}_t$. Then, there exists a probability measure μ on the measurable space $\left(\Omega := \prod_{t\in T} \Omega_t, \mathcal{B} := \bigotimes_{t\in T} \mathcal{B}_t\right)$ such that the image of μ under the natural projection from Ω to Ω_Λ is μ_Λ for each finite set $\Lambda \subset T$.*

B.2 Random variables

If $(X, \mathcal{A}, \mathbb{P})$ is a probability space and $(Y, \mathcal{B})$ is a measurable space, then a *random variable* is simply a measurable function $f : X \to Y$. It can be interpreted as some observable quantity of the probability space. If μ is a measure on Y, we may write $f \sim \mu$ to mean that $f_*\mathbb{P} = \mu$, and we say that f has distribution μ.

Definition B.2. We say the collection $\xi = (\xi_i)_{i\in\mathcal{I}} \subseteq X^Y$ of random variables is *independent* if $\xi_*\mathbb{P}$ is a product measure, that is, if

$$\xi_*\mathbb{P} = \bigotimes_{i\in\mathcal{I}} (\xi_i)_*\mathbb{P}.$$

This is equivalent the usual notion that ξ is independent if for every finite subset $J \subseteq \mathcal{I}$ and for every collection of measurable sets $A_j \in \mathcal{B}$, we have

$$\mathbb{P}\left(\bigcap_{j\in J} \xi_j^{-1}(A_j)\right) = \prod_{j\in J} \mathbb{P}\left(\xi_j^{-1}(A_j)\right).$$

B.3 Weak convergence

Proposition B.3 (Portmanteau)**.** *The following are equivalent:*

i. $\mu_n \xrightarrow{w} \mu$;

ii. $\lim_{n\to\infty} \mu_n(\phi) = \mu(\phi)$ *for all Lipschitz functions* ϕ;

iii. $\liminf_{n\to\infty} \mu_n(U) \geq \mu(U)$ *for all open sets* U;

iv. $\limsup_{n\to\infty} \mu_n(F) \leq \mu(F)$ *for all closed sets* F;

v. $\lim_{n\to\infty} \mu_n(A) = \mu(A)$ *for all continuity sets* A *of the measure* μ.

B.4 Brownian motion

The rigorous mathematical foundation of Brownian motion was laid by Norbert Wiener in 1923, who constructed Brownian motion as a measure on the space of continuous functions. This construction is why the Brownian motion is often called the Wiener process, and defined as the *Wiener measure.*

We shall denote by $\mathcal{N}(\mu, \nu) \in \mathcal{M}_1(\mathbb{R})$ the probability measure corresponding to the normal distribution with mean μ and variance ν. It can be defined as $\mathcal{N}(\mu, \nu) = \rho\,\mathrm{Leb}$, where ρ is the probability density function

$$\rho(x) = \frac{1}{\sqrt{2\pi\nu}} \exp\left(-\frac{(x-\mu)^2}{2\nu}\right).$$

Definition B.4. Let $\mathcal{C}$ be the product σ-algebra on $C([0,1])$. The Wiener measure W is the measure on $(C([0,1]), \mathcal{C})$ defined by the following properties: if we denote by $W_t \colon C([0,1]) \to \mathbb{R}$ the evaluation at time t, seen as a random variable on $(C([0,1]), \mathcal{C}, W)$, then

(i) $W_0 = 0$ almost surely;

(ii) for $t > 0$, $W_t \sim \mathcal{N}(0, t)$;

(iii) if $0 \leq t_1 < t_2 < \cdots < t_n$, then the increments $W_{t_1}, W_{t_2} - W_{t_1}, \ldots, W_{t_n} - W_{t_{n-1}}$ are independent random variables.

The space of càdlàg functions

The space $C([0,1])$ of continuous functions does not let us talk about processes that contain discontinuities, or even about how discontinuous processes can converge in a suitable way to continuous ones. This is important in our setting to deal with the convergence of certain processes induced by discrete dynamics, which consist of step functions, to the Brownian motion, which is continuous almost surely.

The space of *càdlàg* functions, or Skorohod space, is the space of functions whose discontinuities are of the *first kind*, that is, both lateral limits exist, and whose value at the point is equal to the right limit:

$$\mathbb{D}([0,1]) = \left\{ f\colon [0,1] \to \mathbb{R} \,\middle|\, \forall t, \lim_{s\to t^-} x(s) \text{ exists and } \lim_{s\to t^+} x(s) = x(t) \right\}.$$

We want to put a metric on this space with the property that functions are close together if their values are close to each other uniformly, but also allowing small perturbations in the parametrization of time.

Definition B.5 (Skorohod topology)**.** Let Λ be the set of strictly increasing continuous functions from $[0,1]$ to itself. For $f, g \in \mathbb{D}([0,1])$, we define

$$d(f,g) = \inf_{\lambda\in\Lambda} \max\left\{ \|\lambda - \mathrm{id}\|_\infty, \|f - g\circ\lambda\|_\infty \right\}. \tag{B.1}$$

The distance above defines a metric on $\mathbb{D}([0,1])$, whose topology is called the Skorohod topology. Since $d(f,g) \le \|f-g\|_\infty$, the Skorohod topology is coarser than the uniform topology.

Proposition B.6. *The restriction of the Skorohod topology to $C([0,1])$ coincides with the uniform topology. In fact, if $f\colon [0,1] \to \mathbb{R}$ is continuous, then any ball*

$$B_\delta(f) = \{g \in \mathbb{D}([0,1]) \mid \|f-g\|_\infty < \delta\},$$

of the uniform topology contains an open neighborhood of f in the Skorohod topology.

Proof. Since the Skorohod topology is coarser, it suffices to show the second part of the statement. Since f is uniformly continuous, there is $\delta_2 < \delta/2$ such that if $\|\lambda - \mathrm{id}\|_\infty < \delta_2$, then

$$\|f - f\circ\lambda\|_\infty < \delta/2.$$

Let

$$B'(f) = \{g \in \mathbb{D}([0,1]) \mid d(g,f) < \delta_2/2\}.$$

The set $B'(f)$ is a ball of the Skorohod topology, so it suffices to show $B'(f) \subseteq B_\delta(f)$. Let $g \in B'(f)$. Since $\delta_2/2 < \delta_2$, there exists λ such that

$$\max\left\{ \|\lambda - \mathrm{id}\|_\infty, \|g - f\circ\lambda\|_\infty \right\} < \delta_2 < \delta/2.$$

Therefore,

$$\|g - f\|_\infty \le \|g - f\circ\lambda\|_\infty + \|f\circ\lambda - f\|_\infty < \delta.$$

□

Proposition B.7. *If $f_n \to f$ in the Skorohod topology, then $f_n(x) \to f(x)$ for all x in a set E whose complement is at most countable.*

This is not necessary for lemma B.9 because we can take E to be a subset of continuous functions there. The continuous functions $C([0,1])$ are closed in $\mathbb{D}([0,1])$. This could be removed.

Proof. Since $f \in \mathbb{D}([0,1])$, the set of discontinuities of f is at most countable[1]. Let E be a cocountable set such that $f|_E$ is continuous, and let $x \in E$ and $\varepsilon > 0$. Pick δ so that $|x-y| < \delta$ implies $|f(x) - f(y)| < \varepsilon$, and let N be big enough so that $n \geq N$ implies $d(f, f_n) < \min\{\delta, \varepsilon\}/2$. If $n \geq N$, and if we pick a $\lambda \in \Lambda$ that makes the right side of eq. (B.1) $\min\{\delta, \varepsilon\}/2$-close to the infimum, then

$$\begin{aligned} |f(x) - f_n(x)| &\leq |f(x) - f \circ \lambda^{-1}(x)| + |f \circ \lambda^{-1}(x) - f_n(x)| \\ &\leq \varepsilon + \|f - f_n \circ \lambda\|_\infty \leq 2\varepsilon, \end{aligned}$$

because $|\lambda^{-1}(x) - x| \leq \|\lambda - \mathrm{id}\|_\infty < \delta$.

□

The arcsine law

Definition B.8. We shall denote by $\tau^+ \colon \mathbb{D}([0,1]) \to \mathbb{R}$ the observable that returns the proportion of time its function argument is positive, that is,

$$\tau^+(f) = \mathrm{Leb}\{x \in [0,1] : f(x) > 0\} = \int_{[0,1]} 1_{(0,\infty)} \circ f \, d\,\mathrm{Leb}\,.$$

Lemma B.9. *τ^+ is measurable and continuous in a full W-measure set.*

Proof. Since the normal distributions are non-atomic and W_t has normal distribution for every $t \in (0,1]$, we have that

$$W(\{f \in \mathbb{D}([0,1]) : f(t) = 0\}) = 0$$

for all $t \in (0,1]$. Therefore, by Fubini,

$$\mathrm{Leb}(\{t \in [0,1] : f(t) = 0\}) = 0$$

for all f in a set $E \subseteq \mathbb{D}([0,1])$ of full W-measure. It remains to check that τ^+ is continuous at such points $f \in E$. Indeed, 0 is the only discontinuity of the indicator function $1_{(0,\infty)}$, so if f is in this set and $f_n \to f$ in the Skorohod space, then $1_{(0,\infty)} \circ f_n \to 1_{(0,\infty)} \circ f$ almost everywhere by proposition B.7. By the dominated convergence theorem,

$$\tau^+(f) = \int_{[0,1]} 1_{(0,\infty)} \circ f \, d\,\mathrm{Leb} = \lim_{n\to\infty} \int_{[0,1]} 1_{(0,\infty)} \circ f_n \, d\,\mathrm{Leb} = \lim_{n\to\infty} \tau^+(f_n).$$

□

[1]Proof: let t be a discontinuity point of f, and let U be a neighborhood of t where the value of the function is close to the lateral limits at t. Then, f is continuous at $U \setminus \{t\}$, which is open, and therefore contains a rational point q_t. With some care, this will define an injection from discontinuity points to rationals.

Theorem B.10 (Arcsine law)**.** *For $\alpha \in [0,1]$,*

$$(\tau^+)_* W([0,\alpha]) = \frac{2}{\pi} \arcsin \sqrt{\alpha}.$$

Proof. See [19]. Find number of the theorem. □

Theorem B.11 (Forgery Theorem, or Support Theorem)**.** *If $f \in C([0,1])$ and $f(0) = 0$, then every neighborhood of f has positive Wiener measure. Note that by proposition B.6, this is true for neighborhoods both in the uniform and in the Skorohod topologies.*

Proof. See, for instance, [20]. This is known as the Stroock-Varadhan Support Theorem or Lévy's Forgery Theorem. □

Weak invariance principles

We will fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ for the rest of this section. Given a sequence $\xi = (\xi_1, \xi_2, \dots)$ of random variables defined on Ω, we can define their partial sums $S_n^\xi = \sum_{i=1}^n \xi_i$ and a sequence of processes $D_n^\xi \colon \Omega \to \mathbb{D}([0,1])$ given by

$$D_n^\xi(\omega)(t) = \frac{S_{\lfloor nt \rfloor}^\xi(\omega)}{\sqrt{n}},$$

obtained by appropriately scaling the partial sums of the random variables up to some iterate n.

Remarkably, for many classes of sequences ξ, their corresponding processes D_n^ξ converge in some way to the Brownian motion — we denote by an *invariance principle* a result of this form. Perhaps the best known invariance principle is the Donsker's theorem, which is a *weak* invariance principle since it asserts convergence in the weak topology.

Theorem B.12 (Donsker)**.** *If $\xi \subseteq L^2(\mathbb{P})$ is a sequence of i.i.d. variables with mean zero and variance 1, then the measures $(D_n^\xi)_* \mathbb{P}$ converge weakly to the Wiener measure. In other words, for every measurable continuous function $\varphi \colon \mathbb{D}([0,1]) \to \mathbb{R}$,*

$$\int \varphi \circ D_n^\xi \, d\mathbb{P} \to \int \varphi \, dW.$$

The motivation for looking into invariance principles is that they allow transferring certain results from Brownian motions to the sequence of partial sums S_n^ξ. In our case, the result we are interested in is the arcsine law. We know from lemma B.9 that τ^+ is continuous a.e., so weak convergence is enough to transfer its conclusions:

Lemma B.13. *Suppose $\xi \subseteq L^2(\mathbb{P})$ is such that $(D_n^\xi)_* \mathbb{P} \xrightarrow{w} W$. Then the average of the time in which the random walk S_n^ξ is above $M \in \mathbb{R}$ converges to an arcsine distribution, in the sense that*

$$\lim_{n\to\infty} \mathbb{P}\left(\left\{ \omega \in \Omega \;\middle|\; \frac{1}{n} \#\{1 \le i \le n \mid S_n^\xi(\omega) > M\} \le \alpha \right\} \right) = \frac{2}{\pi} \arcsin \sqrt{\alpha}.$$

Proof. Writing $\tau_n^+(\omega) = \frac{1}{n}\#\{1 \leq i \leq n \mid S_n^\xi(\omega) > M\}$, the left-hand side can be rewritten as $\lim_{n\to\infty}(\tau_n^+)_*\mathbb{P}([0,\alpha])$. Set $c_n = M/\sqrt{n}$, and note that

$$\tau_n^+(\omega) = \frac{1}{n}\#\left\{1 \leq i \leq n \,\middle|\, \frac{S_i^\xi(\omega)}{\sqrt{n}} - \frac{M}{\sqrt{n}} > 0\right\} = \tau^+(D_n^\xi(\omega) - c_n).$$

Since by theorem B.10 we know $Z = (\tau^+)_*W$ is equivalent to the Lebesgue measure on $[0,1]$, intervals are continuity sets for it. By the weak convergence assumption and continuity of τ^+, we know $(\tau^+ \circ D_n^\xi)_*\mathbb{P} \xrightarrow{w} Z$, and since $c_n \to 0$ (as a sequence in $\mathbb{D}([0,1])$), in fact

$$(\tau^+ \circ (D_n^\xi - c_n))_*\mathbb{P} \xrightarrow{w} Z.$$

By Portmanteau, this means

$$\lim_{n\to\infty}(\tau_n^+)_*\mathbb{P}([0,\alpha]) = \lim_{n\to\infty}(\tau^+ \circ (D_n^\xi - c_n))_*\mathbb{P}([0,\alpha]) = \frac{2}{\pi}\arcsin\sqrt{\alpha}.$$

□

Ergodic Martingale Differences

Donsker's theorem is not the most general result of its kind. In our case, we are interested in a version of the weak invariance principle for ergodic martingale difference sequences. Intuitively, these are the sequences that behave as the time-one steps of a martingale, in the sense that even if the value of some previous steps are known, the expected value of the next step is still zero.

Definition B.14. A sequence $\xi = (\xi_1, \xi_2 \dots) \in L^1(\mathbb{P})^{\mathbb{N}}$ of random variables is said to be a *martingale difference sequence* if for all n,

$$\mathbb{E}_{\mathbb{P}}(\xi_n \mid \sigma(\xi_1, \dots, \xi_{n-1})) = 0.$$

Stationarity and two-sided processes

In the previous definition we considered an one-sided sequence of random variables. As is often the case in ergodic theory, the sequences we shall be dealing with have the additional property of *stationarity*, which is a sort of time-invariance. This property will also allow us to lift one-sided sequences of variables to two-sided ones, that is, sequences with indices in $\mathbb{Z}$, due to the existence of the invertible extensions exposed in section 1.3.

Definition B.15. A sequence $\xi = (\xi_1, \xi_2 \dots)$ of random variables taking values in Y is said to be *stationary* if $\xi_*\mathbb{P}$ is an invariant measure for the left shift $\sigma\colon Y^{\mathbb{N}} \to Y^{\mathbb{N}}$. Similarly, we can define stationarity for two-sided sequences by replacing $\mathbb{N}$ by $\mathbb{Z}$ in the definition.

Definition B.16. A two-sided sequence $\xi \in L^1(\mathbb{P})^{\mathbb{Z}}$ of random variables is said to be a *martingale difference sequence* if for all n,

$$\mathbb{E}_{\mathbb{P}}(\xi_n \mid \sigma(\{\xi_i \mid i < n\})) = 0.$$

Lemma B.17. *Every stationary one-sided ergodic martingale difference sequence ξ is associated to a two-sided stationary ergodic martingale difference sequence ζ with the same marginal distributions on positive indices. That is, if $\pi\colon Y^{\mathbb{Z}} \to Y^{\mathbb{N}}$ is defined by $\pi(\dots x_{-1}, x_0, x_1, \dots) = (x_1, x_2, \dots)$, we have $(\pi \circ \zeta)_*\mathbb{P} = \xi_*\mathbb{P}$.*

Proof. write □

Invariance principle for ergodic sequences

Definition B.18. A sequence $\xi = (\xi_1, \xi_2 \dots)$ of random variables is said to be *ergodic* if $\xi_*\mathbb{P}$ is an ergodic measure for the left shift $\sigma\colon Y^{\mathbb{N}} \to Y^{\mathbb{N}}$.

Example B.19. Let (X, f, μ) be a measure-preserving system with an ergodic measure μ. Any function $\varphi\colon X \to Y$ defines a sequence of random variables

$$\xi = (\varphi \circ f^{n-1})_{n\geq 1},$$

and this sequence can also be seen as a factor map for the extension $\xi\colon (X, f, \mu) \to (Y^{\mathbb{N}}, \sigma, \xi_*\mu)$. By applying lemma 1.11, we see that the sequence ξ is an ergodic sequence.

The next result can be found in [19, theorem 18.3].

Theorem B.20. *If $\xi \subseteq L^2(\mathbb{P})$ is a stationary ergodic martingale difference sequence for which $\sigma = \|\xi_1\|_2 > 0$, then the measures $(D_n^{\xi})_*\mathbb{P} \xrightarrow{w} \sigma W$.*

Proof. the idea here is to... . For a complete proof, see the aforementioned reference. □

Reverse martingale difference sequences

In ergodic dynamics, the processes we usually care about are the values $\xi = (\varphi \circ f^{n-1})_{n\geq 1}$ of some observable $\varphi \in L^1(\mu)$ along orbits of f. This process, however, cannot possibly be a martingale difference sequence unless $\varphi \equiv 0$, because $\varphi \circ f^n$ is measurable for the σ-algebra $f^{-n}(\mathcal{F})$, which is non-increasing due to measurability of f. In other words, we always have $\mathbb{E}_\mu(\varphi \circ f^n \mid \sigma(\varphi, \dots, \varphi \circ f^{n-1})) = \varphi \circ f^n \neq 0$ in general. This motivates the following definition.

Definition B.21. A sequence $\xi = (\xi_1, \xi_2 \dots)$ of random variables is said to be a *reverse martingale difference sequence* if for all n,

$$\mathbb{E}_{\mathbb{P}}(\xi_n \mid \sigma(\xi_{n+1}, \xi_{n+2}, \dots)) = 0.$$

In the case of a stationary sequence (as is the case for $\xi = (\varphi \circ f^{n-1})_{n\geq 1}$ and $\mathbb{P} = \mu$ an invariant measure for f), results for martingale difference sequences can be stated dually for reverse martingale difference sequences. The following discussion expands on comments and text present in [19, Appendix M22 and Section 19].

Corollary B.22. *If $\xi \in L^2(\mathbb{P})^{\mathbb{N}}$ is a stationary ergodic reverse martingale difference sequence for which $\sigma = \|\xi_1\|_2 > 0$, then the measures $(D_n^{\xi})_*\mathbb{P} \xrightarrow{w} \sigma W$.*

Proof. Consider the space $\mathbb{R}^{\mathbb{N}}$ with the product σ-algebra $\mathcal{H} = \mathcal{B}^{\otimes \mathbb{N}}$ and the measure $\nu = \xi_* \mathbb{P}$. This measure is invariant by the left shift $\sigma \colon \mathbb{R}^{\mathbb{N}} \to \mathbb{R}^{\mathbb{N}}$ by the definition of stationarity. Since $\mathcal{H}$ coincides with the Borel σ-algebra, the construction of the invertible extension (see section 1.3) gives us a stationary measure $\tilde{\nu} \in \mathcal{M}(\mathbb{R}^{\mathbb{Z}})$ for the invertible shift, which we shall call $\tilde{\sigma} \colon \mathbb{R}^{\mathbb{Z}} \to \mathbb{R}^{\mathbb{Z}}$, whose projection $\pi_* \tilde{\nu}$ by $\pi \colon \mathbb{R}^{\mathbb{Z}} \to \mathbb{R}^{\mathbb{N}}$ is equal to ν. The measure $\tilde{\nu}$ is also ergodic for the invertible shift, since ν is.

We now construct an (usual) martingale difference sequence on this space. Let $\mathcal{F}_n = \pi^{-1}(\mathcal{G}_n)$.

Finish □

Re-visit this section.

Appendix C

A categorical lens

C.1 A category

A *category* C consists of a collection of *objects* Ob_C, and for each pair of objects $A, B \in \mathrm{Ob}_C$, there is a collection of *morphisms* $\mathrm{Hom}_C(A, B)$, also called arrows, that possess the following properties:

1. given $A, B, C \in \mathrm{Ob}_C$, there exists an operation

$$\circ : \mathrm{Hom}_C(A, B) \times \mathrm{Hom}_C(B, C) \longrightarrow \mathrm{Hom}_C(A, C)$$

   called categorical composition (note that it does not need to be the composition of functions). If $f : A \longrightarrow B$ and $g : B \longrightarrow C$, we write $g \circ f := \circ(f, g)$. The composition function must be associative; that is, if $h \in \mathrm{Hom}_C(C, D)$, then $(h \circ g) \circ f = h \circ (g \circ f)$;

2. for every $A \in \mathrm{Ob}_C$, there exists an identity morphism $1_A \in \mathrm{Hom}_C(A, A)$, and for all $f, g \in \mathrm{Hom}_C$ with $\mathrm{dom}(f) = \mathrm{cod}(g) = A$, it holds that $f \circ 1_A = f$ and $1_A \circ g = g$.

C.2 Functors

C.3 Natural transformations

C.4 Limits and colimits

Objects

W 33

Bibliography

[1] S. Moss and P. Perrone, "A category-theoretic proof of the ergodic decomposition theorem," *Ergodic Theory and Dynamical Systems*, pp. 1–27, 2023.

[2] A. Katok and B. Hasselblatt, *Introduction to the Modern Theory of Dynamical Systems* (Encyclopedia of Mathematics and Its Applications). Cambridge: Cambridge University Press, 1995.

[3] F. Przytycki and M. Urbański, *Conformal Fractals: Ergodic Theory Methods*, 1st ed. Cambridge University Press, 2010.

[4] M. Andersson, P. D. Carrasco, and R. Saghin, "Non-uniformly hyperbolic endomorphisms," *Compositio Mathematica*, vol. 161, no. 6, pp. 1313–1356, 2025.

[5] M. Andersson, P. D. Carrasco, and R. Saghin. "Non-uniformly hyperbolic endomorphisms." (2024), [Online]. Available: `http://arxiv.org/abs/2206.08295` (visited on 10/15/2025), pre-published.

[6] A. Bonifant and J. Milnor, "Schwarzian derivatives and cylinder maps," *American Mathematical Society*, pp. 1–24, 2008.

[7] V. Ovsienko and S. Tabachnikov, "The Schwarzian Derivative?,"

[8] B. Tortolini, *Annali di scienze matematiche e fisiche*. Tip. delle bella arti, 1855, 520 pp.

[9] R. Ures and C. H. Vásquez, "On the non-robustness of intermingled basins," *Ergodic Theory and Dynamical Systems*, vol. 38, no. 1, pp. 384–400, 2018.

[10] I. Kan, "Open sets of diffeomorphisms having two attractors, each with an everywhere dense basin," *Bulletin of the American Mathematical Society*, vol. 31, no. 1, pp. 68–74, 1994.

[11] C. Bonatti, L. J. Diaz, and M. Viana, *Dynamics Beyond Uniform Hyperbolicity* (Encyclopaedia of Mathematical Sciences). Berlin/Heidelberg: Springer-Verlag, 2005, vol. 102.

[12] D. Ruelle, "Historical behaviour in smooth dynamical systems," in *Global Analysis of Dynamical Systems*, CRC Press, 2001.

[13] F. Takens, "Orbits with historic behaviour, or non-existence of averages," *Nonlinearity*, vol. 21, no. 3, T33, 2008.

[14] F. Hofbauer and G. Keller, "Quadratic maps without asymptotic measure," *Communications in Mathematical Physics*, vol. 127, no. 2, pp. 319–337, 1990.

[15] M. Carvalho and P. Varandas. "Genericity of historic behavior for maps and flows." (2021), [Online]. Available: `http://arxiv.org/abs/2107.01200` (visited on 04/19/2024), pre-published.

[16] M. Viana, *Stochastic Dynamics of Deterministic Systems* (Colóquios Brasileiros de Matemática). Rio de Janeiro, RJ: Associação Instituto Nacional de Matemática Pura e Aplicada, 2021, 197 pp.

[17] V. I. Bogachev, *Measure Theory*. Berlin, Heidelberg: Springer, 2007.

[18] G. Botelho, *Fundamentos de Análise Funcional.*

[19] P. Billingsley, *Convergence of Probability Measures* (Wiley Series in Probability and Statistics. Probability and Statistics Section), 2nd ed. New York: Wiley, 1999, 277 pp.

[20] Y. Peres. "Answer to "The Wiener measure of an open set"," MathOverflow. (2022), [Online]. Available: `https://mathoverflow.net/a/413993/575852` (visited on 12/01/2025).