

**Analysis and development of finite
volume methods for the new generation
of cubed sphere dynamical cores for the
atmosphere**

Luan da Fonseca Santos

REPORT PRESENTED TO THE
INSTITUTE OF MATHEMATICS AND STATISTICS
OF THE UNIVERSITY OF SÃO PAULO
FOR THE DOCTOR OF SCIENCE
QUALIFYING EXAMINATION

Program: Applied Mathematics

Advisor: Prof. Pedro da Silva Peixoto

During the development of this work the author was supported by CAPES and FAPESP (grant number 20/10280-4)

São Paulo
August, 2022

**Analysis and development of finite
volume methods for the new generation
of cubed sphere dynamical cores for the
atmosphere**

Luan da Fonseca Santos

This is the original version of the
qualifying text prepared by candidate
Luan da Fonseca Santos, as submitted
to the Examining Committee.

Resumo

Luan da Fonseca Santos. **Análise e desenvolvimento de métodos de volumes finitos para modelos da nova geração da dinâmica atmosférica baseados na esfera cubada**. Exame de Qualificação (Doutorado). Instituto de Matemática e Estatística, Universidade de São Paulo, São Paulo, 2022.

[illegible]

Palavras-chave: Palavra-chave1. Palavra-chave2. Palavra-chave3.

Abstract

Luan da Fonseca Santos. **Analysis and development of finite volume methods for the new generation of cubed sphere dynamical cores for the atmosphere.** Qualifying Exam (Doctorate). Institute of Mathematics and Statistics, University of São Paulo, São Paulo, 2022.

[illegible]

Keywords: Keyword1. Keyword2. Keyword3.

Contents

Introduction	1
1 One-dimensional finite-volume methods	3
1.1 One-dimensional conservation laws in integral form	3
1.2 The finite-volume approach	7
1.3 The piecewise parabolic method	8
2 Two-dimensional finite-volume method	9
2.1 Two-dimensional consevation laws in integral form	9
2.2 Dimension splitting	9
3 Cubed-sphere grids	11
3.1 Equiangular	11
References	13
Index	15

Introduction

To be written!

Chapter 1

One-dimensional finite-volume methods

1.1 One-dimensional conservation laws in integral form

In this section, we are going to present the derivation of one-dimensional conservation laws in the integral form. The derivation presented here follows [LEVEQUE \(1990\)](#) and [LEVEQUE \(2002\)](#) closely and will be useful to fix some notation. Let us assume that x and t represent the spatial and time coordinate, respectively. Given $[x_1, x_2] \subset \mathbb{R}$, $x_1 \leq x_2$, and a time interval $[t_1, t_2] \subset]0, +\infty[$, $t_1 \leq t_2$, our aim is to describe how m state variables given by functions $q_1, \dots, q_m : \mathbb{R} \times [0, +\infty[\rightarrow \mathbb{R}$ evolve within time in the considered time interval, assuming that we have neither sinks nor sources for the mass of each state variable and also assuming that the mass flow rate is known for all the state variables.

To set the problem in more mathematical terms, let us denote by $\mathbf{q} : \mathbb{R} \times [0, \infty[\rightarrow \mathbb{R}^m$, $\mathbf{q} = \mathbf{q}(x, t)$, the vector of state variables, *i.e.*, $\mathbf{q}_k = q_k$ for $k = 1, \dots, m$. The mass of \mathbf{q} in $[x_1, x_2]$ at time t is defined by:

$$\mathbf{M}_{[x_1, x_2]}(t) := \int_{x_1}^{x_2} \mathbf{q}(x, t) dx \in \mathbb{R}^m. \quad (1.1)$$

Thus, the mass in $[x_1, x_2]$ of the k -th state variable q_k is equal to $(\mathbf{M}_{[x_1, x_2]}(t))_k$, $\forall k = 1, \dots, m$. We are going to assume the following physical constraints concerning the total mass of each state variable:

1. No mass is created;
2. No mass is destroyed.

Also, let us assume that the mass flow rate in a point x and at a time $t > 0$ is given by $\mathbf{f}(\mathbf{q}(x, t))$, where $\mathbf{f} : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is a continuously differentiable (C^1) function. This function \mathbf{f} is known as flux function. With the physical constraints that we imposed, the following

equation must hold for the mass:

$$\frac{d}{dt} \left(\int_{x_1}^{x_2} \mathbf{q}(x, t) dx \right) = \mathbf{f}(\mathbf{q}(x_1, t)) - \mathbf{f}(\mathbf{q}(x_2, t)). \quad (1.2)$$

Equation (1.2) is known as a conservation law written in integral form and tell us how the mass $\mathbf{M}_{[x_1, x_2]}(t)$ varies with time. Another integral form of the conservation law may be obtained integrating Equation (1.2) with respect to time in $[t_1, t_2]$ leading to:

$$\int_{x_1}^{x_2} \mathbf{q}(x, t_2) dx = \int_{x_1}^{x_2} \mathbf{q}(x, t_1) dx + \int_{t_1}^{t_2} \mathbf{f}(\mathbf{q}(x_1, t)) dt - \int_{t_1}^{t_2} \mathbf{f}(\mathbf{q}(x_2, t)) dt. \quad (1.3)$$

Assuming that \mathbf{q} is a C^1 function, we may write:

$$\int_{t_1}^{t_2} \frac{\partial}{\partial t} \mathbf{q}(x, t) dt = \mathbf{q}(x, t_2) - \mathbf{q}(x, t_1), \quad (1.4)$$

and

$$\int_{x_1}^{x_2} \frac{\partial}{\partial x} \mathbf{f}(\mathbf{q}(x, t)) dx = \mathbf{f}(\mathbf{q}(x_2, t)) - \mathbf{f}(\mathbf{q}(x_1, t)). \quad (1.5)$$

Replacing Equations (1.4) and (1.5) in (1.3) we get the differential form of the conservation law:

$$\int_{t_1}^{t_2} \int_{x_1}^{x_2} \left(\frac{\partial}{\partial t} \mathbf{q}(x, t) + \frac{\partial}{\partial x} \mathbf{f}(\mathbf{q}(x, t)) \right) dx dt = 0. \quad (1.6)$$

Since Equation (1.6) must hold for all x_1, x_2, t_1 and t_2 such that $[x_1, x_2] \times [t_1, t_2] \subset \mathbb{R} \times]0, +\infty[$, we obtain the differential form of the conservation law:

$$\frac{\partial}{\partial t} \mathbf{q}(x, t) + \frac{\partial}{\partial x} \mathbf{f}(\mathbf{q}(x, t)) = 0, \quad \forall (x, t) \in \mathbb{R} \times]0, +\infty[. \quad (1.7)$$

We shall assume that the eigenvalues of the Jacobian matrix of the flux function $D\mathbf{f}(\mathbf{q})$ are all real and that $D\mathbf{f}(\mathbf{q})$ is a diagonalizable matrix, $\forall \mathbf{q} \in \mathbb{R}^m$, so that Equation (1.7) is a hyperbolic partial differential equation (LEVEQUE, 1990). As we will specify latter, some initial condition will also be supposed to be known as well.

Many physical relevant equations may be written as Equation (1.7). Some examples are the Euler equations for gas dynamics, obtained when $m = 3$, and the one-dimensional shallow-water equations, obtained $m = 2$. Another relevant equations are the Burgers equation, which is obtained when $m = 1$ and $f(q) = q^2$. The Burgers equation is well known for developing shocks, even for smooth initial conditions and is a simple prototype to study shock formation. At last, the linear advection equation is another interesting example, which is obtained when $m = 1$ and $f(q(x, t)) = u(x, t)q(x, t)$, where $u(x, t)$ is a given velocity. Strictly speaking, the linear advection is not in the form given by Equation (1.7) since f depends on q but also on (x, t) . But, one may check that Equation (1.7) is still hyperbolic in this case. The linear advection equation will play a key role in this work due to its importance to development of atmospheric dynamical cores.

We say that \mathbf{q} is a strong or classical solution to the conservation law (1.7) if it is C^1 and satisfies the Equation (1.7). Applying the steps from Equation (1.3) to Equation (1.7) in a reverse order, one may check that if \mathbf{q} is a strong solution, then it satisfies the integral form (1.3) for all x_1, x_2, t_1 and t_2 such that $[x_1, x_2] \times [t_1, t_2] \subset \mathbb{R} \times]0, +\infty[$. Therefore, Equations (1.3) and (1.7) are equivalent when \mathbf{q} is C^1 . However, the problem (1.3) can be formulated to functions that are not C^1 and have discontinuities. More generally speaking, we say that $\mathbf{q} \in L^\infty(\Omega, \mathbb{R}^m)$ ¹ if it satisfies the Equation (1.3) for all x_1, x_2, t_1 and t_2 such that $[x_1, x_2] \times [t_1, t_2] \subset \mathbb{R} \times]0, +\infty[$. It can be shown that this notion of weak solution is equivalent to requiring that (LEVEQUE, 1990):

$$\int_{-\infty}^{+\infty} \int_0^{+\infty} \left(\frac{\partial}{\partial t} \phi(x, t) \mathbf{q}(x, t) + \frac{\partial}{\partial x} \phi(x, t) \mathbf{f}(\mathbf{q}(x, t)) \right) dt dx = \int_{-\infty}^{+\infty} \phi(x, 0) \mathbf{q}(x, 0) dx, \quad (1.8)$$

$\forall \phi \in C_0^1(\mathbb{R} \times [0, +\infty[)$ where $C_0^1(\mathbb{R} \times [0, +\infty[)$ denotes the set of all continuously differentiable functions with compact support in $\mathbb{R} \times [0, +\infty[$. This formulation of weak solution is more common employed on the construction of Discontinuous Galerkin methods (NAIR *et al.*, 2011).

In order to develop finite-volume methods for conservation laws, it is useful to define the vector of average values of the state variable vector \mathbf{q} in the interval $[x_1, x_2]$ at a time t by:

$$\mathbf{Q}(t) = \frac{1}{\Delta x} \int_{x_1}^{x_2} \mathbf{q}(x, t) dx \in \mathbb{R}^m, \quad (1.9)$$

where $\Delta x = x_2 - x_1$. The Equation (1.2) may be rewritten in terms of \mathbf{Q} as:

$$\frac{d}{dt} \mathbf{Q}(t) = \frac{1}{\Delta x} (\mathbf{f}(\mathbf{q}(x_1, t)) - \mathbf{f}(\mathbf{q}(x_2, t))), \quad (1.10)$$

and so is Equation (1.3):

$$\mathbf{Q}(t_2) = \mathbf{Q}(t_1) + \frac{1}{\Delta x} \left(\int_{t_1}^{t_2} \mathbf{f}(\mathbf{q}(x_1, t)) dt - \int_{t_1}^{t_2} \mathbf{f}(\mathbf{q}(x_2, t)) dt \right). \quad (1.11)$$

To move towards finite volume schemes, we will restrict our attention to a conservation law in a bounded domain of the form $\Omega = [a, b] \times [0, T]$, $a < b$, $T > 0$. However, we must impose some boundary condition. One possible way and that we will adopted in text are the periodic boundary conditions:

$$\mathbf{q}(a, t) = \mathbf{q}(b, t), \quad \forall t \in [0, T]. \quad (1.12)$$

Also, we assume that an initial condition $q_0(x) = q(x, 0)$, $q_0 \in L^\infty([a, b], \mathbb{R}^m)$, is given. Thus, we have specified a Cauchy problem. We notice that Equations (1.10) and (1.11) hold for all x_1, x_2, t_1 and t_2 such that $[x_1, x_2] \times [t_1, t_2] \subset \Omega$. So, let us discretize the domain Ω and write Equations (1.10) and (1.11) in terms of this discretization. Given a positive integer N_T , we define the time step $\Delta t = \frac{T}{N_T}$, $t_n = n\Delta t$, for $n = 0, 1, \dots, N_T$. For the spatial discretization,

¹ $L^\infty(\Omega, \mathbb{R}^m) = \{q : \Omega \rightarrow \mathbb{R}^m \text{ such that } q \text{ is bounded.}\}$

we consider a partition of $[a, b]$ given by:

$$[a, b] = \bigcup_{i=1}^N X_i, \text{ where } X_i = [x_{i-\frac{1}{2}}, x_{i+\frac{1}{2}}] \text{ and } a = x_{\frac{1}{2}} < x_{\frac{3}{2}} < \dots < x_{N-\frac{1}{2}} < x_{N+\frac{1}{2}} = b. \quad (1.13)$$

Each interval X_i is referred to as control volume. We shall use the notations $|X_i| = x_{i+\frac{1}{2}} - x_{i-\frac{1}{2}}$ and $x_i = \frac{1}{2}(x_{i+\frac{1}{2}} + x_{i-\frac{1}{2}})$, $\forall i = 1, \dots, N$, to define the control volume length and midpoint, respectively. We also denote by $\mathbf{Q}_i(t) \in \mathbb{R}^m$ as the vector of average values of state variable vector at time t in the control volume X_i , $\forall i = 1, \dots, N$. Replacing t_1, t_2, x_1 and x_2 by $t_n, t_{n+1}, x_{i-\frac{1}{2}}$ and $x_{i+\frac{1}{2}}$, respectively, in Equation 1.10, we get:

$$\frac{d}{dt} \mathbf{Q}_i(t) = \frac{1}{|X_i|} (\mathbf{f}(\mathbf{q}(x_{i-\frac{1}{2}}, t)) - \mathbf{f}(\mathbf{q}(x_{i+\frac{1}{2}}, t))), \quad \forall i = 1, \dots, N. \quad (1.14)$$

Similarly, Equation (1.11) becomes:

$$\mathbf{Q}_i(t_{n+1}) = \mathbf{Q}_i(t_n) + \frac{1}{|X_i|} \left(\int_{t_n}^{t_{n+1}} \mathbf{f}(\mathbf{q}(x_{i-\frac{1}{2}}, t)) dt - \int_{t_n}^{t_{n+1}} \mathbf{f}(\mathbf{q}(x_{i+\frac{1}{2}}, t)) dt \right), \quad (1.15)$$

$$\forall i = 1, \dots, N, \quad \forall n = 1, \dots, N_T.$$

In order to use a more compact notation, it is helpful to use the following centered difference notation:

$$\delta_x \mathbf{g}(x_i, t) = \mathbf{g}(x_{i+\frac{1}{2}}, t) - \mathbf{g}(x_{i-\frac{1}{2}}, t), \quad (1.16)$$

for an arbitrary vector valued function \mathbf{g} . Using this notation, Equations (1.14) and (1.15) lead to:

$$\frac{d}{dt} \mathbf{Q}_i(t) = -\frac{1}{|X_i|} \delta_x \mathbf{f}(\mathbf{q}(x_i, t)) \quad \forall i = 1, \dots, N, \quad (1.17)$$

and

$$\mathbf{Q}_i(t_{n+1}) = \mathbf{Q}_i(t_n) - \frac{\Delta t}{|X_i|} \delta_x \left(\frac{1}{\Delta t} \int_{t_n}^{t_{n+1}} \mathbf{f}(\mathbf{q}(x_i, t)) dt \right), \quad \forall i = 1, \dots, N, \quad \forall n = 1, \dots, N_T, \quad (1.18)$$

respectively. It is worth pointing out that we have made no approximation in Equations (1.17) and (1.18). Indeed, if \mathbf{q} satisfies Equation (1.2), $\forall [x_1, x_2] \subset [a, b]$ and $\forall t \in [0, T]$, then Equation (1.17) is just Equation (1.2) evaluated in the control volumes and written in terms of the average values \mathbf{Q} . Similarly, if \mathbf{q} satisfies Equation (1.3), $\forall [x_1, x_2] \times [t_1, t_2] \subset \Omega$, then Equation (1.18) is just Equation (1.3) evaluated in the control volumes, at the time instants t_n , and written in terms of the average values \mathbf{Q} .

Notice that in Equation (1.18) we divided and multiplied by Δt , so that we can interpret $\frac{1}{\Delta t} \int_{t_n}^{t_{n+1}} \mathbf{f}(\mathbf{q}(x_i, t)) dt$ as a mean-time average flux. This interpretation is very handy for the derivation of finite-volume schemes.

The formulations given by Equations (1.17) and (1.18) are the cornerstone of the development of finite volume methods for conservation laws. On the right-hand side of Equation (1.17), the flux function \mathbf{f} may be discretized leading to an ordinary differential

equation (ODE) that might be solved using classical ODE integrators. These methods are known as semi-discrete methods (LEVEQUE, 2002), since only the spatial coordinate is discretized. In this work we shall restrict our attention to methods based on Equation (1.18).

1.2 The finite-volume approach

We summarize the problem of the conservation law in the integral form discussed in Section 1.1 in Problem 1.

Problem 1: Given $\Omega = [a, b] \times [0, T]$, a C^1 flux function $\mathbf{f} : \mathbb{R}^m \rightarrow \mathbb{R}^m$, $m \geq 1$, we would like to find the weak solution $\mathbf{q} \in L^\infty(\Omega, \mathbb{R}^m)$ of the conservation law in the integral form:

$$\int_{x_1}^{x_2} \mathbf{q}(x, t_2) dx = \int_{x_1}^{x_2} \mathbf{q}(x, t_1) dx + \int_{t_1}^{t_2} \mathbf{f}(\mathbf{q}(x_1, t)) dt - \int_{t_1}^{t_2} \mathbf{f}(\mathbf{q}(x_2, t)) dt,$$

$\forall [x_1, x_2] \times [t_1, t_2] \subset \Omega$, given the initial condition $\mathbf{q}(x, 0) = \mathbf{q}_0(x)$, $\forall x \in [a, b]$, and assuming periodic boundary conditions, i.e., $\mathbf{q}(a, t) = \mathbf{q}(b, t)$, $\forall t \in [0, T]$.

We point out that, for Problem 1, the total mass in $[a, b]$ satisfies:

$$\mathbf{M}_{[a,b]}(t) = \mathbf{M}_{[a,b]}(0), \quad \forall t \in [0, T]. \quad (1.19)$$

This is the conservation of total mass proprierty and is highly desirable for any numerical scheme that intends to give a robust approximation of the conservation law solution.

In Section 1.1 we introduced a version of Problem 1 considering a discretization of the domain Ω . This idea is summarized in Problem 2.

Problem 2: Assume the framework of Problem 1. We consider positive integers N and N_T , a spatial discretization of $[a, b]$ given by $X_i = [x_{i-\frac{1}{2}}, x_{i+\frac{1}{2}}]$, $\forall i = 1, \dots, N$, $a = x_{\frac{1}{2}} < x_{\frac{3}{2}} < \dots < x_{N-\frac{1}{2}} < x_{N+\frac{1}{2}} = b$, a time discretization $t_n = n\Delta t$, $\Delta t = \frac{T}{N_T}$, $\forall n = 1, \dots, N_T$. Since we are in the framework of Problem 1, it follows that:

$$\mathbf{Q}_i(t_{n+1}) = \mathbf{Q}_i(t_n) - \frac{\Delta t}{|X_i|} \delta_x \left(\frac{1}{\Delta t} \int_{t_n}^{t_{n+1}} \mathbf{f}(\mathbf{q}(x_i, t)) dt \right), \quad \forall i = 1, \dots, N, \quad \forall n = 1, \dots, N_T,$$

where $\mathbf{Q}_i(t) = \frac{1}{|X_i|} \int_{x_{i-\frac{1}{2}}}^{x_{i+\frac{1}{2}}} \mathbf{q}(x, t) dx$.

Our problem now consists of finding the values $\mathbf{Q}_i(t_n)$, $\forall i = 1, \dots, N$, $\forall n = 1, \dots, N_T$, given the initial values $\mathbf{Q}_i(0)$, $\forall i = 1, \dots, N$. In other words, we would like to find the average values of \mathbf{q} in each control volume X_i at the considered time instants.

Finally, we define the one-dimensional (1D) finite-volume (FV) scheme problem as follows in Problem 3.

Problem 3 (1D FV scheme): Assume the framework defined in Problem 2. The finite-

volume approach of Problem 2 consists of a finding a scheme of the form:

$$Q_i^{n+1} = Q_i^n - \frac{\Delta t}{|X_i|} (F_{i+\frac{1}{2}}^n - F_{i-\frac{1}{2}}^n), \quad \forall i = 1, \dots, N, \quad \forall n = 1, \dots, N_T,$$

where $Q_i^n \in \mathbb{R}^m$ is intended to be an approximation of $Q_i(t_n)$ in some sense. The term $F_{i+\frac{1}{2}}^n$ approximates $\frac{1}{\Delta t} \int_{t_n}^{t_{n+1}} \mathbf{f}(\mathbf{q}(x_{i+\frac{1}{2}}, t)) dt$ and the term $F_{i-\frac{1}{2}}^n$ approximates $\frac{1}{\Delta t} \int_{t_n}^{t_{n+1}} \mathbf{f}(\mathbf{q}(x_{i-\frac{1}{2}}, t)) dt$, or, in other words, they estimate the time-averaged fluxes at the control volume X_i boundaries.

1.3 The piecewise parabolic method

Chapter 2

Two-dimensional finite-volume method

2.1 Two-dimensional consevation laws in integral form

2.2 Dimension splitting

Chapter 3

Cubed-sphere grids

3.1 Equiangular

References

- [LEVEQUE 1990] Randall J. LEVEQUE. *Numerical Methods for Conservation Laws*. Birkhäuser Basel, 1990. DOI: [10.1007/978-3-0348-5116-9](https://doi.org/10.1007/978-3-0348-5116-9) (cit. on pp. 3–5).
- [LEVEQUE 2002] Randall J. LEVEQUE. *Finite Volume Methods for Hyperbolic Problems*. Cambridge Texts in Applied Mathematics. Cambridge University Press, 2002. DOI: [10.1017/CBO9780511791253](https://doi.org/10.1017/CBO9780511791253) (cit. on pp. 3, 7).
- [NAIR *et al.* 2011] Ramachandran D. NAIR, Michael N. LEVY, and Peter H. LAURITZEN. “Emerging numerical methods for atmospheric modeling”. In: *Numerical Techniques for Global Atmospheric Models*. Ed. by Peter LAURITZEN, Christiane JABLONOWSKI, Mark TAYLOR, and Ramachandran NAIR. Berlin, Heidelberg: Springer Berlin Heidelberg, 2011, pp. 251–311. ISBN: 978-3-642-11640-7. DOI: [10.1007/978-3-642-11640-7_9](https://doi.org/10.1007/978-3-642-11640-7_9). URL: https://doi.org/10.1007/978-3-642-11640-7_9 (cit. on p. 5).

Index

A

Advection

equation, [9](#), [11](#)

C

Conservation law, [3](#)

in differential form, [4](#)

in integral form, [4](#)

Control volume, [6](#)