$$\frac{1}{1^2} + \frac{1}{2^2} + \cdots = \sum_{n=1}^\infty \frac{1}{n^s} = \zeta(2) = \frac{\pi^2}{6} = \frac{1}{6} \cdot \pi \times \pi, \qquad a_1, \ldots, a_n.$$

$$\hbar$$

$$\bigcirc \times \bigcirc \not\equiv \times \not\equiv \times \vec{\lambda} \text{ "λ "λ "λ "λ ~W 数式外では和文文字扱いのままになっている.}$$

$$\delta \text{いうλ \Rightarrow babcde}$$

$$\text{mathrm: } \delta \text{いうλ \Rightarrow babcde}$$

$$\text{mathhg: } \alpha \delta \text{いうλ \Rightarrow babcde}$$

$$\text{mathg: } \alpha \delta \text{いうλ \Rightarrow babcde}$$

$$\text{mathg: } \alpha \delta \text{いうλ \Rightarrow babcde}$$

$$\text{protected} \text{long macro:->\DLMfontsw@standard {\protect \mathrm } {\protected \ng macro:->\DLMfontsw@standard {\mathtesta } {\protect \mathrm } }$$

$$\text{protected} \text{long macro:->\DLMfontsw@standard {\mathtesta } {\mathtestb } }$$

$$\text{protected} \text{long macro:->\DLMfontsw@standard {\mathtesta } {\mathtestb } }$$

$$\text{protected} \text{long macro:->\DLMfontsw@standard {\mathtesta } {\mathtestb } }$$

$$\text{protected} \text{long macro:->\DLMfontsw@standard } \text{mathtestb } {\mathtestb} \text{ }$$

$$\text{protected} \text{long macro:->\DLMfontsw@standard } \text{mathtestb} \text{ } \text{ }$$

$$\text{protected} \text{ macro:->\Udelimiter } \text{ 4} \text{ symoperators "0230A} \text{ scan_stop: } \text{ ltjsetmathletter}$$

 $\lceil x \rceil + \lfloor x \rfloor + \hat{a} - \sqrt{a} - \lceil \hbar \rceil$

}

{"0230A}