## SHC 798 Assignment 1, 2025

#### Richard Lubega

2025-07-14

## SHC 798 Assignment 1, 2025

#### Part 3: Simple regression

Question 4

Validating the Hand Calculations

```
age <- c(5, 10, 15, 20, 25, 30)

strength <- c(48, 42, 37, 30, 27, 21)


a_beams <- data.frame(age, strength)
```


```
par(mfrow = c(2,2))
hist(a_beams$age, main = "Histogram for Age", xlab = "Age (years)", col = "lightblue")
hist(a_beams$strength, main = "Histogram for Strength", xlab = "Strength (MPa)", col = "lightgreen")

#Applying natural logarithms
a.age <- log(a_beams$age)
a.strength <- log(a_beams$strength)
hist(a.age, main = "Histogram for log(Age)", xlab = "log(Age)", col = "skyblue")
hist(a.strength, main = "Histogram for log(Strength)", xlab = "log(Strength)", col = "lightpink")</pre>
```




## Histogram for Strength






### **Histogram for log(Age)**

## **Histogram for log(Strength)**





(a) and (b)

```
par(mfrow = c(1,1))
```

```
log_beam <- lm(log(strength) ~ log(age), data = a_beams)
summary(log_beam)</pre>
```

(c)

```
##
## Call:
## lm(formula = log(strength) ~ log(age), data = a_beams)
##
## Residuals:
##
                2
                        3
                                       5
  ##
## Coefficients:
            Estimate Std. Error t value Pr(>|t|)
##
             4.66776
                       0.21304 21.911 2.57e-05 ***
## (Intercept)
## log(age)
             -0.43393
                       0.07683 -5.648 0.00484 **
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
##
```

```
## Residual standard error: 0.1138 on 4 degrees of freedom
## Multiple R-squared: 0.8886, Adjusted R-squared: 0.8607
## F-statistic: 31.9 on 1 and 4 DF, p-value: 0.004841

summary(log_beam)$r.squared

(d)

## [1] 0.8885725

summary(log_beam)$coefficients["log(age)", "Pr(>|t|)"]


(e)

## [1] 0.00484069

#####(f)

# Tukey-Anscombe Plot
plot(log_beam$fitted.values, log_beam$residuals, xlab="Fitted", ylab="Residuals", pch = 19, col = "blue title("Residuals vs. Fitted Values") +
lines(loess.smooth(log_beam$fitted.values, log_beam$residuals),col="red") +
abline(h=0, col="grey")
```

# Residuals vs. Fitted Values



## integer(0)