

MATH 350 (Fall 2024): Final Exam Review Session with Luc :)

Throughout these problems, let A and B be commutative rings with $1 \neq 0$.

Problem 1.

- (a) Sort the following by inclusion: fields, Euclidean domains, PIDs, UFDs, and integral domains.
- (b) Give an example of a Euclidean domain that isn't a field and a UFD that isn't a PID.¹
- (c) If A is an integral domain and B is a subring of A , is B also an integral domain?

Problem 2.

Let $\varphi : A \rightarrow B$ be a ring homomorphism.

- (a) Show that the restriction of φ to A^\times , denoted $\varphi|_{A^\times}$, is a group homomorphism from A^\times to B^\times .
- (b) Deduce that if A and B are isomorphic as rings, then A^\times and B^\times are isomorphic as groups.
- (c) Conclude that \mathbb{R} and \mathbb{C} are not isomorphic as rings.²

Problem 3.

Let $\varphi : A \rightarrow B$ be a ring homomorphism. Let J be a subset of B , and let $I := \varphi^{-1}(J)$. Answer the following true-or-false questions with either a proof or a counterexample:

- (a) If A is a field, then φ is injective.
- (b) If J is a subring of B , then I is a subring of A .
- (c) If J is a subring of B , then $\ker \varphi$ is an ideal in I .
- (d) If J is an ideal in B , then I is an ideal in A .
- (e) If J is a prime ideal in B , then I is a prime ideal in A .
- (f) If J is an ideal in B and B/J is an integral domain, then A/I is also an integral domain.
- (g) If J is a maximal ideal in B , then I is a maximal ideal in A .
- (h) If J is an ideal in B and B/J is a field, then A/I is also a field.
- (i) Write $B\varphi(I) := \{bj \mid b \in B, j \in \varphi(I)\}$. If J is an ideal in B , then $B\varphi(I) \subset J$.
- (j) If J is an ideal in B , then $J \subset B\varphi(I)$.

Problem 4.

Let C be the ring of Cauchy sequences³ of rational numbers with respect to the Euclidean metric $d(x, y) = |x - y|$, and let I be the ideal of C whose elements converge to 0.

- (a) Convince yourself that C is a commutative ring. (No need for a proof here—this is just to make sure you remember the ring axioms.)
- (b) Verify that I is an ideal in C . (Hint: Cauchy sequences are bounded.)
- (c) Prove that C/I and \mathbb{R} are isomorphic as rings.⁴ (Hint: Since \mathbb{Q} is dense in \mathbb{R} and \mathbb{R} is a complete metric space⁵ with respect to the Euclidean metric, there is a natural surjection from C to \mathbb{R} .)
- (d) Let A be a ring with $1 \neq 0$, and let \mathbb{F} be a field. Show that if A and \mathbb{F} are isomorphic as rings, then A is a field.
- (e) Deduce that I is a maximal ideal of C .

¹For an example of an integral domain that isn't a UFD, see Problem 5. For an example of a PID that isn't a Euclidean domain, see p. 282 of Dummit and Foote, but it isn't anything you'll need to know for the final.

²Nevertheless, \mathbb{R} and \mathbb{C} are isomorphic as groups. This is because they're isomorphic as vector spaces over \mathbb{Q} .

³A sequence (a_n) is called *Cauchy* if, for all $\varepsilon > 0$, there exists some $N \in \mathbb{N}$ such that $d(a_m - a_n) < \varepsilon$ for all $m, n > N$.

⁴This is Cantor's construction of the real numbers. Note how different it is from the construction by Dedekind cuts!

⁵A metric space X is called *complete* if every Cauchy sequence of elements of X converges to an element of X . A subset $Y \subset X$ is called *dense* in X if, for all $x \in X$, there exists a sequence (y_n) in Y that converges to x .

Problem 5. Consider the commutative ring $R := \mathbb{Z}[\sqrt{-5}]$ with norm $N : R \rightarrow \mathbb{Z}_{\geq 0}$ defined⁶ by $N(a + b\sqrt{-5}) = a^2 + 5b^2$. Note that for all $\alpha, \beta \in R$, we have $N(\alpha\beta) = N(\alpha)N(\beta)$.

- (a) Is R an integral domain? Why or why not?
- (b) Show that if $\lambda \in R$, then $\lambda \in R^\times$ if and only if $N(\lambda) = 1$.
- (c) Show that if $\lambda \in R$ and $N(\lambda) = 9$, then λ is irreducible.
- (d) By considering the equalities $(2 + \sqrt{-5})(2 - \sqrt{-5}) = 9 = 3(3)$, conclude that R is not a UFD.
- (e) Also, conclude that (3) , the ideal in R generated by 3, is not a prime ideal.

Problem 6. Let A be an integral domain with $1 \neq 0$, let $\alpha \in A$, and let (α) be the ideal generated by α . Answer the following true-or-false questions:

- (a) (0) is a prime ideal in A .
- (b) If (α) is maximal, then (α) is prime.
- (c) If (α) is prime, then (α) is maximal.
- (d) If (α) is prime and $\alpha \neq 0$, then (α) is maximal.
- (e) If (α) is prime, $\alpha \neq 0$, and A is a PID, then (α) is maximal. (*Hint: Check the following item.*)
- (f) If (α) is prime, then α is irreducible.
- (g) If α is irreducible, then (α) is prime. (*Hint: Check the previous problem.*)
- (h) If α is irreducible and A is a PID, then (α) is both prime and maximal.

Problem 7. Let's do a little number theory!⁷ Let $\varphi : \mathbb{N} \rightarrow \mathbb{N}$ be the totient function from HW10 #6.

- (a) Let m and n be relatively prime integers. Show that $m\mathbb{Z} \cap n\mathbb{Z} = mn\mathbb{Z}$ and $m\mathbb{Z} + n\mathbb{Z} = \mathbb{Z}$. (*Hint: For any nonzero integers a, b , there exist $x, y \in \mathbb{Z}$ such that $\gcd(a, b) = xa + yb$.*⁸)
- (b) Let k_1, \dots, k_n be pairwise relatively prime integers, and let $K := \prod_{i=1}^n k_i$ be their product. Prove that there exists a ring isomorphism

$$\mathbb{Z}/K \cong \mathbb{Z}/k_1 \times \mathbb{Z}/k_2 \times \cdots \times \mathbb{Z}/k_n.$$

(*Hint: Use induction on n and Sun Zi's theorem, which you proved in HW10 #2 as the "Chinese remainder theorem."*)

- (c) Give an example showing that (b) is false when the k_i 's aren't relatively prime.
- (d) Deduce that if $n = \prod_{i=1}^k p_i^{\alpha_i}$ is the prime factorization of n , then there exists a group isomorphism

$$(\mathbb{Z}/n)^\times \cong (\mathbb{Z}/p_1^{\alpha_1})^\times \times (\mathbb{Z}/p_2^{\alpha_2})^\times \times \cdots \times (\mathbb{Z}/p_k^{\alpha_k})^\times.$$

(*Hint: Use problem 1(b) on this worksheet.*)

- (e) Let $n \in \mathbb{Z}$. Show that $|(\mathbb{Z}/n)^\times| = \varphi(n)$. (*Hint: HW10 #6(a) and HW9 #6(c) might help.*)
- (f) Deduce that if $n = \prod_{i=1}^k p_i^{\alpha_i}$ is the prime factorization of n , then $\varphi(n) = \prod_{i=1}^k \varphi(p_i^{\alpha_i})$.⁹

⁶This is actually the square of the *modulus* function $|\cdot| : \mathbb{C} \rightarrow \mathbb{R}$.

⁷This problem is actually closely related to the *classification of finitely generated abelian groups*, which you should look up if you plan on taking Math 370 (and you should, because I'll be one of the ULAs for it next semester :). I suggest Section 5.2 of Dummit and Foote as a reference.

⁸This is actually a consequence of the fact that \mathbb{Z} is a Euclidean domain. I highly suggest referring to p. 5 of Dummit and Foote for details!

⁹In other words, the totient function is a multiplicative function!

Problem 8. In class, you showed that if \mathbb{F} is a field, then the polynomial ring $\mathbb{F}[x]$ is a Euclidean domain. Prove a strengthened version of the converse: if A is a commutative ring and $A[x]$ is a PID, then A is a field. (*Hint: Check Problems 1(c) and 6(e) from earlier.*)

These next few problems (along with Problem 4(c) from earlier) use the *first isomorphism theorem for rings*, which you proved in HW9 #2.

Problem 9. Let $\mathbb{C}[x, y, z]$ be the ring of polynomials in three variables with complex coefficients, and let $(xz - y)$ be the ideal generated by $xz - y$. Show there exists a ring isomorphism

$$\mathbb{C}[x, y, z]/(xz - y) \cong \mathbb{C}[x, z].$$

Problem 10. In this problem, we prove the *second isomorphism theorem for rings*. Let S be a subring of A , and let I be an ideal in A .

- (a) Show that $S + I$ is a subring of A .
- (b) Show that $S \cap I$ is an ideal in S .
- (c) Prove that there exists a ring isomorphism

$$S/(S \cap I) \cong (S + I)/I.$$

Problem 11. Now, we prove the *third isomorphism theorem for rings*. Let $I \subset J$ be ideals in A .

- (a) Show that J/I is an ideal of A/I .
- (b) Prove that there exists a ring isomorphism

$$(A/I)/(J/I) \cong A/J.$$

- (c) Deduce that J is prime (resp. maximal) in A if and only if J/I is prime (resp. maximal) in A/I .

Problem 12. *This problem is just for fun!* How many continuous ring automorphisms are there from \mathbb{R} to \mathbb{R} ? from \mathbb{C} to \mathbb{C} ? (*Hint 1: How many ring homomorphisms are there from \mathbb{Q} to \mathbb{R} ? from \mathbb{Q} to \mathbb{C} ?*) (*Hint 2: \mathbb{Q} is a dense subset of \mathbb{R} , and $\mathbb{Q}[i]$ is a dense subset of \mathbb{C} .*)

You're doing great! Good luck on the final—you've got this! :)