

MODULAR COMPACTIFICATIONS OF $\mathcal{M}_{2,n}$ I

LUCA BATTISTELLA

ABSTRACT. We study the geometry of Gorenstein curve singularities of genus two, and of their semistable tails. There are two families of such singularities, corresponding to either Weierstrass or conjugate points on a curve of genus two. For every $1 \leq m < n$, a stability condition - using one of the markings as a reference point, and therefore not \mathfrak{S}_n -symmetric - defines proper Deligne-Mumford stacks $\overline{\mathcal{M}}_{2,n}^{(m)}$ containing the locus of smooth curves as a dense open.

CONTENTS

1. Introduction	1
2. Gorenstein curve singularities of genus two	6
3. Tangent sheaf, crimping space, and automorphisms	10
4. Dualising line bundle and semistable tails	17
5. The new moduli functors	29
References	38

1. INTRODUCTION

1.1. From the Deligne-Mumford space to the Hassett-Keel program.

One of the most fascinating and influential results of modern algebraic geometry is the construction of a modular compactification of the stack of smooth pointed curves $\mathcal{M}_{g,n}$, due to P. Deligne, D. Mumford, and F. Knudsen [DM69, Knu83], via the introduction of *stable* pointed curves.

Definition 1.1. A connected, reduced, complete curve C with distinct markings (p_1, \dots, p_n) lying in the smooth locus of C is *stable* if:

- (1) C admits only nodes as singularities;
- (2) every rational component of C has at least three special points (markings or nodes), and every elliptic component has at least one.

Theorem 1.2. Assume $2g - 2 + n > 0$. The moduli stack of stable pointed curves $\overline{\mathcal{M}}_{g,n}$ is a smooth and proper connected Deligne-Mumford stack over $\mathrm{Spec}(\mathbb{Z})$, with projective coarse moduli space $\overline{\mathbf{M}}_{g,n}$. The boundary $\overline{\mathcal{M}}_{g,n} \setminus \mathcal{M}_{g,n}$, representing nodal curves, is a normal crossing divisor.

On one hand, the Deligne-Mumford compactification has nearly every desirable property one could hope for; on the other, it is certainly not the unique modular compactification of $\mathcal{M}_{g,n}$. Classifying all of them is a difficult problem, which has so far only found a partial, beautiful answer in work of D.I. Smyth [Smy13].

Even though the existence of $\overline{\mathcal{M}}_{g,n}$ can be deduced from nowadays standard general theory of stacks [KM97], this moduli space was first constructed as a quotient, prompting the development of the powerful techniques of Geometric Invariant Theory [Gie82, MFK94, BS08]. The study of alternative compactifications of $\mathcal{M}_{g,n}$ is motivated as well by an interest in the birational geometry of $\overline{\mathcal{M}}_{g,n}$, and it is not by chance that the first steps in this direction were moved from a GIT perspective - by changing the invariant theory problem or the stability condition under consideration, and then realising that the resulting quotient still enjoyed a modular interpretation [Sch91, Has05]. The consequent program that goes under the name of B. Hassett and S. Keel aims to describe all the quotients arising in this way, and to determine whether every step of a minimal model program for $\overline{\mathcal{M}}_{g,n}$ can be provided with a modular interpretation in terms of curves with worse than nodal singularities [CTV18, CTV19]. Since the early stages of this program, it has developed into a fascinating playground for implementing ideas that originated from (v)GIT into a general structure theory of Artin stacks [AK16, AFSvdW17, AFS17a, AFS17b]. See for instance [Mor11, FS13] for more comprehensive and instructive accounts.

Only few steps of the Hassett-Keel program have been carried out in full generality. On the other hand, the program has been completed to a larger extent in low genus: with the introduction of weighted pointed curves [Has03] in genus zero, and with Smyth's pioneering work in genus one [Smy11a, Smy11b, Smy18], extending in a direction an earlier work of D. Schubert. The philosophy is that an alternative compactification is defined by allowing a reasonably larger class of curve singularities (*local condition*), identifying their (semi)stable models, and disallowing the latter by imposing a stronger stability condition (*global condition*, usually combinatorial); this ensures that the resulting moduli problem be again separated and universally closed, by the valuative criterion.

A useful notion in this respect is that of the *genus* of an isolated curve singularity: let (C, x) be (the germ of) a reduced curve over an algebraically closed field \mathbf{k} at its unique singular point x , with normalisation $\nu: \tilde{C} \rightarrow C$.

Definition 1.3. Let C have m branches at x , and δ be the \mathbf{k} -dimension of $\nu_*\mathcal{O}_{\tilde{C}}/\mathcal{O}_C$, a skyscraper sheaf supported at x . The genus of (C, x) is then:

$$g = \delta - m + 1.$$

It can be thought of as the number of conditions that a function must satisfy in order to descend from the seminormalisation (the initial object in the category of universal homeomorphism $C' \rightarrow C$, see [Sta19, Tag 0EUS], or a curve with the same topological space as C and an ordinary m -fold point at x) to C . The node, for example, has genus zero, since it coincides with its own seminormalisation. It

is a notion that behaves well in families, in the sense that a complete, reduced curve C with only one singularity at x , of δ -invariant δ , and normalisation a union of m copies of \mathbb{P}^1 , will appear in a family of curves of arithmetic genus g .

Smyth found that, for every fixed number of branches m , there is a unique germ of Gorenstein singularity of genus one up to isomorphism, namely:

$m = 1$: the cusp, $V(y^2 - x^3) \subseteq \mathbb{A}_{x,y}^2$;

$m = 2$: the tacnode, $V(y^2 - yx^2) \subseteq \mathbb{A}_{x,y}^2$;

$m \geq 3$: the union of m general lines through the origin of \mathbb{A}^{m-1} .

Singularities of this kind, with up to m branches, together with nodes, form a deformation-open class of singularities. Furthermore, the elliptic m -fold point can be obtained by contracting a smooth elliptic curve with m rational tails in a one-parameter smoothing, and, roughly speaking, all stable models look like this.

Definition 1.4. [Smy11a] A connected, reduced, complete curve C of arithmetic genus one with smooth distinct markings (p_1, \dots, p_n) is m -stable, $1 \leq m < n$, if:

- (1) it has only nodes and elliptic l -fold points, $l \leq m$, as singularities;
- (2) for every connected subcurve $E \subseteq C$ of arithmetic genus one, its *level*:
 $|E \cap \overline{C} \setminus \overline{E}| + |\{i: p_i \in E\}|$ is strictly larger than m ;
- (3) $H^0(C, \Omega_C^\vee(-\sum_i p_i)) = 0$.

The latter can be taken for a decency condition on the moduli stack. The second one, instead, is essential in guaranteeing the uniqueness of m -stable limits, as per the discussion above. Smyth's main result is the following.

Theorem 1.5. [Smy11a, Smy11b] *The moduli stack of m -stable curves $\overline{\mathcal{M}}_{1,n}(m)$ is a proper irreducible Deligne-Mumford stack over $\mathrm{Spec} \mathbb{Z}[1/6]$. It is not smooth for $m \geq 6$. The coarse moduli spaces $\overline{\mathbf{M}}_{1,n}(m)$ arise as birational models of $\overline{\mathbf{M}}_{1,n}$ for the big line bundles $D(s) = s\lambda + \psi - \Delta$, where ψ is the sum of the ψ -classes, Δ is a boundary class, and s and m are explicitly related.*

1.2. Experimenting on a genus two tale. The following is classically known.

Fact. *There are two unibranch singularities of genus two, namely the ramphoid cusp or A_4 -singularity $V(y^2 - x^5) \subseteq \mathbb{A}_{x,y}^2$, and the singularity $\mathrm{Spec}(\mathbf{k}[t^3, t^4, t^5])$ in 3-space. The former is Gorenstein, and its stable model is a Weierstrass tail (a genus two curve attached to a rational one at a Weierstrass point), while the latter is not Gorenstein, and its stable model is a non-Weierstrass genus two tail.*

See Lemma 2.1 and Proposition 4.2 below. Recall that every smooth curve of genus two is hyperelliptic, i.e. it can be realised as a $2 : 1$ cover of \mathbb{P}^1 , in a unique way up to projectivities. The deck transformation is called the hyperelliptic involution σ ; ramification points (fixed points of σ) are called Weierstrass, while in general $\{p, \sigma(p)\}$ are called conjugate points. See Remark 4.1 below.

Let us try to apply Smyth's approach to genus two curves, starting from $\overline{\mathcal{M}}_{2,2}^{(1)}$. If we are going to require the level of a genus two subcurve to be at least 2, it

seems we need to include non-Gorenstein singularities in order to keep our moduli space proper. This might lead us into trouble; for example, the (log) dualising line bundle is classically used to construct canonical polarisations on stable curves, which in turn are essential in the proof that $\overline{\mathcal{M}}_{g,n}$ is an algebraic stack (or in the construction of $\overline{\mathcal{M}}_{g,n}$). Yet, there is a way around the awful singularity $\mathbf{k}[t^3, t^4, t^5]$.

Fact. *The A_5 -singularity $V(y^2 - yx^3) \subseteq \mathbb{A}_{x,y}^2$ is a Gorenstein singularity of genus two with two branches. Its stable model is a genus two bridge, with the two nodes being conjugate points. A marked union of two \mathbb{P}^1 along an A_5 -singularity has no non-trivial automorphisms as soon as one of the two branches contains at least two markings.*

See Proposition 2.2 and Corollary 3.6 below. Let us go back to $\overline{\mathcal{M}}_{2,2}$. Suppose there is a rational tail R on which the two markings lie, so as to contravene the requirement that the level of any genus two subcurve is at least 2. If R is attached to a Weierstrass point of the minimal subcurve of genus two Z , we may simply contract the latter (in a one-parameter smoothing), thus producing an irreducible ramphoid cusp with two markings. If instead R is attached to a non-Weierstrass point q_1 of Z , we may blow up the one-parameter family at the conjugate point $\sigma(q_1)$ on the central fiber, and then contract Z to get a *dangling* A_5 -singularity (meaning that one of the branches is unmarked), which nonetheless has trivial automorphism group.

To complete the picture, note that, in order to fix a deformation-open class of singularities, we need to allow cusps and tacnodes as well.

Fact. *The singularities appearing in the miniversal family of an A_m -singularity are all and only the A_l -singularities with $l \leq m$.*

See Theorem 2.5 below for a more general statement - valid for all ADE singularities - due to A. Grothendieck. For the sake of separatedness, we should at the same time require that the level of a genus one subcurve be at least 3. Note that hybrid situation may emerge: e.g. an elliptic curve with a cusp, or an irreducible tacnode; what is worth mentioning is that, since we want to allow a tacnode and a cusp sharing a branch, the level condition for genus one subcurves should be applied only to *nodally attached* subcurves. In particular, there cannot be any nodally attached subcurve of genus one. Besides, in the last example we need to break the \mathfrak{S}_2 -symmetry (relabelling the markings) in order to have a unique limit: we do so by declaring that p_1 lie on the cuspidal branch. See Figure 1.

We now have a plausible definition of $\overline{\mathcal{M}}_{2,2}^{(1)}$.

Definition 1.6. A connected, reduced, complete curve of arithmetic genus two C over $\mathbf{k} = \bar{\mathbf{k}}$, with smooth and disjoint markings p_1, p_2 , is 1-stable if:

- (1) C has only A_1- , \dots , A_4- and dangling A_5- singularities.
- (2) Every subcurve of genus two has level 2 at least.
- (3) There is no nodally attached subcurve of genus one; if there are two genus one singularities sharing a branch, p_1 is connected to the one of lowest level.




FIGURE 1. Examples of 2-pointed stable curves and their 1-stable limits.

The main result of the paper is that $\overline{\mathcal{M}}_{2,2}^{(1)}$ is a proper Deligne-Mumford stack, and the generalisations to an arbitrary number of markings that we are going to discuss in the next subsection.

Let us note in passing that the birational map $\overline{\mathcal{M}}_{2,2} \dashrightarrow \overline{\mathcal{M}}_{2,2}^{(1)}$ is not defined everywhere. The reason boils down to the following:

Fact. *There is only one isomorphism class of 2-pointed curves whose normalisation is $(\mathbb{P}^1, q_1) \sqcup (\mathbb{P}^1, q_2, p_1, p_2)$ and having an A_5 -singularity at $q_1 = q_2$. On the other hand, the moduli space of 2-pointed irreducible curves of geometric genus zero and having an A_4 -singularity is isomorphic to \mathbb{A}^1 .*

The second statement can be motivated as follows: the pointed normalisation of such a curve is $(\mathbb{P}^1, q, p_1, p_2)$, which has neither automorphisms, nor deformations. To produce an A_4 -singularity at q we may first collapse a non-zero tangent vector at q (for which we have no choice), producing a cusp, and then collapse a line in the tangent space to the cusp, different from the tangent cone ℓ of the cusp itself (the moduli space is therefore $\mathbb{P}^1 \setminus \{\ell\} = \mathbb{A}^1$). See Lemma 3.2 and the discussion thereafter.

Let $\mathcal{W} \subseteq \overline{\mathcal{M}}_{2,2}$ the codimension 2 locus of Weierstrass tails. From the fact above, we conjecture that there is a factorisation:

$$\begin{array}{ccc} & \text{Bl}_{\mathcal{W}}(\overline{\mathcal{M}}_{2,2}) & \\ \swarrow & & \searrow \\ \overline{\mathcal{M}}_{2,2} & \dashrightarrow & \overline{\mathcal{M}}_{2,2}^{(1)} \end{array}$$

We plan to come back to this point in future work.

1.3. Outline of results and plan of the paper. Here are the main novelties of our construction. It is not a semistable compactification (see [Smy13, Definition 1.2]), in that we allow dangling singularities of type *II* (see ??? below for a definition), whose normalisation has a one-pointed rational component. The necessity to do so was prefigured in [AFS16] already.

The dualising line bundle of a minimal (with no separating nodes) Gorenstein curve of genus one is trivial [Smy11a, Lemma 3.3]. This plays a fundamental role in the study of semistable tails, as well as in the boundedness of the moduli functor. On the other hand the dualising line bundle of a minimal Gorenstein curve of genus two Z has degree two. For this reason we had to desymmetrise the problem and use the first marking as a reference point, in order to write $\omega_Z = \mathcal{O}_Z(q_1 + \bar{q}_1)$ whenever needed, where q_1 is the point of Z closest to p_1 , and \bar{q}_1 is conjugate to q_1 (it is not always defined uniquely).

1.4. Future directions.

- (1) Resolve the indeterminacy of the rational map $\overline{\mathcal{M}}_{2,n}(m_1) \dashrightarrow \overline{\mathcal{M}}_{2,n}(m_2)$; we expect the construction to rely on a semistable compactification of the crimping spaces of the genus two singularities, as in [vdW10, §1.10] and [Smy18]. It would be interesting to put this work in the context of the Hassett-Keel program, as in [Smy11b]. More generally, a question outstanding to our knowledge is whether the whole program fits in the theoretical framework developed in [Hal14].
- (2) Our main motivation to start this project arose from Gromov-Witten theory. In genus one, the link between reduced Gromov-Witten invariants (see for example [VZ08, Zin09, LZ09]) and maps from singular curves (see [Vis12]) was partially uncovered in [BCM18], and brought in plain light in [RSW17a, RSW17b]. With F. Carocci we are investigating whether similar techniques may serve to desingularise the main component of the space of genus two maps to projective space. If this is the case, they will provide a clear definition of reduced invariants, and hopefully ease the access to comparison (standard vs. reduced) results. We expect there will be a(n iso)morphism to the modular blow-up constructed in [HLN12]. This might lead to a mathematical definition of all-genera Gopakumar-Vafa invariants, one day.

1.5. Acknowledgements.

2. GORENSTEIN CURVE SINGULARITIES OF GENUS TWO

In this and the next sections we work over an algebraically closed field of characteristic different from 2, 3, 5. We provide an algebraic classification of the (complete) local rings of Gorenstein curve singularities of genus two.

Let (C, q) be the germ of a curve singularity, and let (R, \mathfrak{m}) denote $(\hat{\mathcal{O}}_{C,q}, \mathfrak{m}_q)$, with normalisation $(\tilde{R}, \tilde{\mathfrak{m}}) \simeq (\mathbf{k}[[t_1]] \oplus \dots \oplus \mathbf{k}[[t_m]], \langle t_1, \dots, t_m \rangle)$. Here m is the number of branches of C at q . Recall from Definition 1.3 that the genus is:

$$g = \delta - m + 1,$$

so, for genus two, $\delta = m + 1$. Following [Smy11a, Appendix A], we consider \tilde{R}/R as a \mathbb{Z} -graded module with:

$$(\tilde{R}/R)_i := \tilde{\mathfrak{m}}^i / (\tilde{\mathfrak{m}}^i \cap R) + \tilde{\mathfrak{m}}^{i+1};$$

furthermore, Smyth (*loc.cit.*) notices that:

- (1) $m + 1 = \delta(p) = \sum_{i \geq 0} \dim_{\mathbf{k}}(\tilde{R}/R)_i$;
- (2) $2 = g = \sum_{i \geq 1} \dim_{\mathbf{k}}(\tilde{R}/R)_i$;
- (3) if $(\tilde{R}/R)_i = (\tilde{R}/R)_j = 0$ then $(\tilde{R}/R)_{i+j} = 0$.

We will also find the following observations to be useful:

- (4) $\sum_{i \geq k} (\tilde{R}/R)_i$ is a grading of $\tilde{\mathfrak{m}}^k / (\tilde{\mathfrak{m}}^k \cap R)$;
- (5) there is an exact sequence of $R/\mathfrak{m} = \mathbf{k}$ -modules:

$$0 \rightarrow A_k := \frac{\tilde{\mathfrak{m}}^k \cap R}{\tilde{\mathfrak{m}}^{k+1} \cap R} \rightarrow \frac{\tilde{\mathfrak{m}}^k}{\tilde{\mathfrak{m}}^{k+1}} \rightarrow \left(\tilde{R}/R \right)_k \rightarrow 0$$

Lemma 2.1. *There are two unibranch curve singularities of genus two; only one of them is Gorenstein, the A_4 -singularity or ramphoid cusp: $V(y^2 - x^5) \subseteq \mathbb{A}_{x,y}^2$.*

Proof. In the unibranch case $\dim_{\mathbf{k}}(\tilde{R}/R)_1 \leq 1$, hence equality must hold (by observation (3) above). We are left with two cases:

- Either $\dim_{\mathbf{k}}(\tilde{R}/R)_2 = 1$ and $\dim_{\mathbf{k}}(\tilde{R}/R)_i = 0$ for all $i \geq 3$: in this case $\tilde{\mathfrak{m}}^3 \subseteq \mathfrak{m}$ by observation (4). From (5) we see that $\tilde{\mathfrak{m}}^3 = \mathfrak{m}$, hence $R \simeq \mathbf{k}[[t^3, t^4, t^5]]$, a spatial non-Gorenstein singularity.
- Or $\dim_{\mathbf{k}}(\tilde{R}/R)_3 = 1$ and $\dim_{\mathbf{k}}(\tilde{R}/R)_i = 0$ for $i = 2$ and for all $i \geq 4$: in this case $\tilde{\mathfrak{m}}^4 \subseteq \mathfrak{m}$ by observation (4). On the other hand from $\dim_{\mathbf{k}}(\tilde{\mathfrak{m}}^2 \cap R / \tilde{\mathfrak{m}}^3 \cap R) = 1$ we deduce that there is a generator of degree 2, and from $\dim_{\mathbf{k}}(\tilde{\mathfrak{m}}^3 \cap R / \tilde{\mathfrak{m}}^4 \cap R) = 0$ there is none of degree 3. We may write the generator as $x = t^2 + ct^3$, and $\mathfrak{m} = \langle x \rangle + \tilde{\mathfrak{m}}^4$. Up to a coordinate change (i.e. automorphism of $\mathbf{k}[[t]]$), we may take $x = t^2$, and

$$\mathfrak{m}/\mathfrak{m}^2 = \langle t^2, t^5 \rangle,$$

so $R \simeq \mathbf{k}[[x, y]]/(x^5 - y^2)$, as anticipated. □

From now on, we are going to study only Gorenstein singularities. With notation as above, let $I = (R : \tilde{R}) = \text{Ann}_R(\tilde{R}/R)$ be the conductor ideal of the singularity. Recall (see e.g. [Hun99, Theorem 6.4]) that (C, q) is Gorenstein iff :

$$\dim_{\mathbf{k}}(R/I) = \dim_{\mathbf{k}}(\tilde{R}/R)(= \delta).$$

Recall from [Ste96, Definition 2-1] that a curve singularity (C, q) is *decomposable* if C is the union of two curves C_1 and C_2 that lie in distinct smooth spaces intersecting each other transversely in q . Given a parametrisation $x_i = x_i(t_1, \dots, t_m)$, $i = 1, \dots, n$, this means that we can find a partition $S \sqcup S' = \{1, \dots, m\}$ such that x_i only depends on $t_s, s \in S$, or $s \in S'$, for all i . Recall from

[AFS16, Proposition 2.1] that, aside from the node, Gorenstein singularities are never decomposable.

Proposition 2.2. *For every fixed integer $m \geq 2$, there are exactly two Gorenstein curve singularities of genus two with m branches.*

Proof. We wish to find a basis for $\mathfrak{m}/\mathfrak{m}^2$, because a map of complete local rings that is surjective on cotangent spaces is surjective. From observation (3) again, we find three possibilities for the vector (d_1, d_2, d_3) , $d_k = \dim_{\mathbf{k}}(\tilde{R}/R)_k$.

Case $(2, 0, 0)$. We see that $\tilde{\mathfrak{m}}^2 \subseteq I$, so (by the Gorenstein assumption) $m+1 = \delta = \dim_{\mathbf{k}}(R/I) \leq \dim_{\mathbf{k}}(R/\tilde{\mathfrak{m}}^2) = \dim_{\mathbf{k}} A_0 + \dim_{\mathbf{k}} A_1 = m-1$, contradiction. We note that in this case the singularity is decomposable.

Case $(1, 1, 0)$. We have $\tilde{\mathfrak{m}}^3 \subseteq I$. We are going to write down the $m-1$ generators of $A_1 \pmod{\tilde{\mathfrak{m}}^3}$ ¹. The first generator, call it, x_1 , has a non-trivial linear term in at least one of the variables, wlog t_1 . By scaling x_1 and possibly adding a multiple of x_1^2 , we can make it into the form: $x_1 = t_1 \oplus p_{1,2}(t_2) \oplus \dots \oplus p_{1,m}(t_m) \pmod{\tilde{\mathfrak{m}}^3}$. Now we can use x_1 and x_1^2 to make sure the second generator does not involve t_1 at all. It will still have a linear term independent of t_1 , say non-trivial in t_2 . By scaling and adding a multiple of x_1^2 , we can write $x_2 = 0 \oplus t_2 \oplus \dots \oplus p_{2,m}(t_m) \pmod{\tilde{\mathfrak{m}}^3}$. By taking a linear combination of x_1 with x_2 and x_2^2 , we may now reduce x_1 to the form $t_1 \oplus 0 \oplus p_{1,3}(t_3) \oplus \dots \oplus p_{1,m}(t_m) \pmod{\tilde{\mathfrak{m}}^3}$. Therefore, by Gaussian elimination with the generators and their squares, we may assume that

$$\begin{aligned} x_1 &= t_1 \oplus 0 \oplus \dots \oplus \alpha_{1,m} t_m + \beta_{1,m} t_m^2 \\ x_2 &= 0 \oplus t_2 \oplus \dots \oplus \alpha_{2,m} t_m + \beta_{2,m} t_m^2 \\ &\dots \\ x_{m-1} &= 0 \oplus \dots \oplus t_{m-1} \oplus \alpha_{m-1,m} t_m + \beta_{m-1,m} t_m^2 \pmod{\tilde{\mathfrak{m}}^3} \end{aligned}$$

We must have $R/I = \langle 1, x_1, \dots, x_{m-1}, y \rangle$ by the Gorenstein condition (if $x_i \in I$, then $t_i \in R$, and it is then easy to see that the singularity would be decomposable). Hence $x_i^2 \in I$ for all but at most one i , say $i = 1$. Then $t_i^2 \in R$ for $i = 2, \dots, m-1$. If $\alpha_{i,m} \neq 0$ for some i in this range, then $t_m^2 \in R$ as well, so $t_1^2 = x_1^2 - O(t_m^2) \in R$, contradicting $\dim_{\mathbf{k}}(\tilde{R}/R)_2 = 1$. Therefore $\alpha_{i,m} = 0$ for $i \in \{2, \dots, m-1\}$. If $\alpha_{1,m} = 0$, then we need a further generator of $\mathfrak{m}/\mathfrak{m}^2$, namely $z = 0 \oplus \dots \oplus t_m^3$. In this case, though, both x_1^2 and z belong to I , so $\dim_{\mathbf{k}}(R/I) = m$, and the

¹To make them into the simplest possible form, we allow ourselves to perform linear algebra operations at first, and only at the end we may change coordinates - the benefit of this two-step process will become apparent in the next section.

singularity cannot be Gorenstein. We have then:

$$\begin{aligned}
 (1) \quad & x_1 = t_1 \oplus 0 \oplus \dots \oplus \alpha_{1,m} t_m + \beta_{1,m} t_m^2 \\
 & x_2 = 0 \oplus t_2 \oplus \dots \oplus \beta_{2,m} t_m^2 \\
 & \dots \\
 & x_{m-1} = 0 \oplus \dots \oplus t_{m-1} \oplus \beta_{m-1,m} t_m^2 \pmod{\tilde{\mathfrak{m}}^3},
 \end{aligned}$$

with $\beta_{1,m} \in \mathbf{k}$ and $\alpha_{1,m}, \beta_{i,m} \in \mathbf{k}^\times$, $i = 2, \dots, m-1$ (by indecomposability). Finally, we may change coordinates in t_m and scale the other t_i to obtain

$$\begin{aligned}
 (2) \quad & x_1 = t_1 \oplus 0 \oplus \dots \oplus t_m \\
 & x_2 = 0 \oplus t_2 \oplus \dots \oplus t_m^2 \\
 & \dots \\
 & x_{m-1} = 0 \oplus \dots \oplus t_{m-1} \oplus t_m^2 \pmod{\tilde{\mathfrak{m}}^3}.
 \end{aligned}$$

We may check that $R/I = \langle 1, x_1, \dots, x_{m-1}, x_1^2 \rangle$ and \tilde{R}/R is of type $(1, 1, 0)$. In the case $m = 2$, we need an extra generator $y = t_2^3$. Equations are given by

- $y(y - x_1^3)$ if $m = 2$ (A_5 -singularity);
- $x_1 x_2 (x_2 - x_1^2)$ if $m = 3$ (D_6 -singularity);
- $\langle x_3(x_1^2 - x_2), x_i(x_j - x_k) \rangle_{1 \leq i < j < k \leq m-1 \text{ or } 1 < j < k < i \leq m-1}$ if $m \geq 4$.

Case $(1, 0, 1)$. We have $\tilde{\mathfrak{m}}^4 \subseteq I$. By an argument similar to the above, we write generators for A_1 as $x_i = \dots \oplus t_i \oplus \dots \oplus \alpha_{i,m} t_m + \beta_{i,m} t_m^2 + \gamma_{i,m} t_m^3$, for $i = 1, \dots, m-1$. Then $R/I = \langle 1, x_1, \dots, x_{m-1}, y \rangle$. For all but at most one i , $x_i^2 \in I$, but definitely $x_i^3 \in I$ for all i . On the other hand $t_m^3 \notin R$, because otherwise $t_i^3 = x_i^3 - \alpha_{i,m}^3 t_m^3 + O(t_m^4)$ would belong to R as well, contradicting $\dim_{\mathbf{k}}(\tilde{R}/R)_3 = 1$. From this we deduce that $\alpha_{i,m} = 0$ for all $i = 1, \dots, m-1$. By $\dim_{\mathbf{k}}(\tilde{R}/R)_2 = 0$ there has to be another generator of degree two in t_m , which we may write as $x_m = t_m^2 + \gamma_{m,m} t_m^3$ of $\mathfrak{m}/\mathfrak{m}^2$. We can use x_m to remove all the t_m^2 pieces from x_1, \dots, x_{m-1} , so we are reduced to

$$\begin{aligned}
 (3) \quad & x_1 = t_1 \oplus 0 \oplus \dots \oplus \gamma_{1,m} t_m^3 \\
 & x_2 = 0 \oplus t_2 \oplus \dots \oplus \gamma_{2,m} t_m^3 \\
 & \dots \\
 & x_{m-1} = 0 \oplus \dots \oplus t_{m-1} \oplus \gamma_{m-1,m} t_m^3 \\
 & x_m = 0 \oplus \dots \oplus t_m^2 + \gamma_{m,m} t_m^3 \pmod{\tilde{\mathfrak{m}}^4},
 \end{aligned}$$

with $\gamma_{m,m} \in \mathbf{k}$ and $\gamma_{i,m} \in \mathbf{k}^\times$, $i = 1, \dots, m-1$ (by indecomposability). Finally, we may change coordinates in t_m and scale the other t_i to obtain

$$\begin{aligned}
 (4) \quad & x_1 = t_1 \oplus 0 \oplus \dots \oplus t_m^3 \\
 & x_2 = 0 \oplus t_2 \oplus \dots \oplus t_m^3 \\
 & \dots \\
 & x_{m-1} = 0 \oplus \dots \oplus t_{m-1} \oplus t_m^3 \\
 & x_m = 0 \oplus \dots \oplus t_m^2 \pmod{\tilde{\mathfrak{m}}^4}.
 \end{aligned}$$

We may check that $R/I = \langle 1, x_1, \dots, x_{m-1}, x_m \rangle$ and \tilde{R}/R is of type $(1, 0, 1)$. For $m = 1$ we recover the unique Gorenstein singularity of Lemma 2.1. Equations are

- $x^5 - y^2$ if $m = 1$ (A_4 -singularity or *ramphoid cusp*);
- $y(y^3 - x^2)$ if $m = 2$ (D_5 -singularity);
- $\langle x_3(x_1 - x_2), x_3^3 - x_1x_2 \rangle$ if $m = 3$;
- $\langle x_i(x_j - x_k), x_m(x_i - x_j), x_m^3 - x_1x_2 \rangle_{i,j,k \in \{1, \dots, m-1\} \text{ all different}}$ if $m \geq 4$.

□

Definition 2.3. In case $(1, 1, 0)$, we shall call the singularities of *type II*, and the branches parametrised by t_1 and t_m *twin*; in case $(1, 0, 1)$, we say the singularity is of *type I*, and the branch parametrised by t_m is called the *singular* branch. We shall refer to them as *special* or *distinguished* branches; all other branches are *axes*. *Branch* remains a generic name, indicating any of the previous ones.

Remark 2.4. Singularities of type I do appear in the miniversal family of singularities of type II, and viceversa. For low values of m - which serves as the playground for our speculations -, this follows neatly from a beautiful result of A. Grothendieck that we have learnt from [CML13] (see also [Arn72, Dem75]):

Theorem 2.5. *Let (C, p) be a curve singularity of ADE type. Singularities that appear in the miniversal deformation of (C, p) are all and only those ADE, whose Dynkin diagram can be obtained as a full subgraph of the diagram of (C, p) .*

3. TANGENT SHEAF, CRIMPING SPACE, AND AUTOMORPHISMS

In this section we analyse the tangent sheaf of a genus two singularity. For a complete Gorenstein curve of genus two with markings, we translate the absence of infinitesimal automorphisms into a (mostly) combinatorial condition. The crimping space naturally makes its appearance in the process.

Lemma 3.1. *Let (C, p) be a Gorenstein curve singularity of genus two, with pointed normalisation $\nu: (\tilde{C}, \{p_i\}_{i=1, \dots, m}) \rightarrow (C, p)$, and assume $\text{char}(\mathbf{k}) \neq 2, 3, 5$. There is a diagram of exact sequences of sheaves*

$$\begin{array}{ccccccc}
0 & \longrightarrow & \nu_* \Omega_{\tilde{C}}^\vee(-\sum_i 3p_i) & \longrightarrow & \nu_* \Omega_{\tilde{C}}^\vee(-\sum_i p_i) & \longrightarrow & \nu_* \bigoplus_i \Omega_{\tilde{C}}^\vee(-p_i)|_{2p_i} \longrightarrow 0 \\
& & \parallel & & \uparrow & & \square & \uparrow \phi \\
0 & \longrightarrow & \nu_* \Omega_{\tilde{C}}^\vee(-\sum_i 3p_i) & \longrightarrow & \Omega_C^\vee & \longrightarrow & \mathbf{k}^{\oplus m} \longrightarrow 0
\end{array}$$

The right-most vertical map admits an explicit description in local coordinates.

Proof. Let $K(\tilde{C})$ denote the constant sheaf of rational functions on \tilde{C} . A section of $\Omega_{\tilde{C}}^\vee \otimes K(\tilde{C})$ is contained in Ω_C^\vee iff its image under the push-forward map

$$\nu_*: \nu_* \mathcal{H}om(\Omega_{\tilde{C}}, K(\tilde{C})) \rightarrow \mathcal{H}om(\Omega_C, K(\tilde{C}))$$

lies in the subspace $\mathcal{H}om(\Omega_C, \mathcal{O}_C)$. We may work locally around the singular point in the coordinates studied in the previous section.

A₄: In the coordinates $x = t^2 + ct^3, y = t^4, z = t^5$ (they are redundant, but it will not matter in what follows), the section $f(t) \frac{d}{dt} \in \nu_* \Omega_{\tilde{C}}^\vee \otimes K(\tilde{C})$ pushes forward to

$$\nu_* \left(f(t) \frac{d}{dt} \right) = (2t + 3ct^2) f(t) \frac{d}{dx} + 4t^3 f(t) \frac{d}{dy} + 5t^4 f(t) \frac{d}{dz},$$

from which, writing $f(t) = f_0 + f_1 t + f_2 t^2 + O(t^3)$, we see that

$$(2t + 3ct^2) f(t), 4t^3 f(t), 5t^4 f(t) \in \hat{\mathcal{O}}_{C,p} \Leftrightarrow f_0 = 0, cf_1 + 2f_2 = 0.$$

A₅: In the coordinates $x = t_1 \oplus at_2 + bt_2^2, y = t_1^3$, the section $f_1(t_1) \frac{d}{dt_1} \oplus f_2(t_2) \frac{d}{dt_2}$ pushes forward to

$$\nu_* \left(f_1(t_1) \frac{d}{dt_1} \oplus f_2(t_2) \frac{d}{dt_2} \right) = (f_1(t_1) \oplus (a + 2bt_2) f_2(t_2)) \frac{d}{dx} + 3t_1^2 f_1(t_1) \frac{d}{dy},$$

from which, writing $f_i(t_i) = f_{i0} + f_{i1} t_i + f_{i2} t_i^2 + O(t_i^3), i = 1, 2$, we see that

$$f_1(t_1) \oplus (a + 2bt_2) f_2(t_2), 3t_1^2 f_1(t_1) \in \hat{\mathcal{O}}_{C,p} \Leftrightarrow \begin{cases} f_{10} = f_{20} = 0, \\ f_{11} = f_{21}, \\ 2bf_{21} + af_{22} = a^2 f_{12}. \end{cases}$$

$\Pi_{m \geq 3}$: In the coordinates of (1),

$$\begin{aligned}
\nu_* \left(\sum_{i=1}^m f_i(t_i) \frac{d}{dt_i} \right) &= (f_1(t_1) \oplus (\alpha_{1,m} + 2\beta_{1,m} t_m) f_m(t_m)) \frac{d}{dx_1} + \\
&\quad \sum_{i=2}^m (f_i(t_i) \oplus 2\beta_{i,m} t_m f_m(t_m)) \frac{d}{dx_i},
\end{aligned}$$

hence we deduce that

$$\nu_* \left(\sum_{i=1}^m f_i(t_i) \frac{d}{dt_i} \right) \in \Omega_C^\vee \otimes \hat{\mathcal{O}}_{C,p} \Leftrightarrow \begin{cases} f_{i0} = 0 & \text{for } i = 1, \dots, m, \\ 2f_{11} = f_{i1} = 2f_{m1}, & \text{for } i = 2, \dots, m-1, \\ \beta_{1,m}f_{m1} + \alpha_{1,m}f_{m2} = \alpha_{1,m}^2 f_{12}. \end{cases}$$

$III_{m \geq 2}$: In the coordinates of (3),

$$\begin{aligned} \nu_* \left(\sum_{i=1}^m f_i(t_i) \frac{d}{dt_i} \right) &= \sum_{i=1}^{m-1} (f_i(t_i) \oplus 3\gamma_{i,m} t_m^2 f_m(t_m)) \frac{d}{dx_i} + \\ &\quad (2t_m + 3\gamma_{m,m} t_m^2) f_m(t_m) \frac{d}{dx_m}, \end{aligned}$$

hence we deduce that

$$\nu_* \left(\sum_{i=1}^m f_i(t_i) \frac{d}{dt_i} \right) \in \Omega_C^\vee \otimes \hat{\mathcal{O}}_{C,p} \Leftrightarrow \begin{cases} f_{i0} = 0 & \text{for } i = 1, \dots, m, \\ f_{i1} = 3f_{m1}, & \text{for } i = 1, \dots, m-1, \\ 3\gamma_{m,m} f_{m1} + 2f_{m2} = 0. \end{cases}$$

□

From this description we see that the letters α, β and γ will play a role in determining the automorphism group of a complete curve with markings. We recall some key concepts from F. van der Wyck's thesis.

We work over \mathbf{k} . We can consider the stack \mathcal{S} of reduced one-dimensional \mathbf{k} -algebras R , and the stack \mathcal{T} of reduced 1d algebras with resolution $(R \hookrightarrow (S, J))$, where S is a smooth one-dimensional \mathbf{k} -algebra, and J is the radical of the conductor of $R \subseteq S$. Basically, R is the (local) ring of a reduced curve with one singular point, S is its normalisation, and J is the ideal of the reduced fiber over the singular point of $\text{Spec}(R)$. \mathcal{S} and \mathcal{T} are limit-preserving stacks over $\text{Spec}(\mathbf{k})$ [vdW10, Proposition 1.21]. Furthermore, we may fix a reduced 1d algebra with resolution $\tau_0 : (R_0 \hookrightarrow (S_0, J_0))$, and consider the substack $\mathcal{T}(\tau_0)$ of reduced 1d algebras with singularity type τ_0 (i.e. isomorphic to τ_0 locally on both the base and the curve, see [vdW10, Definition 1.64]; that various notions of “locally” coincide is proved in [vdW10, Proposition 1.50]). There is a forgetful morphism $\mathcal{T} \rightarrow \mathcal{S}$, and the *crimping space* of τ_0 is defined to be the fiber over R_0 of the restriction of such morphism to $\mathcal{T}(\tau_0)$. The crimping space is a smooth \mathbf{k} -scheme [vdW10, Theorems 1.70 and 1.73]; indeed, it is isomorphic to the quotient of $\text{Aut}_{(S_0, J_0)/\mathbf{k}}$ by $\text{Aut}_{(S_0, J_0)/R_0}$, the latter being the subgroup of automorphisms of the normalisation preserving the subalgebra of the singularity; moreover, the quotient can be computed after modding out the lowest power of J that is contained in R [vdW10, Theorem 1.53]. Crimping spaces should be thought of as moduli for the normalisation map.

Lemma 3.2. *The crimping space of a genus two singularity with m branches is (a number - depending on the type - of copies of) $\mathbb{A}^1 \times (\mathbb{A}^1 \setminus \{0\})^{m-1}$.*

Proof. We resume notation from the previous section. We are going to fix the subalgebra τ_0 given in coordinates by (2) and (4) above respectively.

Type II: recall that in this case $\tilde{\mathfrak{m}}^3 \subseteq R$. For a \mathbf{k} -algebra A , let

$$G_i(A) = \{t_i \mapsto g_{i1}t_i + g_{i2}t_i^2, t_j \mapsto t_j \mid g_{i1} \in A^\times, g_{i2} \in A\},$$

and notice that

$$\text{Aut}_{(\tilde{R}, \tilde{\mathfrak{m}})}^{\text{mod } \tilde{\mathfrak{m}}^3}(A) = (G_1 \times \dots \times G_m) \rtimes \mathfrak{S}_m(A).$$

Consider now the action of a group element of the form $(g_1, \dots, g_m; \text{id}_{\mathfrak{S}_m})$ on the given generators of R :

$$\begin{aligned} x_1 &\mapsto g_{11}t_1 + g_{12}t_1^2 \oplus \dots \oplus g_{m1}t_m + g_{m2}t_m^2; \\ x_i &\mapsto \dots \oplus g_{i1}t_i + g_{i2}t_i^2 \oplus \dots \oplus g_{m1}^2t_m^2, \quad \text{for } i = 2, \dots, m-1. \end{aligned}$$

The former belongs to R iff $g_{11} = g_{m1}$ and $g_{12} = g_{m2}$; the latter does iff $g_{i1} = g_{m1}^2$. Thus, such elements span a subgroup isomorphic to $\mathbb{G}_m \times \mathbb{G}_a^{m-1}(A)$. On the other hand, all branches are isomorphic to one another, but there is a pair of distinguished ones (parametrised by t_1 and t_m respectively). We conclude that

$$\text{Aut}_{\tau_0}^{\text{mod } \tilde{\mathfrak{m}}^3}(A) = (\mathbb{G}_m \times \mathbb{G}_a^{m-1}) \rtimes (\mathfrak{S}_2 \times \mathfrak{S}_{m-2})(A).$$

The quotient is then isomorphic to $\binom{m}{2}$ copies of $\mathbb{A}^1 \times (\mathbb{A}^1 \setminus \{0\})^{m-1}$.

Type III: in this case $\tilde{\mathfrak{m}}^4 \subseteq R$. For a \mathbf{k} -algebra A , let

$$G_i(A) = \{t_i \mapsto g_{i1}t_i + g_{i2}t_i^2 + g_{i3}t_i^3, t_j \mapsto t_j \mid g_{i1} \in A^*, g_{i2}, g_{i3} \in A\},$$

and notice that

$$\text{Aut}_{(\tilde{R}, \tilde{\mathfrak{m}})}^{\text{mod } \tilde{\mathfrak{m}}^4}(A) = (G_1 \times \dots \times G_m) \rtimes \mathfrak{S}_m(A).$$

Consider now the action of a group element of the form $(g_1, \dots, g_m; \text{id}_{\mathfrak{S}_m})$ on the given generators of R :

$$\begin{aligned} x_i &\mapsto \dots \oplus g_{i1}t_i + g_{i2}t_i^2 + g_{i3}t_i^3 \oplus \dots \oplus g_{m1}^3t_m^3, \quad \text{for } i = 1, \dots, m-1; \\ x_m &\mapsto \dots \oplus g_{m1}^2t_m^2 + 2g_{m1}g_{m2}t_m^3. \end{aligned}$$

The former belongs to R iff $g_{i1} = g_{m1}^3$; the latter does iff $g_{m2} = 0$. Thus such elements span a subgroup isomorphic to $\mathbb{G}_m \times \mathbb{G}_a^{m-1} \times \mathbb{G}_a^m(A)$. On the other hand, there is a special (singular) branch, parametrised by t_m . We conclude that

$$\text{Aut}_{\tau_0}^{\text{mod } \tilde{\mathfrak{m}}^4}(A) = (\mathbb{G}_m \times \mathbb{G}_a^{m-1} \times \mathbb{G}_a^m) \rtimes (\mathfrak{S}_{m-1})(A).$$

The quotient is therefore isomorphic to m copies of $\mathbb{A}^1 \times (\mathbb{A}^1 \setminus \{0\})^{m-1}$. \square

It is now clear that the benefit of a two-step classification where at first we do not allow ourselves to change coordinates (i.e. act by automorphisms of the normalisation) is that it makes the crimping space apparent already from the expressions (1) and (3) for the generators of the singularity subalgebra.

There is a more geometric way to see the crimping spaces. It is well-known that a cusp can be obtained by collapsing (*push-out*) a generic (i.e. non-zero)

tangent vector at $p \in \mathbb{P}^1$. More generally, a Gorenstein singularity of genus one and m branches can be obtained by collapsing a generic (not contained in any coordinate linear subspace) tangent line at an ordinary (i.e. rational) m -fold point [Smy11a, Lemma 2.2]. Hence, we recover the crimping space of the elliptic m -fold point, which is isomorphic to $(\mathbb{A}^1 \setminus \{0\})^{m-1}$, as the maximal torus inside $\mathbb{P}(T_p R_m) \simeq \mathbb{P}^{m-1}$, where (R_m, p) is the rational m -fold point. Besides, this gives a natural compactification of the crimping space supporting a universal family of curves - in fact, two: either we collapse non-generic tangent vectors, obtaining non-Gorenstein singularities along the boundary (this family \mathcal{C} has a common normalisation, that is the trivial family $\tilde{\mathcal{C}} = R_m \times \mathbb{P}(T_p R_m)$); or we blow-up $\tilde{\mathcal{C}}$ along the boundary (*sprouting*), and we replace the non-Gorenstein singularities by more elliptic m -fold points, this time with strictly semistable branches [Smy11b, §2.2-3].

Similarly, a Gorenstein singularity of genus two can be obtained by collapsing a generic tangent line to a genus one non-Gorenstein singularity. Indeed, type τ_0^{II} admits a partial normalisation by σ_0^{II} , which is the decomposable union of a tacnode in the (t_1, t_m) -plane together with $m-2$ axes, by adjoining the generator t_m^2 ; while type τ_0^{III} by σ_0^{III} , which is the decomposable union of a cusp (parametrised by t_m) together with $m-1$ axes, by adjoining t_m^3 . These fit together nicely in the following picture: if we restrict \mathcal{C} from the previous paragraph to the union of the coordinate lines in $\mathbb{P}(T_p R_m)$, we obtain m copies of σ_0^{III} over the points, together with $\binom{m}{2}$ copies of the universal curve of type σ_0^{II} over its crimping space - which is isomorphic to $\mathbb{A}^1 \setminus \{0\}$ -, identified with the open lines. Let $P = \mathbb{P}(T_p \mathcal{C}_{|\cup \text{lines}})$ be the projectivised tangent space at the singular point in the fiber of this family of genus one singularities. For each of the $\binom{m}{2}$ coordinate lines, P has one component P_i^{II} that is a \mathbb{P}^{m-1} -bundle over such line; besides, P has m components P_j^{III} isomorphic to \mathbb{P}^m and supported over the points. The crimping space of the genus two singularities (of type II and III together) with m branches can be seen to be an open subscheme of P : it is obtained by removing from the \mathbb{P}^{m-1} -fibers of P^{II} the $m-1$ hyperplanes generated by (a) the tangent line to the tacnode and the $m-2$ axes, and (b) the plane containing the tacnode and all but one of the $m-2$ axes; and from each component of P^{III} the m planes generated by (a) the tangent cone of the cusp and the $m-1$ axes, and (b) the plane containing the cusp and all but one of the $m-1$ axes.

Remark 3.3. The crimping space is related to the moduli of arrows ϕ as in the diagram of Lemma 3.1, satisfying a number of requirements.

We note that $H^0(\Omega_{\mathbb{P}^1}^\vee(-p)_{|2p})$ is the tangent space to the subgroup of automorphisms of \mathbb{P}^1 fixing one point p , so it is isomorphic as a Lie algebra to the only non-abelian Lie algebra of dimension two V . It has a basis $e_1 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, $e_2 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$ with $[e_1, e_2] = -2e_2$. The vector (φ, ψ) therefore corresponds to the

infinitesimal automorphism:

$$t \mapsto \frac{1 + \epsilon \varphi t}{1 - \epsilon(\varphi t + \psi)} = t + \epsilon(2\varphi t - \psi t^2).$$

We require ϕ to be the embedding (i.e. a point of $\text{Gr}(m, V^{\oplus m})$) of a Lie subalgebra, such that the corresponding group of infinitesimal automorphisms fixes uniquely the subalgebra of a singularity of genus two in $\mathbf{k}[[t_1]] \oplus \dots \oplus \mathbf{k}[[t_m]]$.

We start with some heuristics. The unibranch case goes as follows: the subalgebra of $\mathbf{k}[[t]]$ generated by $x = t^2 + ct^3$ is preserved by (φ, ψ) iff

$$(1 + 2\varphi)^2 t^2 - 2\psi(1 + 2\varphi)t^3 + c(1 + 2\varphi)^3 t^3 \text{ is a multiple of } t^2 + ct^3,$$

which reduces to $\varphi(1 + 2\varphi)c = \psi$. This furthermore determines c iff $\varphi \neq 0$. Note that in this case (dimension one) the Lie subalgebra condition is automatically satisfied. We have found $(\varphi, \psi) \in \mathbf{k}^\times \times \mathbf{k}$.

The case of type II_2 -algebras is more interesting. Let $x = (t_1, \alpha t_2 + \beta t_2^2)$ be the generator of such an algebra. The image of x under $(\varphi_1, \psi_1, \varphi_2, \psi_2)$ is:

$$((1 + 2\varphi_1)t_1 - \psi_1 t_1^2, \alpha(1 + 2\varphi_2)t_2 + (\beta(1 + 4\varphi_2) - \alpha\psi_2)t_2^2),$$

from which we deduce:

$$(5) \quad \varphi_1 = \varphi_2 \quad \text{and} \quad 2\beta\varphi_2 - \alpha\psi_2 = -\alpha^2\psi_1.$$

Now let $\phi: \mathbf{k}^2 \rightarrow V^{\oplus 2}$ be given by $\begin{pmatrix} \varphi_{11} & \psi_{11} & \varphi_{12} & \psi_{12} \\ \varphi_{21} & \psi_{21} & \varphi_{22} & \psi_{22} \end{pmatrix}$, with Plücker coordinates w_{ij} for the minor of the i -th and j -th columns. The first condition in (5) immediately implies

$$(6) \quad w_{13} = 0 \quad \text{and} \quad w_{12} = -w_{23}, w_{14} = w_{34}.$$

The second condition in (5) implies

$$(7) \quad \alpha w_{12} = w_{14} \quad \text{and} \quad 2\beta w_{12} = \alpha w_{24}$$

so that $(\alpha, \beta) \in \mathbf{k}^\times \times \mathbf{k}$ is determined as soon as $w_{12}, w_{14} \neq 0$. Notice that (6) implies in particular the Plücker equation

$$w_{12}w_{34} - w_{13}w_{24} + w_{14}w_{23}.$$

It is easy to see that the condition for ϕ to be a sub-Lie algebra is

$$\text{rk} \begin{pmatrix} \varphi_{11} & \psi_{11} & \varphi_{12} & \psi_{12} \\ \varphi_{21} & \psi_{21} & \varphi_{22} & \psi_{22} \\ 0 & w_{12} & 0 & w_{34} \end{pmatrix} = 2,$$

translating into

$$\begin{aligned} w_{12}w_{13} &= 0 & w_{12}(w_{34} - w_{14}) &= 0 \\ w_{13}w_{34} &= 0 & (w_{12} + w_{23})w_{34} &= 0 \end{aligned}$$

which also are automatically satisfied after (6). We see the various equations above cut inside $\mathbb{P}_{[w_{ij}]}^5$ the locus

$$(\mathbb{A}_{w_{14}/w_{12}}^1 \setminus \{0\}) \times \mathbb{A}_{w_{24}/w_{12}}^1.$$

More generally, given a subalgebra $R_{\alpha,\beta}$ of $W = \bigoplus_{i=1}^m \mathbf{k}[[t_i]]/(t_i^2)$, with generators of the form described in (1), the subalgebra of $V_{(\varphi_i, \psi_i)_{i=1, \dots, m}}^{\oplus m}$ preserving $R_{\alpha,\beta}$ is isomorphic to $\mathbf{k}^{\oplus m}$ with equations (see Lemma 3.1):

$$\begin{cases} 2\varphi_1 = \varphi_i = 2\varphi_m, & \text{for } i = 2, \dots, m-1, \\ 2\beta_{1,m}\varphi_m - \alpha_{1,m}\psi_m = -\alpha_{1,m}^2\psi_1; \end{cases}$$

it is easily seen that such a subalgebra of $V^{\oplus m}$ does not determine $R_{\alpha,\beta}$, but it does determine $(\alpha_{1,m}, \beta_{1,m})$. The situation of type III is analogous.

We are going to use the preceding discussion in order to study the automorphism group of complete marked curves with a genus two singularity. The only case that actually requires such a discussion is when every component of the normalisation is rational and contains one extra marking, besides the preimage of the singularity: in this case it does make a difference what point of the crimping space we are looking at. The concept has been formalised again in van der Wyck's thesis, see [vdW10, Proposition 1.102, Theorem 1.105 and Corollary 1.106], where he introduces the concept of reduced pointed curve with resolution of type T (encoding the amount and type of the singular points, the distribution of genus and markings among the components of the normalisation, and the adjacency data between components and singular points), and the algebraic stack \mathcal{N}_T of such objects. In the case at hand, such stack is isomorphic to $[\mathbb{A}^1/\mathbb{G}_m]$ (see also [vdW10, Examples 1.111-112]), and it therefore has two points: one with \mathbb{G}_m , and the other with trivial stabiliser.

Definition 3.4. The *atom* of type II_m is obtained by gluing the subalgebra of $\mathbf{k}[t_1] \oplus \dots \oplus \mathbf{k}[t_m]$ generated by x_1, \dots, x_{m-1} (and y) as in (2) (and following lines) with m copies of $(\mathbf{k}[s], (s))$ under the identification $s_i = t_i^{-1}$. There is a \mathbb{G}_m -action on the type II atom by $\lambda.t_i = \lambda t_i$ for $i = 1, m$ and $\lambda.t_i = \lambda^2 t_i$ for $i = 2, \dots, m-1$.

Similarly, the atom of type III_m is obtained by gluing the subalgebra of $\mathbf{k}[t_1] \oplus \dots \oplus \mathbf{k}[t_m]$ generated by x_1, \dots, x_m as in (4) with m copies of $(\mathbf{k}[s], (s))$ under the identification $s_i = t_i^{-1}$. There is a \mathbb{G}_m -action on the type III atom by $\lambda.t_i = \lambda t_i$ for $i = m$ and $\lambda.t_i = \lambda^3 t_i$ for $i = 1, \dots, m-1$.

The curve with a genus two singularity and one marked point for every branch that has trivial automorphism group will be called the *non-atom*.

Following the previous discussion, there is a more geometric way to realise the dicotomy between the atom and the non-atom. The non-Gorenstein genus one singularity of type σ_0^{II} (resp. σ_0^{III}), with every branch rational and one-marked, has automorphism group \mathbb{G}_m^{m-1} (resp. \mathbb{G}_m^m); the latter group therefore acts on the

tangent space at the singular point, and it can be checked that of the lines fixed by this action only one sits inside the open subset corresponding to the crimping space, while all other lines in the open are identified under the group action - by collapsing them, they give rise to the atom and non-atom respectively.

As a third viewpoint, automorphisms can be studied by twisting the exact sequences of Lemma 3.1 by the ideal of the markings and then taking global sections. It appears that the dicotomy arises from the map ϕ : if the last condition imposed on infinitesimal automorphisms interweaves first and second order non-trivially (i.e. when $\beta_{1,m}$, resp. $\gamma_{m,m}$, are non-zero) then it is enough that automorphisms be trivial to second order on all branches for them to be trivial tout-court.

Finally, we shall turn the condition that the automorphism group be finite into a combinatorial one. For this, recall Smyth's description of Gorenstein curves of genus one with no automorphisms [Smy11a, Proposition 2.3, Corollary 2.4].

Definition 3.5. Let (C, p_1, \dots, p_n) be a pointed reduced curve. A connected subcurve $D \subseteq C$ is said to be *nodally attached* if $D \cap \overline{C \setminus D}$ consists of nodes only. We say that C is *residually DM* (rDM) if every nodal and nodally attached subcurve D of C , marked by $\{p_i \in D\} \cup D \cap \overline{C \setminus D}$, is Deligne-Mumford stable.

Special points are either nodes or markings.

Corollary 3.6. Let (C, p_1, \dots, p_n) be a pointed Gorenstein curve of arithmetic genus two. $H^0(C, \Omega_C^\vee(-\sum_{i=1}^n p_i)) = 0$ is equivalent to either of the following:

- (1) C has a singularity of type $II_{m \geq 2}$: either all branches contain exactly one special point and C is not atomic; or at least one of its twin branches contains a special point, each of its axes contains at least one special point, and at least one branch has at least two. Furthermore C is rDM.
- (2) C has a singularity of type $III_{m \geq 1}$: either all branches contain exactly one special point and C is not atomic; or each of its axes contains at least one special point, and at least one branch has at least two. Furthermore C is rDM.
- (3) C has two elliptic m -fold points: each of their branches contains at least one special point, and either they share a branch, or at least one branch of each singular point contains at least two special points. Furthermore C is rDM.
- (4) C has one elliptic m -fold point: one of its branches is a genus one curve, or two of its branches coincide, and each of the other ones contains at least one special point; otherwise, all branches contain at least one special point, and at least one branch has at least two. Furthermore C is rDM.
- (5) C contains only nodes and is Deligne-Mumford stable.

4. DUALISING LINE BUNDLE AND SEMISTABLE TAILS

Given a family of prestable (pointed) curves of genus two over the spectrum of a discrete valuation ring $\mathcal{C} \rightarrow \Delta$, with smooth generic fiber \mathcal{C}_η and regular total space, we classify the subcurves of the central fiber \mathcal{C}_0 that can be contracted

to produce a Gorenstein singularity of genus two. In the genus one case, Smyth answered the analogous question by identifying the class of *balanced* subcurves: subcurves of arithmetic genus one, such that, when breaking them into a *core* (minimal subcurve of genus one, i.e. not containing any separating node) and a number of rational trees (with root corresponding to the component adjacent to the core, and leaves corresponding to the components adjacent to the portion of \mathcal{C}_0 that is not contracted), the distance between any leaf and the root for any such tree is constant. In the case at hand, the answer turns out to be slightly more complicated: first, the special branch(es) of a type I (resp. II) singularity correspond through a rational chain to a Weierstrass (resp. two conjugate) point(s) of the core, and the special branches are always the closest to the core. Second, if the core is reducible, the lengths of the rational trees may vary according to where their attaching points lie on the core, but they are determined by the length of the special chains and the configuration of the attaching points on the core.

Remark 4.1. While there are no special points on a smooth curve of genus zero or one, the simplest instance of Brill-Noether theory involves smooth curves of genus two. Every such C is *hyperelliptic*: it admits a unique (up to reparametrisation) $2:1$ cover $\phi: C \rightarrow \mathbb{P}^1$, induced by the complete canonical linear system, i.e. $|K_C|$ is the unique \mathfrak{g}_2^1 on C ; said otherwise, there is a unique element $\sigma \in \text{Aut}(C)$, called the *hyperelliptic involution*, such that $C/\langle\sigma\rangle \simeq \mathbb{P}^1$. A point $x \in C$ is called *Weierstrass* if it is a ramification point for ϕ (or, equivalently, a fixed point for σ); from the Riemann-Hurwitz formula it follows that there are six Weierstrass points on every smooth curve of genus two. Two points x_1, x_2 are said to be conjugate (write $x_2 = \overline{x_1}$) if there exists a point $z \in \mathbb{P}^1$ such that $\phi^{-1}(z) = \{x_1, x_2\}$ (or, equivalently, $\sigma(x_1) = x_2$). These notions may be extended to nodal curves by declaring (C, x) to be Weierstrass if its stabilisation lies in the closure of

$$\mathcal{W} = \{(C, x) \mid C \text{ smooth and } x \text{ Weierstrass}\} \subseteq \overline{\mathcal{M}}_{2,1},$$

and similarly for conjugate points. We then need to study the limiting behaviour of Weierstrass points when a smooth curve degenerates to a nodal one. This is a difficult problem when it comes to higher genus curves; it has received considerable attention since the '70s, in work of Arbarello, Eisenbud-Harris, and many others. In our case it boils down to understanding admissible covers [HM82] of degree two with a branch locus of degree six; said otherwise, up to the involution action, the Weierstrass locus is isomorphic to $\overline{\mathcal{M}}_{0,6}/\mathfrak{S}_5$, and the conjugate locus is isomorphic to $\overline{\mathcal{M}}_{0,7}/\mathfrak{S}_6$. We remark that (C, x) being Weierstrass is an intrinsic notion if C is of compact type (or, more generally, tree-like), but it may depend on the smoothing otherwise (i.e. the fiber of $\mathcal{W} \rightarrow \overline{\mathcal{M}}_2$ may have positive dimension); we have benefitted from the exposition in [Dia85, Appendix 2], [Cuk89, Proposition (3.0.6)], and [HM98, Theorem 5.45].

- If x belongs to a component of genus one E , which is attached to another component of genus one at a node y , then x is Weierstrass iff $2x \sim 2y \in \text{Pic}(E)$;




FIGURE 2. Admissible covers and Weierstrass points.

if instead E has a self-node that glues y_1 with y_2 , then x is Weierstrass iff $2x \sim y_1 + y_2 \in \text{Pic}(E)$.

If x is on a rational component R , x is Weierstrass if either R is attached to a genus one curve at two distinct points, or R has a self-node gluing y_1 and y_2 and is attached to a genus one tail at y_3 , in which case we require $\phi(y_1) = \phi(y_2)$ for a double cover $\phi: R \rightarrow \mathbb{P}^1$ ramified at x and y_3 , or R has two self-nodes gluing y_1 with y_2 , and y_3 with y_4 , in which case we require x to be a ramification point for a double cover $\phi: R \rightarrow \mathbb{P}^1$ such that $\phi(y_1) = \phi(y_2)$ and $\phi(y_3) = \phi(y_4)$. See Figure 2.

- If x_1 and x_2 are conjugate, they have to map to the same component of the target of the admissible cover. The description of the previous point works by replacing every condition on $2x$ by its analogue for $x_1 + x_2$. There are a few more situations to take into account: x_1 and x_2 could belong to a rational component R bubbling off from a Weierstrass point of a genus two curve; or bridging between two distinct curves of genus one; or x_1 and x_2 could lie on two distinct rational components R_1 and R_2 intersecting at one node and meeting a curve of genus one in two distinct points (\dagger); or R_1 and R_2 intersecting each other in three points. See Figure 3.

We observe here that in case (\dagger), the singularity of the total space of a smoothing $\mathcal{C} \rightarrow \Delta$ at the two distinguished nodes (separating the elliptic component from the rational chain) are both A_k for the same k , because they map to the same




FIGURE 3. Admissible covers and conjugate points.

node of the target in the admissible cover picture. This fact is stable under base change, and it determines a symmetry of the rational chain in the model with regular total space.

Proposition 4.2. *Let $\phi: \mathcal{C} \rightarrow \overline{\mathcal{C}}$ be a contraction over the spectrum of a discrete valuation ring Δ , where: $\mathcal{C} \rightarrow \Delta$ is a family of prestable (reduced, nodal) curves of arithmetic genus two, with regular total space and smooth generic fiber \mathcal{C}_η ; and $\overline{\mathcal{C}} \rightarrow \Delta$ is a family of Gorenstein curves of arithmetic genus two, with $\overline{\mathcal{C}}_\eta$ smooth, and $\overline{\mathcal{C}}_0$ containing a genus two singularity of type I_m at q . Denote by $(Z; q_1, \dots, q_m)$ the exceptional locus $\text{Exc}(\phi) = \phi^{-1}(q)$, marked with $Z \cap \overline{\mathcal{C}}_0 \setminus Z$, where q_m corresponds to the special branch of $\overline{\mathcal{C}}_0$. Then:*

- (1) *The stabilisation of (Z, q_m) is Weierstrass.*
- (2) *Let x_m be the point of the core of Z closest to q_m , and let k be the length of R_m , the rational chain separating q_m from x_m . With similar notation, for every $i = 1, \dots, m-1$, R_i has length*

$$\begin{cases} 3k + 1 + \text{dist}(x_m, x_i) & \text{if } x_i \neq x_m, \\ k + 2 \text{dist}(q_m, r_i) & \text{if } x_i = x_m, \end{cases}$$

where r_i is the point of R_m closest to q_i , and $\text{dist}(a, b)$ is the number of irreducible components between the points a and b (so for example it is 1 if a and b lie on the same irreducible component but $a \neq b$).

Proposition 4.3. *Same as above with $\overline{\mathcal{C}}_0$ containing a genus two singularity of type II_m , and q_1, q_m corresponding to the special branches. Then:*

(1) The stabilisation of (Z, q_1, q_m) is conjugate.

(2) R_1 and R_m have the same length k , and, for $i = 2, \dots, m-1$, R_i has length

$$\begin{cases} 2k + \min_{\epsilon \in \{1, m\}} \text{dist}^*(x_\epsilon, x_i) & \text{if } x_1 \neq x_m, \text{ and } x_i \notin \{x_1, x_m\}, \\ k + \text{dist}(q_1, r_i) & \text{if } x_1 \neq x_m, \text{ and } x_i = x_1 (+\text{symm. } 1 \leftrightarrow m), \\ 2k + \text{dist}(x_1, r_m) + \text{dist}(x_1, x_i) & \text{if } x_1 = x_m, \text{ and } x_i \neq x_1, \\ k + \text{dist}(q_1, r_i) + \delta(\text{dist}(r_m, r_i) - 1) & \text{if } x_1 = x_m = x_i, \text{ and } \delta = \begin{cases} 1 & r_m \in [q_1, r_i] \\ 0 & \text{otherwise} \end{cases} \end{cases}$$

where we set $\text{dist}^*(x_\epsilon, x_i) = 1$ if the core consists of a genus one curve with a rational bridge, and x_i lies between x_1 and x_m on this rational bridge, and $\text{dist}^*(x_\epsilon, x_i) = \text{dist}(x_\epsilon, x_i)$ otherwise.

Proposition 4.4. *Let $(\mathcal{C}, \Sigma_1, \dots, \Sigma_n) \rightarrow \Delta$ be a family of pointed semistable curves of arithmetic genus two such that \mathcal{C} has regular total space and smooth generic fiber, and $(\mathcal{C}, \Sigma_1) \rightarrow \Delta$ is Weierstrass. Let (Z, q_1, \dots, q_m) be a genus two subcurve of \mathcal{C}_0 containing none of the $\Sigma_i(0)$, marked by $Z \cap \mathcal{C}_0 \setminus \overline{Z}$ so that the tail containing Σ_1 is attached to Z at q_1 , and satisfying all the shape prescriptions of Proposition 4.2(2). There exists a contraction $\phi: \mathcal{C} \rightarrow \overline{\mathcal{C}}$ over Δ , with exceptional locus Z , such that $\overline{\mathcal{C}} \rightarrow \Delta$ is a family of Gorenstein curves containing a type I_m singularity in the central fiber.*

Proposition 4.5. *Same as above with $(\mathcal{C}, \Sigma_1, \Sigma_2) \rightarrow \Delta$ conjugate, (Z, q_1, \dots, q_m) shaped as prescribed by Proposition 4.3(2), and the resulting $\overline{\mathcal{C}} \rightarrow \Delta$ containing a type II_m singularity in the central fiber.*

Proof. (of Proposition 4.2) By blowing down all the rational trees on $\overline{\mathcal{C}}_0$, we can assume that the latter does not contain any separating node. Consider then the hyperelliptic cover $\tau: \overline{\mathcal{C}} \rightarrow \mathbb{P}(\pi_* \omega_{\overline{\mathcal{C}}/\Delta})$; restricting to the central fiber, τ contracts all axes, and gives a $2:1$ covering of \mathbb{P}^1 by the special branch, ramified at the singularity and at another point; in fact, we can extend the image of this point to a section of $\mathbb{P}(\pi_* \omega_{\overline{\mathcal{C}}/\Delta})$ lying inside the branch locus of τ . By pulling this back to \mathcal{C} via $\tau \circ \phi$ we get a horizontal divisor Δ' ; clearly, the stable model of (\mathcal{C}, Δ') is Weierstrass, and its central fiber coincides with the stabilisation of (Z, q_1) . This proves the first claim. (The proof of Proposition 4.3(1) is entirely analogous, by noticing that the preimage of a generic hyperplane section of $\mathbb{P}(\pi_* \omega_{\overline{\mathcal{C}}/\Delta})$ will mark the two special branches of $\overline{\mathcal{C}}_0$.)

We now come to a proof of the more combinatorial claim (2) of the Proposition. Since $\overline{\mathcal{C}} \rightarrow \Delta$ is Gorenstein, and ϕ is assumed to be an isomorphism outside Z , because the dualising sheaf behaves well under restriction to open subschemes, we have an equality of line bundles:

$$\phi^* \omega_{\overline{\mathcal{C}}/\Delta} = \omega_{\mathcal{C}/\Delta}(D),$$

for some effective (Cartier) divisor D on \mathcal{C} supported on Z . The next lemma will help us determine the coefficients of D along the components of Z containing q_i .

Lemma 4.6. *Let $\nu: C \rightarrow \bar{C}$ be the normalisation of a Gorenstein singularity of genus two, with $\nu^{-1}(q) = \{q_1, \dots, q_m\}$. Then $\nu^*\omega_{\bar{C}} = \omega_C(2q_1 + \dots + 2q_{m-1} + 4q_m)$ (type I) or $\nu^*\omega_{\bar{C}} = \omega_C(3q_1 + 2q_2 + \dots + 2q_{m-1} + 3q_m)$ (type II).*

Proof. The dualising sheaf of a reduced curve admits an explicit description (due to Rosenlicht, see e.g. [AK70, Proposition VIII.1.16]) in terms of residues:

$$\omega_{\bar{C}}(U) = \{\eta \in \Omega_C \otimes K(\nu^{-1}(U)) \mid \sum_{p_i \in \nu^{-1}(p), p \in U} \text{Res}_{p_i}((\nu^*f)\eta) = 0, \forall f \in \mathcal{O}_{\bar{C}}(U)\}.$$

We are going to use the explicit coordinates in (1) and (3). In case I, we know that $\tilde{\mathfrak{m}}^4 \subseteq R$, therefore we have poles of fourth order at most. It is enough to study the possible polar tails. On the other hand, $t_i^2 \in R$ for all i implies the part of order three is trivial. So let

$$\eta = c_1 \frac{dt_1}{t_1^4} + b_1 \frac{dt_1}{t_1^2} + a_1 \frac{dt_1}{t_1} \oplus \dots \oplus c_m \frac{dt_m}{t_m^4} + b_m \frac{dt_m}{t_m^2} + a_m \frac{dt_m}{t_m}.$$

From looking at $1 \cdot \eta$ we deduce $\sum_{i=1}^m a_i = 0$; from $x_i \cdot \eta$ we see $b_i + c_m = 0$ for all i , and from $x_i^3 \cdot \eta$ we have $c_i = 0$ for all i . (The statement about third order poles can be evinced from $x_i^2 \cdot \eta$ or from $z \cdot \eta$ indifferently.) Therefore $\omega_C/\nu_*\omega_{\bar{C}}$ is spanned by

$$\begin{aligned} & \frac{dt_1}{t_1} - \frac{dt_m}{t_m}, \dots, \frac{dt_{m-1}}{t_{m-1}} - \frac{dt_m}{t_m}, \frac{dt_m}{t_m^2} \\ & \bar{\eta} = \frac{dt_1}{t_1^2} + \dots + \frac{dt_{m-1}}{t_{m-1}^2} - \frac{dt_m}{t_m^4}. \end{aligned}$$

In particular ω_C is generated by $\bar{\eta}$ as an \mathcal{O}_C -module.

In case II, we know that $\tilde{\mathfrak{m}}^3 \subseteq R$, so we have poles of third order at most. Let

$$\eta = c_1 \frac{dt_1}{t_1^3} + b_1 \frac{dt_1}{t_1^2} + a_1 \frac{dt_1}{t_1} \oplus \dots \oplus c_m \frac{dt_m}{t_m^3} + b_m \frac{dt_m}{t_m^2} + a_m \frac{dt_m}{t_m}.$$

From looking at $1 \cdot \eta$ we deduce $\sum_{i=1}^m a_i = 0$; from $x_i \cdot \eta$ we see $b_1 + b_m = 0$ (if $i = 1$), and $b_i + c_m = 0$ (if $i = 2, \dots, m-1$); finally from $x_i^2 \cdot \eta$ we have $c_1 + c_m = 0$ (if $i = 1$), and $c_i = 0$ (if $i = 2, \dots, m-1$). Therefore $\omega_C/\nu_*\omega_{\bar{C}}$ is spanned by

$$\begin{aligned} & \frac{dt_1}{t_1} - \frac{dt_m}{t_m}, \dots, \frac{dt_{m-1}}{t_{m-1}} - \frac{dt_m}{t_m}, \frac{dt_1}{t_1^2} - \frac{dt_m}{t_m^2}, \\ & \bar{\eta} = \frac{dt_1}{t_1^3} + \frac{dt_2}{t_2^2} + \dots + \frac{dt_{m-1}}{t_{m-1}^2} - \frac{dt_m}{t_m^3}. \end{aligned}$$

In particular ω_C is generated by $\bar{\eta}$ as an \mathcal{O}_C -module. □

Corollary 4.7. *The dualising sheaf has multi-degree $(0, \dots, 0, 2)$ (case I) and $(1, 0, \dots, 0, 1)$ (case II) respectively.*

Remark 4.8. It follows from this computation and Corollary 3.6 that the finiteness condition on automorphism groups, $H^0(\bar{C}, \Omega_{\bar{C}}^\vee(-\sum_{i=1}^n p_i)) = 0$, implies ampleness of $\omega_{\bar{C}}(\sum_{i=1}^n p_i)$.

Let us now go back to the proof of Proposition 4.2. Because $\phi|_{\mathcal{C}_0 \setminus Z}$ is the normalisation of $\bar{\mathcal{C}}_0$ at q , and, letting T_i be the tail of $\mathcal{C}_0 \setminus Z$ attached to Z at q_i , we know that $\omega_{\mathcal{C}/\Delta|T_i} = \omega_{T_i}(q_i)$ by adjunction, it follows from Lemma 4.6 that D has multiplicity 3 at the component of Z containing q_m in case I (resp. 2 at the components containing q_1 and q_m in case II), and 1 at all other components containing a q_i . Set $\mathcal{L} = \omega_{\mathcal{C}/\Delta}(D) = \phi^* \omega_{\bar{\mathcal{C}}/\Delta}$; we shall analyse the consequences of $\mathcal{L}|_Z = \mathcal{O}_Z$. We think of Z as being the union of a core K and a number ($\leq m$ by semistability) of rational trees.

Let d_A denote the multiplicity of the divisor D along the component A of Z . First, we claim that no component can appear with $d_A = 0$. Assume that this occurred along one of the rational trees. Call S ($S \simeq \mathbb{P}^1$) a component furthest from the core such that $d_S = 0$; R the one that precedes it, and T_1, \dots, T_h the ones that follow it (when sweeping the tree from the core) - so that $h \geq 1$ by the previous paragraph, and $d_{T_i} \geq 1$ by inductive assumption. Then, by adjunction,

$$\deg(\mathcal{L}|_S) = -2 + (h+1) + d_R + \sum d_{T_i} = 0,$$

which necessarily implies $h = 1$, and $d_R = d_{T_i} = 0$, contradicting the assumption. The case that S belongs to the core is similar (ω_S might only be more positive).

Let us now consider $d_S = 1$. We stick to the notation above; furthermore, there may be a number k of q_i , $i \in \{(1), 2, \dots, m-1\}$ in case II (resp. I), lying on S . Then, again by adjunction,

$$\deg(\mathcal{L}|_S) = -2 + d_R + \sum d_{T_i} = 0.$$

so either $d_R = 2$, $h = 0$ and $k \geq 1$ arbitrary, i.e. S is adjacent to $\overline{C \setminus Z}$; or $d_R = 1$, $h = 1$, and $d_{T_1} = 1$ (with k arbitrary). In the latter case, though, by repeating the same argument on T_1 etc., we would find an infinite chain in Z .

Remark 4.9. More generally, an analogous computation shows that, when balancing a component A of multiplicity d_A , all neighbouring components of multiplicity $d_A - 1$ can be safely ignored (at the same time, the number of such components is bounded only by m , due to the semistability of Z).

We now prove that $d_R > d_S$ holds in general for S on a rational tree. The preceding paragraphs deal with the cases $d_S = 0, 1$; we may therefore assume $d_S > 1$ (which in particular implies $0 \leq k \leq 2$). We have

$$\deg(\mathcal{L}|_S) = -2 + d_R - (d_S - 1)(h + k + 1) + \sum d_{T_i} = 0.$$

By proceeding inductively from leaves to root, we can assume that $d_S > d_{T_i}$, $i = 1, \dots, h$. We may therefore rewrite the previous equality as

$$d_R = (d_S - 1)(h + k + 1) - \sum d_{T_i} + 2 \geq (d_S - 1)(k + 1) + 2 = d_S + 1 + k(d_S - 1) > d_S.$$

In fact, we can prove as on [Smy11a, p.893] that $d_R = d_S + 1$, unless $d_S = 3$ and $q_m \in S$ (type I), or $d_S = 2$ and either q_1 or q_m (or both) are on S (type II). We introduce some terminology to describe the weighted dual graph of D .

Definition 4.10. A g -chain is a weighted graph that is a chain and such that the weight of two adjacent vertices differ by g . We call g the growth rate; the vertex with highest (resp. lowest) weight is called the root (resp. leaf) of the chain. An (a, g) -chain is a g -chain with leaf weight a . The chain C_1 can be attached to the chain C_0 by identifying the root of C_1 with a vertex of C_0 having the same weight. A 1-tree is obtained by attaching a number of $(1, 1)$ -chains among themselves.

Let us now look at a component S with $d_S = 2$ and at least one of q_1 and q_m attached to it. The balancing equation is

$$\deg(\mathcal{L}|_S) = -2 - (h + k + 1) + d_R + \sum d_{T_i} = 0,$$

with $k \in \{1, 2\}$. The preceding discussion implies that $d_{T_i} = 1$ for all $i = 1, \dots, h$, so $d_R = 3 + k$. If $k = 2$, i.e. both q_1 and q_m are on S - in which case they are indeed equidistant from the core -, then $d_R = 5$, and it can be shown inductively that the multiplicity of D increases by 3 for every step we make from S towards the core. The same holds in case I, with q_m attached to S and $d_S = 3$.

Finally, say $d_S = 2$ and only $q_1 \in S$. Then $d_R = 4$, and the growth rate along the chain that connects S to the core is 2, unless there is a component S' at which two 2-chains meet.

Definition 4.11. A 2-tree is obtained by attaching a number of $(1, 1)$ -chains to a $(2, 2)$ -chain. A 3-tree is obtained by attaching a number of $(1, 1)$ -chains either to a $(3, 3)$ -chain, or to a weighted graph itself obtained by attaching two $(2, 2)$ -chains to the leaf of a 3-chain.

From the preceding discussion it is clear that the weighted dual graph of D is obtained by attaching a number of 1-trees, and either (a) one 3-tree or (b) two 2-trees to the dual graph of the core K , weighted in an appropriate fashion.

Finally, let us look at the core K . Consider it as a one-pointed (case (a)), resp. two-pointed (case (b)) curve of genus two, by ignoring all the attachment points of the 1-trees (which works by Remark 4.9), and let $\bar{K} \in \bar{\mathcal{M}}_{2,1}$ (resp. $\bar{\mathcal{M}}_{2,2}$) be its stable model. The following can happen:

- (1) K is a smooth curve of genus two. In case (a), let R be the component adjacent to the core along the 3-tree, and let $x = R \cap K$; then $d_K = d_R + 3$ by balancing R . Now balancing K gives

$$\omega_{\mathcal{C}/\Delta}(d_R R + d_K K)|_K = \omega_K(d_R x - (d_R + 2)x) \simeq \mathcal{O}_K,$$

which admits a solution if and only if K is Weierstrass. Similarly case (b) can be balanced if and only if K is conjugate.

- (2) K contains two distinct subcurves of genus one E_1 and E_2 . We start by solving the balancing equation on one of them, say $E = E_1$. If all but one




FIGURE 4. Examples of Z with two genus one components (the white dots). The numbers indicate the multiplicity of D . On the left, one 3-tree attached to E_1 ; on the right, two 2-trees attached to the same bead of the rational chain separating E_1 from E_2 .

of the neighbouring components have multiplicity $d_E - 1$, then the last one is forced to have multiplicity $d_E - 1$ as well (by degree reasons). The case that all but two neighbouring components have multiplicity $d_E - 1$ occurs when either one 2-tree or one 3-tree (and exactly one) is attached to E at x ; let F be the other component with undetermined multiplicity, which lies between E_1 and E_2 (possibly $F = E_2$), and let $E \cap F = \{y\}$. The case of a 2-tree forces $d_F = d_E$ by degree reason, but then we are left to solve $x \sim y$ in $\text{Pic}(E)$, which is impossible; on the other hand, the case of a 3-tree imposes $d_F = d_E + 1$ and $2x \sim 2y$ in $\text{Pic}(E)$, i.e. K is Weierstrass. By the same token, the two 2-trees have two hit the same genus one curve, say E_1 , in nodes x_1, x_2 such that $x_1 + x_2 \sim 2y$ and $d_F = d_E + 1$.

Assume now that there is a chain of rational curves S_i lying between E_1 and E_2 in K , and one of the special trees connects to one of the S_i ; in case (b), then, both 2-trees must connect to (possibly different) S_i , by the previous paragraph. Furthermore, the growth rate along the rational chains at E_1 and E_2 has to be 1. This in turn implies that, in case (b), the growth rate along the chain separating the two 2-trees is 0. In particular, the two 2-trees are attached to components with the same multiplicity for D , so q_1 and q_m are equidistant from the core. See Figure 4 for some examples.

- (3) $\bar{K} \in \Delta_{irr}$, i.e. K contains an irreducible subcurve of arithmetic genus one E , with two points y_1 and y_2 on E that are joined in K by a (possibly empty) rational chain. We see as above that either a 3-tree is attached to a point $x \in E$ satisfying $2x \sim y_1 + y_2$ in $\text{Pic}(E)$, or two 2-trees are attached to $x_1, x_2 \in E$ satisfying $x_1 + x_2 \sim y_1 + y_2$ in $\text{Pic}(E)$, or the rational chain is not empty and all the distinguished trees are attached to it. In this case, solve the balancing equation on E : let $d = d_E$, d_1 and d_2 be the multiplicities of the rational components attached to y_1 and y_2 respectively; then either $d_1 = d_2 = d - 1$, or $d_1 = d - 1 + k$, $d_2 = d - 1 - k$ and $r_1 - r_2$ is k -torsion in $\text{Pic}(E)$. But, by chasing the balancing equation along the rational necklace, we find that, if $k \geq 1$, then the growth rate increases when passing through





FIGURE 5. K is a genus one curve E with a rational bridge R . Left column, one 3-tree; right, two 2-trees. Above: the special trees affere to E ; below: they affere to R - note that R is necessarily symmetric in this case.

a distinguished bead, so that ultimately $d - 1 - k = d_2 > d_1 d - 1 + k$, which is absurd. So the only possibility is to have a rational chain symmetric with respect to the distinguished beads, namely: in case (a) the two pieces of the rational chain standing between the special bead and E have the same length, and in case (b) the distance shortest path between a special bead and E is the same for the two special beads. See Figure 5.


- (4) Finally, we consider the case that K has geometric genus 0. It is easy to see that, if a special tree afferes to a stable component, the only chance that the balancing condition may be satisfied is that there are two 2-trees and they affere to different stable components; the semistable chains have arbitrary length and D has the same multiplicity along every component of the core; see the upper left corner of Figure 6. There remain three possibilities for the dual graph, according to how the distinguished components (denoted by R) and the other stable components (denoted by A and B) distribute themselves. Denoting by t the growth rate along various rational chains, we find that in case (a) and (b).1 balancing along A or B is equivalent to $\sum_i t_i = 1$. Assume $t_1 \geq 0$; then $d_A > d_B$, therefore $t_2, t_3 > 0$, which contradicts $\sum_i t_i = 1$.

FIGURE 6. When K consists of rational curves.

Similarly, if $t_1 \leq -2$, then $d_A < d_B$, therefore $t_2, t_3 < 0$, which makes $\sum_i t_i = 1$ again impossible. We find only one solution with $t_1 = -1$ and $t_2 = t_3 = 0$ - notice that it is a degeneration of the case considered in the previous point. On the other hand, in case (b).2, we find $\sum_i t_i = 1$ when balancing B , and $\sum_i t_i = -1$ when balancing A , which is a contradiction.

This concludes the proof of Propositions 4.2(2) and 4.3(2). \square

Remark 4.12. There is a stable 2-pointed curve that arises as a solution of the balancing equation, yet is not conjugate, namely:



In this case, the line bundle $\omega_{\mathcal{C}/\Delta}(D)$ trivial along Z might not be semiample.

Proof. (of Proposition 4.4) By blowing down some rational tails outside Z , we can assume that $\mathcal{C}_0 \setminus Z = \sqcup_{i=1}^m T_i$ with each $T_i \simeq \mathbb{P}^1$. The image of $\Sigma_i(0)$ and $\Sigma_j(0)$ may now coincide for $i \neq j$. The total space of the curve can still be assumed to be smooth. By abuse of notation, we denote the resulting family of pointed

curves by $(\mathcal{C}, \Sigma_1, \dots, \Sigma_n)$. By assumption on the shape of Z , we can find an effective Cartier D supported on Z such that $\mathcal{L} = \omega_{\mathcal{C}/\Delta}(D + \Sigma)$ be trivial on Z and relatively ample elsewhere (both on T_i and on the generic fiber). Consider a second line bundle $\mathcal{L}' = \mathcal{O}(2\Sigma_1 + \Sigma)$. Since we assumed Σ_1 to be Weierstrass, $\mathcal{L}_\eta \simeq \mathcal{L}'_\eta$. On the other hand it is easy to see that the multi-degrees of \mathcal{L}_0 and \mathcal{L}'_0 coincide; it follows from the separatedness of $\text{Pic}_{\mathcal{C}/\Delta}^0 \rightarrow \Delta$ (see [?, p. 136] or [BLR90, §9.4]) that \mathcal{L} and \mathcal{L}' are isomorphic line bundles, so that in particular \mathcal{L} is trivial on a neighbourhood of Z . Observe now that

$$R^1\pi_*\mathcal{L}(-D) = R^1\pi_*\omega_{\mathcal{C}/\Delta}(\Sigma) = 0$$

by semistability, hence $\pi_*\mathcal{L} \rightarrow \pi_*(\mathcal{L}|_D) = \pi_*\mathcal{O}_D$ which contains the constants, showing that \mathcal{L} is semiample along Z (that it is along the T_i is easier). We therefore have a contraction

$$\mathcal{C} \xrightarrow{\phi} \overline{\mathcal{C}} = \underline{\text{Proj}}_\Delta \left(\bigoplus_{n \geq 0} \pi_*\mathcal{L}^{\otimes n} \right) \rightarrow \Delta$$

associated to \mathcal{L} . The proof that $\overline{\mathcal{C}} \rightarrow \Delta$ is a flat family of Gorenstein curves goes along the lines of [Smy11a, Lemma 2.13] or [RSW17a, Proposition 3.7.3.1]. It is clear from the classification that it contains a type I_m -singularity in the central fiber. The proof of Proposition 4.5 is entirely analogous. \square

Caveat 1. It seems not to be true in general that, if we have a family of semistable curves $\pi: \mathcal{C} \rightarrow \Delta$ over a discrete valuation ring, and a line bundle \mathcal{L} on \mathcal{C} that is trivial on a higher genus subcurve Z of \mathcal{C}_0 , and π -ample elsewhere, then \mathcal{L} is π -semiample, i.e. relatively generated by global sections. It will rather depend on the family, and, in particular, the assumption that the total space of \mathcal{C} is regular along Z seems to be essential. This seems not to be an issue in positive characteristic, thanks to a result of S. Keel [Kee99], so assume $\text{char}(\mathbf{k}) = 0$; we give a counterexample using the theory of limit linear series: we produce a linear series that can be smoothed while having basepoints along a Weierstrass tail.² For any such smoothing, the corresponding line bundle \mathcal{L} is not globally generated along Z - though we do not know how the powers of \mathcal{L} behave.

Let X_0 be the nodal curve obtained by attaching $R \simeq \mathbb{P}^1$ to a Weierstrass point q of a smooth genus two curve Z . Choose $d \gg 0$ ($d \geq 5$ is enough), and let us study the moduli space of complete linear systems of degree d on (smoothings of) X_0 ; with $r = d - 2$, the Brill-Noether number is $\rho = 2$ (the dimension of the Jacobian). On the other hand, assume that the R -aspect of the lls has $\mathcal{L}_Y|_Z \simeq \mathcal{O}_Z$; then the Z -aspect has $\mathcal{L}_Z|_Z \simeq \mathcal{O}_Z(dq)$, whose vanishing sequence is $\alpha_Z(q) = \{0, 1, \dots, d-4, d-2, d\}$, from which we deduce for the complementary aspect $\alpha_R(q) \geq \{0, 2, 4, 5, \dots, d\}$. We want to show that all such aspects are

²It has been pointed out to us by F. Carocci that a similar but easier computation can be carried out for a genus one tail as well. This shows that the regularity assumption of [Smy11a, Lemma 2.13] is necessary.

smoothable, by appealing to the Regeneration Theorem [HM98, Theorem 5.41]. Notice that in the case at hand we have a choice of a two-dimensional subspace of $\langle 1, t, t^2, t^3 \rangle_{\mathbf{k}} \subseteq H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(d))$ meeting the subspace $\langle t^2, t^3 \rangle_{\mathbf{k}}$ non-trivially, i.e. the locus in $\mathrm{Gr}(1, \mathbb{P}^3)$ of lines meeting a fixed line ℓ , which is a Schubert cycle of dimension 3. We therefore need to put X_0 in a family over a base B of dimension 1 at least. We shall do so by considering the family X obtained by attaching R to a moving point of Z , so that X_0 is the fiber of X over $q \in Z$.

Let us start by examining the other possibilities for $\mathcal{G}_d^{d-2}(X_0)$: the R -aspect can in fact restrict to any line bundle of degree 0 on Z , which we are going to write as $\mathcal{O}_Z(p_1 + p_2 - 2q)$ for two moving points p_1, p_2 on Z (think of them as coordinates on $\mathrm{Pic}(Z)$). Then $\mathcal{L}_Z = \mathcal{O}_Z((d-2)q + p_1 + p_2)$.

$\subseteq \mathrm{Pic}(Z)$	dim	$\alpha_{Z;d-3,d-2}$	$\alpha_{R;0,1}$	$\subseteq \mathbb{P}H^0(R, \mathcal{O}_R(d))$	dim
$p_1 + p_2 \sim 2q$	0	$\{d-2, d\}$	$\geq \{0, 2\}$	$\{\ell' \in \mathrm{Gr}(1, \mathbb{P}^3) \ell' \cap \ell \neq \emptyset\}$	3
$2q \approx p_1 + p_2 \geq q$	1	$\{d-3, d-1\}$	$\geq \{1, 3\}$	\mathbb{P}^1	1
$p_1 + p_2 \not\approx q$	2	$\{d-3, d-2\}$	$\{2, 3\}$	pt	0

If we now let q vary in $B \simeq Z$, we may generically assume that it is not Weierstrass. We then find the following:

$\subseteq \mathrm{Pic}(X)$	dim	$\alpha_{Z;d-3,d-2}$	$\alpha_{R;0,1}$	$\subseteq \mathbb{P}H^0(R, \mathcal{O}_R(d))$	dim
$p_1 + p_2 \sim \omega_Z$	$0+1$	$\{d-2, d-1\}$	$\geq \{1, 2\}$	$(\mathbb{P}^2)^*$	2
$p_1 + p_2 \sim 2q$	$0+1$	$\{d-3, d\}$	$\geq \{0, 3\}$	\mathbb{P}^2	2
$\omega_Z, 2q \approx p_1 + p_2 \geq q$	$1+1$	$\{d-3, d-1\}$	$\geq \{1, 3\}$	\mathbb{P}^1	1
$p_1 + p_2 \not\approx q$	$2+1$	$\{d-3, d-2\}$	$\{2, 3\}$	pt	0

We conclude that $\mathcal{G}_d^{d-2}(X/B)$ has pure dimension 3, and we may therefore apply the Regeneration Theorem to deduce that all lls with $\mathcal{L}_{R|Z} \simeq \mathcal{O}_Z$ - in particular those with base-locus Z - are smoothable.

5. THE NEW MODULI FUNCTORS

The following generalises [Smy11a, Definition 3.4].

Definition 5.1. Let (C, p_1, \dots, p_n) be a reduced curve, marked by smooth points. For a nodally attached subcurve $D \subseteq C$, we define its *level* as

$$\mathrm{lev}(D) = |D \cap \overline{C \setminus D}| + |\{p_1, \dots, p_n\} \cap D|.$$

We say a Gorenstein curve C is *minimal* if it contains no node x such that the normalisation of C at x consists of two connected components, one of which has genus zero. When C has arithmetic genus one, this is the same as saying that C contains no separating nodes. Recall [Smy11a, Lemma 3.3].

Lemma 5.2. *A minimal Gorenstein curve Z of arithmetic genus one can be: a smooth elliptic curve; a ring of $r \geq 1$ copies of \mathbb{P}^1 ; or an elliptic m -fold point whose normalisation is the disjoint union of m copies of \mathbb{P}^1 . In any case $\omega_Z \simeq \mathcal{O}_Z$.*

We may similarly describe minimal (sub)curves of genus two.

Lemma 5.3. *A minimal Gorenstein curve Z of arithmetic genus two can be:*

- (1) a smooth curve of genus two;
- (2) the union of two minimal Gorenstein curves of genus one, E_1 and E_2 , nodally separated by a (possibly empty) rational chain R ;
- (3) the union of a minimal Gorenstein curve of genus one E , and a (possibly empty) rational chain R , along two distinct nodes;
- (4) the union of two copies of $(\mathbb{P}^1, 0, 1, \infty)$ with three (possibly empty) rational chains R_0, R_1, R_∞ joining the homonymous points;
- (5) an elliptic m -fold point x whose normalisation is the disjoint union of either $m-1$ copies of \mathbb{P}^1 (i.e. two branches of x coincide), or $m-1$ copies of \mathbb{P}^1 and a minimal Gorenstein curve of genus one (i.e. Z contains two genus one subcurves sharing a branch);
- (6) or a singularity of genus two with m -branches, whose normalisation is the disjoint union of m copies of \mathbb{P}^1 .

Remark 5.4. In both cases (5) and (6) there are special branches supporting the degree of ω_Z (compare with Definition 2.3 and Corollary 4.7; recall that the restriction of the dualising sheaf to a component introduces a twist by the conductor ideal, see Noether's formula [Cat82, Proposition 1.2]). Notice that the notion of conjugate points is not always intrinsic to the curve.

Similarly to [Smy11a, Corollary 3.2, Lemma 3.5] we can prove the following.

Lemma 5.5. *Let (C, p_1, \dots, p_n) be a pointed semistable curve of arithmetic genus two, with minimal genus two subcurve Z . For every subcurve $Z' \subseteq C$ of genus two, we have an inclusion $Z \subseteq Z'$ and $\text{lev}(Z) \leq \text{lev}(Z')$.*

We finally come to the definition of m -stability for curves of genus two.

Definition 5.6. Fix positive integers $m < n$. Let (C, p_1, \dots, p_n) be a connected, reduced, complete curve of arithmetic genus two, marked by smooth distinct points. We say that C is m -stable if:

- (1) C is Gorenstein with only: nodes; elliptic l -fold points, $l \leq m+1$; type $I_{\leq m}$, type $II_{\leq m}$, and dangling II_{m+1} singularities of genus two, as singular points.
- (2) If Z is a connected subcurve of arithmetic genus two, then $\text{lev}(Z) > m$.
- (3) If E is a nodally attached subcurve of arithmetic genus one, $\text{lev}(E) > m+1$.
- (4) $H^0(C, \Omega_C^\vee(-\sum_{i=1}^n p_i)) = 0$.
- (5) If C contains a singularity of genus two, or an elliptic l -fold with a self-branch, p_1 is connected (through a rational chain) to one of the special branches (see Remark 5.4). If C contains two genus one subcurves sharing a branch, and E_1 has level less than $m+2$, then p_1 is connected to E_1 .

Remark 5.7. The definition is not \mathfrak{S}_n -symmetric. In the arguments below, we exploit the asymmetry to write the dualising line bundle of a genus two (sub)curve Z as $\omega_Z \simeq \mathcal{O}_Z(q_1 + \bar{q}_1)$, where q_1 is the point of Z closest to p_1 , and \bar{q}_1 its conjugate, possibly depending on a one-parameter smoothing. Compare with the situation in genus one, where the dualising line bundle of a minimal Gorenstein curve is trivial (all smooth points are non-special).

Remark 5.8. If there is a subcurve of genus one, condition (3) and condition (4) jointly imply condition (2). Indeed, from Corollary 3.6 we see that $\text{lev}(Z) \geq \text{lev}(E) - 1$. The only cases (up to relabelling) in which the level drops by one are: when $Z = (E, p_1, \dots, p_{l-2}, q_1, q_2) \sqcup_{\{q_1, q_2\}} (\mathbb{P}^1, q_1, q_2, p_{l-1})$; and when $Z = (E_1, p_1, \dots, p_{l-1}, q) \sqcup_q (E_2, q)$, where all the E have genus one.

Lemma 5.9 (boundedness). *If (C, p_1, \dots, p_n) is an m -stable curve of genus two, the N -th power of $A = \omega_C(\sum_{i=1}^n p_i)$ is very ample for every $N > 2 + 8(m+1)$.*

Proof. It is enough to show that, for every pair of points $p, q \in C$ (possibly equal):

- (1) *basepoint-freeness*: $H^1(C, A^{\otimes N} \otimes I_p) = 0$;
- (2) *separating points and tangent vectors*: $H^1(C, A^{\otimes N} \otimes I_p I_q) = 0$.

By Serre duality we may equivalently show that $H^0(C, \omega_C \otimes A^{-N} \otimes (I_p I_q)^\vee) = 0$. Let $\nu: \tilde{C} \rightarrow C$ be the normalisation, and let $\nu^{-1}(p) = \{p_1, \dots, p_h\}$, $\nu^{-1}(q) = \{q_1, \dots, q_k\}$, with $h, k \leq m+1$. It follows from Proposition 2.2 (and [Smy11a, Proposition A.3]) that $\nu_* \mathcal{O}_{\tilde{C}}(-D) \subseteq I_p I_q$ for $D = 4(\sum_{i=1}^h p_i + \sum_{j=1}^k q_j)$ (note that $\deg(D) \leq 8(m+1)$); furthermore, the quotient is torsion, therefore, by applying $\mathcal{H}om(-, \mathcal{O}_C)$ and adjunction, we find $(I_p I_q)^\vee \subseteq \nu_* \mathcal{O}_{\tilde{C}}(D)$. It is thus enough to show that $H^0(\tilde{C}, \mathcal{O}_{\tilde{C}}(D) \otimes \nu^*(\omega_C \otimes A^{-N})) = 0$. Finally, $\nu^* \omega_C$ has degree at most two, and $\nu^* A$ has degree at least one on any branch of \tilde{C} , hence it is enough to take $N > 2 + 8(m+1)$. \square

Lemma 5.10 (deformation openness). *Let $(\mathcal{C}, \sigma_1, \dots, \sigma_n) \rightarrow S$ be a family of curves over a Noetherian base scheme with n sections. The locus*

$$\{s \in S \mid (\mathcal{C}_s, \sigma_1(s), \dots, \sigma_n(s)) \text{ is } m\text{-stable}\}$$

is Zariski-open in S .

Proof. Having connected fibers which are Gorenstein curves of arithmetic genus two is an open condition (see for example [Sta19, Tag 0E1M]). Only singularities of genus zero (nodes), one (elliptic l -folds), and two may then occur.

The case $m = 1$ deserves special attention. In this case, that condition (1) is open follows from acknowledging that $I_1 = A_4$, $I_2 = A_5$, while tacnodes, cusps, and nodes are A_3 , A_2 , and A_1 -singularities respectively, and from Grothendieck's results on the deformation theory of ADE singularities (see Theorem 2.5 above).

The case $m \geq 2$ simply follows from upper semicontinuity of embedded dimension and the fact that we have exhausted all possible Gorenstein singularities of genus ≤ 2 , and embedding dimension $\leq m+1$.

Condition (4) translates to: the locus where the automorphism group is unramified is open in the base. Homogeneity can be used to prove that being unramified, which is open in the source, is also open in the target, for the structural morphism of a group scheme; see the end of the proof of [Smy11a, Lemma 3.10].

The other conditions are topological, hence constructible. With Noetherian assumptions, it is enough to check their openness over the spectrum of a discrete valuation ring. Assume that the geometric generic fiber $C_{\bar{\eta}}$ contains two genus

one subcurve $E_{1,\bar{\eta}}$ and $E_{2,\bar{\eta}}$; their closures E_1 and E_2 in \mathcal{C} are then flat families of genus one curves over Δ . If $E_{1,\bar{\eta}}$ and $E_{2,\bar{\eta}}$ are disconnected, then so are E_1 and E_2 , by local constancy of the number of connected components (from the Zariski decomposition and [Sta19, Tag 0E0D]). If $E_{1,\bar{\eta}}$ and $E_{2,\bar{\eta}}$ are joined by a disconnecting node $q_{\bar{\eta}}$, then so are $E_{1,0}$ and $E_{2,0}$; indeed, the unique limit of $q_{\bar{\eta}}$ must be a singular point of the projection, but cannot be any worse than a node (because we have already exhausted all the available genus). Finally, if $E_{1,\bar{\eta}}$ and $E_{2,\bar{\eta}}$ share a branch, then so do $E_{1,0}$ and $E_{2,0}$; on the other hand, if $E_{i,\bar{\eta}}$ has more than one branch, then so does E_i . Similarly, if $C_{\bar{\eta}}$ contains only one subcurve of genus one, with two nodes joined by a rational chain, so does C_0 . The upshot of this discussion is that

$$|E_{i,\bar{\eta}} \cap \overline{C_{\bar{\eta}} \setminus E_{i,\bar{\eta}}}| = |E_{i,0} \cap \overline{C_0 \setminus E_{i,0}}|.$$

The number of markings on E_i is also constant. Hence we can deduce condition (3) for $C_{\bar{\eta}}$ from the same condition on C_0 . Condition (2) follows in this case from Remark 5.8; it can be proved analogously when there is no subcurve of genus one.

Finally, suppose that $C_{\bar{\eta}}$ has a genus two singularity, then so does C_0 . The (union of the) distinguished branch(es) $E_{\bar{\eta}}$ of $C_{\bar{\eta}}$ is a genus one singularity, and so is its limit E_0 in C_0 . It has to contain the distinguished branch(es) of C_0 , because any subcurve not containing them has genus zero; therefore, by assumption, E_0 contains $p_{1,0}$. Then also $E_{\bar{\eta}}$ contains $p_{1,\bar{\eta}}$, because the markings are contained in the non-singular locus of the curve. Similarly, $C_{\bar{\eta}}$ has a genus one singularity with a self-branch, the limit of such a branch is a genus one subcurve E_0 of C_0 ; the latter may very well acquire a genus two singularity, but E_0 will contain the special branches of it, so it will be connected to p_1 . We conclude as above. The case that $C_{\bar{\eta}}$ contains a genus one subcurve of low level is analogous. We have thus proved that condition 5 is open. \square

Definition 5.11. We shall denote by $\overline{\mathcal{M}}_{2,n}^{(m)}$ the moduli functor of n -pointed m -stable curves of genus two.

It follows from the previous lemmas and standard arguments that $\overline{\mathcal{M}}_{2,n}^{(m)}$ is represented by a Deligne-Mumford stack of finite type.

Proposition 5.12 (Valuative criterion of properness for $\overline{\mathcal{M}}_{2,n}^{(m)}$). *Given a smooth n -pointed curve of genus two C_{η} over a discrete valuation field $\eta = \text{Spec}(K) \hookrightarrow \Delta$, there exists a finite base-change $\Delta' \rightarrow \Delta$ after which C_{η} can be completed to an m -stable curve over Δ' . Two such models are always dominated by a third one.*

Proof. Existence of limits. We start with a smooth n -pointed curve of genus two over a discrete valuation field. By the semistable reduction theorem [DM69, Corollary 2.7], we may find a finite base-change $\Delta' \rightarrow \Delta$ and a semistable curve $\mathcal{C}' \rightarrow \Delta'$ with regular total space, such that its generic fiber is isomorphic to the pullback of the curve we started with. By Castelnuovo's criterion, we may further assume that the central fiber contains no rational tails.

We check whether p_1 is Weierstrass or not: in the former case, change base with $\pi'' \mapsto (\pi')^3$ and resolve; in the latter, mark \mathcal{C}' with an extra section \bar{p}_1 given by the closure of the conjugate point $\bar{p}_1(\eta)$ (unless it coincides with one of p_1, \dots, p_n ; if it coincides with p_1 , we have a Weierstrass point indeed), then change base with $\pi'' \mapsto (\pi')^2$ and resolve. We drop the primes from notation. \mathcal{C}_0 is now marked with a(n extra) smooth point \bar{p}_1 . The base-change is a technical expedient we find useful in the forthcoming construction.

We claim there is a unique genus two subcurve Z of \mathcal{C}_0 that satisfies the shape requirements of Proposition 4.2 (resp. 4.3) - or consists of two disjoint balanced subcurves of genus one - guaranteeing the curve we obtain by contracting Z has bounded above singularities and bounded below level as in conditions (1),(2), and (3) of Definition 5.6.

We think of this process as drawing a family of expanding circles on the dual graph (except, they are neither exactly circles, nor always expanding). Note that we may at any point blow-up the curve at a marking on the central fiber, and consider the strict transform of the corresponding section; thus markings can effectively be considered as infinite legs in the dual graph. We are going to contract the strict interior of the circle; note that the number of branches of the resulting singularity is determined by the inner valence of the circle, and the level by the outer valence.

For simplicity, we start by examining the case that the core of \mathcal{C}_0 is irreducible, and p_1 is Weierstrass. Step 0: if the core K has level $\geq m+1$, then the curve is already m -stable. Otherwise, draw a first circle comprising K , and reaching every second closest rational component along any rational tree attached to K , except for the tree containing p_1 . Note that the inner valence is exactly $\text{lev}(K)$, thanks to the base-change we have performed earlier, and in particular it is no larger than m ; on the other hand, semistability implies that the outer valence is non-decreasing - it is important to consider markings as infinite legs for this -: if it is $\geq m+1$ we stop, otherwise we repeat the process. Calling K_1 the union of the components strictly inside the first circle, \mathcal{C}_0 is the union of K_1 and a number of rational trees; if the outer valence at the first step is still $\leq m$, we enlarge the circle by reaching one step further along the rational tree containing p_1 , and three steps along all the other ones. Because a circle of very large radius has both inner and outer valence equal to $n > m$, by increasing the radius step by step we will eventually reach level $m+1$ or higher. If we stop at the l -th step, the line bundle we will use to perform the contraction is

$$\begin{aligned} \mathcal{L} = \omega_{\mathcal{C}/\Delta}(3lK + \sum_{R \in [p_1, K]} [3l - 3 \text{dist}(R, K)]_+ R + \sum_{R \notin T_1} [3l - \text{dist}(R, K)]_+ R + \\ \sum_{R \in T_1, R \notin [p_1, K]} [3l - 3 \text{dist}(T_1 \wedge T_R, K) - \text{dist}(R, T_1 \wedge T_R)]_+ R + \sum_{i=1}^n p_i) \end{aligned}$$

where T_1 is the rational tree (connected component of $\mathcal{C}_0 \setminus K$) containing p_1 , T_R the one containing R , $T_1 \wedge T_R$ their common component furthest from the core, dist is the distance on the dual graph, and $[k]_+ = \max\{0, k\}$ for any integer k . If \mathcal{L} differs from $\omega_{\mathcal{C}/\Delta}^{\log}$ by a vertical divisor whose support is Z , the shape prescription being satisfied by construction, it follows from Proposition 4.4 that \mathcal{L} is π -semiample, and the contraction it produces contains a singularity of type I with p_1 afferring to the special branch. Note that \mathcal{L} contracts as well the semistable rational components that are disjoint from Z , hence the resulting curve has no (infinitesimal) automorphisms. The level condition is satisfied by construction, therefore $\overline{\mathcal{C}}_0$ is m -stable.

The case that the core is irreducible and p_1 is not Weierstrass is dealt with in a similar fashion. Remember that in this case we have constructed a conjugate section \bar{p}_1 ; this is an auxiliary marking that will be forgotten in the end, and should not be taken into account when computing the level. At every step we draw a larger circle by including one more component along the trees containing p_1 and \bar{p}_1 , and two more along every other tree; the inner valence of the new circle is the same as the outer valence of the old one, thanks to the $2 : 1$ base-change. At the l -th step we are going to use the line bundle

$$\begin{aligned} \mathcal{L} = \omega_{\mathcal{C}/\Delta}(2lK + \sum_{\substack{R \in [p_1, K] \\ \text{or } [\bar{p}_1, K]}} [2l - 2 \text{dist}(R, K)]_+ R + \sum_{R \notin T_1, \bar{T}_1} [2l - \text{dist}(R, K)]_+ R + \\ \sum_{\substack{R \in T_1, R \notin [p_1, K] \text{ or} \\ R \in \bar{T}_1, R \notin [\bar{p}_1, K]}} [2l - 2 \text{dist}(T_1 \wedge T_R, K) - \text{dist}(R, T_1 \wedge T_R)]_+ R + \sum_{i=1}^n p_i + \bar{p}_1) \end{aligned}$$

to contract the strict interior of the circle. It follows from Proposition 4.5 that \mathcal{L} is π -semiample, and the associated contraction contains a singularity of type II with p_1 afferring to one of the twin branches. Note that a II_{m+1} singularity will occur only if the level is m and \bar{p}_1 does not coincide with any other special point, so that one of the twin branches remains dangling after forgetting \bar{p}_1 . Again, the stability condition is satisfied by construction.

Suppose next that the minimal subcurve of genus two Z contains two subcurves of genus one; call E_1 and E_2 the minimal such, and assume that p_1 afferes to E_1 in a point that is 2-torsion with respect to the node separating E_1 from E_2 . We start by drawing expanding circles around E_2 until the level condition for it is satisfied, and then we do the same for E_1 , so that (5) holds; observe, though, that as soon as the two circles touch, the contraction will not have two distinguished genus one subcurves anymore, therefore we need to check (2) rather than (3).

- If $\text{lev}(E_2) \geq m + 2$ is attained before the circle around E_2 gets to touch E_1 , take the next $l_2 \equiv 2 \pmod{3}$ (thus “undoing” the $3 : 1$ base-change, which was




FIGURE 7. An instance of the critical step, when we pass from contracting the two genus one subcurves separately, to contracting the genus two subcurve as a whole. On the left, the contraction produces a tacnode with a genus one branch, having $\text{lev}_2 = 3$. On the right, the contraction produces a singularity of type I_3 . Note that in the meantime we performed a blow-up at p_1 .

unnecessary in this case), then contract the inner disc by the line bundle

$$\mathcal{L}_2 = \omega_{\mathcal{C}/\Delta}((l_2 + 1)E_2 + \sum [l_2 + 1 - \text{dist}(E_2, R)]_+ R + \sum_{i=1}^n p_i).$$

Smyth’s contraction lemma [Smy11a, Lemma 2.13] applies, so that E_2 is contracted to an elliptic l -fold point q_2 ($l \leq m + 1$). Consider now E_1 . If $\text{lev}(E_1) \leq m + 1$, start drawing expanding circles around it. Either level $\geq m + 2$ can be reached before touching the singularity at q_2 , or, by contracting the maximal balanced subcurve of genus one containing E_1 and not q_2 , we produce a curve having two genus one singularities that share a branch. Note that p_1 affers to the only genus one subcurve that may have level $\leq m + 1$.

- Otherwise, one step before including E_1 , we may contract the disc around E_2 to produce a genus one singularity with a genus one branch. If $\text{lev}_2 \leq m$ at this point, we need to contract the whole genus two subcurve. What happens at this critical step is that the multiplicity of D along E_1 grows from 0 to 3, hence D will be supported three steps further along each rational tail departing from E_2 . Because the critical step is $\equiv 1 \pmod{3}$, the length of the rational tails is and remains $\equiv 2 \pmod{3}$, hence we perform only one “meaningful” step forward (i.e. on the original central fiber, before changing base), thanks to the preliminary base-change. See Figure 7. In particular, the inner valence of the disc will be $\leq m$. We may now proceed as in the irreducible case, expanding the circle at every step by 1 along T_1 and by 3 along all other rational tails.

The case that the central fiber contains two subcurves of genus one, and p_1 affers to a non-Weierstrass point, is analogous: it is enough to replace the number 3 by the number 2, and 2 by 1, in the previous argument. The only novelty is, it can happen that p_1 is equidistant from E_1 and E_2 , afferring to the rational chain joining them. In this case we start by expanding a circle around the one with

the lowest level; if they have the same level, expand them simultaneously. If at a later stage p_1 becomes closer to one of the two circles, proceed as above.

When the core consists of an elliptic curve E with a rational bridge R , and p_1 affers to a Weierstrass point - either on E or on R -, we have noticed above (see the end of Remark 4.1), that R comprises an odd number $k = 2h + 1$ of rational components. We start as above by enlarging a balanced circle around E in order to establish the level condition (3); once again there is a critical step (after which we contract a genus two subcurve, and check (2) instead) when the circle touches itself along R , and we observe that this happens after the $(3h+2)$ -th step, therefore extending the circle by 1 along the tail containing p_1 , and by 3 on all the other ones, only makes one meaningful step. The case of a non-Weierstrass point on the elliptic bridge is analogous.

Finally, the case that the central fiber has geometric genus zero is dealt with as if the core was irreducible, since we do not introduce any genus zero singularity other than the node (the remaining ones are not Gorenstein).

Uniqueness of limits. By the semistable reduction theorem, there is a diagram

$$\begin{array}{ccc}
 & \mathcal{C}^{ss} & \\
 \phi \swarrow & & \searrow \phi' \\
 \mathcal{C} & & \mathcal{C}' \\
 & \searrow & \swarrow \\
 & \Delta &
 \end{array}$$

extending the isomorphism between the generic fibers, where \mathcal{C}^{ss} has semistable central fiber and regular total space.

Claim 1: If \mathcal{C}'_0 has only singularities of genus $\leq i$ ($i = 0, 1$), then so does \mathcal{C}_0 . First, assume that \mathcal{C}'_0 has only nodes. If \mathcal{C}_0 has a singular point x of genus one, $E := \phi^{-1}(x)$ is an *unmarked* subcurve of arithmetic genus one and level $\leq m + 1$ of \mathcal{C}_0^{ss} . Then so is $\phi'(E)$: indeed, ϕ' being a contraction, it has connected fibers, which excludes the possibility that ϕ' lowers the genus of E by realising a finite cover of a line. This contradicts the m -stability of \mathcal{C}' . We may argue similarly if x is a genus two singularity with $\leq m$ branches. On the other hand, if x is dangling Π_{m+1} , there is a -1 -curve R adjacent to $\phi^{-1}(x)$; ϕ' must contract R by DM stability of \mathcal{C}' , hence $\phi'(\phi^{-1}(x))$ is again a genus two curve of level $\leq m$.

Assume now that \mathcal{C}'_0 has at worst singularities of genus one, while \mathcal{C}_0 has a singularity x of genus two; the case of a dangling Π_{m+1} can be excluded as above. Then $\mathcal{C}_0^{ss} = Z \cup R_1 \cup \dots \cup R_l$, with $Z = \phi^{-1}(x)$ and $l \leq m$. If Z has geometric genus two, or is irreducible of geometric genus one, $\phi'(Z)$ violates the m -stability of \mathcal{C}' . If Z contains a unique subcurve E of genus one, with a rational bridge R , then at least one of R_1, \dots, R_l must be connected to R , for otherwise \mathcal{C}'_0 - which is obtained by contracting a balanced subcurve around E , not including the entire R - would have a positive dimensional automorphism group (scaling a semistable

component of R). Therefore $\text{lev}(E) \leq (l-1) + 2 \leq m+1$. Similarly, if Z contains two subcurves of genus one E_1 and E_2 , then $(\text{lev}(E_1) - 1) + (\text{lev}(E_2) - 1) \leq l$, hence at least one of the two has level $\leq m+1$. In any case, $\phi'(E)$ contradicts the m -stability of \mathcal{C}' .

Claim 2: We may assume that \mathcal{C}^{ss} contains either no -1 -curve, or only one, which is contracted by neither ϕ nor ϕ' .

If there is a -1 -curve contracted by both, ϕ and ϕ' factor through a smaller regular model. Assume there is a -1 -curve not contracted by ϕ . Then, by condition (4), its image has to be one of the special branches of a genus two singularity; on the other hand, by condition (5), the only special branch of a singularity of type I must contain some special points; therefore we conclude that the singular point x of \mathcal{C}_0 is dangling of type III_{l+1} , $l \leq m$. If we let $Z = \phi^{-1}(x)$, we may write $\mathcal{C}_0 = Z \cup R_0 \cup \dots \cup R_l$, with $R_0 = R$, and R_l the tail containing p_1 . By Claim 1, ϕ' has to contract a genus two subcurve Z' as well. If Z' is of the shape described in Proposition 4.5 and it contains R , then $Z \subsetneq Z'$ is easily seen, which implies \mathcal{C}'_0 has a singularity of type II with more than $m+1$ branches, by the level condition (2) on \mathcal{C}_0 ; on the other hand, if $Z' \subsetneq Z$ were disjoint from R , then \mathcal{C}'_0 would not satisfy condition (2). Similarly, if Z' is of the shape described in Proposition 4.4, then R_0 and R_l must meet in a “trunk” T attached to a Weierstrass point of the core of \mathcal{C}_0^{ss} , and it can be argued as above that $Z' = Z$ must hold.

Claim 3: The exceptional loci of ϕ and ϕ' coincide.

If \mathcal{C}_0 has only nodes, then so does \mathcal{C}'_0 by Claim 1, and we can conclude by the uniqueness part of the stable reduction theorem.

If \mathcal{C}_0 has a genus one singularity x , it cannot have a genus two singularity as well, so neither can \mathcal{C}'_0 by Claim 1. If \mathcal{C}_0 has a second genus one singularity y , let $E_1 = \phi^{-1}(x)$ and $E_2 = \phi^{-1}(y)$; they are disjoint balanced subcurves of genus one and level $\leq m+1$ in \mathcal{C}_0^{ss} , therefore ϕ' must contract them. Enlarging the contraction radius of any one of them would produce a singularity with at least $m+2$ branches (by condition (3) on \mathcal{C}_0), unless by enlarging we make them touch, in which case we would contract to a genus two singularity; but this is not possible, by Claim 1. The case of a single genus one singularity with a genus one branch, or with a disjoint subcurve of genus one, or with two branches joined by a (possibly empty) rational chain, is similar.

Finally, the case that \mathcal{C}_0 has a genus two singularity, has already been discussed at the end of Claim 2. To summarise, writing $\mathcal{C}_0^{ss} = Z \cup R_1 \cup \dots \cup R_l$, with $Z = \phi^{-1}(x)$ and $l \leq m$ - the case of a dangling III_{m+1} was dealt with before -, $\phi'(Z)$ must be a point x' , by stability considerations. Call $Z' = (\phi')^{-1}(x')$ and note that $Z \subseteq Z'$. If x and x' are singularities of the same type, $Z = Z'$ is easily deduced by level/singularity (i.e. outer/inner valence) considerations, the key point being that the shape of the curve has only one parameter (the “radius” of the circle), which is determined by m -stability. On the other hand, if x were of type II and x' of type I , the two special trees determined by x would have to share a trunk attached to a Weierstrass point of the core, and $Z \subseteq Z'$ would

imply $Z \subsetneq Z'$, which together with condition (2) for \mathcal{C}' would make x' into a singularity with too many branches.

The claim follows from observing that the exceptional locus of ϕ (resp. ϕ') is the union of the fibers F over higher genus singularities of \mathcal{C}_0 (resp. \mathcal{C}'_0), and the rational components with only two special points that are disjoint from F .

Claim: The generic isomorphism between \mathcal{C} and \mathcal{C}' extends over Δ .

Follows from [Deb01, Lemma1.13]. \square

Summing up, we have proved the following:

Theorem 5.13. *For every $1 \leq m < n$, m -stability defines a proper Deligne-Mumford stack $\overline{\mathcal{M}}_{2,n}^{(m)}$ over \mathbf{k} , containing $\mathcal{M}_{2,n}$ as a dense open substack.*

REFERENCES

- [AFS16] Jarod Alper, Maksym Fedorchuk, and David Ishii Smyth. Singularities with \mathbb{G}_m -action and the log minimal model program for $\overline{\mathcal{M}}_g$. *J. Reine Angew. Math.*, 721:1–41, 2016.
- [AFS17a] Jarod Alper, Maksym Fedorchuk, and David Ishii Smyth. Second flip in the Hassett-Keel program: existence of good moduli spaces. *Compos. Math.*, 153(8):1584–1609, 2017.
- [AFS17b] Jarod Alper, Maksym Fedorchuk, and David Ishii Smyth. Second flip in the Hassett-Keel program: projectivity. *Int. Math. Res. Not. IMRN*, (24):7375–7419, 2017.
- [AFSvdW17] Jarod Alper, Maksym Fedorchuk, David Ishii Smyth, and Frederick van der Wyck. Second flip in the Hassett-Keel program: a local description. *Compos. Math.*, 153(8):1547–1583, 2017.
- [AK70] Allen Altman and Steven Kleiman. *Introduction to Grothendieck duality theory*. Lecture Notes in Mathematics, Vol. 146. Springer-Verlag, Berlin-New York, 1970.
- [AK16] Jarod Alper and Andrew Kresch. Equivariant versal deformations of semistable curves. *Michigan Math. J.*, 65(2):227–250, 2016.
- [Arn72] V. I. Arnol'd. Normal forms of functions near degenerate critical points, the Weyl groups A_k, D_k, E_k and Lagrangian singularities. *Funkcional. Anal. i Priložen.*, 6(4):3–25, 1972.
- [BCM18] Luca Battistella, Francesca Carocci, and Cristina Manolache. Reduced invariants from cuspidal maps. *arXiv e-prints*, page arXiv:1801.07739, Jan 2018.
- [BLR90] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud. *Néron models*, volume 21 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1990.
- [BS08] Elizabeth Baldwin and David Swinarski. A geometric invariant theory construction of moduli spaces of stable maps. *Int. Math. Res. Pap. IMRP*, (1):Art. ID rp. 004, 104, 2008.
- [Cat82] Fabrizio Catanese. Pluricanonical-Gorenstein-curves. In *Enumerative geometry and classical algebraic geometry (Nice, 1981)*, volume 24 of *Progr. Math.*, pages 51–95. Birkhäuser Boston, Boston, MA, 1982.
- [CML13] Sebastian Casalaina-Martin and Radu Laza. Simultaneous semi-stable reduction for curves with ADE singularities. *Trans. Amer. Math. Soc.*, 365(5):2271–2295, 2013.

- [CTV18] Giulio Codogni, Luca Tasin, and Filippo Viviani. On the first steps of the minimal model program for the moduli space of stable pointed curves. *arXiv e-prints*, page arXiv:1808.00231, Aug 2018.
- [CTV19] Giulio Codogni, Luca Tasin, and Filippo Viviani. On some modular contractions of the moduli space of stable pointed curves. *arXiv e-prints*, page arXiv:1904.13212, Apr 2019.
- [Cuk89] Fernando Cukierman. Families of Weierstrass points. *Duke Math. J.*, 58(2):317–346, 1989.
- [Deb01] Olivier Debarre. *Higher-dimensional algebraic geometry*. Universitext. Springer-Verlag, New York, 2001.
- [Dem75] Michel Demazure. Classification des germes à point critique isolé et à nombres de modules 0 ou 1 (d’après V. I. Arnol’d). pages 124–142. *Lecture Notes in Math.*, Vol. 431, 1975.
- [Dia85] Steven Diaz. Exceptional Weierstrass points and the divisor on moduli space that they define. *Mem. Amer. Math. Soc.*, 56(327):iv+69, 1985.
- [DM69] P. Deligne and D. Mumford. The irreducibility of the space of curves of given genus. *Inst. Hautes Études Sci. Publ. Math.*, (36):75–109, 1969.
- [FS13] Maksym Fedorchuk and David Ishii Smyth. Alternate compactifications of moduli spaces of curves. In *Handbook of moduli. Vol. I*, volume 24 of *Adv. Lect. Math. (ALM)*, pages 331–413. Int. Press, Somerville, MA, 2013.
- [Gie82] D. Gieseker. *Lectures on moduli of curves*, volume 69 of *Tata Institute of Fundamental Research Lectures on Mathematics and Physics*. Published for the Tata Institute of Fundamental Research, Bombay; Springer-Verlag, Berlin-New York, 1982.
- [Hal14] Daniel Halpern-Leistner. On the structure of instability in moduli theory. *arXiv e-prints*, page arXiv:1411.0627, Nov 2014.
- [Has03] Brendan Hassett. Moduli spaces of weighted pointed stable curves. *Adv. Math.*, 173(2):316–352, 2003.
- [Has05] Brendan Hassett. Classical and minimal models of the moduli space of curves of genus two. In *Geometric methods in algebra and number theory*, volume 235 of *Progr. Math.*, pages 169–192. Birkhäuser Boston, Boston, MA, 2005.
- [HLN12] Yi Hu, Jun Li, and Jingchen Niu. Genus Two Stable Maps, Local Equations and Modular Resolutions. *arXiv e-prints*, page arXiv:1201.2427, Jan 2012.
- [HM82] Joe Harris and David Mumford. On the Kodaira dimension of the moduli space of curves. *Invent. Math.*, 67(1):23–88, 1982. With an appendix by William Fulton.
- [HM98] Joe Harris and Ian Morrison. *Moduli of curves*, volume 187 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1998.
- [Hun99] Craig Huneke. Hyman Bass and ubiquity: Gorenstein rings. In *Algebra, K-theory, groups, and education (New York, 1997)*, volume 243 of *Contemp. Math.*, pages 55–78. Amer. Math. Soc., Providence, RI, 1999.
- [Kee99] Seán Keel. Basepoint freeness for nef and big line bundles in positive characteristic. *Ann. of Math. (2)*, 149(1):253–286, 1999.
- [KM97] Seán Keel and Shigefumi Mori. Quotients by groupoids. *Ann. of Math. (2)*, 145(1):193–213, 1997.
- [Knu83] Finn F. Knudsen. The projectivity of the moduli space of stable curves. II. The stacks $M_{g,n}$. *Math. Scand.*, 52(2):161–199, 1983.
- [LZ09] Jun Li and Aleksey Zinger. On the genus-one Gromov-Witten invariants of complete intersections. *J. Differential Geom.*, 82(3):641–690, 2009.
- [MFK94] D. Mumford, J. Fogarty, and F. Kirwan. *Geometric invariant theory*, volume 34 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (2) [Results in Mathematics and Related Areas (2)]*. Springer-Verlag, Berlin, third edition, 1994.

- [Mor11] Ian Morrison. Mori theory of moduli spaces of stable curves. <http://www.projectivepress.com/moduli/moristablecurves.pdf>, 2011.
- [RSW17a] Dhruv Ranganathan, Keli Santos-Parker, and Jonathan Wise. Moduli of stable maps in genus one and logarithmic geometry I. *arXiv e-prints*, page arXiv:1708.02359, August 2017.
- [RSW17b] Dhruv Ranganathan, Keli Santos-Parker, and Jonathan Wise. Moduli of stable maps in genus one and logarithmic geometry II. *arXiv e-prints*, page arXiv:1709.00490, September 2017.
- [Sch91] David Schubert. A new compactification of the moduli space of curves. *Compositio Math.*, 78(3):297–313, 1991.
- [Smy11a] David Ishii Smyth. Modular compactifications of the space of pointed elliptic curves I. *Compos. Math.*, 147(3):877–913, 2011.
- [Smy11b] David Ishii Smyth. Modular compactifications of the space of pointed elliptic curves II. *Compos. Math.*, 147(6):1843–1884, 2011.
- [Smy13] David Ishii Smyth. Towards a classification of modular compactifications of $\mathcal{M}_{g,n}$. *Invent. Math.*, 192(2):459–503, 2013.
- [Smy18] David Ishii Smyth. Intersections of psi-classes on moduli spaces of m-stable curves. *arXiv e-prints*, page arXiv:1808.03214, Aug 2018.
- [Sta19] The Stacks Project Authors. *Stacks Project*. <https://stacks.math.columbia.edu>, 2019.
- [Ste96] Jan Stevens. On the classification of reducible curve singularities. In *Algebraic geometry and singularities (La Rábida, 1991)*, volume 134 of *Progr. Math.*, pages 383–407. Birkhäuser, Basel, 1996.
- [vdW10] Frederick van der Wyck. *Moduli of singular curves and crimping*. PhD thesis, Harvard University, 2010.
- [Vis12] Michael Viscardi. Alternate compactifications of the moduli space of genus one maps. *Manuscripta Math.*, 139(1-2):201–236, 2012.
- [VZ08] Ravi Vakil and Aleksey Zinger. A desingularization of the main component of the moduli space of genus-one stable maps into \mathbb{P}^n . *Geom. Topol.*, 12(1):1–95, 2008.
- [Zin09] Aleksey Zinger. Reduced genus-one Gromov-Witten invariants. *J. Differential Geom.*, 83(2):407–460, 2009.

Luca Battistella

Max Planck Institut für Mathematik - Bonn

battistella@mpim-bonn.mpg.de