

A QUANTUM LEFSCHETZ THEOREM FOR QUASIMAP INVARIANTS VIA RELATIVE QUASIMAPS

LUCA BATTISTELLA AND NAVID NABIJOU

ABSTRACT. We define moduli spaces of relative toric quasimaps in genus zero in the spirit of A. Gathmann. When X is a smooth toric variety and Y is a very ample hypersurface in X we construct a virtual class on the moduli space of relative quasimaps to (X, Y) which can be used to define relative quasimap invariants of the pair. We obtain a recursion formula which expresses each relative invariant in terms of invariants of lower multiplicity. Finally we apply this formula to derive a quantum Lefschetz theorem expressing the absolute quasimap invariants of Y in terms of those of X . We include several appendices collecting proofs of standard results in quasimap theory.

CONTENTS

1. Introduction	1
2. Relative stable quasimaps	5
3. Recursion formula for \mathbb{P}^N relative a hyperplane	13
4. Recursion formula in the general case	19
5. The quasimap mirror formula	24
Appendix A. The comparison morphism	30
Appendix B. Notes on quasimaps	37
Appendix C. Some intersection-theoretic lemmas	53
References	54

1. INTRODUCTION

In this paper we construct moduli spaces of relative quasimaps as substacks of moduli spaces of (absolute) quasimaps. This provides a common generalisation of two different theories: stable quasimaps on the one hand, and relative stable maps (in the sense of A. Gathmann) on the other. In this introductory section we briefly recall these, providing the context for our work.

1.1. Stable quasimaps. The moduli space of *stable toric quasimaps*

$$\overline{\mathcal{Q}}_{g,n}(X, \beta)$$

was constructed by I. Ciocan-Fontanine and B. Kim [CFK10] as a compactification of the moduli space of smooth curves in a smooth and complete toric variety X . Roughly speaking, the objects are rational maps $C \dashrightarrow X$ where C is a nodal curve, subject to a stability condition. The precise definition relies on the description of X as a GIT quotient.

The space $\overline{\mathcal{Q}}_{g,n}(X, \beta)$ is a proper Deligne–Mumford stack of finite type, and admits a virtual fundamental class, which is used to define curve-counting invariants for X , called *quasimap invariants*.

This theory agrees with that of stable quotients [MOP11] when both are defined, namely when X is a projective space. There is a common generalisation given by the theory of stable quasimaps to GIT quotients [CFKM14]. For simplicity, however, we will work mostly in the toric setting (though this restriction is probably not essential for our arguments). Thus in this paper when we say “quasimaps” we are implicitly talking about toric quasimaps.

The quasimap invariants provide an alternative system of curve counts to the more well-known Gromov–Witten invariants. These latter invariants are defined via moduli spaces of stable maps, and as such we will often refer to them as *stable map invariants* in this paper.

The genus zero quasimap invariants are expected to coincide with the stable map invariants when X is a toric Fano variety [CM]; this has been proven (in all genera) for $X = \mathbb{P}^N$ [MOP11, Theorem 3] [Man12b, §5.4]. More generally, the case of a projective complete intersection of Fano index at least 2 can be obtained by combining A. Givental’s mirror theorem [Giv98, Theorem 0.1] with the wall-crossing formulae of I. Ciocan-Fontanine and B. Kim [CFK10, Conjecture 7.2.10] [CFK14, §5.5 and Conjecture 6.3.1].

In general, however, the invariants differ, the difference being encoded by certain wall-crossing formulae [CFK14]. The motivation for this comes from mirror symmetry: the idea is that the quasimap invariants of X should correspond to the B -side theory of X^\vee (this is in contrast to the Gromov–Witten invariants, which live on the A -side); see [CFK10, §7].

1.2. Relative stable maps. In [Gat02] A. Gathmann constructs a moduli space of relative stable maps to the pair (X, Y) as a closed substack of the moduli space of (absolute) stable maps to X

$$\overline{\mathcal{M}}_{0,\alpha}(X|Y, \beta) \hookrightarrow \overline{\mathcal{M}}_{0,n}(X, \beta)$$

parametrising stable maps with prescribed tangencies to Y at the marked points.

Unfortunately this space does not admit a natural perfect obstruction theory. Nevertheless in the case where Y is very ample it is still possible to construct a virtual fundamental class by intersection-theoretic methods, and hence one can define relative Gromov–Witten invariants.

There is a recursion formula for these virtual classes which allows one to express any relative invariant of (X, Y) in terms of absolute invariants of Y and relative invariants with lower contact multiplicities.

By successively increasing the contact multiplicities from zero to the maximum possible value, Gathmann obtains an algorithm expressing the absolute invariants of Y in terms of those of X : see [Gat02, Corollary 5.7]. In [Gat03] this result is applied to produce an alternative proof of the mirror theorem for projective hypersurfaces [Giv96a] [LLY97].

1.3. Relative stable quasimaps. In this paper we construct moduli spaces of relative quasimaps in genus zero. We prove a recursion relation similar to Gathmann's formula, and use this to derive a quantum Lefschetz formula for quasimap invariants.

This construction carries over to the setting of ϵ -stable quasimaps; since for $\epsilon > 1$ these moduli spaces agree with the space of stable maps, one can view our construction as giving a common generalisation of the two stories outlined above.

The plan of the paper is as follows. In §§2.1 and 2.2 we provide a brief review of the theories of stable quasimaps and relative stable maps. Then in §2.3 we define the moduli space of relative quasimaps as a substack of the moduli space of (absolute) quasimaps:

$$\overline{\mathcal{Q}}_{0,\alpha}(X|Y, \beta) \subseteq \overline{\mathcal{Q}}_{0,n}(X, \beta)$$

Here X is a smooth toric variety, Y is a smooth hypersurface and $\alpha = (\alpha_1, \dots, \alpha_n)$ encodes the orders of tangency of the marked points to Y . Note that we *do not* require Y to be toric.

In §3 we examine the special case of a hyperplane $H \subseteq \mathbb{P}^N$. We find that the moduli space is irreducible of the expected dimension (in fact, more than this: it is the closure of the so-called “nice locus” consisting of maps from a \mathbb{P}^1 whose image is not contained in H). Thus it has an actual fundamental class, which we can use to define relative quasimap invariants.

Another useful fact about this special case is that there exists a birational comparison morphism:

$$\chi : \overline{\mathcal{M}}_{0,n}(\mathbb{P}^N, d) \rightarrow \overline{\mathcal{Q}}_{0,n}(\mathbb{P}^N, d)$$

This restricts to a birational morphism between the relative spaces, which we use to push down Gathmann's formula to obtain a recursion formula for relative stable quasimaps. The stronger stability condition for quasimaps significantly simplifies the boundary terms which appear.

In §4 we deal with the case of an arbitrary pair (X, Y) with Y very ample. We use the embedding $X \hookrightarrow \mathbb{P}^N$ defined by $\mathcal{O}_X(Y)$ to construct a virtual class $[\overline{\mathcal{Q}}_{0,\alpha}(X|Y, \beta)]^{\text{virt}}$.

We then prove the recursion formula for (X, Y) by pulling back the formula for (\mathbb{P}^N, H) . This requires several comparison theorems for virtual classes, extending results in Gromov–Witten theory to the setting of quasimaps.

Finally in §5 we apply the recursion formula of §4 to obtain a quantum Lefschetz formula for quasimap invariants, i.e. a formula expressing the invariants of Y in terms of those of X . This recovers Corollary 5.5.1 in [CFK14], and can be interpreted as a mirror theorem for Y . The argument is similar in spirit to the one given in [Gat03], however the stronger stability condition considerably simplifies both the proof and the final result.

We also include several appendices, collecting together results which are presumably well-known to experts. Appendix A discusses the comparison morphism; Appendix B contains foundational results in quasimap theory, including functoriality and the splitting axiom; Appendix C contains a number of intersection-theoretic lemmas.

Acknowledgements. The authors wish to thank Tom Coates, Cristina Manolache, Andrea Petracci and Richard Thomas for many helpful discussions. L.B. is supported by [REF] and N.N. is supported by [REF]

1.4. Table of notation. We will use the following notation, most of which is introduced in the main body of the paper.

X	a smooth projective toric variety
Y	a smooth very ample hypersurface in X
Σ	the fan of X
$\Sigma(1)$	the set of 1-dimensional cones of Σ
ρ	an element of $\Sigma(1)$
D_ρ	the toric divisor in X corresponding to ρ
$\overline{\mathcal{M}}_{g,n}(X, \beta)$	the moduli space of stable maps to X
$\overline{\mathcal{M}}_{0,\alpha}(X Y, \beta)$	the moduli space of relative stable maps to (X, Y) ; see §2.2
$\overline{\mathcal{Q}}_{g,n}(X, \beta)$	the moduli space of toric quasimaps to X ; see §2.1
$\mathcal{Q}_{0,\alpha}^\circ(X Y, \beta)$	the nice locus of relative quasimaps to (X, Y) ; see §3.1
$\overline{\mathcal{Q}}_{0,\alpha}(X Y, \beta)$	the moduli space of relative quasimaps to (X, Y) ; see §2.3
$\mathcal{D}_{\alpha,k}^Q(X Y, \beta)$	the quasimap comb locus; see §3.2
$\mathcal{D}^Q(X Y, A, B, M)$	(a component of) the comb locus; see §3.2
$\mathcal{E}^Q(X Y, A, B, M)$	the total product for the comb locus; see §4.3
$\mathcal{D}^Q(X, A, B)$	the quasimap centipede locus; see Appendix B.3
$\mathcal{E}^Q(X, A, B)$	the total product for the centipede locus; see Appendix B.3
$\mathfrak{M}_{g,n}^{\text{wt}}$	the moduli stack of weighted prestable curves; see Appendix B.3
$\mathfrak{Pic}_{g,n}^{d,\text{st}}$	the usual open substack of the relative Picard stack of the universal curve over $\mathfrak{M}_{g,n}$; see Appendix A
\mathfrak{Bun}_G	the moduli stack of principal G -bundles on the universal curve over $\mathfrak{M}_{g,n}$; see Appendix B.4

$\overline{Q}(f)$	the push-forward morphism between quasimap spaces; see Appendix B.1
χ	the comparison morphism from stable maps to quasimaps; see Appendix A
$f^!$	Gysin morphism for f a regular embedding
$f_v^!$	virtual pull-back for f virtually smooth; see Appendix C
$f_\Delta^!$	diagonal pull-back; see Appendix C

2. RELATIVE STABLE QUASIMAPS

2.1. Review of absolute stable quasimaps. We briefly recall the definition and basic properties of the moduli space of toric quasimaps; see [CFK10] for more details.

Definition 2.1 ([CFK10, Definition 3.1.1]). Let $X = X_\Sigma$ be a smooth and projective toric variety with fan $\Sigma \subseteq N_{\mathbb{Q}}$, let $M = N^\vee = \text{Hom}(N, \mathbb{Z})$ and let $\mathcal{O}_{X_\Sigma}(1)$ be a fixed polarisation, which we can write (non-uniquely) in terms of the T -invariant divisors as:

$$\mathcal{O}_{X_\Sigma}(1) = \otimes_{\rho \in \Sigma(1)} \mathcal{O}_{X_\Sigma}(D_\rho)^{\otimes \alpha_\rho}$$

for some $\alpha_\rho \in \mathbb{Z}$. We fix the following numerical invariants: a genus $g \geq 0$, number of marked points $n \geq 0$ and effective curve class $\beta \in H_2^+(X)$. Then a *stable (toric) quasimap* is given by the data

$$((C, x_1, \dots, x_n), (L_\rho, u_\rho)_{\rho \in \Sigma(1)}, (\varphi_m)_{m \in M})$$

where:

- (1) (C, x_1, \dots, x_n) is a prestable curve of genus g with n marked points;
- (2) the L_ρ are line bundles on C of degree $d_\rho = D_\rho \cdot \beta$;
- (3) the u_ρ are global sections of L_ρ ;
- (4) $\varphi_m: \bigotimes_{\rho \in \Sigma(1)} L_\rho^{\otimes \langle \rho, m \rangle} \rightarrow \mathcal{O}_C$ are isomorphisms, such that $\varphi_m \otimes \varphi_{m'} = \varphi_{m+m'}$ for all $m, m' \in M$.

These are required to satisfy the following two conditions:

- (1) *nondegeneracy*: there is a finite (possibly empty) set of smooth and non-marked points $B \subseteq C$, called the *basepoints* of the quasimap, such that for all $x \in C \setminus B$ there exists a maximal cone $\sigma \in \Sigma_{\max}$ with $u_\rho(x) \neq 0$ for all $\rho \notin \sigma$;
- (2) *stability*: if we let $L = \otimes_\rho L_\rho^{\otimes \alpha_\rho}$ then the following \mathbb{Q} -divisor is ample

$$\omega_C(x_1 + \dots + x_n) \otimes L^{\otimes \epsilon}$$

for every rational $\epsilon > 0$.

It can be shown that this definition does not depend on the choice of polarisation.

Remark 2.2. This definition is motivated by D. Cox's description of the functor of points of a toric variety in terms of Σ -collections [Cox95a]; see

also Appendix B.1. A quasimap defines a rational map $C \dashrightarrow X$ with base locus equal to B . (This can be expressed in a more generalisable manner as follows: a quasimap is a map to the stack quotient $[\mathbb{A}^{\Sigma(1)}/(\mathbb{G}_m)^r]$ such that B is the preimage of the unstable locus.)

In particular a quasimap without any basepoints defines a morphism $C \rightarrow X$. Thus the basepoints appear in the (virtual) boundary of the moduli space, in much the same way as the locus of stable maps with rational tails appears in the boundary of the moduli space of stable maps. This is something more than just a vague analogy; these loci correspond to each other under the comparison morphism when $X \simeq \mathbb{P}^N$; see Appendix A.

More generally, one can define the notion of a family of quasimaps over a base scheme S , and what it means for two such families to be isomorphic; one thus obtains a moduli space

$$\overline{\mathcal{Q}}_{g,n}(X, \beta)$$

of stable (toric) quasimaps to X , which is a proper Deligne–Mumford stack of finite type [CFK10, §3]

As with the case of stable maps, there is a combinatorial characterisation of stability which is much easier to check in practice; a prestable quasimap is stable if and only if the following conditions hold:

- (1) the line bundle L defined above must have strictly positive degree on any rational component with fewer than three special points, and on any elliptic component with no special points;
- (2) C cannot have any rational components with fewer than two special points (called *rational tails*).

Condition (1) is analogous to the ordinary stability condition for stable maps. Condition (2) is new, however, and gives quasimaps a distinctly different flavour to stable maps; we shall sometimes refer to it as the *strong stability condition*.

Remark 2.3. Unlike in Gromov–Witten theory, $\overline{\mathcal{Q}}_{g,n+1}(X, \beta)$ is *not* the universal curve over $\overline{\mathcal{Q}}_{g,n}(X, \beta)$ since markings cannot be basepoints. In fact there is not even a morphism between these spaces in general.

The moduli space $\overline{\mathcal{Q}}_{g,n}(X, \beta)$ admits a perfect obstruction theory relative to the moduli space $\mathfrak{M}_{g,n}$ of source curves ([CFK10, §5], and hence one can construct a virtual class

$$[\overline{\mathcal{Q}}_{g,n}(X, \beta)]^{\text{virt}} \in A_{\text{vdim } \overline{\mathcal{Q}}_{g,n}(X, \beta)}(\overline{\mathcal{Q}}_{g,n}(X, \beta))$$

where the virtual dimension is the same as for stable maps:

$$\text{vdim } \overline{\mathcal{Q}}_{g,n}(X, \beta) = (\dim X - 3)(1 - g) - (K_X \cdot \beta) + n$$

Since the markings are not basepoints there exist evaluation maps

$$\text{ev}_i : \overline{\mathcal{Q}}_{g,n}(X, \beta) \rightarrow X$$

and there are ψ -classes defined in the usual way by pulling back the relative dualising sheaf of the universal curve

$$\psi_i = c_1(x_i^* \omega_{C/\overline{\mathcal{Q}}})$$

where $C \rightarrow \overline{\mathcal{Q}} = \overline{\mathcal{Q}}_{g,n}(X, \beta)$ is the universal curve and $x_i : \overline{\mathcal{Q}} \rightarrow C$ is the section defining the i th marked point. Putting all these pieces together, we can define *quasimap invariants*:

$$\langle \gamma_1 \psi^{k_1}, \dots, \gamma_n \psi^{k_n} \rangle_{g,n,\beta}^X = \int_{[\overline{\mathcal{Q}}_{g,n}(X, \beta)]^{\text{virt}}} \prod_{i=1}^n \text{ev}_i^*(\gamma_i) \psi_i^{k_i}$$

We use the same correlator notation as in Gromov–Witten theory; this should not cause any confusion.

Example 2.4. Consider $\overline{\mathcal{Q}}_{0,2}(\mathbb{P}^2, 1)$. What are its objects? By the strong stability condition (2) above, we see that the source curve must be irreducible. On the other hand since \mathbb{P}^2 has Picard rank 1 we may exploit the isomorphisms φ_m to reduce ourselves to considering one line bundle equipped with three sections. Thus the data of the quasimap is $((C, x_1, x_2), L, u_0, u_1, u_2)$ where $(C, L) \cong (\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1))$.

Pick coordinates $[s : t]$ on \mathbb{P}^1 such that the marked points are $[1 : 0]$ and $[0 : 1]$. We can express the sections as $u_i = a_i s + b_i t$; the requirement that the markings are not basepoints then translates into the following stability condition:

$$(a_0, a_1, a_2) \neq (0, 0, 0) \quad \text{and} \quad (b_0, b_1, b_2) \neq (0, 0, 0).$$

The group $\text{Aut}(C; x_1, x_2) \cong \mathbb{G}_m$ acts by rotation $\lambda : [s : t] \mapsto [s : \lambda t]$, while $\text{Aut}(L) \cong \mathbb{G}_m$ acts by scalar multiplication on \underline{a} and \underline{b} . Thus the \mathbb{G}_m^2 action on $\mathbb{A}_{\underline{a}, \underline{b}}^6$ is encoded by the following weight matrix:

$$\begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 & 1 \end{bmatrix}$$

It is now clear that the quotient is $\mathbb{P}^2 \times \mathbb{P}^2$; in fact, we see that the evaluation map

$$(\text{ev}_1, \text{ev}_2) : \overline{\mathcal{Q}}_{0,2}(\mathbb{P}^2, 1) \rightarrow \mathbb{P}^2 \times \mathbb{P}^2$$


is an isomorphism. It is given in the above notation by:

$$((\mathbb{P}^1; [1 : 0], [0 : 1]); \mathcal{O}_{\mathbb{P}^1}(1); u_0, u_1, u_2) \mapsto ([a_0 : a_1 : a_2], [b_0 : b_1 : b_2])$$

Notice that the locus where $(a_0, a_1, a_2) = \mu(b_0, b_1, b_2)$, i.e. the diagonal in $\mathbb{P}^2 \times \mathbb{P}^2$ is precisely the locus of quasimaps which have a basepoint. The point $[a_0 : a_1 : a_2] = [b_0 : b_1 : b_2] \in \mathbb{P}^2$ is the image of the underlying “residual map” of degree 0, obtained by dividing all the sections by a local equation

of the basepoint (equivalently, by extending the rational map $C \dashrightarrow \mathbb{P}^2$ to a morphism $C \rightarrow \mathbb{P}^2$).

On the other hand, $(ev_1, ev_2): \overline{\mathcal{M}}_{0,2}(\mathbb{P}^2, 1) \rightarrow \mathbb{P}^2 \times \mathbb{P}^2$ is *not* an isomorphism. Off the diagonal, the images of the two marked points determine uniquely the image of the stable map, i.e. the line through them. On the diagonal however, the following maps with a rational tail appear:



The image of the degree 1 component under f can be any line passing through the point of \mathbb{P}^2 to which the other component is contracted. Hence $\overline{\mathcal{M}}_{0,2}(\mathbb{P}^2, 1) \simeq \text{Bl}_\Delta(\mathbb{P}^2 \times \mathbb{P}^2)$. The comparison morphism χ (see Appendix A) can be interpreted as the blow-down map, and it induces an isomorphism of the rational tails-free locus with the basepoint-free locus.

Remark 2.5. There is a more general notion of ϵ -stable quasimap [CFKM14, §7.1]. Here the stability condition, namely that the line bundle

$$\omega_C(x_1 + \dots + x_n) \otimes L^{\otimes \epsilon}$$

is ample, is only required to hold for a fixed $\epsilon \in \mathbb{Q}_{>0}$ (instead of for arbitrary ϵ , as was the case with ordinary quasimaps).

This has the effect of allowing some rational tails to appear, as long as their degree is high enough with respect to ϵ . In order to keep the moduli space separated, one has to bound the multiplicity of the basepoints that can occur.

By boundedness and the fact that the degree is an integer-valued function, there exist finitely many critical values of ϵ which divide $\mathbb{Q}_{>0}$ into chambers inside which the moduli spaces $\overline{\mathcal{Q}}_{g,n}^\epsilon(X, \beta)$ do not change.

Furthermore for ϵ sufficiently small we recover the space of (ordinary) quasimaps, and for ϵ sufficiently large we obtain the moduli space of stable maps. Thus one can view the spaces of ϵ -stable quasimaps as interpolating between these two extremes, and they have proven successful as a tool for comparing quasimap invariants to stable map invariants [CFK14].

Remark 2.6. There is another variant which is going to play a role in later sections: that of *parametrised quasimaps* [CFK10, §7]. The idea is that a parametrised quasimap comes with a preferred rational component (by introducing the extra data of an isomorphism with \mathbb{P}^1) and the stability condition is imposed *on all but the preferred component*. This mimics the construction of graph spaces in Gromov-Witten theory and induces a \mathbb{G}_m -action on $\overline{\mathcal{Q}}_{g,n}^\epsilon(X, \beta)$ by rotating the preferred component. The fixed loci

and their equivariant normal bundles are well-understood, at least in the toric setting [CFK10, §7].

Note that we no longer require $2g - 2 + n \geq 0$, due to the modified stability condition. In particular it makes sense, and turns out to be very useful, to consider unmarked parametrised quasimaps $\overline{QG}_{0,0}(X, \beta)$. In this case the source curve needs to be irreducible.

Example 2.7. $\overline{QG}_{0,0}(\mathbb{P}^r, d) = \mathbb{P}^N$ with $N = (r + 1)(d + 1) - 1$. Indeed, the curve and line bundle must be $(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(d))$ and we are left with choosing $r + 1$ sections of $\mathcal{O}_{\mathbb{P}^1}(d)$ (not all zero) up to automorphisms of $\mathcal{O}_{\mathbb{P}^1}(d)$, i.e. up to scaling. For an early appearance of such spaces, see for instance [Ber00].

2.2. Review of relative stable maps. Given a smooth projective variety X and a smooth divisor Y , Gathmann's moduli space of relative stable maps parametrises stable maps to X with specified tangencies to Y at the marked points; see [Gat02] for details.

Remark 2.8. There are many other approaches to defining relative stable maps besides Gathmann's: the moduli space of maps to expanded degenerations of J. Li [Li01] [Li02], the twisted stable maps of D. Abramovich and B. Fantechi [AF16], the logarithmic stable maps with expansions of B. Kim [Kim10] and the logarithmic stable maps (without expansions) of M. Gross and B. Siebert [GS13] [GS16], Q. Chen [Che14] and D. Abramovich and Q. Chen [AC14]. However, the invariants defined via these theories are all known to coincide [AMW14], so the choice of which variant to work with mainly depends on one's intended application.

Definition 2.9 ([Gat02, Definition 1.1]). Let X be a smooth projective variety and $Y \subseteq X$ a smooth divisor. Fix a number $n \geq 0$ of marked points, an effective curve class $\beta \in H_2^+(X)$ and an n -tuple $\alpha = (\alpha_1, \dots, \alpha_n)$ of non-negative integers such that $\sum_i \alpha_i \leq Y \cdot \beta$. Then the moduli space

$$\overline{\mathcal{M}}_{0,\alpha}(X|Y, \beta)$$

of relative stable maps to (X, Y) is defined to be the locus in $\overline{\mathcal{M}}_{0,n}(X, \beta)$ of stable maps $(C \rightarrow S, (x_i : S \rightarrow C)_{i=1}^n, f : C \rightarrow X)$ satisfying the following two conditions:

- (1) if x_i is a marked point such that $\alpha_i > 0$ then $f(x_i) \in Y$;
- (2) if we consider $f^*[Y] \in A_0(f^{-1}Y)$ then the difference $f^*[Y] - \sum_i \alpha_i x_i$ is an effective class.

This forms a closed substack of $\overline{\mathcal{M}}_{0,n}(X, \beta)$. Condition (1) is required in order for the class $\sum_i \alpha_i x_i$ to make sense in $A_0(f^{-1}Y)$.

Remark 2.10. When $\alpha = (0, \dots, 0)$, condition (2) becomes $Y \cdot \beta \geq 0$, so $\overline{\mathcal{M}}_{0,(0,\dots,0)}(X|Y, \beta) = \overline{\mathcal{M}}_{0,n}(X, \beta)$ as long as Y is nef.

Remark 2.11. The definition given above works in families; however there is a more combinatorial definition for individual maps which is more useful in

practice (see [Gat02, Remark 1.4]): a stable map (C, x_1, \dots, x_n, f) is a relative stable map if and only if, for each connected component Z of $f^{-1}(Y) \subseteq C$:

- (1) if Z is a point and is equal to a marked point x_i , then the multiplicity of f to Y at x_i is greater than or equal to α_i ;
- (2) if Z is one-dimensional (hence a union of irreducible components of C) and if we let $C^{(i)}$ for $1 \leq i \leq r$ denote the irreducible components of C adjacent to Z , and $m^{(i)}$ denote the multiplicity of $f|_{C^{(i)}}$ to Y at the node $Z \cap C^{(i)}$, then we must have:

$$(1) \quad Y \cdot f_*[Z] + \sum_{i=1}^r m^{(i)} \geq \sum_{x_i \in Z} \alpha_i$$

Remark 2.12. In the second case above we call Z an *internal* component and the $C^{(i)}$ *external* components. Note that Z is not necessarily irreducible: the term “component” is justified by the fact that it is a *connected* component of $f^{-1}(Y)$.

Remark 2.13. In the case of maximal multiplicity $\sum_i \alpha_i = Y \cdot \beta$, all the inequalities in the above definition must actually be equalities.

In the case where $X = \mathbb{P}^N$ and $Y = H$ is a hyperplane, one can show that $\overline{\mathcal{M}}_{0,\alpha}(\mathbb{P}^N|H, d)$ is irreducible with dimension equal to the expected dimension:

$$\text{vdim } \overline{\mathcal{M}}_{0,\alpha}(X|Y, \beta) = \text{vdim } \overline{\mathcal{M}}_{0,n}(X, \beta) - \sum_{i=1}^n \alpha_i$$

Hence it has a fundamental class from which one can define relative Gromov–Witten invariants.

More generally if $Y \subseteq X$ is very ample one can use the embedding $X \hookrightarrow \mathbb{P}^N$ corresponding to $|\mathcal{O}_X(Y)|$ to obtain a cartesian diagram:

$$\begin{array}{ccc} \overline{\mathcal{M}}_{0,\alpha}(X|Y, \beta) & \longrightarrow & \overline{\mathcal{M}}_{0,\alpha}(\mathbb{P}^N|H, d) \\ \downarrow & \square & \downarrow \\ \overline{\mathcal{M}}_{0,n}(X, \beta) & \xrightarrow{\varphi} & \overline{\mathcal{M}}_{0,n}(\mathbb{P}^N, d) \end{array}$$

Then the fact that $\overline{\mathcal{M}}_{0,n}(\mathbb{P}^N, d)$ is smooth allows us to define a virtual class on $\overline{\mathcal{M}}_{0,\alpha}(X|Y, \beta)$ by diagonal pull-back (see Appendix C of the current paper):

$$[\overline{\mathcal{M}}_{0,\alpha}(X|Y, \beta)]^{\text{virt}} = \varphi^! [\overline{\mathcal{M}}_{0,\alpha}(\mathbb{P}^N|H, d)]$$

Thus one can define relative Gromov–Witten invariants by capping the virtual class with products of evaluation classes and psi classes.

In [Gat02, §§2-4] Gathmann proves a recursion relation inside the Chow group of $\overline{\mathcal{M}}_{0,\alpha}(X|Y, \beta)$. This describes what happens when we increase

the multiplicity of one of the marked points by one. Let us therefore fix a marked point $x_k \in \{x_1, \dots, x_n\}$ and let $e_k = (1, \dots, 0, \dots, 0)$. We then have

$$(\alpha_k \psi_k + \text{ev}_k^*[Y])[\overline{\mathcal{M}}_{0,\alpha}(X|Y, \beta)]^{\text{virt}} = [\overline{\mathcal{M}}_{0,\alpha+e_k}(X|Y, \beta)]^{\text{virt}} + [\mathcal{D}_{\alpha,k}(X, \beta)]^{\text{virt}}$$

where $\mathcal{D}_{\alpha,k}(X, \beta)$ is an appropriate *comb locus*. Repeated application of this result shows that both the relative Gromov–Witten invariants of (X, Y) and the Gromov–Witten invariants of Y are completely determined by the Gromov–Witten invariants of X [Gat02, Corollary 5.7]. This relation is then applied in [Gat03] to obtain a new proof of the mirror theorem for projective hypersurfaces.

2.3. Definition of relative stable quasimaps. We now give the main definition of the paper. From here on X will denote a smooth projective toric variety and $Y \subseteq X$ a very ample hypersurface. We do not require that Y is toric.

Consider the line bundle $\mathcal{O}_X(Y)$ and the section s_Y cutting out Y . By [Cox95b] we have a natural isomorphism of \mathbb{C} -vector spaces

$$H^0(X, \mathcal{O}_X(Y)) = \left\langle \prod_{\rho} z_{\rho}^{a_{\rho}} : \sum_{\rho} a_{\rho} D_{\rho} = Y \right\rangle_{\mathbb{C}}$$

where the z_{ρ} for $\rho \in \Sigma(1)$ are the generators of the Cox ring of X and the a_{ρ} are non-negative integers. We can therefore write s_Y as

$$s_Y = \sum_{\underline{a}=(a_{\rho})} \lambda_{\underline{a}} \prod_{\rho} z_{\rho}^{a_{\rho}}$$

for some scalars $\lambda_{\underline{a}} \in \mathbb{C}$. The idea is that a quasimap

$$((C, x_1, \dots, x_n), (L_{\rho}, u_{\rho})_{\rho \in \Sigma(1)}, (\varphi_m)_{m \in M})$$

should “map” a point $x \in C$ into Y if and only if the section

$$(2) \quad u_Y := \sum_{\underline{a}} \lambda_{\underline{a}} \prod_{\rho} u_{\rho}^{a_{\rho}}$$

vanishes at x . We now explain how to make sense of expression (2). For each \underline{a} we have a well-defined section

$$u_{\underline{a}} := \lambda_{\underline{a}} \prod_{\rho} u_{\rho}^{a_{\rho}} \in H^0(C, \otimes_{\rho} L_{\rho}^{\otimes a_{\rho}})$$

and if we have \underline{a} and \underline{b} such that $\sum_{\rho} a_{\rho} D_{\rho} = Y = \sum_{\rho} b_{\rho} D_{\rho}$ then these differ by an element m of M . Thus the isomorphism φ_m allows us to view the sections $u_{\underline{a}}$ and $u_{\underline{b}}$ as sections of the same bundle, which we denote by L_Y . Then we can sum these together to obtain u_Y . There is a choice involved here, but up to isomorphism it does not matter; see the proof of functoriality in Appendix B.1 for more details.

The upshot is that we obtain a line bundle L_Y on C , which plays the role of the “pull-back” of $\mathcal{O}_X(Y)$ along the “map” $C \rightarrow X$, and a global section

$$u_Y \in H^0(C, L_Y)$$

which plays the role of the “pull-back” of s_Y .

Definition 2.14. With notation as above, let $n \geq 2$ be a number of marked points, $\beta \in H_2^+(X)$ be an effective curve class and $\alpha = (\alpha_1, \dots, \alpha_n)$ a collection of non-negative integers such that $\sum_i \alpha_i \leq Y \cdot \beta$. Then we define the *moduli space of relative stable quasimaps*

$$\overline{\mathcal{Q}}_{0,\alpha}(X|Y, \beta) \subseteq \overline{\mathcal{Q}}_{0,n}(X, \beta)$$

to be the locus of quasimaps

$$((C \rightarrow S, (x_i : S \rightarrow C)_{i=1}^n), (L_\rho, u_\rho)_{\rho \in \Sigma(1)}, (\varphi_m)_{m \in M})$$

such that:

- (1) if x_i is a marking such that $\alpha_i > 0$, then $x_i^* u_Y = 0$;
- (2) if we let $u_Y^*(0) \in A_0(u_Y^{-1}(0))$ denote the class defined by the Gysin map for L_Y , then the difference $u_Y^*(0) - \sum_i \alpha_i x_i$ is an effective class.

The class $u_Y^*(0)$ is defined as follows. Consider the cartesian diagram

$$\begin{array}{ccc} u_Y^{-1}(0) & \longrightarrow & C \\ \downarrow & \square & \downarrow u_Y \\ C & \xrightarrow{0_Y} & L_Y \end{array}$$

where 0_Y is the zero section. We have a Gysin map

$$0_Y^! : A_*(C) \rightarrow A_*(u_Y^{-1}(0))$$

and we define $u_Y^*(0) := 0_Y^!([C])$.

Remark 2.15. As in the case of relative stable maps (see §2.2) there is an alternative definition which is easier to check in practice: a quasimap is a relative quasimap if and only if for every connected component Z of $u_Y^{-1}(0)$ we have that:

- (1) if Z is a point and is equal to a marked point x_i , then the order of vanishing of u_Y at x_i is greater than or equal to α_i ;
- (2) if Z is one-dimensional (hence a union of irreducible components) and if we let $C^{(i)}$ for $1 \leq i \leq r$ denote the irreducible components of C adjacent to Z , and $m^{(i)}$ the order of vanishing of u_Y at the node $Z \cap C^{(i)}$, then we must have:

$$(3) \quad \deg L_Y|_Z + \sum_{i=1}^r m^{(i)} \geq \sum_{x_i \in Z} \alpha_i$$

As it stands we do not know much about this moduli space. In the following section we will examine the case $X = \mathbb{P}^N$ and $Y = H$ a hyperplane in detail. We will then apply the results obtained there to deduce facts about the general case.

3. RECURSION FORMULA FOR \mathbb{P}^N RELATIVE A HYPERPLANE

3.1. Basic properties of the moduli space. In this section we will show that the moduli space

$$\overline{\mathcal{Q}}_{0,\alpha}(\mathbb{P}^N|H, d)$$

is irreducible of the expected dimension, and thus admits a fundamental class. We then prove a recursion formula for these fundamental classes by pushing forward Gathmann's recursion formula along the comparison morphism:

$$\chi : \overline{\mathcal{M}}_{0,n}(\mathbb{P}^N, d) \rightarrow \overline{\mathcal{Q}}_{0,n}(\mathbb{P}^N, d)$$

Let us briefly recall what this morphism does. Every stable map defines a quasimap which is stable except for the fact that it may have rational tails. χ has the effect of contracting these rational tails and introducing a basepoint at the corresponding node, with multiplicity equal to the degree of the rational tail; see Appendix A for more details.

For the remainder of this section we set $X = \mathbb{P}_{z_0 \dots z_N}^N$ and $Y = H = \{z_0 = 0\}$. Given a quasimap

$$((C, x_1, \dots, x_n), L, u_0, \dots, u_N) \in \overline{\mathcal{Q}}_{0,n}(\mathbb{P}^N, d)$$

the line bundle L_Y of the previous section is equal to L and the section u_Y is equal to u_0 .

Lemma 3.1. *The comparison morphism restricts to a morphism*

$$\chi : \overline{\mathcal{M}}_{0,\alpha}(\mathbb{P}^N|H, d) \rightarrow \overline{\mathcal{Q}}_{0,\alpha}(\mathbb{P}^N|H, d)$$

Proof. We need to verify that a relative stable map is sent to a relative stable quasimap by χ . Since the contraction of a rational tail R always occurs away from the markings, we only need to examine the internal components Z of the quasimap. To be more precise, we have to show that the inequality (3) is satisfied, using the fact that the inequality (1) is satisfied by the stable map that we started with.

Let us describe this stable map around Z . For each basepoint x on Z there is a rational tail R of the stable map attached to Z at x . This is either internal (mapped into H) or external (not mapped into H).

If R is internal then both R and Z live inside the same connected component Z' of $f^{-1}(H)$. Applying χ has the effect of contracting R and increasing the degree of the line bundle on Z by $H \cdot f_*[R]$. Thus the left hand side of inequality (1) is left unchanged, and since the right hand side is also unaltered we obtain inequality (3).

On the other hand if R is external then the multiplicity $m^{(R)}$ of $R \cap Z$ satisfies:

$$m^{(R)} \leq H \cdot f_*[R]$$

Since applying χ has the effect of replacing $m^{(R)}$ by $H \cdot f_*[R]$ in the left hand side of (1), inequality (3) holds for the quasimap. Thus we obtain a morphism from the relative stable map space to the relative quasimap space, as claimed. \square

Let us denote by

$$\mathcal{Q}_{0,\alpha}^\circ(\mathbb{P}^N|H, d) \subseteq \overline{\mathcal{Q}}_{0,\alpha}(\mathbb{P}^N|H, d)$$

the *nice locus*, consisting of those quasimaps with irreducible source curve $C \cong \mathbb{P}^1$ and no basepoints (so that we have an actual map $u : C \rightarrow \mathbb{P}^N$) such that the curve is not mapped inside H and u has tangency at least α_i to H at the marking x_i .

This is an irreducible, locally closed substack of $\overline{\mathcal{Q}}_{0,n}(\mathbb{P}^N, d)$ of codimension $\sum_i \alpha_i$, by essentially the same argument as in [Gat02, Lemma 1.8]. In fact it is isomorphic to the nice locus inside the stable map space, denoted $\mathcal{M}_{0,\alpha}(\mathbb{P}^N|H, d)$ by Gathmann (see [Gat02, Def. 1.6]); the stricter stability condition has no effect when the source curve is irreducible).

Lemma 3.2. $\overline{\mathcal{Q}}_{0,\alpha}(\mathbb{P}^N|H, d)$ is equal to the closure of the nice locus $\mathcal{Q}_{0,\alpha}^\circ(\mathbb{P}^N|H, d)$ inside $\overline{\mathcal{Q}}_{0,n}(\mathbb{P}^N, d)$.

Proof. $\overline{\mathcal{Q}}_{0,\alpha}(\mathbb{P}^N|H, d) \subseteq \overline{\mathcal{Q}_{0,\alpha}^\circ(\mathbb{P}^N|H, d)}$: we show that any relative stable quasimap can be infinitesimally deformed to a relative stable quasimap with no basepoints. This is in particular a relative stable map; we then appeal to [Gat02, Prop. 1.14] to deform this stable map and obtain a point in the nice locus. Since this deformation does not introduce any rational tails, this is also a deformation of quasimaps, and the statement follows.

We induct on the number of components containing at least one basepoint. Suppose this number is non-zero (otherwise there is nothing to prove) and pick such a component C_0 , with base-points y_1, \dots, y_k . Recall that this means that $u_i(y_j) = 0$ for all i and j . We will deform the section $u_N|_{C_0}$ to a new section $u'_N|_{C_0}$ in such a way that $u'_N|_{C_0}(y_j) \neq 0$ and in such a way that we do not introduce any new basepoints. Notice that since the relative condition only depends on u_0 , the resulting deformed quasimap will still be a relative quasimap.

Now, by nondegeneracy and the fact that there exists a basepoint, we must have $\deg(L|_{C_0}) > 0$, and since $C_0 \cong \mathbb{P}^1$ we can find a section w_0 of $L|_{C_0} \cong \mathcal{O}_{\mathbb{P}^1}(d_0)$ not vanishing at any of the base-points y_i .

We then set

$$u'_N|_{C_0} := u_N|_{C_0} + \epsilon w_0$$

and $u'_i|_{C_0} = u_i|_{C_0}$ for all other i . Notice that $u'_N|_{C_0}(y_j) \neq 0$ for all j as claimed. It is also clear that we do not introduce any new basepoints, since

$u'_N|_{C_0}(y) = 0$ implies $u_N|_{C_0}(y) = 0$ (put differently: being a basepoint is a closed condition).

It remains to extend the section $u'_N|_{C_0}$ to a section u'_N on the whole curve. Let C_1, \dots, C_r be the components of C adjacent to C_0 and let $q_i = C_0 \cap C_i$. We need to modify the sections $u_N|_{C_i}$ in such a way that $u'_N|_{C_i}(q_i) = u'_N|_{C_0}(q_i)$.

By nondegeneracy, we can choose a section w_i of $L|_{C_i}$ such that $w_i(q_i) \neq 0$. Then set:

$$u'_N|_{C_i} := u_N|_{C_i} + \epsilon \left(\frac{w_0(q_i)}{w_i(q_i)} \right) \cdot w_i$$

Then indeed we have:

$$u'_N|_{C_i}(q_i) = u_N(q_i) + \epsilon \left(\frac{w_0(q_i)}{w_i(q_i)} \right) \cdot w_i(q_i) = u_N(q_i) + \epsilon w_0(q_i) = u'_N|_{C_0}(q_i)$$

We can continue this process, replacing C_0 by C_i ; since the genus of the curve is zero there are no cycles in the dual intersection graph, and so we will never come to the same component twice. In this way we obtain a new quasimap

$$((C, x_1, \dots, x_n), L, u_0, \dots, u_{N-1}, u'_N)$$

over $\text{Spec } \mathbb{C}[\epsilon]/(\epsilon^2)$ which has no basepoints on C_0 . We can repeat this process for all the components of C (using higher powers of ϵ each time in order to ensure that we never introduce additional basepoints) and thus we obtain an infinitesimal deformation of our original quasimap which has no basepoints, as required.

$\overline{\mathcal{Q}}_{0,\alpha}^\circ(\mathbb{P}^N|H, d) \subseteq \overline{\mathcal{Q}}_{0,\alpha}(\mathbb{P}^N|H, d)$: consider a family of stable quasimaps over a smooth curve S , such that the generic fibre lies in the nice locus. We may blow-up the source curve (a fibered surface over S) in the locus of basepoints (which consists of finitely many smooth points of the central fiber) and repeat this process a finite number of times in order to obtain an actual morphism to \mathbb{P}^N . This has the effect of adding rational tails at the basepoints in the central fibre. If the morphism is constant on any of these rational tails we may contract them, and thus we obtain a family of stable maps which pushes down along χ to our original family of quasimaps.

The general fibre is not modified at all, and so is still in the nice locus. By [Gat02, Lemma 1.9] it follows that the central fibre is a relative stable map, and then by applying χ and appealing to Lemma 3.1 it follows that the same is true for the central fibre of the family of quasimaps. \square

Corollary 3.3. *The moduli space $\overline{\mathcal{Q}}_{0,\alpha}(\mathbb{P}^N|H, d)$ is irreducible of the expected dimension. Hence it has a fundamental class.*

Proof. This holds because the moduli space is equal to the closure of the nice locus, which is irreducible of the expected dimension. \square

Since it has a fundamental class, we can define *relative quasimap invariants* for the pair (\mathbb{P}^N, H) :

$$\langle \text{ev}_1^* \gamma_1 \psi_1^{k_1}, \dots, \text{ev}_n^* \gamma_n \psi_n^{k_n} \rangle_{0, \alpha, d}^{\mathbb{P}^N | H} := \int_{[\overline{\mathcal{Q}}_{0, \alpha}(\mathbb{P}^N | H, \beta)]} \prod_{i=1}^n \text{ev}_i^* \gamma_i \psi_i^{k_i}$$

We will now prove a number of properties about the fundamental class. These immediately imply corresponding properties about the relative invariants.

Corollary 3.4. *The comparison morphism from relative stable maps to relative quasimaps is birational. In particular it sends the fundamental class to the fundamental class.*

Proof. This follows because the comparison morphism restricts to an isomorphism on the nice locus, which by the lemma above is a dense open subset of both spaces. \square

3.2. Proof of the recursion formula. We wish to obtain a recursion formula relating the quasimap invariants of multiplicity α with the quasimap invariants of multiplicity $\alpha + e_k$, as in [Gat02, Theorem 2.6]. This process of “increasing the multiplicities” can be naively performed in the same way as Gathmann: for $m = \alpha_k + 1$ the following section (of the pull-back of the jet bundle of the universal line bundle)

$$\sigma_k^m := x_k^* d_{C/\overline{\mathcal{Q}}}^m(u_0) \in H^0(\overline{\mathcal{Q}}, x_k^* \mathcal{P}_{C/\overline{\mathcal{Q}}}^m(\mathcal{L}))$$

cuts out $\overline{\mathcal{Q}}_{0, \alpha + e_k}(\mathbb{P}^N | H, d)$ inside $\overline{\mathcal{Q}} = \overline{\mathcal{Q}}_{0, \alpha}(\mathbb{P}^N | H, d)$, along with a number of degenerate contributions (called the *comb loci*) parametrising quasimaps for which x_k belongs to an internal component $Z \subseteq C$ (a component on which u_0 vanishes), such that

$$\deg(L|_Z) + \sum_{i=1}^r m^{(i)} = \sum_{x_i \in Z} \alpha_i$$

(here by “component” we really mean “connected component of the vanishing locus of u_0 ”). Quasimap stability means that these degenerate contributions cannot contain any rational tails; this is really the only difference with the case of stable maps.

Indeed, we can actually push forward Gathmann’s formula along the comparison morphism

$$\chi: \overline{\mathcal{M}}_{0, \alpha}(\mathbb{P}^N | H, d) \rightarrow \overline{\mathcal{Q}}_{0, \alpha}(\mathbb{P}^N | H, d)$$

and due to Corollary 3.4 above, the only terms which change are the comb loci containing rational tails. In fact these disappear, since the restriction of the comparison map to these loci has positive-dimensional fibres:

Lemma 3.5. *Consider a rational tail component in the comb locus of the moduli space of stable maps, i.e. a moduli space of the form:*

$$\overline{\mathcal{M}}_{0,(m^{(i)})}(\mathbb{P}^N|H, d)$$

Then (assuming that $Nd > 1$) we have

$$\dim \left([\overline{\mathcal{M}}_{0,(m^{(i)})}(\mathbb{P}^N|H, d)] \cap \text{ev}_1^*(\text{pt}_H) \right) > 0$$

where $\text{pt}_H \in A^{N-1}(H)$ is a point class. Thus the pushforward along χ of any comb locus with a rational tail is 0.

Proof. This is a simple dimension count. We have

$$\begin{aligned} \dim \left([\overline{\mathcal{M}}_{0,(m^{(i)})}(\mathbb{P}^N|H, d)] \cap \text{ev}_1^*(\text{pt}_H) \right) &= (N-3) + d(N+1) + (1-m^{(i)}) - (N-1) \\ &= (Nd-1) + (d-m^{(i)}) \end{aligned}$$

from which the lemma follows because $m^{(i)} \leq d$. \square

Remark 3.6. With an eye to the future, we remark that these rational tail components contribute nontrivially to the Gromov–Witten invariants of a Calabi–Yau hypersurface in projective space, and so their absence from the quasimap recursion formula accounts for the divergence between Gromov–Witten and quasimap invariants in the Calabi–Yau case [Gat03, Rmk. 1.6].

Since we wish to apply the projection formula to Gathmann’s recursion relation, we should express the cohomological terms which appears as pull-backs:

Lemma 3.7. *We have:*

$$\begin{aligned} \chi^*(\psi_k) &= \psi_k \\ \chi^*(\text{ev}_k^* H) &= \text{ev}_k^* H \end{aligned}$$

Proof. We will actually show that:

$$\begin{aligned} \chi^* x_k^* \omega_{C/\overline{Q}} &= x_k^* \omega_{C/\overline{M}} \\ \chi^* x_k^* \mathcal{L} &= \text{ev}_k^* \mathcal{O}_{\mathbb{P}^N}(H) \end{aligned}$$

This follows by considering the following diagram:

$$\begin{array}{ccccc} & & & \mathbb{P}^N & \\ & \nearrow f & & \nwarrow & \\ C_{\overline{M}} & \xrightarrow{\sigma^{ss}} & \chi^* C_{\overline{Q}} & \xrightarrow{\quad} & C_{\overline{Q}} \\ & \searrow x_k & \downarrow x_k & \square & \downarrow x_k \\ & & \overline{\mathcal{M}}_{0,\alpha}(\mathbb{P}^N|H, d) & \xrightarrow{\chi} & \overline{\mathcal{Q}}_{0,\alpha}(\mathbb{P}^N|H, d) \end{array}$$

where σ^{ss} is the strong stabilisation map which contracts the rational tails, and so is an isomorphism near the markings. \square

Proposition 3.8. *Define the quasimap comb locus $\mathcal{D}_{\alpha,k}^Q(\mathbb{P}^N|H, d)$ as the union of the moduli spaces*

$$\mathcal{D}^Q(\mathbb{P}^N|H, A, B, M) := \overline{\mathcal{Q}}_{0,|\alpha^{(0)}|+r}(H, d_0) \times_{H^r} \prod_{i=1}^r \overline{\mathcal{Q}}_{0,(m^{(i)}) \cup \alpha^{(i)}}(\mathbb{P}^N|H, d_i)$$

where the union runs over all splittings $A = (\alpha^{(0)}, \dots, \alpha^{(r)})$ of the markings, $B = (d_0, \dots, d_r)$ of the degree and all valid multiplicities $M = (m^{(1)}, \dots, m^{(r)})$ such that the above spaces are all well-defined (in particular $|\alpha^{(0)}| + r$ and $|\alpha^{(i)}| + 1$ are all ≥ 2) and such that

$$d_0 + \sum_{i=1}^r m^{(i)} = \sum \alpha^{(0)}$$

Equip this with the sum of the (product) fundamental classes, where each term is weighted by:

$$\frac{m^{(1)} \dots m^{(r)}}{r!}$$

Then the following formula holds

$$(\alpha_k \psi_k + \text{ev}_k^* H) \cdot [\overline{\mathcal{Q}}_{0,\alpha}(\mathbb{P}^N|H, d)] = [\overline{\mathcal{Q}}_{0,\alpha+e_k}(\mathbb{P}^N|H, d)] + [\mathcal{D}_{\alpha,k}^Q(\mathbb{P}^N|H, d)].$$

Proof. This follows from [Gat02, Thm. 2.6] by pushforward along χ , using the projection formula and Lemmas 3.4, 3.5 and 3.7. \square

Remark 3.9. In the discussion above we have implicitly used the fact that there exists a commuting diagram of comb loci:

$$\begin{array}{ccc} \mathcal{D}^M(\mathbb{P}^N|H, A, B, M) & \longrightarrow & \overline{\mathcal{M}}_{0,\alpha}(\mathbb{P}^N|H, d) \\ \downarrow & & \downarrow \\ \mathcal{D}^Q(\mathbb{P}^N|H, A, B, M) & \longrightarrow & \overline{\mathcal{Q}}_{0,\alpha}(\mathbb{P}^N|H, d) \end{array}$$

The vertical arrow on the left is a product of comparison morphisms. The horizontal arrow at the top is the glueing morphism which glue together the various pieces of the comb to produce a single relative stable map. Here we explain how to define the corresponding glueing morphism for quasimaps (i.e. the bottom horizontal arrow).

Suppose for simplicity that $r = 1$ and consider an element of the quasimap comb locus, consisting of two quasimaps:

$$\begin{aligned} &((C^0, x_1^0, \dots, x_{n_0}^0, q), L^0, u_0^0, \dots, u_N^0) \\ &((C^1, x_1^1, \dots, x_{n_1}^1, q), L^1, u_0^1, \dots, u_N^1) \end{aligned}$$

such that $u^0(q) = u^1(q)$ in \mathbb{P}^N . We want to glue these quasimaps together along q . The definition of the curve is obvious; we simply take:

$$C = C^0 \cup_q C^1$$

On the other hand, glueing together the line bundles L^0 and L^1 to obtain a line bundle L over C requires a choice of scalar $\lambda \in \mathbb{G}_m$, in order to match up the fibres over q . Furthermore if the sections are to extend as well then this scalar must be chosen in such a way that it takes $(u_0^0(q), \dots, u_N^0(q)) \in (L_q^0)^{\oplus(N+1)}$ to $(u_0^1(q), \dots, u_N^1(q)) \in (L_q^1)^{\oplus(N+1)}$. Since q is not a basepoint of either quasimap (being a marking) these tuples are nonzero, and so λ is unique if it exists. Furthermore it exists if and only if these tuples belong to the same \mathbb{G}_m -orbit in \mathbb{C}^{N+1} . This is precisely the statement that $u^0(q) = u^1(q) \in \mathbb{P}^N$.

This justifies description of this space as a “comb locus.” Similar arguments apply for more general toric varieties.

4. RECURSION FORMULA IN THE GENERAL CASE

4.1. The virtual class on the space of relative quasimaps. Let X be an arbitrary toric variety (smooth and proper) and $Y \subseteq X$ a very ample hypersurface (not necessarily toric). The complete linear system associated to $\mathcal{O}_X(Y)$ defines an embedding $i : X \hookrightarrow \mathbb{P}^N$ such that $i^{-1}(H) = Y$ (for a certain hyperplane H). By the functoriality property of quasimap spaces (see Appendix B.1) we have a map:

$$k := \overline{Q}(i) : \overline{Q}_{0,n}(X, \beta) \rightarrow \overline{Q}_{0,n}(\mathbb{P}^N, d)$$

where $d = i_*\beta$. Since i is a closed embedding it follows that k is as well. Furthermore k admits a compatible perfect obstruction theory (see Appendix B.2), so we have a notion of virtual pull-back along k (which coincides with the diagonal pull-back according to Lemma C.1).

It is easy to show that k restricts to a morphism between the relative spaces, and thus we have a diagram of embeddings

$$\begin{array}{ccc} \overline{Q}_{0,\alpha}(X|Y, \beta) & \xhookrightarrow{g} & \overline{Q}_{0,\alpha}(\mathbb{P}^N|H, d) \\ \downarrow f & \square & \downarrow j \\ \overline{Q}_{0,n}(X, \beta) & \xhookrightarrow{k} & \overline{Q}_{0,n}(\mathbb{P}^N, d) \end{array}$$

which one can show is cartesian. As such we can define a virtual class on $\overline{Q}_{0,\alpha}(X|Y, \beta)$ by pullback along k (virtual or diagonal):

$$[\overline{Q}_{0,\alpha}(X|Y, \beta)]^{\text{virt}} := k^![\overline{Q}_{0,\alpha}(\mathbb{P}^N|H, d)]$$

We use this class to define relative quasimap invariants in general:

$$\langle \gamma_1 \psi_1^{k_1}, \dots, \gamma_n \psi_n^{k_n} \rangle_{0,\alpha,\beta}^{X|Y} := \int_{[\overline{Q}_{0,\alpha}(X|Y,\beta)]^{\text{virt}}} \prod_{i=1}^n \text{ev}_i^*(\gamma_i) \cdot \psi_i^{k_i}$$

These invariants will play a role in our proof of the mirror formula in §5.

4.2. Relative spaces pull back. The idea is to prove the recursion formula for (X, Y) by pulling back the formula for (\mathbb{P}^N, H) along k . In order to do this, we need to understand how the various virtual classes involved in the formula pull back along this map. The first two terms pull back by the very definition of the virtual class:

Lemma 4.1. $k^![\overline{\mathcal{Q}}_{0,\alpha}(\mathbb{P}^N|H, d)] = [\overline{\mathcal{Q}}_{0,\alpha}(X|Y, \beta)]^{\text{virt}}$

It thus remains to consider the third term, namely the virtual class of the comb locus. This is the technical heart of the proof.

4.3. Comb loci pull back. Recall that we can write $\mathcal{D}_{\alpha,k}^Q(X|Y, \beta)$ as a union of comb loci

$$\mathcal{D}^Q(X|Y, A, B, M) := \overline{\mathcal{Q}}_{0,A_0 \cup \{q_1, \dots, q_r\}}(Y, \beta_0) \times_{Y^r} \prod_{i=1}^r \overline{\mathcal{Q}}_{0,\alpha^{(i)} \cup (m_i)}(X|Y, \beta_i)$$

where A and B are partitions of the marked points and curve class respectively, and $M = (m_1, \dots, m_r)$ records the intersection multiplicity with Y at the nodes connecting the internal component to the external components (the spine of the comb to the teeth). Since the virtual class on $\mathcal{D}_{\alpha,k}^Q(X|Y, \beta)$ is equal to the sum of the virtual classes of the $\mathcal{D}^Q(X|Y, A, B, M)$, we can deal with each of these comb loci separately.

Remark 4.2. Note that Y is not in general toric, and so we should clarify what we mean by:

$$\overline{\mathcal{Q}}(Y) = \overline{\mathcal{Q}}_{0,A_0 \cup \{q_1, \dots, q_n\}}(Y, \beta_0)$$

There are two possibilities here: one is to *define* this space as the cartesian product

$$\begin{array}{ccc} \overline{\mathcal{Q}}(Y) & \longrightarrow & \overline{\mathcal{Q}}(H) \\ \downarrow & \square & \downarrow \\ \overline{\mathcal{Q}}(X) & \xrightarrow{k} & \overline{\mathcal{Q}}(\mathbb{P}^N) \end{array}$$

and equip it with the virtual class pulled back along k :

$$[\overline{\mathcal{Q}}(Y)]^{\text{virt}} := k^![\overline{\mathcal{Q}}(H)]$$

Using this definition, $\overline{\mathcal{Q}}(Y)$ consists of those quasimaps in $\overline{\mathcal{Q}}(X)$ for which $u_Y \equiv 0$.

This has obvious advantages from the point of view of our computations, but is conceptually unsatisfying. On the other hand, $Y \subseteq X$ defines a $(\mathbb{C}^*)^r$ -invariant subvariety in the prequotient of X , which we refer to (by analogy with the case $X = \mathbb{P}^N$) as the *cone* of Y :

$$C(Y) \subseteq \mathbb{A}^{\Sigma_X(1)}$$

Then Y is equal to the GIT quotient

$$Y = C(Y) // (\mathbb{C}^*)^r$$

and so we may use the more general theory of quasimaps to GIT quotients [CFKM14] to define $\overline{\mathcal{Q}}(Y)$ and its virtual class.

We should then show that these two definitions of $\overline{\mathcal{Q}}(Y)$ agree (i.e. that there exists an isomorphism between these moduli spaces which preserves the virtual class). This is carried out in Appendix B.4.

Now, the comb locus sits inside the full product

$$\mathcal{E}^Q(X|Y, A, B, M) := \overline{\mathcal{Q}}_{0, A_0 \cup \{q_1, \dots, q_r\}}(Y, \beta_0) \times \prod_{i=1}^r \overline{\mathcal{Q}}_{0, \alpha^{(i)} \cup (m_i)}(X|Y, \beta_i)$$

which we may endow with the product virtual class (with weighting as before):

$$[\mathcal{E}^Q(X|Y, A, B, M)]^{\text{virt}} := \left(\frac{m^{(1)} \cdots m^{(r)}}{r!} \right) \cdot \left([\overline{\mathcal{Q}}_{0, A_0 \cup \{q_1, \dots, q_r\}}(Y, \beta_0)]^{\text{virt}} \times \prod_{i=1}^r [\overline{\mathcal{Q}}_{0, \alpha^{(i)} \cup (m_i)}(X|Y, \beta_i)]^{\text{virt}} \right)$$

We have the following cartesian diagram

$$\begin{array}{ccc} \mathcal{D}^Q(X|Y, A, B, M) & \longrightarrow & \mathcal{E}^Q(X|Y, A, B, M) \\ \downarrow & \square & \downarrow \\ X^r & \xrightarrow{\Delta_{X^r}} & X^r \times X^r \end{array}$$

and we can use this to define a *product virtual class* on the comb locus:

$$[\mathcal{D}^Q(X|Y, A, B, M)]^{\text{virt}} := \Delta_{X^r}^! [\mathcal{E}^Q(X|Y, A, B, M)]^{\text{virt}}$$

Remark 4.3. This is the same definition of the virtual class of the comb locus that we gave in §3.2 in the case $(X, Y) = (\mathbb{P}^N, H)$.

On the other hand, there is another cartesian diagram defining the comb locus:

$$\begin{array}{ccc} \mathcal{D}^Q(X|Y, A, B, M) & \xrightarrow{k} & \mathcal{D}^Q(\mathbb{P}^N|H, A, i_*B, M) \\ \downarrow & \square & \downarrow \\ \overline{\mathcal{Q}}_{0,n}(X, \beta) & \xrightarrow{k} & \overline{\mathcal{Q}}_{0,n}(\mathbb{P}^N, d) \end{array}$$

Remark 4.4. Technically this is not quite correct: the fibre product is actually the union of comb loci over all partitions B' such that $i_*B' = i_*B$. But this subtlety makes no difference to our arguments.

Remember that we are trying to show that the virtual class of the comb locus pulls back nicely along k . The result we need is:

Lemma 4.5. *For any α we have:*

$$k^! [\mathcal{D}^Q(\mathbb{P}^N|H, A, i_*B, M)]^{\text{virt}} = [\mathcal{D}^Q(X|Y, A, B, M)]^{\text{virt}}$$

Let us introduce the following shorthand notation: we fix the the data of A, B, M and set:

$$\begin{aligned}\mathcal{D}(X|Y) &:= \mathcal{D}^Q(X|Y, A, B, M) \\ \mathcal{E}(X|Y) &:= \mathcal{E}^Q(X|Y, A, B, M) \\ \mathcal{D}(X) &:= \mathcal{D}^Q(X, A, B) \\ \mathcal{E}(X) &:= \mathcal{E}^Q(X, A, B) \\ \overline{\mathcal{Q}}(X) &:= \overline{\mathcal{Q}}_{0,n}(X, \beta)\end{aligned}$$

and similarly for (\mathbb{P}^N, H) . Here $\mathcal{D}(X)$ and $\mathcal{E}(X)$ are the centipede loci introduced in Appendix B.3; they are defined in the same way as the comb loci, except that we replace both the quasimaps to Y and the relative quasimaps to (X, Y) by quasimaps to X . There is a cartesian diagram

$$\begin{array}{ccc}\mathcal{E}(X|Y) & \longrightarrow & \mathcal{E}(\mathbb{P}^N|H) \\ \downarrow & \square & \downarrow \theta \\ \mathcal{E}(X) & \longrightarrow & \mathcal{E}(\mathbb{P}^N)\end{array}$$

and since $\mathcal{E}(\mathbb{P}^N)$ is smooth and there is a natural fundamental class on $\mathcal{E}(\mathbb{P}^N|H)$, we have a diagonal pull-back morphism $\theta^! = \theta^!_{\Delta}$ (see Appendix C).

Lemma 4.6. $[\mathcal{E}(X|Y)]^{\text{virt}} = \theta^![\mathcal{E}(X)]^{\text{virt}}$

Proof. It suffices to check that in the following cartesian diagram

$$\begin{array}{ccc}\overline{\mathcal{Q}}(Y) & \longrightarrow & \overline{\mathcal{Q}}(H) \\ \downarrow & \square & \downarrow \theta \\ \overline{\mathcal{Q}}(X) & \longrightarrow & \overline{\mathcal{Q}}(\mathbb{P}^N)\end{array}$$

we have $\theta^![\overline{\mathcal{Q}}(X)]^{\text{virt}} = [\overline{\mathcal{Q}}(Y)]^{\text{virt}}$. Depending on one's definition of $\overline{\mathcal{Q}}(Y)$ (see Remark 4.2 above) this is either true by definition or is a standard result whose proof can be found in Appendix B.4. \square

Now consider the following cartesian diagram

$$\begin{array}{ccccc}\mathcal{D}(X) & \longrightarrow & \mathcal{D}(\mathbb{P}^N) & \longrightarrow & \mathfrak{M}_{A,B}^{\text{wt}} \\ \downarrow \varphi_X & \square & \downarrow \varphi_{\mathbb{P}^N} & \square & \downarrow \psi \\ \overline{\mathcal{Q}}(X) & \xrightarrow{k} & \overline{\mathcal{Q}}(\mathbb{P}^N) & \longrightarrow & \mathfrak{M}_{0,n,\beta}^{\text{wt}}\end{array}$$

where $\mathfrak{M}_{0,n,\beta}^{\text{wt}}$ is the moduli space of prestable curves weighted by the class β [Cos06, §2] and:

$$\mathfrak{M}_{A,B}^{\text{wt}} := \mathfrak{M}_{0,A_0 \cup \{q_1, \dots, q_r\}, \beta_0}^{\text{wt}} \times \prod_{i=1}^r \mathfrak{M}_{0,A_i \cup \{q_i\}, \beta_i}^{\text{wt}}$$

The maps $\mathcal{D}(X) \rightarrow \mathfrak{M}_{A,B}^{\text{wt}}$ and $\overline{\mathcal{Q}}(X) \rightarrow \mathfrak{M}_{0,n,\beta}^{\text{wt}}$ admit relative perfect obstruction theories which are the same as the usual ones relative to the moduli spaces of *unweighted* curves. Furthermore the morphism ψ admits a perfect obstruction theory; see Appendix B.3 for details. This induces virtual pull-back morphisms, and by the splitting axiom (see Lemma B.7) we have

$$[\mathcal{D}(X)]^{\text{virt}} := \Delta_{X^r}^! [\mathcal{E}(X)]^{\text{virt}} = \psi^! [\overline{\mathcal{Q}}(X)]^{\text{virt}}$$

from which it follows by commutativity of virtual pull-backs that:

$$(4) \quad [\mathcal{D}(X)]^{\text{virt}} = \psi^! [\overline{\mathcal{Q}}(X)]^{\text{virt}} = \psi^! k^! [\overline{\mathcal{Q}}(\mathbb{P}^N)] = k^! \psi^! [\overline{\mathcal{Q}}(\mathbb{P}^N)] = k^! [\mathcal{D}(\mathbb{P}^N)]$$

Proof of Lemma 4.5. Putting all the preceding results together, we consider the cartesian digram:

$$\begin{array}{ccccc} \mathcal{D}(X|Y) & \longrightarrow & \mathcal{E}(X|Y) & \longrightarrow & \mathcal{E}(\mathbb{P}^N|H) \\ \downarrow & \square & \downarrow & \square & \downarrow \theta \\ \mathcal{D}(X) & \longrightarrow & \mathcal{E}(X) & \longrightarrow & \mathcal{E}(\mathbb{P}^N) \\ \downarrow & \square & \downarrow & & \\ X^r & \xrightarrow{\Delta_{X^r}} & X^r \times X^r & & \end{array}$$

We then have:

$$\begin{aligned} [\mathcal{D}(X|Y)]^{\text{virt}} &= \Delta_{X^r}^! [\mathcal{E}(X|Y)]^{\text{virt}} && \text{by definition} \\ &= \Delta_{X^r}^! \theta^! [\mathcal{E}(X)]^{\text{virt}} && \text{by Lemma 4.6} \\ &= \theta^! \Delta_{X^r}^! [\mathcal{E}(X)]^{\text{virt}} && \text{by commutativity} \\ &= \theta^! [\mathcal{D}(X)]^{\text{virt}} && \text{by definition} \\ &= \theta^! k^! [\mathcal{D}(\mathbb{P}^N)] && \text{by formula (4) above} \\ &= \theta^! k^! \Delta_{(\mathbb{P}^N)^r}^! [\mathcal{E}(\mathbb{P}^N)] && \text{by definition} \\ &= k^! \Delta_{(\mathbb{P}^N)^r}^! \theta^! [\mathcal{E}(\mathbb{P}^N)] && \text{by commutativity} \\ &= k^! \Delta_{\mathbb{P}^{N^r}}^! [\mathcal{E}(\mathbb{P}^N|H)] && \text{by Lemma 4.6} \\ &= k^! [\mathcal{D}(\mathbb{P}^N|H)] && \text{by definition} \end{aligned}$$

Summing over all the components of $\mathcal{D}_{\alpha,k}^Q(\mathbb{P}^N|H, d)$ we obtain the result. \square

Theorem 4.7. *Let X be a smooth and proper toric variety and let $Y \subseteq X$ be a very ample hypersurface (not necessarily toric). Then, with the set-up as in the preceding discussion, we have an equality*

$$(\alpha_k \psi_k + e v_k^*[Y])[\overline{\mathcal{Q}}_{0,\alpha}(X|Y, \beta)]^{\text{virt}} = [\overline{\mathcal{Q}}_{0,\alpha+e_k}(X|Y, \beta)]^{\text{virt}} + [\mathcal{D}_{\alpha,k}^{\mathcal{Q}}(X|Y, \beta)]^{\text{virt}}$$

in the Chow group of $\overline{\mathcal{Q}}_{0,\alpha}(X|Y, \beta)$.

Proof. Apply $k^!$ to Proposition 3.8, using Lemmas 4.1 and 4.5. \square

5. THE QUASIMAP MIRROR FORMULA

We are going to reproduce Gathmann's proof of the mirror formula with relative stable maps [Gat03] in the context of quasimaps, thanks to the extension of his formula to this setting that we have proved in the previous sections. We have chosen to work with unparametrised quasimaps, hence the minimum number of markings is two; this minimal choice turns out to be extremely convenient because it determines the shape of the source curve to a high degree, so to grant a great level of control on degenerate contributions appearing in Gathmann's algorithm. The absence of rational tails in the quasimap moduli space makes the recursion much simpler, even in the CY case.

We would like to think of this as a Lefschetz-type theorem, in that it expresses certain (restricted) quasimap invariants of a hypersurface Y in terms of those of the ambient space X . As it turns out, we have also retrieved a formula of Ciocan-Fontanine and Kim [CFK14, Corollary 5.5.1] (but with more restrictive assumptions on the target); under this new light, the formula can be simply interpreted as a relation between some residues for the \mathbb{G}_m -action on the space of 0-pointed and 1-pointed *parametrised* quasimap invariants of the hypersurface Y . It is remarkable how, knowing only about a small sector (i.e. invariants with few insertions), it is possible to formally reconstruct the full quasimap potential; a point which was greatly clarified to us by the discussion in [CFK14, Section 5.5].

We are going to be interested in the following **setup**: X is a smooth projective toric variety and Y is a smooth *very ample* hypersurface in it, satisfying the following *semi-positivity assumption*, that $-K_Y$ is nef. Notice that, by adjunction, it follows from our hypotheses that $-K_X$ is positive (at least) on every effective curve class *coming from* Y . Let us denote by r the dimension of X and assume it is *at least* 3. Then, in fact, every curve class on X comes from Y (by Lefschetz's hyperplane theorem) and X is *Fano*.

Denote dual bases for $H^*(X; \mathbb{Q})$ by η^i and η_i ($i = 0, \dots, k$), with $\eta^0 = \mathbb{1}_X$ and $\eta^1 = Y$, which induce bases $\rho_i = i^* \eta_i$ for $i^* H^*(X)$ (extend it to a basis of $H^*(Y)$ by adding $\rho_{k+1}, \dots, \rho_{k'}$) and dually $\rho^i, i = 1, \dots, k'$; notice that the class of a point on Y is given by restricting the dual of η^1 , i.e. it is ρ_1 , while the class of a point on X is annihilated when restricted to Y , i.e. $\rho_0 = 0$. Furthermore, remark that, everytime we look at a relative space

$\overline{\mathcal{Q}}_{0,(m,0)}(X|Y, \beta)$ with $m > 0$, the evaluation map $\text{ev}_1: \overline{\mathcal{Q}}_{0,(m,0)}(X|Y, \beta) \rightarrow X$ factors through Y (so all the insertions can be first pulled back to Y).

Definition 5.1. Let X be a smooth projective toric variety (or a complete intersection in a toric variety, or more generally any GIT quotient for which the quasimap spaces are defined), and consider

$$S_0^X(z, \beta) = (\text{ev}_1)_* \left(\frac{1}{z - \psi_1} [\overline{\mathcal{Q}}_{0,2}(X, \beta)]^{\text{vir}} \right)$$

for every effective curve class $\beta \in H_2^+(X, \mathbb{Z})$. Set $S_0^X(z, 0) = \mathbb{1}_X$ and

$$S_0^X(z, q) = \sum_{\beta \geq 0} S_0^X(z, \beta) q^\beta.$$

Theorem 5.2. Let X be a toric Fano variety of dimension at least 3, and $i: Y \subseteq X$ a very ample hypersurface such that $-K_Y$ is nef. Then

$$(5) \quad \frac{\sum_{\beta \geq 0} q^\beta \prod_{j=0}^{Y \cdot \beta} (Y + jz) S_0^X(z, \beta)}{P_0(q)} = i_* S_0^Y(z, q)$$

where

$$\begin{aligned} P_0(q) &= 1 + \sum_{\substack{\beta > 0: \\ K_Y \cdot \beta = 0}} (Y \cdot \beta) q^\beta \langle [pt_Y], \mathbb{1}_X \rangle_{\overline{\mathcal{Q}}_{0,(Y \cdot \beta, 0)}(X|Y, \beta)} \\ &= 1 + \sum_{\substack{\beta > 0: \\ K_Y \cdot \beta = 0}} q^\beta (Y \cdot \beta)! \langle \psi_1^{Y \cdot \beta - 1} [pt_X], \mathbb{1}_X \rangle_{\overline{\mathcal{Q}}_{0,2}(X, \beta)}. \end{aligned}$$

Proof. Define

$$S_{0,(m)}^{X|Y}(z, \beta) = (\text{ev}_1)_* \left(\frac{1}{z - \psi_1} [\overline{\mathcal{Q}}_{0,(m,0)}(X|Y, \beta)]^{\text{vir}} \right),$$

which coincides with the absolute S_0 -function defined above for $m = 0$, and

$$T_{(m)}^{X|Y}(z, \beta) = (\text{ev}_1)_* \left(m [\overline{\mathcal{Q}}_{0,(m,0)}(X|Y, \beta)]^{\text{vir}} + \frac{1}{z - \psi_1} [D_m^Q(X|Y, \beta)]^{\text{vir}} \right).$$

Then, by Gathmann's formula, we can prove that


$$(6) \quad (Y + mz) S_{0,(m)}^{X|Y}(z, \beta) = S_{0,(m+1)}^{X|Y}(z, \beta) + T_{(m)}^{X|Y}(z, \beta),$$

from which it follows that

$$\prod_{j=0}^{Y \cdot \beta} (Y + jz) S_0^X(z, \beta) = \sum_{m=0}^{Y \cdot \beta} \prod_{j=m+1}^{Y \cdot \beta} (Y + jz) T_{(m)}^{X|Y}(z, \beta).$$

It is now a matter of evaluating the RHS. Notice that $T_{(m)}^{X|Y}(z, \beta)$ is made of two parts:

- the *boundary terms*: since there are only two markings and the first one is required to lie in Y , the strong stability condition for quasimaps forces the shape of the source curve to be that of a snake which the hypersurface cuts into two pieces, the internal one of degree $\beta^{(0)}$, and the external one of degree $\beta^{(1)}$ and multiplicity $m^{(1)}$ of contact with Y , with the first marked point belonging to the internal component and the second to the external one.



The invariants which we need to consider will hence be of the form

$$\langle i^* \eta_i \psi_1^j, \rho^h \rangle_{\overline{\mathcal{Q}}_{0,2}(Y, \beta^{(0)})} \langle \rho_h, \mathbb{1}_X \rangle_{\overline{\mathcal{Q}}_{0,(m^{(1)},0)}(X|Y, \beta^{(1)})}, \quad h \in \{1, \dots, k'\}$$

Consider the following dimensional computation:

$$\begin{aligned} 0 \leq \text{codim } \rho^h &= \dim Y - \text{codim } \rho_h \\ &= \dim Y - \text{vdim } \overline{\mathcal{Q}}_{0,(m^{(1)},0)}(X|Y, \beta^{(1)}) \\ &= \dim Y - (\dim X - 3 - K_X \cdot \beta^{(1)} + 2 - m^{(1)}) \\ &= K_Y \cdot \beta^{(1)} - Y \cdot \beta^{(1)} + m^{(1)} \leq 0 \end{aligned}$$

where the last equality follows from adjunction, and the inequality follows from $K_Y \leq 0$ and $m^{(1)} \leq Y \cdot \beta^{(1)}$. This shows that the only non-trivial contributions are due to the classes $\beta^{(1)}$ such that $K_Y \cdot \beta^{(1)} = 0$, and the order of tangency is forced to be maximal, i.e. $m^{(1)} = Y \cdot \beta^{(1)}$. Furthermore, the only relevant insertions are $\rho^1 = \mathbb{1}_Y$ and $\rho_1 = [pt_Y]$. Finally, $m^{(1)} = Y \cdot \beta^{(1)}$ implies that

$$m = \alpha_1 = Y \cdot \beta^{(0)} + m^{(1)} = Y \cdot \beta,$$

hence the boundary contributions do not show up until the very end of the process of “increasing the multiplicity”.

- The remaining term in $T_{(m)}^{X|Y}(z, \beta)$ is $m(\text{ev}_1)_*[\overline{\mathcal{Q}}_{0,(m,0)}(X|Y, \beta)]^{\text{vir}}$; notice that it only gets insertions from the cohomology of X (restricted to Y). On the other hand

$$\text{vdim } \overline{\mathcal{Q}}_{0,(m,0)}(X|Y, \beta) = \dim X - 3 - K_X \cdot \beta + 2 - m \geq r - 1$$

because $m \leq Y \cdot \beta$ and $-(K_X + Y) \cdot \beta \geq 0$, by adjunction, projection formula, and for every effective curve class β (coming from Y , but saying this is superfluous by Lefschetz's hyperplane theorem as we have already remarked); since the restriction of the class $[pt_X]$ to Y vanishes, the only insertion that contributes is η_1 (by definition of a dual basis, all other dimension 1 classes vanish when restricted to Y), forcing the equality $m = Y \cdot \beta$, so that again this correction term is non-trivial only in the last step of the algorithm.

So, in the end, we see that equation 6 reduces to

$$\begin{aligned} \prod_{j=0}^{Y \cdot \beta} (Y + jz) S_0^X(z, \beta) &= T_{(Y, \beta)}^{X|Y}(z, \beta) \\ &= \sum_{i=1, \dots, k; j \geq 0} z^{j+1} \eta^i \langle \rho_i \psi_1^j, \mathbb{1}_Y \rangle_{\overline{\mathcal{Q}}_{0,2}(Y, \beta)} \\ &+ \sum_{\substack{0 < \beta^{(0)} < \beta \\ \beta^{(0)} + \beta^{(1)} = \beta}} z^{j+1} \eta^i \langle \rho_i \psi_1^j, \mathbb{1}_Y \rangle_{\overline{\mathcal{Q}}_{0,2}(Y, \beta^{(0)})} (Y \cdot \beta^{(1)}) \langle [pt_Y], \mathbb{1}_X \rangle_{\overline{\mathcal{Q}}_{0,(Y, \beta^{(1)}, 0)}(X|Y, \beta^{(1)})} \\ &+ \eta^1 (Y \cdot \beta) \langle [pt_Y], \mathbb{1}_X \rangle_{\overline{\mathcal{Q}}_{0,(Y, \beta, 0)}(X|Y, \beta)} \end{aligned}$$

if β is such that $K_Y \cdot \beta = 0$ (which implies $K_Y \cdot \beta^{(1)} = 0$ as well, for every effective decomposition $\beta = \beta^{(0)} + \beta^{(1)}$, due to the semi-positivity assumption on Y); while, if $K_Y \cdot \beta < 0$, it simply reduces to

$$\prod_{j=0}^{Y \cdot \beta} (Y + jz) S_0^X(z, \beta) = \sum_{i=1, \dots, k; j \geq 0} z^{j+1} \eta^i \langle \rho_i \psi_1^j, \mathbb{1}_Y \rangle_{\overline{\mathcal{Q}}_{0,2}(Y, \beta)} = i_* S_0^Y(z, \beta).$$

The proof of the first claim is now evident. We are left with evaluating $P(q)$.

In order to do that, we use again Gathmann's algorithm, this time in the opposite direction, to go all the way back to X ; so it starts:

$$[\overline{\mathcal{Q}}_{0,(Y, \beta, 0)}(X|Y, \beta)]^{\text{vir}} = (Y + (Y \cdot \beta - 1) \psi_1) [\overline{\mathcal{Q}}_{0,(Y, \beta - 1, 0)}(X|Y, \beta)]^{\text{vir}} - [D_{Y, \beta}^Q(X|Y, \beta)]^{\text{vir}}$$

When looking at the boundary, the invariants that come into play are of the form

$$\langle [pt_Y], \rho^h \rangle_{\overline{\mathcal{Q}}_{0,2}(Y, \beta^{(0)})} \langle \rho_h, \mathbb{1}_X \rangle_{\overline{\mathcal{Q}}_{0,(Y, (\beta - \beta^{(0)}) - 1, 0)}(X|Y, \beta - \beta^{(0)})}$$

but notice that they must vanish by dimensional reasons, since

$$\text{codim}(\rho^h) = \dim Y - 3 + 2 - K_Y \cdot \beta^{(0)} - \dim Y = -1.$$

So

$$\begin{aligned}
(Y \cdot \beta) \langle [pt_Y], \mathbb{1}_X \rangle_{\overline{\mathcal{Q}}_{0,(Y,\beta,0)}(X|Y,\beta)} &= \\
= (Y \cdot \beta) \int_{[\overline{\mathcal{Q}}_{0,2}(X,\beta)]^{\text{vir}}} \text{ev}_1^*(\eta_1) \prod_{j=0}^{Y \cdot \beta - 1} (\text{ev}_1^* Y + j \psi_1) &= \\
= (Y \cdot \beta)! \langle [pt_X] \psi_1^{Y \cdot \beta - 1}, \mathbb{1}_X \rangle_{\overline{\mathcal{Q}}_{0,2}(X,\beta)}. &
\end{aligned}$$

the second equality because $Y \cdot \eta_1 = [pt_X]$ and $Y^2 \cdot \eta_1 = 0$. \square

Corollary 5.3. *If Y is itself Fano, then there is no correction term*

$$\sum_{\beta \geq 0} q^\beta \prod_{j=0}^{Y \cdot \beta} (Y + jz) S_0^X(z, \beta) = i_* S_0^Y(z, q)$$

Corollary 5.4. *Let Y_5 be the quintic three-fold in \mathbb{P}^4 . Then*

$$i_* S_0^{Y_5}(z, q) = \frac{I_{\text{small}}^{Y_5}(z, q)}{P^{Y_5}(q)},$$

where

$$I_{\text{small}}^{Y_5}(z, q) = 5H + \sum_{d>0} \frac{\prod_{j=0}^{5d} (H + jz)}{\prod_{j=0}^d (H + jz)^5} q^d$$

and

$$P^{Y_5}(q) = 1 + \sum_{d>0} \frac{(5d)!}{(d!)^5} q^d.$$

Remark 5.5. This formula (and, more generally, formulae for concavex bundles over products of projective spaces) was already obtained in [CZ14, Theorem 1] via equivariant localisation.

5.1. Comparison with the work of Ciocan-Fontanine and Kim. We would like to compare our formula to [CFK14, Corollary 5.5.1].

In [CFK14, Section 5] they introduce (in the more general context of ϵ -stable quasimaps to GIT quotients)

- the J^ϵ -function:

$$J^\epsilon(\mathbf{t}, z) = \sum_{k \geq 0, \beta \geq 0} q^\beta (\text{ev}_\bullet)_* \left(\frac{\prod_{i=1}^k \text{ev}_i^*(\mathbf{t})}{k!} \cap \text{Res}_{F_0} [\overline{\mathcal{Q}} G_{0,k}^\epsilon(Y, \beta)]^{\text{vir}} \right)$$

- the S^ϵ -operator

$$S^\epsilon(z)(\gamma) = \sum_{m \geq 0, \beta \geq 0} \frac{q^\beta}{m!} (\text{ev}_1)_* \left(\frac{[\overline{\mathcal{Q}}_{0,2+m}^\epsilon(Y, \beta)]^{\text{vir}}}{z - \psi} \text{ev}_2^*(\gamma) \prod_{j=3}^{2+m} \text{ev}_j^*(\mathbf{t}) \right)$$

- the P^ϵ -series

$$P^\epsilon(\mathbf{t}, z) = \sum_h \rho^h \sum_{m \geq 0, \beta \geq 0} \frac{q^\beta}{m!} [\overline{QG}_{0,1+m}^\epsilon(Y, \beta)] \cap \text{ev}_1^*(\rho_h p_\infty)$$

where $p_\infty \in H_{\mathbb{G}_m}^*(\mathbb{P}^1)$ is defined via its restrictions to the \mathbb{G}_m -fixed points: $p_{\infty|0} = 0, p_{\infty|\infty} = -z$.

They prove by localisation that [CFK14, Theorem 5.4.1]

$$J^\epsilon(z) = S^\epsilon(z)(P^\epsilon).$$

Furthermore, they prove that, restricting to $\mathbf{t} = 0$ and semi-positive targets, the only class that matches non-trivially with $P^\epsilon|_{\mathbf{t}=0}$ is $[pt_Y]$, and the above formula takes the simpler form of a product [CFK14, Corollary 5.5.1]

$$\frac{J^\epsilon|_{\mathbf{t}=0}}{\langle [pt_Y], P^\epsilon|_{\mathbf{t}=0} \rangle} = \mathbb{1}_Y + \sum_h \rho^h \left(\sum_{\beta \neq 0} q^\beta \left\langle \frac{\rho_h}{z - \psi}, \mathbb{1}_Y \right\rangle_{0,2,\beta}^\epsilon \right).$$

Notice that the restriction of $S^\epsilon(z)(\mathbb{1}_Y)$ to $\mathbf{t} = 0$ that appears on the RHS of this formula coincides with what we have called $S_0^Y(z, q)$ above.

They also observe that, if we write the $\frac{1}{z}$ -expansion of $J_{\mathbf{t}=0}^\epsilon$ as

$$J_{\mathbf{t}=0}^\epsilon = J_0^\epsilon(q) \mathbb{1}_Y + O\left(\frac{1}{z}\right)$$

then $\langle [pt_Y], P^\epsilon|_{\mathbf{t}=0} \rangle = J_0^\epsilon(q)$.

Let us look more closely at $J_{\mathbf{t}=0}^\epsilon = \sum_{\beta \geq 0} q^\beta (\text{ev}_\bullet)_* (\text{Res}_{F_0} [\overline{QG}_{0,0}^\epsilon(Y, \beta)]^{\text{vir}})$. Recall that in our context $Y \subseteq X$ is a very ample hypersurface and X is toric Fano. Furthermore, set $\epsilon = 0^+$. We have the following diagram:

$$\begin{array}{ccccc} \overline{QG}_{0,0}(Y, \beta) & \longleftrightarrow & F_0^Y & \xrightarrow{\text{ev}_\bullet} & Y \\ \downarrow i & & \downarrow & & \downarrow i \\ \overline{QG}_{0,0}(X, \beta) & \longleftrightarrow & F_0^X & \xrightarrow{\text{ev}_\bullet} & X \end{array}$$

- By a slight generalisation of [CFKM14, Propositions 6.2.2 and 6.2.3], $\iota_* [\overline{QG}_{0,0}(Y, \beta)]^{\text{vir}} = e(\pi_* E_{0,0,\beta}^Y(z)) \cap [\overline{QG}_{0,0}(X, \beta)]^{\text{vir}}$ as \mathbb{G}_m -equivariant classes, where π is the universal curve on $\overline{QG}_{0,0}(X, \beta)$ and $E_{0,0,\beta}^Y(z)$ is the equivariant line bundle on it associated to $\mathcal{O}_X(Y)$. This is analagous to the bundle L_Y used in the definition of relative quasimaps (see §2.3).
- Since the fibers of π are irreducible (by the stability condition and the fact that there are no markings, there can only be the parametrised component), the following splitting holds:

$$e(\pi_* E_{0,0,\beta}^Y(z)) = \prod_{j=0}^{Y \cdot \beta} c_1(\sigma_0^* E_{0,0,\beta}^Y(z) \otimes \omega_\pi^{\otimes j})$$

coming from evaluating at (the j -th order infinitesimal thickening of) the zero section σ_0 and the jet bundles exact sequence:

$$0 \longrightarrow \pi_*(E_{0,0,\beta}^Y(-j\sigma_0)) \longrightarrow \pi_*E_{0,0,\beta}^Y \longrightarrow \sigma_0^*P^{j-1}(E_{0,0,\beta}^Y) \longrightarrow 0$$

$$0 \longrightarrow \Omega_\pi^{\otimes j} \otimes E_{0,0,\beta}^Y \longrightarrow P^j(E_{0,0,\beta}^Y) \longrightarrow P^{j-1}(E_{0,0,\beta}^Y) \longrightarrow 0$$

which, restricting to F_0^X , gives:

$$\iota_*[F_0^Y]^{\text{vir}} = \prod_{j=0}^{Y \cdot \beta} (Y + iz)[F_0^X]^{\text{vir}}.$$

- The small J^{0+} -function for toric varieties has been evaluated by Givental [Giv96b][CFK10, Definition 7.2.8]:

$$(\text{ev}_\bullet)_* \frac{[F_0^X]^{\text{vir}}}{e(N_{F_0/\overline{Q}G_{0,0}(X,\beta)})} = \prod_{\rho \in \Sigma_X(1)} \frac{\prod_{j=-\infty}^0 (D_\rho + jz)}{\prod_{j=-\infty}^{\int_\beta D_\rho} (D_\rho + jz)} = \frac{\prod_{\rho \in \Sigma_X(1): D_\rho \cdot \beta \leq 0} (D_\rho + jz)}{\prod_{\rho \in \Sigma_X(1): D_\rho \cdot \beta > 0} (D_\rho + jz)}$$

So, using $\sum_{\rho \in \Sigma_X(1)} D_\rho = -K_X$ and $(Y + K_X) \cdot \beta = 0$, we see that

$$J_0^Y(q) = \sum_{\beta \geq 0} q^\beta (Y \cdot \beta)! \frac{\prod_{\rho \in \Sigma_X(1): D_\rho \cdot \beta < 0} (-1)^{-D_\rho \cdot \beta} (-D_\rho \cdot \beta)!}{\prod_{\rho \in \Sigma_X(1): D_\rho \cdot \beta > 0} (D_\rho \cdot \beta)!}$$

- Since X is Fano, $J_{|t=0}^X = S_{|t=0}^X(\mathbb{1}_X)$.
- The coefficient $\langle [pt_X] \psi_1^{Y \cdot \beta - 1}, \mathbb{1}_X \rangle_{\overline{Q}_{0,2}(X,\beta)}$ that appears in our P -series (multiplied by $(Y \cdot \beta)!$), can be deduced from the expansion of $S_{|t=0}^X(\mathbb{1}_X)$ given above, and it turns out to be

$$\langle [pt_X], S_{|t=0}^X(\mathbb{1}_X) \rangle [z^{Y \cdot \beta}] = \frac{\prod_{\rho \in \Sigma_X(1): D_\rho \cdot \beta < 0} (-1)^{-D_\rho \cdot \beta} (-D_\rho \cdot \beta)!}{\prod_{\rho \in \Sigma_X(1): D_\rho \cdot \beta > 0} (D_\rho \cdot \beta)!}.$$

So we may conclude that the i_* of [CFK14, Corollary 5.5.1] coincides with our Equation 5.


APPENDIX A. THE COMPARISON MORPHISM

We summarise the existence of the comparison morphism for \mathbb{P}^r and how it implies that GW and quasimap invariants of projective space coincide. This has been proven in [MOP11, Theorem 3] and [Man12b, Section 4.3] (but see also [Ber00, Proposition 4.1] and [PR03, Theorem 7.1] for inspiration). We shall try to clarify as many details as possible, for our own benefit and, hopefully, that of the novice reader.

In order to give a morphism $\chi: \overline{\mathcal{M}}_{g,n}(\mathbb{P}^r, d) \rightarrow \overline{\mathcal{Q}}_{g,n}(\mathbb{P}^r, d)$ we need to be able to canonically associate a family of quasimaps on a base S to any family of stable maps on the same base.


The pointwise construction is the following: a stable map has no base points, so the only thing that might prevent it from being a stable quasimap is the presence of rational tails (of positive degree, by the stable maps stability condition). Let $C = C^{(0)} \sqcup_{q_i} R_i$ be the source curve; the rational tail R_i has degree d_i and is joined to the permanent curve $C^{(0)}$ at the node q_i , which is the only special point on R_i ; hence all the markings belong to $C^{(0)}$. The map to \mathbb{P}^r is equivalent to the data of a line bundle $L = f^* \mathcal{O}_{\mathbb{P}^r}(1)$ on C and $r + 1$ sections s_0, \dots, s_r thereof. We associate to such a stable map the quasimap $(C^{(0)}, \mathbf{x}; L|_{C^{(0)}} \otimes \mathcal{O}_{C^{(0)}}(\sum_i d_i q_i); \hat{s}_0, \dots, \hat{s}_r)$, where \hat{s}_j is the restriction of s_j to $C^{(0)}$, seen as a section of $L|_{C^{(0)}} \otimes \mathcal{O}_{C^{(0)}}(\sum_i d_i q_i)$ through the inclusion $L|_{C^{(0)}} \hookrightarrow L|_{C^{(0)}} \otimes \mathcal{O}_{C^{(0)}}(\sum_i d_i q_i)$. Notice that the resulting quasimap has a base-point of order d_i at q_i .

The construction in families requires us to find a line bundle on the universal curve that is trivial on the rational tails and relatively ample elsewhere. This can be performed at the level of Picard stacks: let $\mathfrak{Pic}_{g,n}^{d, \text{st}}$ be the open substack of $\mathfrak{Pic}(\pi: \mathfrak{C}_{g,n} \rightarrow \mathfrak{M}_{g,n})$ obtained by requiring that the total degree of the line bundle is d , the multi-degree is nonnegative and $\mathcal{L} \otimes \omega_\pi^{\log}$ is ample relative to π , where \mathcal{L} is the universal line bundle. Let T^δ be the locus in the universal curve over $\mathfrak{Pic}_{g,n}^{d, \text{st}}$ spanned by rational tails on which \mathcal{L} has degree δ ; this is a Cartier divisor by deformation theory and smoothness of the stack $\mathfrak{C}_{\mathfrak{Pic}}$. Notice that T^{δ_0} and T^{δ_1} (say $\delta_0 < \delta_1$) do intersect in a stratum of codimension 1 in both of them, where the rational tail splits into two rational components, the furthest from $C^{(0)}$ having degree δ_0 .



Claim: the line bundle $\mathcal{M} = \mathcal{L} \otimes \omega_\pi^{\log} \otimes \bigotimes_{0 < \delta \leq d} \mathcal{O}_{\mathfrak{C}}((\delta - 1)T^\delta)$ on $\mathfrak{C}_{\mathfrak{Pic}}$ has degree 0 on every component of every rational tail, and is π -relatively ample elsewhere.

Proof. Consider a curve $C^{(0)} \sqcup_q R$ with a rational tail of degree δ , such that R consists of n many components $R^{(1)}, \dots, R^{(n)}$, each of degree $\delta^{(1)}, \dots, \delta^{(n)}$ respectively, numbered from the closest to the farthest from $C^{(0)}$; set $T_i = \bigcup_{j=i}^n R_j$ and $\epsilon_i = \delta - 1 - \sum_{j=1}^{i-1} \delta_j$.



A general one-parameter family in $\mathfrak{Pic}_{g,n}^{d,st}$ will give us a smoothing of such a curve; the universal curve over such a family is a normal surface S ; we can compute the degree of the restriction of \mathcal{M} to components of the central fiber of this family by first restricting \mathcal{M} to S , and then using intersection theory on this normal surface.

Notice that restricting $\bigotimes_{0 < \delta \leq d} \mathcal{O}_{\mathbb{C}}((\delta-1)T^\delta)$ to this family gives $\mathcal{O}_S(\sum_{j=1}^n \epsilon_j T_j)$. Since $R^{(i)}$ is a (-2) -curve for $i = 1, \dots, n-1$, and $R^{(n)}$ is a (-1) -curve, we get

$$R^{(i)} \cdot T_j = \begin{cases} 0, & \text{for } j < i \\ -1, & \text{for } j = i \\ 1, & \text{for } j = i+1 \\ 0 & \text{for } j > i+1 \end{cases}$$

hence $\deg(\mathcal{M}|_{R^{(i)}}) = \delta^{(i)} - \epsilon_i + \epsilon_{i+1} = 0$ for $i = 1 \dots, n-1$, while for $i = n$ it is $\delta^{(n)} - 1 - \epsilon_n = 0$, as ω^{\log} is trivial on the (-2) curves and has degree -1 on $R^{(n)}$. The last assertion of the claim follows from the stability condition and the fact that $\mathcal{O}_{\mathbb{C}}(T^\delta)$ is effective when restricted to $C^{(0)}$. \square

By taking the relative Proj construction we obtain another curve $\hat{\mathbb{C}} = \underline{\text{Proj}}_{\mathfrak{Pic}} \left(\bigoplus_{k \geq 0} \pi_* \mathcal{M}^{\otimes k} \right)$ over $\mathfrak{Pic}_{g,n}^{d,st}$, with a map ρ that contracts the rational tails

$$\begin{array}{ccc} \mathbb{C}_{\mathfrak{Pic}} & \xrightarrow{\rho} & \hat{\mathbb{C}} \\ & \searrow \pi & \downarrow \pi' \\ & & \mathfrak{Pic}_{g,n}^{d,st} \end{array}$$

It is flat because it is a family of genus g curves over a reduced base. Furthermore, it can be checked by cohomology and base-change [Har77, Theorem 12.11][Knu83, Corollary 1.5] (notice that the fibers of ρ are either points or rational curves) that $\hat{\mathcal{L}} = \rho_* \left(\mathcal{L} \otimes \bigotimes_{0 < \delta \leq d} \mathcal{O}_{\mathfrak{C}}(\delta T^\delta) \right)$ is a line bundle on $\hat{\mathfrak{C}}$ of degree d relative to π' (such that $\rho^* \hat{\mathcal{L}} \simeq \mathcal{L} \otimes \bigotimes_{0 < \delta \leq d} \mathcal{O}_{\mathfrak{C}}(\delta T^\delta)$), hence the universal property gives us a commutative diagram (with Cartesian square)

$$\begin{array}{ccccc} \mathfrak{C}_{\text{Pic}} & \xrightarrow{\rho} & \hat{\mathfrak{C}} & \longrightarrow & \mathfrak{C}_{\text{Pic}} \\ & \searrow \pi & \downarrow \pi' & \square & \downarrow \pi \\ & & \mathfrak{Pic}_{g,n}^{d,\text{st}} & \xrightarrow{\chi'} & \mathfrak{Pic}_{g,n}^{d,\text{st}} \end{array}$$

The very same construction, with the line bundles pulled back from the Picard stack, and the sections of \mathcal{L} seen as sections of $\mathcal{L} \otimes \bigotimes_{0 < \delta \leq d} \mathcal{O}_{\mathfrak{C}}(\delta T^\delta)$ through the inclusion of line bundles ($\mathcal{O}_{\mathfrak{C}}(T^\delta)$ is effective), and descended to sections of $\hat{\mathcal{L}}$ on $\hat{\mathfrak{C}}$ gives us the comparison morphism $\chi: \overline{\mathcal{M}}_{g,n}(\mathbb{P}^r, d) \rightarrow \overline{\mathcal{Q}}_{g,n}(\mathbb{P}^r, d)$, fitting in a commutative diagram

$$\begin{array}{ccc} \overline{\mathcal{M}}_{g,n}(\mathbb{P}^r, d) & \xrightarrow{\chi} & \overline{\mathcal{Q}}_{g,n}(\mathbb{P}^r, d) \\ \downarrow v_{\mathcal{M}} & & \downarrow v_{\mathcal{Q}} \\ \mathfrak{Pic}_{g,n}^{d,\text{st}} & \xrightarrow{\chi'} & \mathfrak{Pic}_{g,n}^{d,\text{st}} \end{array}$$

and, as before,

$$\begin{array}{ccccc} C_{\mathcal{M}} & \xrightarrow{\rho} & \hat{C} = \chi^* C_{\mathcal{Q}} & \longrightarrow & C_{\mathcal{Q}} \\ & \searrow \pi_{\mathcal{M}} & \downarrow \hat{\pi} & \square & \downarrow \pi_{\mathcal{Q}} \\ & & \overline{\mathcal{M}}_{g,n}(\mathbb{P}^r, d) & \xrightarrow{\chi} & \overline{\mathcal{Q}}_{g,n}(\mathbb{P}^r, d) \end{array}$$

The comparison between virtual fundamental classes is best outlined in the arXiv version of [Man12b, Remark 5.20]. Call $v'_{\mathcal{M}} = \chi' \circ v_{\mathcal{M}}$. We may endow it with an obstruction theory by means of

$$\begin{array}{ccccccc} v_{\mathcal{M}}^* \mathbb{L}_{\chi'} & \longrightarrow & \mathbb{E}_{v'_{\mathcal{M}}} & \longrightarrow & \mathbb{E}_{v_{\mathcal{M}}} & \xrightarrow{[1]} & \longrightarrow \\ \downarrow & & \downarrow & & \downarrow & & \\ v_{\mathcal{M}}^* \mathbb{L}_{\chi'} & \longrightarrow & \mathbb{L}_{v'_{\mathcal{M}}} & \longrightarrow & \mathbb{L}_{v_{\mathcal{M}}} & \xrightarrow{[1]} & \longrightarrow \end{array}$$

Notice that χ' is a morphism (not of DM type) between smooth Artin stacks, hence we can only deduce that $\mathbb{L}_{\chi'}$ is supported in $[-1, 1]$. It is therefore easily seen that $\mathbb{E}_{v'_{\mathcal{M}}}$ is also supported in $[-1, 1]$; in order to show that

it is actually a perfect obstruction theory, consider the long exact sequence

$$\begin{aligned} 0 \rightarrow h^{-1}v_{\mathcal{M}}^*\mathbb{L}_{\chi'} &\rightarrow h^{-1}\mathbb{E}_{v_{\mathcal{M}}} \rightarrow h^{-1}\mathbb{E}_{v_{\mathcal{M}}} \\ &\rightarrow h^0v_{\mathcal{M}}^*\mathbb{L}_{\chi'} \rightarrow h^0\mathbb{E}_{v_{\mathcal{M}}} \rightarrow h^0\mathbb{E}_{v_{\mathcal{M}}} \\ &\rightarrow h^1v_{\mathcal{M}}^*\mathbb{L}_{\chi'} \rightarrow h^1\mathbb{E}_{v_{\mathcal{M}}} \rightarrow 0 \end{aligned}$$

and observe that, dually, $h^{-1}v_{\mathcal{M}}^*\mathbb{T}_{\chi'}$ injects into $h^0\mathbb{E}_{v_{\mathcal{M}}}^{\vee} \simeq h^0\mathbb{T}_{v_{\mathcal{M}}}$, because every infinitesimal automorphism of the rational tail induces a nontrivial deformation of the stable map (since the degree of the latter is positive on every component of the rational tail); we conclude that $h^1\mathbb{E}_{v_{\mathcal{M}}} = 0$.

Claim: there is a morphism of obstruction theories $\chi^*\mathbb{E}_{v_Q} \rightarrow \mathbb{E}_{v_{\mathcal{M}}}$ [Man12b, Lemma 4.19].

Dually, $\mathbb{E}_{v_{\mathcal{M}}}^{\vee} = R^{\bullet}\pi_{\mathcal{M}*}\mathcal{L}^{\oplus r+1} = R^{\bullet}\hat{\pi}_*(\rho_*\mathcal{L}^{\oplus r+1})$, while, by cohomology and base-change, $\chi^*\mathbb{E}_{v_Q}^{\vee} = R^{\bullet}\hat{\pi}_*(\hat{\mathcal{L}}^{\oplus r+1})$, where $\hat{\mathcal{L}} = \rho_*(\mathcal{L} \otimes \bigotimes_{0 < \delta \leq d} \mathcal{O}_{\mathbb{C}}(\delta T^{\delta}))$, so $\mathbb{E}_{v_{\mathcal{M}}}^{\vee} \rightarrow \chi^*\mathbb{E}_{v_Q}^{\vee}$ comes from the inclusion of line bundles on $C_{\mathcal{M}}$

$$\mathcal{L} \hookrightarrow \mathcal{L} \otimes \bigotimes_{0 < \delta \leq d} \mathcal{O}_{\mathbb{C}}(\delta T^{\delta}).$$

Claim: this morphism factors through $\mathbb{E}_{v_{\mathcal{M}}}$.

$$\begin{array}{ccccc} & & \chi^*\mathbb{E}_{v_Q} & & \\ & \swarrow \text{dashed} & \downarrow & \searrow \phi & \\ \mathbb{E}_{v_{\mathcal{M}}} & \xrightarrow{\quad} & \mathbb{E}_{v_{\mathcal{M}}} & \xrightarrow{\quad} & v_{\mathcal{M}}^*\mathbb{L}_{\chi'}[1] \end{array}$$

In order to prove that the dashed arrow exists, we need to show that ϕ is the zero map. Dually, we look at $v_{\mathcal{M}}^*\mathbb{T}_{\chi'}[-1] \xrightarrow{\phi^{\vee}} R^{\bullet}\hat{\pi}_*(\hat{\mathcal{L}}^{\oplus r+1})$. Notation: call R the rational tail, joined at the rest of the curve (which we denote by $(C^{(0)}, \mathbf{p})$ as a marked curve), at the node q , which we may occasionally think of as a (smooth) point on $C^{(0)}$. We claim that:

- $h^0(\phi^{\vee})$ is zero because: the LHS involves automorphisms of the rational tail that leave $C^{(0)}$ fixed, while the RHS involves deformations of $C^{(0)}$, so there is no possible interference.
- $h^1(\phi^{\vee})$ is zero because: **this is slightly awkward**. There are two types of possible contributions to the LHS. They correspond to either moving the node q along $C^{(0)}$, or smoothing it. The former appears in the relative tangent of χ' only if the marked curve $(C^{(0)}, \mathbf{p})$ has no automorphisms that may “move q back”, i.e. $(C^{(0)}, \mathbf{p})$ is a stable pointed curve. The latter matters only if $(C^{(0)}, q, \mathbf{p})$ has no moduli, i.e. $(C^{(0)}, \mathbf{p})$ is a rational tail with less than 3 markings. **I will try to justify why the first type vanishes under $h^1(\phi^{\vee})$, and leave the second type because I do not understand it as yet.** Look at the long exact sequence

$$\begin{aligned}
0 \rightarrow \mathrm{Hom}(\Omega_{C^{(0)}}, \mathcal{O}_{C^{(0)}}(-q - \sum p_i)) &\rightarrow \mathrm{Hom}(\Omega_{C^{(0)}}, \mathcal{O}_{C^{(0)}}(-\sum p_i)) \rightarrow \\
T_{C^{(0)}, q} \rightarrow \mathrm{Ext}^1(\Omega_{C^{(0)}}, \mathcal{O}_{C^{(0)}}(-q - \sum p_i)) &\rightarrow \mathrm{Ext}^1(\Omega_{C^{(0)}}, \mathcal{O}_{C^{(0)}}(-\sum p_i)) \rightarrow 0
\end{aligned}$$

We are interested in what happens to

$$\frac{T_{C^{(0)}, q}}{\mathrm{Im}(\mathrm{Hom}(\Omega_{C^{(0)}}, \mathcal{O}_{C^{(0)}}(-\sum p_i)))}$$

under $h^1(\phi^\vee)$. If we can show that $h^1(\phi^\vee)$ factors through $\mathrm{Ext}^1(\Omega_{C^{(0)}}, \mathcal{O}_{C^{(0)}}(-\sum p_i))$ we are in business. Indeed the natural maps

$$\begin{array}{ccccc}
\mathrm{Def}_L & \longrightarrow & \mathrm{Def}_{(C, L)} & \longrightarrow & \mathrm{Def}_C \\
\downarrow & & \downarrow & & \downarrow \\
H^1(\mathcal{O}_C) & \longrightarrow & H^1(L^{\oplus r+1}) & \longrightarrow & H^1(f^*T_{\mathbb{P}^r})
\end{array}$$

show that $h^1(\phi^\vee)$ factors through

$$\mathrm{Ext}^1(\Omega_{C^{(0)}}, \mathcal{O}_{C^{(0)}}(-q - \sum p_i)) \rightarrow \mathrm{Ext}^1(\Omega_{C^{(0)}}, \mathcal{O}_{C^{(0)}}) \rightarrow \mathrm{Ext}^1(f^*\Omega_{\mathbb{P}^r}, \mathcal{O}_{C^{(0)}}) \simeq H^1(f^*T_{\mathbb{P}^r}).$$

- $h^2(\phi^\vee)$ is zero because: $\mathbb{E}_{v_M}^\vee$ is supported in $[0, 1]$.

Now the cone $C(\phi)$ gives an obstruction theory relative to χ . A priori, it is supported in $[-2, 0]$. By the octahedral axiom




$$\begin{array}{ccccc}
\chi^*\mathbb{E}_{v_Q} & & & & \\
\downarrow \phi & \searrow \phi' & & & \\
\mathbb{E}_{v'_M} & \longrightarrow & \mathbb{E}_{v_M} & \longrightarrow & v_M^*\mathbb{L}_{\chi'}[1] \\
& & \searrow & \nearrow & \\
& & C(\phi') & & \\
\downarrow & \nearrow & & & \\
C(\phi) & & & &
\end{array}$$

it is enough to observe that $C(\phi')$ is supported in $[-1, 0]$ [Man12b, Lemma 4.20] and that $v_M^*\mathbb{L}_{\chi'}[1]$ is supported in degrees $[-2, 0]$, in order to conclude that $C(\phi) = \mathbb{E}_\chi$ is a perfect obstruction theory. The conclusion that

$$\chi_*[\overline{\mathcal{M}}_{g,n}(\mathbb{P}^r, d)]^{\mathrm{vir}} = [\overline{\mathcal{Q}}_{g,n}(\mathbb{P}^r, d)]^{\mathrm{vir}}$$

follows from the connectedness of $\overline{\mathcal{M}}_{g,n}(\mathbb{P}^r, d)$ [KP01] (hence of $\overline{\mathcal{Q}}_{g,n}(\mathbb{P}^r, d)$) and an application of the virtual push-forward theorem [Man12b, Proposition 4.21].

We shall now explain with an example the reason why a naive attempt to extend the comparison morphism to a general toric variety fails. The problem in a nutshell is that not all toric divisors are nef: a rational tail

FIGURE 1. Toric fan for \mathbb{F}_1 .FIGURE 2. Nef cone $\text{Nef}(\mathbb{F}_1)$.FIGURE 3. Mori cone $\overline{\text{NE}}(\mathbb{F}_1)$.

contained in a divisor which is not nef may have negative degree $-d$ with respect to the corresponding line bundle; when contracting such a rational tail, we shall take the line bundle $L(-dq)$, but what to do with the sections? We would like to divide them by z^d , where z is a local coordinate around q , but no condition forces such a divisibility to happen. Otherwise said, there is now an inclusion $L_{|C^{(0)}}(-dq) \hookrightarrow L_{|C^{(0)}}$, but the (restriction of the) given sections of L do not necessarily live in the image of $H^0(C^{(0)}, L_{|C^{(0)}}(-dq)) \hookrightarrow H^0(C^{(0)}, L_{|C^{(0)}})$.

A concrete example is found when looking at the Hirzebruch surface $\mathbb{F}_1 = \text{Bl}_p \mathbb{P}^1$.

$\text{Pic}(\mathbb{F}_1)$ is generated by $[D_3]$ and $[D_4]$, with relations $[D_1] = [D_3]$ and $[D_2] = [D_4] - [D_3]$, and the intersection table is given by

$$\begin{cases} D_3^2 = 0 \\ D_3 \cdot D_4 = 0 \\ D_4^2 = 1 \end{cases}$$

When thinking of \mathbb{F}_1 as a \mathbb{P}^1 -bundle over \mathbb{P}^1 , C_1 and C_3 represent the fibers of the bundle (over the toric points of \mathbb{P}^1), while C_4 (resp. C_2) is the zero/positive (resp. infinity/negative) section; when thinking of \mathbb{F}_1 as $\text{Bl}_p \mathbb{P}^1$, C_2 is the exceptional divisor, C_4 is the toric line not passing through p , and C_1, C_3 are the strict transforms of the toric lines through p .

Let us look at $\overline{\mathcal{M}}_{0,2}(\mathbb{F}_1, [C_4])$. Since $[C_4] = [C_2] + [C_3]$, there are going to be maps of the following sort: the source curve is reducible $R_1 \sqcup_q R_2$, R_1 is mapped isomorphically to a fiber (i.e. in class $[C_3]$) and R_2 is mapped isomorphically to C_2 , all the markings belong to R_1 . So R_2 is a rational tail and deserves to be contracted. Notice that the line bundle $\mathcal{O}(D_2)$ has degree -1 on R_2 (and 1 on R_1). In this case everything works well because the corresponding section $u_{2|R_1}$ must vanish at the node, so we can divide it by a chosen (once for all toric line bundles) section of $\mathcal{O}_{R_1}(q)$.

Consider now $\overline{\mathcal{M}}_{0,2}(\mathbb{F}_1, 2[C_2] + [C_3])$. Certainly there are going to be maps similar to the ones described above, with R_2 now covering C_2 2:1. The point is that $\mathcal{O}(D_2)$ has degree -2 on R_2 , but $u_{2|R_1}$ doesn't have to vanish at the node of order 2, so we are in trouble. [Something is going on here: in this case there is a boundary component where the map is of the type that we have just described, and the requirement that \$u_{2|R_1}\$ vanishes of order 2 at the node defines precisely the intersection with the main component. Check this. Could we possibly exploit this phenomenon to define a smaller compactification, possibly even smaller than quasimaps?](#)

APPENDIX B. NOTES ON QUASIMAPS

In this appendix we collect several foundational results in quasimap theory, including:

- (1) *Functoriality* (§B.1): given a morphism $f: Y \rightarrow X$ we describe the induced map:

$$\overline{Q}(f): \overline{Q}_{g,n}(Y, \beta) \rightarrow \overline{Q}_{g,n}(X, f_*\beta)$$

We also discuss (§B.2) when $\overline{Q}(f)$ admits a compatible perfect obstruction theory.

- (2) *Splitting axiom* (§B.3): this gives an equality between two natural virtual classes on boundary strata (i.e. loci where the underlying curve is reducible of a prescribed type).
- (3) *Comparison with the GIT construction* (§B.4): we show that for a (not necessarily toric) hypersurface $Y \hookrightarrow X$, our definition of $\overline{Q}(Y)$ as a

substack of $\overline{Q}(X)$ coincides with the definition of $\overline{Q}(Y)$ given by the description of Y as a GIT quotient (see [CFKM14]).

B.1. Functoriality. In the case of stable maps, a morphism $f : Y \rightarrow X$ induces a morphism between the corresponding moduli spaces

$$\overline{M}(f) : \overline{M}_{g,n}(Y, \beta) \rightarrow \overline{M}_{g,n}(X, f_*\beta)$$

given by (post)composition with f ; in general this induced morphism may involve stabilisation of the source curve. Because of this, we may say that the construction of the moduli space of stable maps is *functorial*.

It is natural to ask whether the same holds for the moduli space of quasimaps, i.e. whether we have a morphism:

$$\overline{Q}(f) : \overline{Q}_{g,n}(Y, \beta) \rightarrow \overline{Q}_{g,n}(X, f_*\beta)$$

Since here the objects of the moduli space are not maps, we cannot simply compose with f , but we should define an operation which agrees with composing with f on the locus of quasimaps without basepoints. In [CFK14, Section 3.1] a definition (in the GIT context) is given when f is an embedding into a projective space; we shall discuss the general situation of a morphism between toric varieties $f : Y \rightarrow X$.

Our approach uses the language of Σ -collections introduced by D. Cox. This approach is natural insofar as a quasimap is a generalisation of a Σ -collection. We will refer extensively to [Cox95b] and [Cox95a], which we recommend as an introduction for any readers unfamiliar with the theory.

Let X and Y be smooth and proper toric varieties with fans $\Sigma_X \subseteq N_X$ and $\Sigma_Y \subseteq N_Y$. Suppose we are given $f : Y \rightarrow X$ (which we do not assume to be a toric morphism). By [Cox95a, Theorem 1.1] the data of such a map is equivalent to a Σ_X -collection on Y :

$$((L_\rho, u_\rho)_{\rho \in \Sigma_X(1)}, (\varphi_{m_x})_{m_x \in M_X})$$

In addition, [Cox95b] allows us to describe line bundles on Y and their global sections in terms of the homogeneous coordinates $(z_\tau)_{\tau \in \Sigma_Y(1)}$. All of these observations are combined into the following theorem, which is so useful that we will state it here in its entirety:

Theorem B.1. [Cox95a, Theorem 3.2] *The data of a morphism $f : Y \rightarrow X$ is the same as the data of homogeneous polynomials*

$$P_\rho \in S_{\beta_\rho}^Y$$

for $\rho \in \Sigma_X(1)$, where $\beta_\rho \in \text{Pic } Y$ and $S_{\beta_\rho}^Y$ is the corresponding graded piece of the Cox ring

$$S^Y = k[z_\tau : \tau \in \Sigma_Y(1)]$$

This data is required to satisfy the following two conditions:

- (1) $\sum_{\rho \in \Sigma_X(1)} \beta_\rho \otimes n_\rho = 0$ in $\text{Pic } Y \otimes N_X$.
- (2) $(P_\rho(z_\tau)) \notin Z(\Sigma_X) \subseteq \mathbb{A}_k^{\Sigma_X(1)}$ whenever $(z_\tau) \notin Z(\Sigma_Y) \subseteq \mathbb{A}_k^{\Sigma_Y(1)}$.

Furthermore, two such sets of data (P_ρ) and (P'_ρ) correspond to the same morphism if and only if there exists a $\lambda \in \text{Hom}_{\mathbb{Z}}(\text{Pic } X, \mathbb{G}_m)$ such that

$$\lambda(D_\rho) \cdot P_\rho = P'_\rho$$

for all $\rho \in \Sigma_X(1)$. Finally, if we define $\tilde{f}(z_\tau) = (P_\rho(z_\tau))$ then this defines a lift of f to the prequotients:

$$\begin{array}{ccc} \mathbb{A}_k^{|\Sigma_Y(1)|} \setminus Z(\Sigma_Y) & \xrightarrow{\tilde{f}} & \mathbb{A}_k^{|\Sigma_X(1)|} \setminus Z(\Sigma_X) \\ \downarrow q_Y & & \downarrow q_X \\ Y & \xrightarrow{f} & X \end{array}$$

Aside B.2. Throughout this section we will stick to the notation established above; in particular we will use ρ to denote a ray in $\Sigma_X(1)$ and τ to denote a ray in $\Sigma_Y(1)$.

Recall our goal: given a map $f : Y \rightarrow X$ we wish to define a “push-forward” map:

$$\overline{Q}(f) : \overline{Q}_{g,n}(Y, \beta) \rightarrow \overline{Q}_{g,n}(X, f_*\beta)$$

Consider therefore a quasimap $((C, x_1, \dots, x_n), (L_\tau, u_\tau)_{\tau \in \Sigma_Y(1)}, (\varphi_{m_Y})_{m_Y \in M_Y})$ with target Y . Pick data $(P_\rho)_{\rho \in \Sigma_X(1)}$ corresponding to the map f , as in the theorem above; we will later see that our construction does not depend on this choice.

The idea of the construction is as follows. Assume for a moment that $C \simeq \mathbb{P}^1$ and that the quasimap is without basepoints, so that we have an actual morphism $C \rightarrow Y$. Then, by the theorem of Cox that we have just recalled, we can lift this morphism to the prequotient as in the following diagram

$$\begin{array}{ccccc} \mathbb{A}_k^2 \setminus \{0\} & \xrightarrow{(u_\tau)} & \mathbb{A}_k^{|\Sigma_Y(1)|} \setminus Z(\Sigma_Y) & \xrightarrow{(P_\rho)} & \mathbb{A}_k^{|\Sigma_X(1)|} \setminus Z(\Sigma_X) \\ \downarrow & & \downarrow & & \downarrow \\ C & \longrightarrow & Y & \longrightarrow & X \end{array}$$

from which it follows that the composition $C \rightarrow Y \rightarrow X$ is given in homogeneous coordinates by:


$$(P_\rho((u_\tau)_{\tau \in \Sigma_Y(1)}))_{\rho \in \Sigma_X(1)}.$$

In general the quasimap to Y will still be given by a bunch of line bundles and their sections on C , and we can still make sense of the expressions $P_\rho(u_\tau)$ as sections of line bundles on C obtained from the given ones by means of the recipe encoded by f ; morally, we can trivialise the given line bundles near a point of C and treat the above polynomial expression as a function on this neighbourhood. This will allow us to define the pushforward along f on the quasimap space.

Let us make this precise now. We shall first address the issue of stabilising the curve: in case $f_*\beta = 0$, observe that

$$\overline{\mathcal{Q}}_{g,n}(X, 0) \simeq \overline{\mathcal{M}}_{g,n} \times X$$

hence the stabilisation morphism is the usual $\overline{\mathcal{Q}}_{g,n}(X, 0) \rightarrow \overline{\mathcal{M}}_{g,n}$, while the map to X is given by postcomposing with f any of the evaluation morphisms. On the other hand it may well be that, even if $f_*\beta \neq 0$, the source curve is reducible and some of its components have a degree β_0 in Y the pushforward of which is trivial in X .



Pick a(ny) polarisation $\mathcal{O}_X(1)$ and write $f^*\mathcal{O}_X(1) = \bigotimes_{\tau} L_{\tau}^{\otimes a_{\tau}}$, with $a_{\tau} \geq 0$ (because $f^*\mathcal{O}_X(1)$ is nef). Notice that the line bundle $\omega_{\pi}(x_1 + \dots + x_n) \otimes \bigotimes_{\tau} \mathcal{L}_{\tau}^{\otimes a_{\tau}}$ on the universal curve of $\overline{\mathcal{Q}}_{g,n}(Y, \beta)$ provides the required contraction

$$\begin{array}{ccc} C_Y & \xrightarrow{\phi} & \overline{C} \\ & \searrow \pi & \swarrow \pi' \\ & \overline{\mathcal{Q}}_{g,n}(Y, \beta) & \end{array}$$

Let the sections of \overline{C} be the ones of C_Y composed with ϕ .

Resuming the notation above, for each $\rho \in \Sigma_X(1)$, P_{ρ} is a polynomial in the z_{τ} ; we can write it as

$$(7) \quad P_{\rho}(z_{\tau}) = \sum_{\underline{a}} P_{\rho}^{\underline{a}}(z_{\tau}) = \sum_{\underline{a}} \mu_{\underline{a}} \prod_{\tau} z_{\tau}^{a_{\tau}}$$

where the sum is over a finite number of multindices $\underline{a} = (a_{\tau}) \in \mathbb{N}^{|\Sigma_Y(1)|}$ and the $\mu_{\underline{a}}$ are nonzero scalars. Observe that, for each \underline{a} , the line bundle $\bigotimes_{\tau} \mathcal{L}_{\tau}^{\otimes a_{\tau}}$ on C_Y is trivial on the ρ -contracted components, which are always rational curves; hence, by cohomology and base-change, the following gives

a line bundle on \overline{C} :

$$\tilde{\mathcal{L}}_\rho^a = \phi_* \bigotimes_{\tau} \mathcal{L}_\tau^{\otimes a_\tau}$$

Then we may take the following section of $\tilde{\mathcal{L}}_\rho^a$:

$$\tilde{u}_\rho^a = P_\rho^a(u_\tau) = \mu_a \prod_{\tau} u_\tau^{a_\tau};$$

notice that such sections must be constant on ϕ -contracted components of C_Y and so descend to \overline{C} . Thus each of the terms P_ρ^a of P_ρ defines a section \tilde{u}_ρ^a of a line bundle $\tilde{\mathcal{L}}_\rho^a$. But what we want is a single section \tilde{u}_ρ of a single line bundle $\tilde{\mathcal{L}}_\rho$. This is where the isomorphisms φ_{m_Y} come in.

Recall that we have a short exact sequence:

$$(8) \quad 0 \longrightarrow M_Y \xrightarrow{\theta} \mathbb{Z}^{\Sigma_Y(1)} \longrightarrow \text{Pic } Y \longrightarrow 0$$

Let \underline{a} and \underline{b} be multindices appearing in the sum (7) above. By the homogeneity of P_ρ we have

$$\sum_{\tau} a_\tau D_\tau = \beta_\rho = \sum_{\tau} b_\tau D_\tau$$

which is precisely the statement that in the above sequence \underline{a} and \underline{b} map to the same element of $\text{Pic } Y$ (namely β_ρ). Hence there exists a unique $m_Y \in M_Y$ such that:

$$\theta(m_Y) = \underline{a} - \underline{b}$$

Now, the isomorphism φ_{m_Y} (contained in the data of our original quasimap) is a map:

$$\varphi_{m_Y} : \bigotimes_{\tau} \mathcal{L}_\tau^{\otimes \langle m_Y, n_\tau \rangle} \cong \mathcal{O}_{C_Y}$$

By definition, $\theta(m_Y) = (\langle m_Y, n_\tau \rangle)_{\tau \in \Sigma_Y(1)}$. But also $\theta(m_Y) = (a_\tau - b_\tau)_{\tau \in \Sigma_Y(1)}$. Hence we have:

$$\varphi_{m_Y} : \bigotimes_{\tau} \mathcal{L}_\tau^{\otimes a_\tau} \cong \bigotimes_{\tau} \mathcal{L}_\tau^{\otimes b_\tau}$$

In other words, we have well-defined canonical isomorphisms

$$\tilde{\mathcal{L}}_\rho^a \cong \tilde{\mathcal{L}}_\rho^b$$

for all \underline{a} and \underline{b} . Let us choose one such \underline{a} (it doesn't matter which); call it \underline{a}^ρ . We define:

$$\tilde{\mathcal{L}}_\rho := \tilde{\mathcal{L}}_\rho^{\underline{a}^\rho}$$

Then for all \underline{b} we can use the above isomorphism to view \tilde{u}_ρ^b as a section of $\tilde{\mathcal{L}}_\rho$. Summing all of these together with the appropriate coefficients we thus obtain a section \tilde{u}_ρ of $\tilde{\mathcal{L}}_\rho$, which we can write (with abuse of notation) as:

$$\tilde{u}_\rho = \sum_{\underline{a}} \mu_{\underline{a}} \prod_{\tau} u_\tau^{a_\tau}$$

Note that if we had made a different choice of \underline{a}^ρ above the result would have been isomorphic.

Thus far we have constructed line bundles and sections $(\tilde{\mathcal{L}}_\rho, \tilde{u}_\rho)_{\rho \in \Sigma_X(1)}$ on \overline{C} . It remains to define the isomorphisms

$$\tilde{\varphi}_{m_X} : \otimes_\rho \tilde{\mathcal{L}}_\rho^{\otimes \langle m_X, n_\rho \rangle} \cong \mathcal{O}_{\overline{C}}$$

for all $m_X \in M_X$. The left hand side is:

$$\otimes_\rho \tilde{\mathcal{L}}_\rho^{\otimes \langle m_X, n_\rho \rangle} = \otimes_\rho \left(\phi_* \otimes_\tau \mathcal{L}_\tau^{\otimes a_\tau^\rho} \right)^{\otimes \langle m_X, n_\rho \rangle} = \phi_* \otimes_\tau \mathcal{L}_\tau^{\otimes \left(\sum_\rho a_\tau^\rho \langle m_X, n_\rho \rangle \right)}$$

Now, for $m_Y \in M_Y$ we have isomorphisms $\varphi_{m_Y} : \otimes_\tau \mathcal{L}_\tau^{\otimes \langle m_Y, n_\tau \rangle} \cong \mathcal{O}_{C_Y}$. Hence, in order to construct $\tilde{\varphi}_{m_X}$ it is tempting to look for an m_Y such that

$$\langle m_Y, n_\tau \rangle = \sum_\rho a_\tau^\rho \langle m_X, n_\rho \rangle$$

for all $\tau \in \Sigma_Y(1)$ (we will then set $\tilde{\varphi}_{m_X} = \varphi_{m_Y}$). Consider therefore the short exact sequence (8). Recall that $\theta(m_Y) = (\langle m_Y, n_\tau \rangle)_{\tau \in \Sigma_Y(1)}$. Hence we need to show that

$$\left(\sum_\rho a_\tau^\rho \langle m_X, n_\rho \rangle \right)_{\tau \in \Sigma_Y(1)}$$

belongs to the image of θ , i.e. that it belongs to the kernel of the second map (notice that m_Y is then unique because θ is injective). This is equivalent to saying that

$$\sum_\tau \sum_\rho a_\tau^\rho \langle m_X, n_\rho \rangle D_\tau = 0 \in \text{Pic } Y$$

Now, we have

$$\sum_\tau a_\tau^\rho D_\tau = \beta_\rho$$

so that the above sum becomes

$$\sum_\rho \langle m_X, n_\rho \rangle \beta_\rho = \left\langle m_X, \sum_\rho \beta_\rho \otimes n_\rho \right\rangle = \langle m_X, 0 \rangle = 0$$

where $\sum_\rho \beta_\rho \otimes n_\rho = 0$ by Condition (1) in Theorem B.1. So there does indeed exist a (unique) $m_Y \in M_Y$ such that $\langle m_Y, n_\tau \rangle = \sum_\rho a_\tau^\rho \langle m_X, n_\rho \rangle$, so that we can set:

$$\tilde{\varphi}_{m_X} = \varphi_{m_Y} : \bigotimes_\rho \tilde{\mathcal{L}}_\rho^{\otimes \langle m_X, n_\rho \rangle} \cong \mathcal{O}_{\overline{C}}$$

Thus, we have produced a quasimap with target X and class $f_*\beta$ on the base $\overline{\mathcal{Q}}_{g,n}(Y, \beta)$:

$$(\overline{C}, (\tilde{\mathcal{L}}_\rho, \tilde{u}_\rho)_{\rho \in \Sigma_X(1)}, (\tilde{\varphi}_{m_X})_{m_X \in M_X})$$

The proof that this construction does not depend on the choice of (P_ρ) is straightforward and is left to the reader.

It remains to demonstrate that the quasimap thus constructed is nondegenerate and stable. Nondegeneracy follows immediately from Condition (2) in Theorem B.1. Put differently: the original quasimap defined a rational map $C \dashrightarrow Y$, whereas the new quasimap defines a rational map which is simply the composition $C \dashrightarrow Y \rightarrow X$. Therefore the set of basepoints is exactly the same.

Stability follows precisely from the construction of ϕ : if $\mathcal{O}_X(1) = \bigotimes_{\rho} L_{\rho}^{\otimes b_{\rho}}$, then $\omega_{\pi'}(\bar{x}_1 + \dots \bar{x}_n) \otimes \bigotimes_{\rho} \tilde{\mathcal{L}}_{\rho}^{\otimes b_{\rho}}$ will be π' -ample on \bar{C} , since we have contracted all the components on which $f^*\mathcal{O}_X(1)$ was trivial without introducing any rational tail.

Finally, by the universal property of $\bar{\mathcal{Q}}_{g,n}(X, \beta)$ there is an induced diagram

$$\begin{array}{ccccc} C_Y & \xrightarrow{\phi} & \bar{C} & \xrightarrow{\quad} & C_X \\ & \searrow \pi & \downarrow \pi' & \square & \downarrow \pi_X \\ & & \bar{\mathcal{Q}}_{g,n}(Y, \beta) & \xrightarrow{\bar{Q}(f)} & \bar{\mathcal{Q}}_{g,n}(X, f_*\beta) \end{array}$$

This completes the proof of the following.

Theorem B.3. *Let X and Y be smooth proper toric varieties and $f : Y \rightarrow X$ a morphism. Assume that f satisfies Condition (1) above. Then there exists a natural push-forward map*

$$\bar{Q}(f) : \bar{\mathcal{Q}}_{g,n}(Y, \beta) \rightarrow \bar{\mathcal{Q}}_{g,n}(X, f_*\beta)$$

which does not modify the underlying prestable curves.

Remark B.4. Theorem B.1 tells us that we can lift any morphism between toric varieties to an equivariant morphism between the prequotients

$$\begin{array}{ccc} \mathbb{A}_k^{|\Sigma_Y(1)|} \setminus Z(\Sigma_Y) & \xrightarrow{\tilde{f}} & \mathbb{A}_k^{|\Sigma_X(1)|} \setminus Z(\Sigma_X) \\ \downarrow q_Y & & \downarrow q_X \\ Y & \xrightarrow{f} & X \end{array}$$

where the torus homomorphism $G_Y = \text{Hom}_{\mathbb{Z}}(\text{Pic}(Y), \mathbb{C}^*) \rightarrow G_X = \text{Hom}_{\mathbb{Z}}(\text{Pic}(Y), \mathbb{C}^*)$ is induced in the obvious way by $f : Y \rightarrow X$. Now, thinking of quasimaps as maps to the quotient stack, functoriality is again clear by postcomposition with \tilde{f} (notice that the preimage of the unstable locus of X is the unstable locus of Y).

Finally, let us describe how this push-forward morphism behaves when f is a nonconstant map $\mathbb{P}^r \rightarrow \mathbb{P}^N$, since we will make use of this later. Write f in homogeneous coordinates as:

$$f[z_0, \dots, z_r] = [f_0(z_0, \dots, z_r), \dots, f_N(z_0, \dots, z_r)]$$

where the f_i are all homogeneous of degree $a > 0$. Then given a quasimap with target \mathbb{P}^r

$$(C, L, u_0, \dots, u_r)$$

the pushed-forward quasimap with target \mathbb{P}^N is:

$$(C, L^{\otimes a}, f_0(u_0, \dots, u_r), \dots, f_N(u_0, \dots, u_r))$$

B.2. Relative obstruction theories for $\overline{Q}(Y) \rightarrow \overline{Q}(X)$. Assume now that $f: Y \rightarrow X$ is a morphism (between projective varieties) satisfying one of the following equivalent conditions:

- (1) f is finite;
- (2) for any ample line bundle $\mathcal{O}_X(1)$ on X , $f^*\mathcal{O}_X(1)$ is ample on Y ;
- (3) for every effective curve class $\beta \in H_2^+(Y)$, $f_*\beta \neq 0$.

Observe then that the induced morphism

$$k = \overline{Q}(f): \overline{Q}_{g,n}(Y, \beta) \rightarrow \overline{Q}_{g,n}(X, f_*\beta)$$

commutes with the projections to $\mathfrak{M}_{g,n}$, i.e. there is no need to stabilise the underlying curve. We are going to describe a situation in which it is possible to define an operation of pullback along k on Chow groups.

Even in the easiest possible case when $Y \hookrightarrow X$ is a regular embedding, k itself is not necessarily a regular embedding, and so the Gysin map in the sense of [Ful98] is not guaranteed to exist.

However, when $\overline{Q}_{g,n}(X, f_*\beta)$ is a smooth stack (or rather, when it is unobstructed, for instance when $X = \mathbb{P}^N$ and $g = 0$ or $(g, n) = (1, 0)$) there is a way around this. In [Man12a] a generalisation of the Gysin map called the *virtual pull-back* is defined for morphisms endowed with a relative perfect obstruction theory. Moreover, a sufficient condition is given [Man12a, Corollary 4.9] for this map to respect the virtual classes.

Lemma B.5. *For a finite morphism of smooth toric varieties $f: Y \rightarrow X$, there exists a relative obstruction theory \mathbf{E}_k for the morphism*

$$k: \overline{Q}_{g,n}(Y, \beta) \rightarrow \overline{Q}_{g,n}(X, f_*\beta)$$

which fits into a compatible triple with the standard obstruction theories for the quasimap spaces over $\mathfrak{M}_{g,n}$. Furthermore, \mathbf{E}_k is perfect if $\overline{Q}_{g,n}(X, f_\beta)$ is unobstructed, so that:*

$$k_{\vee}^![\overline{Q}_{g,n}(X, f_*\beta)] = [\overline{Q}_{g,n}(Y, \beta)]^{\text{virt}}$$

Proof. Note first that, since k does not change the source curve of a quasimap, we indeed have a commuting triangle:

$$\begin{array}{ccc} \overline{Q}_{g,n}(Y, \beta) & \xrightarrow{k} & \overline{Q}_{g,n}(X, f_*\beta) \\ & \searrow & \swarrow \\ & \mathfrak{M}_{g,n} & \end{array}$$

We have perfect obstruction theories $\mathbf{E}_{\overline{Q}(Y)/\mathfrak{M}}$ and $\mathbf{E}_{\overline{Q}(X)/\mathfrak{M}}$ and we want to find a perfect obstruction theory \mathbf{E}_k . Consider the diagram of universal curves

$$\begin{array}{ccc} C_Y & \xrightarrow{\alpha} & C_X \\ \downarrow \pi & \square & \downarrow \rho \\ \overline{Q}_{g,n}(Y, \beta) & \xrightarrow{k} & \overline{Q}_{g,n}(X, f_*\beta) \end{array}$$

which is cartesian because k does not alter the source curve of any quasimap. We have sheaves \mathcal{F}_Y and \mathcal{F}_X on C_Y and C_X respectively such that:

$$\begin{aligned} \mathbf{E}_{\overline{Q}(Y)/\mathfrak{M}}^\vee &= \mathbf{R}^\bullet \pi_* \mathcal{F}_Y \\ \mathbf{E}_{\overline{Q}(X)/\mathfrak{M}}^\vee &= \mathbf{R}^\bullet \rho_* \mathcal{F}_X \end{aligned}$$

It follows (by flatness of ρ) that when we pull back the latter obstruction theory to $\overline{Q}(Y)$ we obtain:

$$k^* \mathbf{E}_{\overline{Q}(X)/\mathfrak{M}}^\vee = \mathbf{R}^\bullet \pi_* \alpha^* \mathcal{F}_X$$

To construct a compatible triple, we require a morphism $k^* \mathbf{E}_{\overline{Q}(X)/\mathfrak{M}} \rightarrow \mathbf{E}_{\overline{Q}(Y)/\mathfrak{M}}$. Dually, it is therefore enough to construct a morphism of sheaves on C_Y

$$\mathcal{F}_Y \rightarrow \alpha^* \mathcal{F}_X$$

and then apply $\mathbf{R}^\bullet \pi_*$. This is analogous to the morphism $f^* T_Y \rightarrow f^* T_X|_Y$ which is used in the stable maps setting. However the construction for quasimaps requires a little more ingenuity, because we do not have access to a universal map f .

The sheaf \mathcal{F}_Y is defined on C_Y by the short exact sequence

$$0 \rightarrow \mathcal{O}_{C_Y}^{\oplus r_Y} \rightarrow \oplus_\tau \mathcal{L}_\tau \rightarrow \mathcal{F}_Y \rightarrow 0$$

where $r_Y = \text{rk Pic } Y$ (implicitly we have chosen a basis for this \mathbb{Z} -module). Similarly \mathcal{F}_X is defined on C_X by:

$$0 \rightarrow \mathcal{O}_{C_X}^{\oplus r_X} \rightarrow \oplus_\rho \mathcal{L}_\rho \rightarrow \mathcal{F}_X \rightarrow 0$$

We will construct our morphism by first constructing a morphism:

$$\oplus_\tau \mathcal{L}_\tau \rightarrow \alpha^*(\oplus_\rho \mathcal{L}_\rho)$$

Recall that $f: Y \rightarrow X$ is given by homogeneous polynomials

$$P_\rho \in S_{\beta_\rho}^Y \subset S^Y = k[z_\tau : \tau \in \Sigma_Y(1)]$$

in the Cox ring of Y , where $\beta_\rho = f^*[D_\rho] \in \text{Pic } Y$. For all monomials appearing in P_ρ , if we look at their exponents $(a_\tau)_{\tau \in \Sigma_Y(1)}$, we have $\sum_{\tau \in \Sigma_Y(1)} a_\tau [D_\tau] = \beta_\rho$ by homogeneity, hence we can use the isomorphisms

parametrised by M_Y as in the proof of Proposition B.3 above in order to interpret the (P_ρ) as a morphism

$$(P_\rho)_{\rho \in \Sigma_X(1)}: \bigoplus_{\tau} \mathcal{L}_\tau \rightarrow \bigoplus_{\rho} \bigotimes_{\tau} \mathcal{L}_\tau^{\otimes a_\tau^\rho} = \bigoplus_{\rho} \tilde{\mathcal{L}}_\rho = \alpha^* \left(\bigoplus_{\rho} \mathcal{L}_\rho \right)$$

where the notation is as in §B.1. Thus we have constructed a morphism $\bigoplus_{\tau} \mathcal{L}_\tau \rightarrow \alpha^*(\bigoplus_{\rho} \mathcal{L}_\rho)$.

On the other hand, $f: Y \rightarrow X$ induces a pullback map on line bundles $\text{Pic}(X) \rightarrow \text{Pic}(Y)$. Since we have implicitly chosen bases for these \mathbb{Z} -modules, this gives rise to a matrix, whose transpose we denote by:

$$Q \in \text{Mat}_{r_X \times r_Y}(\mathbb{Z})$$

It is now clear by the functoriality construction that the square in the following diagram is commutative; hence it induces the (dashed) map of sheaves that we were hoping for

$$(9) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_{C_Y}^{\oplus r_Y} & \longrightarrow & \bigoplus_{\tau} \mathcal{L}_\tau & \longrightarrow & \mathcal{F}_Y \longrightarrow 0 \\ & & \downarrow Q & & \downarrow (P_\rho) & & \downarrow \text{---} \\ & & \mathcal{O}_{C_Y}^{\oplus r_X} & \longrightarrow & \alpha^* \left(\bigoplus_{\rho} \mathcal{L}_\rho \right) & \longrightarrow & \alpha^* \mathcal{F}_X \longrightarrow 0 \end{array}$$

Applying $R^\bullet \pi_*$ and dualising we obtain a morphism between the obstruction theories for the quasimap spaces, and we can complete this to obtain an exact triangle

$$k^* \mathbf{E}_{\overline{Q}(X)/\mathfrak{M}} \rightarrow \mathbf{E}_{\overline{Q}(Y)/\mathfrak{M}} \rightarrow \mathbf{E}_k \xrightarrow{[1]}$$

on $\overline{Q}(Y)$. The axioms of a triangulated category then give a morphism of exact triangles:

$$\begin{array}{ccccc} k^* \mathbf{E}_{\overline{Q}(X)/\mathfrak{M}} & \longrightarrow & \mathbf{E}_{\overline{Q}(Y)/\mathfrak{M}} & \longrightarrow & \mathbf{E}_k \xrightarrow{[1]} \\ \downarrow & & \downarrow & & \downarrow \\ k^* \mathbf{L}_{\overline{Q}(X)/\mathfrak{M}} & \longrightarrow & \mathbf{L}_{\overline{Q}(Y)/\mathfrak{M}} & \longrightarrow & \mathbf{L}_k \xrightarrow{[1]} \end{array}$$

It follows from a simple diagram chase that $\mathbf{E}_k \rightarrow \mathbf{L}_k$ is a relative obstruction theory. On the other hand, assuming that $\overline{Q}_{g,n}(X, f_*\beta)$ is unobstructed, we may look at the long exact sequence in cohomology and find

$$0 \rightarrow h^{-2}(\mathbf{E}_k) \rightarrow h^{-1}(k^* \mathbf{E}_{\overline{Q}(X)/\mathfrak{M}}) = 0$$

Hence $h^{-2}(\mathbf{E}_k) = 0$ and it is easy to show using similar arguments that \mathbf{E}_k is of perfect amplitude contained in $[-1, 0]$. \square

Remark B.6. The short exact sequence defining \mathcal{F}_X should be thought of as the pull-back of the Euler sequence

$$0 \rightarrow \mathcal{O}_X^{\oplus r_X} \rightarrow \bigoplus_{\rho \in \Sigma_X(1)} \mathcal{O}_X(D_\rho) \rightarrow T_X \rightarrow 0$$

along the map $C \rightarrow X$ (if such a map existed). In particular, if we work away from the locus of basepoints then $\mathcal{F}_X = u^* T_X$.

In particular, for every smooth projective variety $i: X \hookrightarrow \mathbb{P}^N$, we have a virtual pull-back morphism

$$k_v^! : A_*(\overline{\mathcal{Q}}_{0,n}(\mathbb{P}^N, d)) \rightarrow A_*(\overline{\mathcal{Q}}_{0,n}(X, \beta))$$

where $d = i_*\beta$, and more generally for any cartesian diagram

$$\begin{array}{ccc} F & \xrightarrow{\quad} & G \\ \downarrow & \square & \downarrow \\ \overline{\mathcal{Q}}_{0,n}(X, \beta) & \xrightarrow{k} & \overline{\mathcal{Q}}_{0,n}(\mathbb{P}^N, d) \end{array}$$

we get an associated virtual pull-back morphism:

$$k_v^! : A_*(G) \rightarrow A_*(F)$$

This is used in §4 to pull-back the recursion formula for the pair (\mathbb{P}^N, H) and obtain a recursion formula in the general case.

B.3. Splitting axiom. In this section we consider certain boundary strata of the moduli space of quasimaps, called *centipede loci*. These are the analogues in the absolute setting of the comb loci which appear in the relative setting (§3.2). The general element of such a locus has a source curve with $r + 1$ irreducible components, one “trunk” of the centipede and r “legs.” Each of these components has a prescribed genus, curve class and set of marked points.

Given such a locus, there are two natural virtual classes with which it can be equipped. One is the product virtual class induced by the absolute product of the $r + 1$ quasimap spaces, and the other is the class pulled back from the ambient moduli space. In this section we show that these classes coincide. This is the quasimap version of the *splitting axiom* from Gromov–Witten theory, called the *cutting edges axiom* in [Beh97].

We first establish notation. Fix a smooth projective toric variety X and numerical invariants g, n, β such that the corresponding quasimap space is defined. Now fix partitions $G = (g_0, \dots, g_r)$ of the genus, $A = (A_0, \dots, A_r)$ of the marked points and $B = (\beta_0, \dots, \beta_r)$ of the curve class and consider the following space (which we call the *centipede locus*):

$$\mathcal{D}^Q(X, G, A, B) := \overline{\mathcal{Q}}_{g_0, A_0 \cup \{q_1, \dots, q_r\}}(X, \beta_0) \times_{X^r} \prod_{i=1}^r \overline{\mathcal{Q}}_{g_i, A_i \cup \{q_i\}}(X, \beta_i)$$

Of course we assume that every element of the partition is in the stable range, so that every factor in the above product makes sense. See Remark 3.9 for a justification of why these are the correct boundary strata to consider. We can equip the centipede locus with the product virtual class in the following way. Set

$$\mathcal{E}^Q(X, G, A, B) := \overline{\mathcal{Q}}_{g_0, A_0 \cup \{q_1, \dots, q_r\}}(X, \beta_0) \times \prod_{i=1}^r \overline{\mathcal{Q}}_{g_i, A_i \cup \{q_i\}}(X, \beta_i)$$

which we endow with the product class:

$$[\mathcal{E}^Q(X, G, A, B)]^{\text{virt}} := [\overline{\mathcal{Q}}_{g_0, A_0 \cup \{q_1, \dots, q_r\}}(X, \beta_0)]^{\text{virt}} \times \prod_{i=1}^r [\overline{\mathcal{Q}}_{g_i, A_i \cup \{q_i\}}(X, \beta_i)]^{\text{virt}}$$

We then consider the cartesian diagram:

$$(10) \quad \begin{array}{ccc} \mathcal{D}^Q(X, G, A, B) & \xrightarrow{h} & \mathcal{E}^Q(X, G, A, B) \\ \downarrow \text{ev}_q & \square & \downarrow \text{ev}_q \\ X^r & \xrightarrow{\Delta_{X^r}} & X^r \times X^r \end{array}$$

Since X is smooth Δ_{X^r} is a regular embedding, so we have a Gysin map which we use to define:

$$[\mathcal{D}^Q(X, G, A, B)]^{\text{virt}} := \Delta_{X^r}^! [\mathcal{E}^Q(X, G, A, B)]^{\text{virt}}$$

Notice that if we set

$$\mathfrak{M}_{G, A, B}^{\text{wt}} := \mathfrak{M}_{g_0, A_0 \cup \{q_1, \dots, q_r\}, \beta_0}^{\text{wt}} \times \prod_{i=1}^r \mathfrak{M}_{g_i, A_i \cup \{q_i\}, \beta_i}^{\text{wt}}$$

then there is a morphism given by forgetting everything except the source curves and their classes

$$\rho_E : \mathcal{E}^Q(X, G, A, B) \rightarrow \mathfrak{M}_{G, A, B}^{\text{wt}}$$

and the virtual class on $\mathcal{E}^Q(X, G, A, B)$ is induced by a perfect obstruction theory $\mathbf{E}_{\rho_E} \rightarrow \mathbf{L}_{\rho_E}$ given by the product of the standard obstruction theories for each factor:

$$\overline{\mathcal{Q}}_{g_i, A_i \cup \{q_i\}}(X, \beta_i) \rightarrow \mathfrak{M}_{g_i, A_i, \beta_i}^{\text{wt}}$$

On the other hand, we have the following cartesian diagram

$$(11) \quad \begin{array}{ccc} \mathcal{D}^Q(X, G, A, B) & \xrightarrow{\varphi} & \overline{\mathcal{Q}}_{0, n}(X, \beta) \\ \downarrow \rho_D & \square & \downarrow \rho_Q \\ \mathfrak{M}_{G, A, B}^{\text{wt}} & \xrightarrow{\psi} & \mathfrak{M}_{g, n, \beta}^{\text{wt}} \end{array}$$

The bottom horizontal map is not a closed immersion: due to the existence of degree-0 rational components, there may be many possible equally valid

ways of breaking up a nodal curve. For instance, consider the following example of two elements which map to the same curve under ψ . [FIGURE]

Nevertheless ψ has a natural perfect obstruction theory, given by \mathbf{L}_ψ : we only need to show that it is supported in $[-1, 0]$. Consider the exact triangle:

$$\psi^* \mathbf{L}_{\mathfrak{M}_{g,n,\beta}^{\text{wt}}} \rightarrow \mathbf{L}_{\mathfrak{M}_{G,A,B}^{\text{wt}}} \rightarrow \mathbf{L}_\psi \xrightarrow{[1]}$$

The first two terms are concentrated in degrees $[0, 1]$, because they are the cotangent complexes of smooth Artin stacks. Therefore \mathbf{L}_ψ is concentrated in degrees $[-1, 1]$. Furthermore, if we examine the long exact cohomology sequence near $h^1(\mathbf{L}_\psi)$ we find

$$h^1(\psi^* \mathbf{L}_{\mathfrak{M}_{g,n,\beta}^{\text{wt}}}) \rightarrow h^1(\mathbf{L}_{\mathfrak{M}_{G,A,B}^{\text{wt}}}) \rightarrow h^1(\mathbf{L}_\psi) \rightarrow 0$$

and hence we must show that the first map is surjective. But this is dual to the map which takes an infinitesimal automorphism of the disconnected curve to an infinitesimal automorphism of the corresponding connected curve (obtained by glueing together the “nodal” marked points). The space of infinitesimal automorphisms of a nodal curve splits into a direct sum of infinitesimal automorphisms of each component; since the glueing does not affect the components, we see that this map is an isomorphism. Hence $h^1(\mathbf{L}_\psi) = 0$ as claimed; morally this follows from the fact that the fibres of ψ are Deligne–Mumford.

Hence there is a virtual pull-back map $\psi^!$ which defines a class

$$\psi^! [\overline{\mathcal{Q}}_{g,n}(X, \beta)]^{\text{virt}}$$

on $\mathcal{D}^Q(X, G, A, B)$. By functoriality of virtual pull-backs, this is induced by the perfect obstruction theory:

$$\varphi^* \mathbf{E}_{\rho_Q} \rightarrow \mathbf{L}_{\rho_D}$$

Finally if we look at (10) we see that $\text{ev}_q^* \mathbf{L}_{\Delta_{X^r}} \rightarrow \mathbf{L}_h$ is a perfect obstruction theory for the map h . To summarise, we have a triangle

$$(12) \quad \begin{array}{ccc} \mathcal{D}^Q(X, G, A, B) & \xrightarrow{h} & \mathcal{E}^Q(X, G, A, B) \\ & \searrow \rho_D & \swarrow \rho_E \\ & \mathfrak{M}_{G,A,B}^{\text{wt}} & \end{array}$$

where all three morphisms are equipped with perfect obstruction theories. We simply need to check that these fit together in a compatible triple

Lemma B.7. *There is a compatible triple*

$$(h^* \mathbf{E}_{\rho_E}, \varphi^* \mathbf{E}_{\rho_Q}, \text{ev}_q^* \mathbf{L}_{\Delta_{X^r}})$$

for the triangle (12). Hence by functoriality of virtual pull-backs we have:

$$\psi^! [\overline{\mathcal{Q}}_{g,n}(X, \beta)]^{\text{virt}} = \Delta_{X^r}^! [\mathcal{E}^Q(X, G, A, B)]^{\text{virt}} = [\mathcal{D}^Q(X, G, A, B)]^{\text{virt}}$$

Proof. We need to construct a morphism of triangles

$$\begin{array}{ccccccc}
h^*E_{\rho_E} & \longrightarrow & \varphi^*E_{\rho_Q} & \longrightarrow & \mathrm{ev}_q^*L_{\Delta_{X^r}} & \xrightarrow{[1]} & \longrightarrow \\
\downarrow & & \downarrow & & \downarrow & & \\
h^*L_{\rho_E} & \longrightarrow & L_{\rho_D} & \longrightarrow & L_h & \xrightarrow{[1]} & \longrightarrow
\end{array}$$

Consider the following diagram:

$$\begin{array}{ccccc}
h^*\tilde{C} & \xrightarrow{\nu} & \varphi^*C & \longrightarrow & C \\
\searrow \eta & & \downarrow & \square & \downarrow \pi \\
& & \mathcal{D}^Q(X, G, A, B) & \xrightarrow{\varphi} & \overline{\mathcal{Q}}_{g,n}(X, \beta)
\end{array}$$

Here \tilde{C} is the universal (disconnected) curve over $\mathcal{E}^Q(X, G, A, B)$, which we have pulled back to $\mathcal{D}^Q(X, G, A, B)$, while φ^*C is the universal curve over $\mathcal{D}^Q(X, G, A, B)$. Therefore the map $\nu : h^*\tilde{C} \rightarrow \varphi^*C$ is (fiberwise) a partial normalisation map given by normalising the nodes which connect the “trunk” of the centipede to the “legs.”

There are natural sheaves \mathcal{F} and $\tilde{\mathcal{F}}$ on C and $h^*\tilde{C}$ respectively, such that

$$\begin{aligned}
\varphi^*E_{\rho_Q}^\vee &= R^\bullet \pi_* \mathcal{F} \\
h^*E_{\rho_E}^\vee &= R^\bullet \eta_* \tilde{\mathcal{F}}
\end{aligned}$$

Furthermore $\nu^*\mathcal{F} \simeq \tilde{\mathcal{F}}$, hence by tensoring the partial normalisation short exact sequence

$$0 \rightarrow \mathcal{O}_{\varphi^*C} \rightarrow \nu_* \mathcal{O}_{h^*\tilde{C}} \rightarrow \mathcal{O}_q \rightarrow 0$$

with \mathcal{F} and applying the projection formula, we obtain

$$0 \rightarrow \mathcal{F} \rightarrow \nu_* \tilde{\mathcal{F}} \rightarrow \mathcal{F}_q \rightarrow 0$$

on φ^*C , where q is the locus of nodes connecting the trunk to the spine. (The fact that the morphism on the left is injective follows by applying the Snake Lemma to the short exact sequence defining \mathcal{F} .) To this we can apply $R^\bullet \pi_*$ to obtain an exact triangle

$$(13) \quad R^\bullet \pi_* \mathcal{F} \rightarrow R^\bullet \eta_* \tilde{\mathcal{F}} \rightarrow R^\bullet \pi_* \mathcal{F}_q \xrightarrow{[1]}$$

Finally, notice that, since quasimaps are required not to have base-points at the nodes, the fibre of the sheaf \mathcal{F} at each of the nodes q can actually be identified with the tangent to the toric variety X at the image of the node itself, i.e. $R^\bullet \pi_* \mathcal{F}_q \cong \mathrm{ev}_q^* T_{X^r} = \mathrm{ev}_q^* T_{\Delta_{X^r}}[-1]$. Dualising sequence (13) we obtain

$$h^*E_{\rho_E} \rightarrow \varphi^*E_{\rho_Q} \rightarrow \mathrm{ev}_q^* E_{\Delta_{X^r}} \xrightarrow{[1]}$$

as required. \square

B.4. Comparison with the GIT construction. Let X be a smooth projective toric variety and $Y \hookrightarrow X$ a smooth very ample hypersurface. The complete linear system $|\mathcal{O}_X(Y)|$ gives an embedding $i : X \hookrightarrow \mathbb{P}^N$ which expresses Y as the intersection inside \mathbb{P}^N of X and a certain hyperplane H : $Y = i^{-1}(H)$. We can *define* the moduli space of quasimaps to Y via the following cartesian diagram:

$$\begin{array}{ccc} \overline{\mathcal{Q}}_{g,n}(Y, \beta) & \hookrightarrow & \overline{\mathcal{Q}}_{g,n}(H, d) \\ \downarrow & \square & \downarrow \\ \overline{\mathcal{Q}}_{g,n}(X, \beta) & \xrightarrow{k} & \overline{\mathcal{Q}}_{g,n}(\mathbb{P}^N, d) \end{array}$$

where $d = i_*\beta$. This moduli space is easy to describe: let s_Y denote the section of $\mathcal{O}_X(Y)$ cutting out Y inside X . Recall from §2.3 that for any quasimap

$$((C, x_1, \dots, x_n), (L_\rho, u_\rho)_{\rho \in \Sigma_X(1)}, (\varphi_m)_{m \in M_X}) \in \overline{\mathcal{Q}}_{g,n}(X, \beta)$$

we can construct a section u_Y of a line bundle L_Y on C , which plays the role of the pull-back of s_Y to C . Then

$$\overline{\mathcal{Q}}_{g,n}(Y, \beta) \subseteq \overline{\mathcal{Q}}_{g,n}(X, \beta)$$

consists of those quasimaps such that $u_Y \equiv 0$.

The cartesian diagram above can also be used to endow $\overline{\mathcal{Q}}_{g,n}(Y, \beta)$ with a virtual class via virtual (or diagonal) pull-back along k . Thus we can define quasimap invariants for Y .

On the other hand, Y has the natural structure of a GIT quotient

$$Y = C(Y) // G$$

where $C(Y) \subseteq \mathbb{A}^{\Sigma_X(1)}$ is the affine cone over Y and $G = \text{Hom}_{\mathbb{Z}}(\text{Pic}(X), \mathbb{G}_m) \cong \mathbb{G}_m^{r_X}$ acts on $C(Y)$ via the natural inclusion

$$\mathbb{G}_m^{r_X} \hookrightarrow \mathbb{G}_m^{\Sigma_X(1)}$$

(here $C(Y) \subseteq \mathbb{A}^{\Sigma_X(1)}$ is preserved by G because it is cut out by a homogeneous equation). In [CFKM14] moduli spaces of quasimaps are constructed for GIT quotient targets (satisfying a number of conditions, all of which hold for Y). There is thus a moduli space

$$\overline{\mathcal{Q}}_{g,n}^{\text{GIT}}(Y, \beta)$$

which admits a virtual class. Hence we have two moduli spaces of quasimaps to Y , each equipped with a virtual class, and we want to check that these definitions agree.

Objects of $\overline{\mathcal{Q}}_{g,n}^{\text{GIT}}(Y, \beta)$ are diagrams of the form

$$\begin{array}{ccc} P & \longrightarrow & C(Y) \\ \downarrow & & \\ C & & \end{array}$$

where C is a prestable curve, P is a principal G -bundle and the map $P \rightarrow C(Y)$ is G -equivariant. Equivalently, an object consists of a prestable curve C , a principal G -bundle P and a section u of the associated $C(Y)$ -bundle:

$$\begin{array}{ccc} P \times_G C(Y) & & \\ p \downarrow \uparrow u & & \\ C & & \end{array}$$

The obstruction theory on this space is defined relative to the stack \mathfrak{Bun}_G parametrising principal G -bundles on the universal curve:

$$C\mathfrak{M}_{g,n} \rightarrow \mathfrak{M}_{g,n}$$

It is given by

$$\mathbf{E}_{\overline{Q}/\mathfrak{Bun}_G}^\vee = \mathbf{R}^\bullet \pi_*(u^* \mathbf{L}_p)$$

where π is the universal curve over $\overline{Q} = \overline{Q}_{g,n}^{\text{GIT}}(Y, \beta)$. There is a natural isomorphism

$$\mathfrak{Bun}_G \cong \times_{\mathfrak{M}_{g,n}}^{r_X} \mathfrak{Pic}_{g,n}$$

given by sending P to the r_X individual factors of the affine bundle $P \times_G \mathbb{A}^{r_X}$. Furthermore there is a G -equivariant embedding

$$\begin{array}{ccc} P \times_G C(Y) & \xhookrightarrow{j} & P \times_G \mathbb{A}^{\Sigma_X(1)} \cong \bigoplus_{\rho \in \Sigma_X(1)} L_\rho \\ p \downarrow \uparrow u & & \swarrow \\ C & & \end{array}$$

which expresses $P \times_G C(Y)$ as the vanishing locus of u_Y in $\bigoplus_{\rho \in \Sigma_X(1)} L_\rho$. This shows that the two definitions of the moduli space agree.

Finally we must compare the virtual classes. Using the normal sheaf sequence for the inclusion j (relative to the base C) we obtain a short exact sequence on C :

$$0 \rightarrow u^* T_p \rightarrow \bigoplus_{\rho \in \Sigma_X(1)} L_\rho \rightarrow u^* \mathbf{N}_{P \times_G C(Y) / \bigoplus_{\rho \in \Sigma_X(1)} L_\rho} \rightarrow 0$$

Since $P \times_G C(Y)$ is defined by the vanishing of u_Y , we see that the final term is isomorphic to the line bundle L_Y discussed above. Thus as elements of the derived category

$$u^* T_p = \left[\bigoplus_{\rho \in \Sigma_X(1)} L_\rho \rightarrow L_Y \right]$$

Applying $\mathbf{R}^\bullet \pi_*$ we obtain on the left hand side the obstruction theory for the GIT moduli space relative \mathfrak{Bun}_G . On the other hand, the first term on

the right hand side is the obstruction theory for $\overline{\mathcal{Q}}(X)$ relative the product of the Picard stacks (isomorphic to \mathfrak{Bun}_G via the discussion above) whereas the second term is the relative obstruction theory for $\overline{\mathcal{Q}}(Y)$ inside $\overline{\mathcal{Q}}(X)$. Thus the virtual classes agree, as claimed.

APPENDIX C. SOME INTERSECTION-THEORETIC LEMMAS

In this appendix we explicitly define the *diagonal pull-back* along a morphism whose target is unobstructed (used in [Gat02]) and verify that this agrees with the virtual pull-back of [Man12a] when both are defined. We also check that it satisfies some expected compatibility properties.

Consider a morphism of DM stacks $f: Y \rightarrow X$ over a smooth base \mathfrak{M} , such that X is smooth over \mathfrak{M} and Y carries a virtual class given by a perfect obstruction theory $\mathbf{E}_{Y/\mathfrak{M}}$. Then, for every Cartesian diagram

$$\begin{array}{ccc} G & \xrightarrow{g} & F \\ \downarrow q & \square & \downarrow p \\ Y & \xrightarrow{f} & X \end{array}$$

and every class $\alpha \in A_*(F)$, we may define

$$f_{\Delta}^!(\alpha) = \Delta_X^!([Y]^{\text{vir}} \times \alpha) \in A_*(G)$$

which we call the *diagonal (virtual) pull-back*. We first show that it coincides with the usual virtual pull-back along f in the presence of a compatible perfect obstruction theory for f .

Lemma C.1. *Assume that there exists a relative obstruction theory \mathbf{E}_f compatible with $\mathbf{E}_{Y/\mathfrak{M}}$ and the standard (unobstructed) obstruction theory for X , i.e:*

$$\begin{array}{ccccc} f^* \mathbf{L}_{X/\mathfrak{M}} & \longrightarrow & \mathbf{E}_{Y/\mathfrak{M}} & \longrightarrow & \mathbf{E}_f \xrightarrow{[1]} \\ \downarrow \text{Id} & & \downarrow & & \downarrow \\ f^* \mathbf{L}_{X/\mathfrak{M}} & \longrightarrow & \mathbf{L}_{Y/\mathfrak{M}} & \longrightarrow & \mathbf{L}_f \xrightarrow{[1]} \end{array}$$

Then for every Cartesian diagram and every class $\alpha \in A_*(F)$ as above,

$$f_v^!(\alpha) = f_{\Delta}^!(\alpha).$$

Proof. Consider the following cartesian diagram:

$$\begin{array}{ccccccc} G & \xrightarrow{q \times g} & Y \times_{\mathfrak{M}} F & \xrightarrow{\text{pr}_1} & Y & & \\ \downarrow g & \square & \downarrow f \times \text{Id} & \square & \downarrow f & & \\ F & \xrightarrow{p \times \text{Id}} & X \times_{\mathfrak{M}} F & \xrightarrow{\text{pr}_1} & X & & \\ \downarrow p & \square & \downarrow \text{Id} \times p & & & & \\ X & \xrightarrow{\Delta_X} & X \times_{\mathfrak{M}} X & & & & \end{array}$$

Then, by commutativity of virtual pull-backs, we have

$$\begin{aligned}
 \Delta_X^!([Y]^{\text{vir}} \times \alpha) &= \Delta^!((f_v^![X]) \times \alpha) \\
 &= \Delta_X^!(f_v^!([X] \times \alpha)) \\
 &= f_v^!(\Delta_X^!([X] \times \alpha)) \\
 &= f_v^!(\alpha)
 \end{aligned}$$

as required. \square

Secondly, we show that the *diagonal* virtual pull-back behaves similarly to an ordinary virtual pull-back (e.g. commutes with other virtual pull-backs) even in the absence of a compatible perfect obstruction theory.

Lemma C.2. *The diagonal virtual pull-back morphism as defined above commutes with ordinary Gysin maps and with virtual pull-backs.*

Proof. First consider the case of ordinary Gysin maps. We must consider a cartesian diagram:

$$\begin{array}{ccccc}
 Y'' & \longrightarrow & X'' & \longrightarrow & S \\
 \downarrow & \square & \downarrow & \square & \downarrow k \\
 Y' & \longrightarrow & X' & \longrightarrow & T \\
 \downarrow & \square & \downarrow & & \\
 Y & \xrightarrow{f} & X & &
 \end{array}$$

with k a regular embedding and $f: Y \rightarrow X$ as before. We need to show that for all $\alpha \in A_*(X')$:

$$k^! f_\Delta^!(\alpha) = f_\Delta^! k^!(\alpha)$$

We form the cartesian diagram:

$$\begin{array}{ccccc}
 Y'' & \longrightarrow & Y \times X'' & \longrightarrow & S \\
 \downarrow & \square & \downarrow & \square & \downarrow k \\
 Y' & \longrightarrow & Y \times X' & \longrightarrow & T \\
 \downarrow & \square & \downarrow & & \\
 X & \xrightarrow{\Delta_X} & X \times X & &
 \end{array}$$

And apply commutativity of usual Gysin morphisms. In the case where k is not a regular embedding but rather is equipped with a relative perfect obstruction theory, the same argument works with $k^!$ replaced by $k_v^!$. \square

REFERENCES

- [AC14] Dan Abramovich and Qile Chen. Stable logarithmic maps to Deligne-Faltings pairs II. *Asian J. Math.*, 18(3):465–488, 2014.
- [AF16] Dan Abramovich and Barbara Fantechi. Orbifold techniques in degeneration formulas. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 16(2):519–579, 2016.

- [AMW14] Dan Abramovich, Steffen Marcus, and Jonathan Wise. Comparison theorems for Gromov-Witten invariants of smooth pairs and of degenerations. *Ann. Inst. Fourier (Grenoble)*, 64(4):1611–1667, 2014.
- [Beh97] K. Behrend. Gromov-Witten invariants in algebraic geometry. *Invent. Math.*, 127(3):601–617, 1997.
- [Ber00] Aaron Bertram. Another way to enumerate rational curves with torus actions. *Invent. Math.*, 142(3):487–512, 2000.
- [CFK10] Ionuț Ciocan-Fontanine and Bumsig Kim. Moduli stacks of stable toric quasimaps. *Adv. Math.*, 225(6):3022–3051, 2010.
- [CFK14] Ionuț Ciocan-Fontanine and Bumsig Kim. Wall-crossing in genus zero quasimap theory and mirror maps. *Algebr. Geom.*, 1(4):400–448, 2014.
- [CFKM14] Ionuț Ciocan-Fontanine, Bumsig Kim, and Daves Maulik. Stable quasimaps to GIT quotients. *J. Geom. Phys.*, 75:17–47, 2014.
- [Che14] Qile Chen. Stable logarithmic maps to Deligne-Faltings pairs I. *Ann. of Math. (2)*, 180(2):455–521, 2014.
- [CM] Tom Coates and Cristina Manolache. In preparation.
- [Cos06] Kevin Costello. Higher genus Gromov-Witten invariants as genus zero invariants of symmetric products. *Ann. of Math. (2)*, 164(2):561–601, 2006.
- [Cox95a] David A. Cox. The functor of a smooth toric variety. *Tohoku Math. J. (2)*, 47(2):251–262, 1995.
- [Cox95b] David A. Cox. The homogeneous coordinate ring of a toric variety. *J. Algebraic Geom.*, 4(1):17–50, 1995.
- [CZ14] Yaim Cooper and Aleksey Zinger. Mirror symmetry for stable quotients invariants. *Michigan Math. J.*, 63(3):571–621, 2014.
- [Ful98] William Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 1998.
- [Gat02] Andreas Gathmann. Absolute and relative Gromov-Witten invariants of very ample hypersurfaces. *Duke Math. J.*, 115(2):171–203, 2002.
- [Gat03] Andreas Gathmann. Relative Gromov-Witten invariants and the mirror formula. *Math. Ann.*, 325(2):393–412, 2003.
- [Giv96a] Alexander B. Givental. Equivariant Gromov-Witten invariants. *Internat. Math. Res. Notices*, (13):613–663, 1996.
- [Giv96b] Alexander B. Givental. Equivariant Gromov-Witten invariants. *Internat. Math. Res. Notices*, (13):613–663, 1996.
- [Giv98] Alexander Givental. *A Mirror Theorem for Toric Complete Intersections*, pages 141–175. Birkhäuser Boston, Boston, MA, 1998.
- [GS13] Mark Gross and Bernd Siebert. Logarithmic Gromov-Witten invariants. *J. Amer. Math. Soc.*, 26(2):451–510, 2013.
- [GS16] M. Gross and B. Siebert. Intrinsic mirror symmetry and punctured Gromov-Witten invariants. *ArXiv e-prints*, September 2016.
- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.
- [Kim10] Bumsig Kim. Logarithmic stable maps. In *New developments in algebraic geometry, integrable systems and mirror symmetry (RIMS, Kyoto, 2008)*, volume 59 of *Adv. Stud. Pure Math.*, pages 167–200. Math. Soc. Japan, Tokyo, 2010.
- [Knu83] Finn F. Knudsen. The projectivity of the moduli space of stable curves. II. The stacks $M_{g,n}$. *Math. Scand.*, 52(2):161–199, 1983.
- [KP01] B. Kim and R. Pandharipande. The connectedness of the moduli space of maps to homogeneous spaces. In *Symplectic geometry and mirror symmetry (Seoul, 2000)*, pages 187–201. World Sci. Publ., River Edge, NJ, 2001.

- [Li01] Jun Li. Stable morphisms to singular schemes and relative stable morphisms. *J. Differential Geom.*, 57(3):509–578, 2001.
- [Li02] Jun Li. A degeneration formula of GW-invariants. *J. Differential Geom.*, 60(2):199–293, 2002.
- [LLY97] Bong H. Lian, Kefeng Liu, and Shing-Tung Yau. Mirror principle. I. *Asian J. Math.*, 1(4):729–763, 1997.
- [Man12a] Cristina Manolache. Virtual pull-backs. *J. Algebraic Geom.*, 21(2):201–245, 2012.
- [Man12b] Cristina Manolache. Virtual push-forwards. *Geom. Topol.*, 16(4):2003–2036, 2012.
- [MOP11] Alina Marian, Dragos Oprea, and Rahul Pandharipande. The moduli space of stable quotients. *Geom. Topol.*, 15(3):1651–1706, 2011.
- [PR03] Mihnea Popa and Mike Roth. Stable maps and Quot schemes. *Invent. Math.*, 152(3):625–663, 2003.