

Dualities for abelian ℓ -groups and vector lattices beyond archimedeanity

Luca Carai, University of Salerno

joint work with S. Lapenta and L. Spada

Ordered Algebras and Logic, Les Diablerets, Switzerland

April 2, 2022

Abelian ℓ -groups and vector lattices

Definition

- An **abelian ℓ -group** is an abelian group A equipped with a lattice order such that $a \leq b$ implies $a + c \leq b + c$ for every $a, b, c \in A$.

Definition

- An **abelian ℓ -group** is an abelian group A equipped with a lattice order such that $a \leq b$ implies $a + c \leq b + c$ for every $a, b, c \in A$.
- A **vector lattice** is an abelian ℓ -group V equipped with a structure of \mathbb{R} -vector space such that $0 \leq r$ and $0 \leq v$ imply $rv \geq 0$ for each $r \in \mathbb{R}$ and $v \in V$.

Abelian ℓ -groups and vector lattices

Definition

- An **abelian ℓ -group** is an abelian group A equipped with a lattice order such that $a \leq b$ implies $a + c \leq b + c$ for every $a, b, c \in A$.
- A **vector lattice** is an abelian ℓ -group V equipped with a structure of \mathbb{R} -vector space such that $0 \leq r$ and $0 \leq v$ imply $rv \geq 0$ for each $r \in \mathbb{R}$ and $v \in V$.

Abelian ℓ -groups and vector lattices form **varieties**.

Congruences in abelian ℓ -groups and vector lattices correspond to ℓ -ideals.

Definition

- An ℓ -ideal in an abelian ℓ -group is a subgroup I that is convex, i.e. $|a| \leq |b|$ and $b \in I$ imply $a \in I$.
- An ℓ -ideal in a vector lattice is a vector subspace that is convex.

Congruences in abelian ℓ -groups and vector lattices correspond to ℓ -ideals.

Definition

- An ℓ -ideal in an abelian ℓ -group is a subgroup I that is convex, i.e. $|a| \leq |b|$ and $b \in I$ imply $a \in I$.
- An ℓ -ideal in a vector lattice is a vector subspace that is convex.

Definition

- A proper ℓ -ideal is called ℓ -maximal if it is maximal wrt inclusion.
- A nontrivial abelian ℓ -group/vector lattice A is ℓ -simple if $\{0\}$ and A are the only ℓ -ideals of A .

Definition

An abelian ℓ -group/vector lattice is **semisimple** if the intersection of all its maximal ℓ -ideals is $\{0\}$.

It is **archimedean** if $na \leq b$ for every $n \in \mathbb{N}$ implies $a \leq 0$.

Definition

An abelian ℓ -group/vector lattice is **semisimple** if the intersection of all its maximal ℓ -ideals is $\{0\}$.

It is **archimedean** if $na \leq b$ for every $n \in \mathbb{N}$ implies $a \leq 0$.

Semisimple \Rightarrow archimedean

Definition

An abelian ℓ -group/vector lattice is **semisimple** if the intersection of all its maximal ℓ -ideals is $\{0\}$.

It is **archimedean** if $na \leq b$ for every $n \in \mathbb{N}$ implies $a \leq 0$.

Semisimple \Rightarrow archimedean

Archimedean \Rightarrow semisimple (if finitely generated)

Definition

An abelian ℓ -group/vector lattice is **semisimple** if the intersection of all its maximal ℓ -ideals is $\{0\}$.

It is **archimedean** if $na \leq b$ for every $n \in \mathbb{N}$ implies $a \leq 0$.

Semisimple \Rightarrow archimedean

Archimedean \Rightarrow semisimple (if finitely generated)

- A/I is simple iff I is maximal.
- A/I is semisimple iff I is intersection of maximal ℓ -ideals.

Baker-Beynon duality

Piecewise linear functions



Definition

A continuous function $f : \mathbb{R}^\kappa \rightarrow \mathbb{R}$ is **piecewise linear** if there exist g_1, \dots, g_n linear homogeneous polynomials in the variables $(x_\alpha)_{\alpha < \kappa}$ such that for each $x \in \mathbb{R}^\kappa$ we have $f(x) = g_i(x)$ for some $i = 1, \dots, n$.

Piecewise linear functions

Definition



A continuous function $f : \mathbb{R}^k \rightarrow \mathbb{R}$ is **piecewise linear** if there exist g_1, \dots, g_n linear homogeneous polynomials in the variables $(x_\alpha)_{\alpha < k}$ such that for each $x \in \mathbb{R}^k$ we have $f(x) = g_i(x)$ for some $i = 1, \dots, n$.



Piecewise linear functions

Definition

A continuous function $f : \mathbb{R}^k \rightarrow \mathbb{R}$ is **piecewise linear** if there exist g_1, \dots, g_n linear homogeneous polynomials in the variables $(x_\alpha)_{\alpha < k}$ such that for each $x \in \mathbb{R}^k$ we have $f(x) = g_i(x)$ for some $i = 1, \dots, n$.



Piecewise linear functions

- The set $\text{PWL}_{\mathbb{R}}(\mathbb{R}^{\kappa})$ of piecewise linear functions on \mathbb{R}^{κ} is a vector lattice with pointwise operations.

Piecewise linear functions

- The set $\text{PWL}_{\mathbb{R}}(\mathbb{R}^{\kappa})$ of piecewise linear functions on \mathbb{R}^{κ} is a vector lattice with pointwise operations.
- The set $\text{PWL}_{\mathbb{Z}}(\mathbb{R}^{\kappa})$ of piecewise linear functions on \mathbb{R}^{κ} such that g_1, \dots, g_n have integer coefficients is an abelian ℓ -group with pointwise operations.

Piecewise linear functions

- The set $\text{PWL}_{\mathbb{R}}(\mathbb{R}^{\kappa})$ of piecewise linear functions on \mathbb{R}^{κ} is a vector lattice with pointwise operations.
- The set $\text{PWL}_{\mathbb{Z}}(\mathbb{R}^{\kappa})$ of piecewise linear functions on \mathbb{R}^{κ} such that g_1, \dots, g_n have integer coefficients is an abelian ℓ -group with pointwise operations.

Theorem

- $\text{PWL}_{\mathbb{R}}(\mathbb{R}^{\kappa})$ is iso to the free vector lattice on κ generators.
- $\text{PWL}_{\mathbb{Z}}(\mathbb{R}^{\kappa})$ is iso to the free abelian ℓ -group on κ generators.

Piecewise linear functions

If $X \subseteq \mathbb{R}^k$, we denote by $\text{PWL}_{\mathbb{R}}(X)$ and $\text{PWL}_{\mathbb{Z}}(X)$ the sets of piecewise linear maps restricted to X .

Piecewise linear functions

If $X \subseteq \mathbb{R}^k$, we denote by $\text{PWL}_{\mathbb{R}}(X)$ and $\text{PWL}_{\mathbb{Z}}(X)$ the sets of piecewise linear maps restricted to X .

Definition

A subset of \mathbb{R}^k is a **cone** if it is closed under multiplication by nonnegative scalars.

Piecewise linear functions

If $X \subseteq \mathbb{R}^\kappa$, we denote by $\text{PWL}_{\mathbb{R}}(X)$ and $\text{PWL}_{\mathbb{Z}}(X)$ the sets of piecewise linear maps restricted to X .

Definition

A subset of \mathbb{R}^κ is a **cone** if it is closed under multiplication by nonnegative scalars.

Theorem (Baker 1968)

- *Every κ -generated semisimple vector lattice is isomorphic to $\text{PWL}_{\mathbb{R}}(C)$ where C is a cone that is closed in \mathbb{R}^κ .*
- *Every κ -generated semisimple abelian ℓ -group is isomorphic to $\text{PWL}_{\mathbb{Z}}(C)$ where C is a cone that is closed in \mathbb{R}^κ .*

Theorem (Beynon 1974)

- *The category of **semisimple vector lattices** is dually equivalent to the category of **closed cones** in \mathbb{R}^κ and piecewise linear maps with real coefficients.*

Theorem (Beynon 1974)

- The category of *semisimple vector lattices* is dually equivalent to the category of *closed cones* in \mathbb{R}^κ and piecewise linear maps with real coefficients.
- The category of *semisimple abelian ℓ -groups* is dually equivalent to the category of *closed cones* in \mathbb{R}^κ and piecewise linear maps with integer coefficients.

Theorem (Beynon 1974)

- The category of *finitely generated archimedean vector lattices* is dually equivalent to the category of *closed cones* in \mathbb{R}^n for $n \in \mathbb{N}$ and piecewise linear maps with real coefficients.
- The category of *finitely generated archimedean abelian ℓ -groups* is dually equivalent to the category of *closed cones* in \mathbb{R}^n for $n \in \mathbb{N}$ and piecewise linear maps with integer coefficients.

General affine duality approach

Basic Galois connection

Let V be the variety of abelian ℓ -groups or the variety of vector lattices. Let $A \in V$, κ a cardinal, and \mathcal{F}_κ be the free algebra in V over κ generators.

Basic Galois connection

Let V be the variety of abelian ℓ -groups or the variety of vector lattices. Let $A \in V$, κ a cardinal, and \mathcal{F}_κ be the free algebra in V over κ generators.

For any $T \subseteq \mathcal{F}_\kappa$ and $S \subseteq A^\kappa$, we define the following operators.

$$\mathbb{V}_A(T) = \{x \in A^\kappa \mid t(x) = 0 \text{ for all } t \in T\}$$

$$\mathbb{I}_A(S) = \{t \in \mathcal{F}_\kappa \mid t(x) = 0 \text{ for all } x \in S\}.$$

$\mathbb{I}_A(S)$ is always an ℓ -ideal.

Basic Galois connection

Let V be the variety of abelian ℓ -groups or the variety of vector lattices. Let $A \in V$, κ a cardinal, and \mathcal{F}_κ be the free algebra in V over κ generators.

For any $T \subseteq \mathcal{F}_\kappa$ and $S \subseteq A^\kappa$, we define the following operators.

$$\mathbb{V}_A(T) = \{x \in A^\kappa \mid t(x) = 0 \text{ for all } t \in T\}$$

$$\mathbb{I}_A(S) = \{t \in \mathcal{F}_\kappa \mid t(x) = 0 \text{ for all } x \in S\}.$$

$\mathbb{I}_A(S)$ is always an ℓ -ideal.

Basic Galois connection

$$T \subseteq \mathbb{I}_A(S) \quad \text{iff} \quad S \subseteq \mathbb{V}_A(T) .$$

Algebraic Nullstellensatz

(Caramello, Marra, and Spada 2012)

- Let I be an ℓ -ideal of \mathcal{F}_κ . We have $I = \mathbb{I}_A(x)$ for some $x \in A^\kappa$ iff \mathcal{F}_κ / I embeds into A .

Algebraic Nullstellensatz

(Caramello, Marra, and Spada 2012)

- Let I be an ℓ -ideal of \mathcal{F}_κ . We have $I = \mathbb{I}_A(x)$ for some $x \in A^\kappa$ iff \mathcal{F}_κ / I embeds into A .
- $\mathbb{I}_A(S) = \bigcap_{x \in S} \mathbb{I}_A(x)$.

Fixpoints of the basic Galois connection

Algebraic Nullstellensatz

(Caramello, Marra, and Spada 2012)

- Let I be an ℓ -ideal of \mathcal{F}_κ . We have $I = \mathbb{I}_A(x)$ for some $x \in A^\kappa$ iff \mathcal{F}_κ / I embeds into A .
- $\mathbb{I}_A(S) = \bigcap_{x \in S} \mathbb{I}_A(x)$.

Definition

The subsets $\mathbb{V}_A(I) = \{x \in A^\kappa \mid t(x) = 0 \text{ for all } t \in I\}$ are the closed subsets of a topology on A^κ called the **Zariski topology**.

Fixpoints of the basic Galois connection

Algebraic Nullstellensatz

(Caramello, Marra, and Spada 2012)

- Let I be an ℓ -ideal of \mathcal{F}_κ . We have $I = \mathbb{I}_A(x)$ for some $x \in A^\kappa$ iff \mathcal{F}_κ / I embeds into A .
- $\mathbb{I}_A(S) = \bigcap_{x \in S} \mathbb{I}_A(x)$.

Definition

The subsets $\mathbb{V}_A(I) = \{x \in A^\kappa \mid t(x) = 0 \text{ for all } t \in I\}$ are the closed subsets of a topology on A^κ called the **Zariski topology**.

The fixpoints of the Galois connection are:

- the intersections of ideals I of \mathcal{F}_κ such that \mathcal{F}_κ / I embeds into A ,
- the Zariski closed subsets of A^κ .

Theorem (Caramello, Marra, and Spada 2012)

The Galois connection induces a dual equivalence between

- *the category of algebras of V that are subdirect products of subalgebras of A , and*

Theorem (Caramello, Marra, and Spada 2012)

The Galois connection induces a dual equivalence between

- *the category of algebras of V that are subdirect products of subalgebras of A , and*
- *the category of Zariski closed subsets C of A^κ where κ ranges over all the cardinal numbers.*

Theorem (Caramello, Marra, and Spada 2012)

The Galois connection induces a dual equivalence between

- *the category of algebras of V that are subdirect products of subalgebras of A , and*
- *the category of Zariski closed subsets C of A^κ where κ ranges over all the cardinal numbers.*

$$\mathcal{F}_\kappa / I \longrightarrow \mathbb{V}_A(I)$$

$$\mathcal{F}_\kappa / \mathbb{I}_A(C) \longleftarrow C$$

Theorem

An abelian ℓ -group embeds into \mathbb{R} iff it is simple or trivial.

Moreover, every simple vector lattice is isomorphic to \mathbb{R} .

Theorem

An abelian ℓ -group embeds into \mathbb{R} iff it is simple or trivial.

Moreover, every simple vector lattice is isomorphic to \mathbb{R} .

- Every semisimple abelian ℓ -group/vector lattice is subdirect product of subalgebras of \mathbb{R} .

Theorem

An abelian ℓ -group embeds into \mathbb{R} iff it is simple or trivial.

Moreover, every simple vector lattice is isomorphic to \mathbb{R} .

- Every semisimple abelian ℓ -group/vector lattice is subdirect product of subalgebras of \mathbb{R} .
- The Zariski closed subsets of \mathbb{R}^{κ} are the closed cones.

Theorem

An abelian ℓ -group embeds into \mathbb{R} iff it is simple or trivial.

Moreover, every simple vector lattice is isomorphic to \mathbb{R} .

- Every semisimple abelian ℓ -group/vector lattice is subdirect product of subalgebras of \mathbb{R} .
- The Zariski closed subsets of \mathbb{R}^κ are the closed cones.
- $\mathcal{F}_\kappa / \mathbb{I}_\mathbb{R}(C) \cong \text{PWL}_\mathbb{R}(C)$ (vector lattices)
 $\mathcal{F}_\kappa / \mathbb{I}_\mathbb{R}(C) \cong \text{PWL}_\mathbb{Z}(C)$ (abelian ℓ -groups)

Applying the general affine duality approach with $A = \mathbb{R}$

Theorem

An abelian ℓ -group embeds into \mathbb{R} iff it is simple or trivial.

Moreover, every simple vector lattice is isomorphic to \mathbb{R} .

- Every semisimple abelian ℓ -group/vector lattice is subdirect product of subalgebras of \mathbb{R} .
- The Zariski closed subsets of \mathbb{R}^κ are the closed cones.
- $\mathcal{F}_\kappa / \mathbb{I}_\mathbb{R}(C) \cong \text{PWL}_\mathbb{R}(C)$ (vector lattices)
 $\mathcal{F}_\kappa / \mathbb{I}_\mathbb{R}(C) \cong \text{PWL}_\mathbb{Z}(C)$ (abelian ℓ -groups)

Thus, this approach yields Baker-Beynon duality.

Beyond Baker-Beynon duality

Definition

An ℓ -ideal I is **prime** if $a \wedge b \in I$ implies $a \in I$ or $b \in I$.

Definition

An ℓ -ideal I is **prime** if $a \wedge b \in I$ implies $a \in I$ or $b \in I$.

- A/I is linearly ordered iff I is prime.
- Every ℓ -ideal is intersection of prime ℓ -ideals.
- Every abelian ℓ -group/vector lattice is subdirect product of linearly ordered ones.

Definition

An ℓ -ideal I is **prime** if $a \wedge b \in I$ implies $a \in I$ or $b \in I$.

- A/I is linearly ordered iff I is prime.
- Every ℓ -ideal is intersection of prime ℓ -ideals.
- Every abelian ℓ -group/vector lattice is subdirect product of linearly ordered ones.

To apply the general affine duality approach we need A such that every linearly ordered abelian ℓ -group/vector lattice embeds into A .

Definition

An ℓ -ideal I is **prime** if $a \wedge b \in I$ implies $a \in I$ or $b \in I$.

- A/I is linearly ordered iff I is prime.
- Every ℓ -ideal is intersection of prime ℓ -ideals.
- Every abelian ℓ -group/vector lattice is subdirect product of linearly ordered ones.

To apply the general affine duality approach we need A such that every linearly ordered abelian ℓ -group/vector lattice embeds into A .

This is not possible for cardinality reasons. However, such an A exists if we impose a bound on the cardinality/number of generators.

Definition

An ℓ -ideal I is **prime** if $a \wedge b \in I$ implies $a \in I$ or $b \in I$.

- A/I is linearly ordered iff I is prime.
- Every ℓ -ideal is intersection of prime ℓ -ideals.
- Every abelian ℓ -group/vector lattice is subdirect product of linearly ordered ones.

To apply the general affine duality approach we need A such that every linearly ordered abelian ℓ -group/vector lattice embeds into A .

This is not possible for cardinality reasons. However, such an A exists if we impose a bound on the cardinality/number of generators.

Theorem

Let γ be a cardinal. There exists an ultrapower \mathcal{U} of \mathbb{R} such that every κ -generated linearly ordered abelian ℓ -group/vector lattice with $\kappa \leq \gamma$ embeds into \mathcal{U} .

Applying the general affine duality approach with $A = \mathcal{U}$

If $\kappa \leq \gamma$, then every κ -generated abelian ℓ -group/vector lattice is subdirect product of subalgebras of \mathcal{U} .

Applying the general affine duality approach with $A = \mathcal{U}$

If $\kappa \leq \gamma$, then every κ -generated abelian ℓ -group/vector lattice is subdirect product of subalgebras of \mathcal{U} .

Theorem (C., Lapenta, and Spada)

Let γ be a cardinal. There exists an ultrapower \mathcal{U} of \mathbb{R} such that:

- *The category of κ -generated vector lattices for some $\kappa \leq \gamma$ is dually equivalent to the category of Zariski closed subsets of \mathcal{U}^κ for some $\kappa \leq \gamma$.*

Applying the general affine duality approach with $A = \mathcal{U}$

If $\kappa \leq \gamma$, then every κ -generated abelian ℓ -group/vector lattice is subdirect product of subalgebras of \mathcal{U} .

Theorem (C., Lapenta, and Spada)

Let γ be a cardinal. There exists an ultrapower \mathcal{U} of \mathbb{R} such that:

- *The category of κ -generated vector lattices for some $\kappa \leq \gamma$ is dually equivalent to the category of Zariski closed subsets of \mathcal{U}^κ for some $\kappa \leq \gamma$.*
- *The category of κ -generated abelian ℓ -groups for some $\kappa \leq \gamma$ is dually equivalent to the category of Zariski closed subsets of \mathcal{U}^κ for some $\kappa \leq \gamma$.*

Applying the general affine duality approach with $A = \mathcal{U}$

If $\kappa \leq \gamma$, then every κ -generated abelian ℓ -group/vector lattice is subdirect product of subalgebras of \mathcal{U} .

Theorem (C., Lapenta, and Spada)

Let γ be a cardinal. There exists an ultrapower \mathcal{U} of \mathbb{R} such that:

- *The category of κ -generated vector lattices for some $\kappa \leq \gamma$ is dually equivalent to the category of Zariski closed subsets of \mathcal{U}^κ for some $\kappa \leq \gamma$.*
- *The category of κ -generated abelian ℓ -groups for some $\kappa \leq \gamma$ is dually equivalent to the category of Zariski closed subsets of \mathcal{U}^κ for some $\kappa \leq \gamma$.*

The Zariski topology on \mathcal{U}^κ depends on whether we work with abelian ℓ -groups or vector lattices.

Enlargements of piecewise linear functions

Every piecewise linear function $f : \mathbb{R} \rightarrow \mathbb{R}$ can be extended to a function ${}^*f : \mathcal{U} \rightarrow \mathcal{U}$ by setting ${}^*f([(r_i)_{i \in I}]) = [(f(r_i))_{i \in I}]$.

Enlargements of piecewise linear functions

Every piecewise linear function $f : \mathbb{R} \rightarrow \mathbb{R}$ can be extended to a function ${}^*f : \mathcal{U} \rightarrow \mathcal{U}$ by setting ${}^*f([(r_i)_{i \in I}]) = [(f(r_i))_{i \in I}]$.

Similarly, we can extend every piecewise linear $f : \mathbb{R}^\kappa \rightarrow \mathbb{R}$ to ${}^*f : \mathcal{U}^\kappa \rightarrow \mathcal{U}$ which is called the **enlargement** of f .

Enlargements of piecewise linear functions

Every piecewise linear function $f : \mathbb{R} \rightarrow \mathbb{R}$ can be extended to a function ${}^*f : \mathcal{U} \rightarrow \mathcal{U}$ by setting ${}^*f([(r_i)_{i \in I}]) = [(f(r_i))_{i \in I}]$.

Similarly, we can extend every piecewise linear $f : \mathbb{R}^\kappa \rightarrow \mathbb{R}$ to ${}^*f : \mathcal{U}^\kappa \rightarrow \mathcal{U}$ which is called the **enlargement** of f .

We define:

$${}^*\text{PWL}_{\mathbb{R}}(\mathcal{U}^\kappa) = \{{}^*f \mid f \in \text{PWL}_{\mathbb{R}}(\mathbb{R}^\kappa)\},$$

$${}^*\text{PWL}_{\mathbb{Z}}(\mathcal{U}^\kappa) = \{{}^*f \mid f \in \text{PWL}_{\mathbb{Z}}(\mathbb{R}^\kappa)\}.$$

Enlargements of piecewise linear functions

Every piecewise linear function $f : \mathbb{R} \rightarrow \mathbb{R}$ can be extended to a function ${}^*f : \mathcal{U} \rightarrow \mathcal{U}$ by setting ${}^*f([(r_i)_{i \in I}]) = [(f(r_i))_{i \in I}]$.

Similarly, we can extend every piecewise linear $f : \mathbb{R}^\kappa \rightarrow \mathbb{R}$ to ${}^*f : \mathcal{U}^\kappa \rightarrow \mathcal{U}$ which is called the **enlargement** of f .

We define:

$${}^*\text{PWL}_{\mathbb{R}}(\mathcal{U}^\kappa) = \{{}^*f \mid f \in \text{PWL}_{\mathbb{R}}(\mathbb{R}^\kappa)\},$$

$${}^*\text{PWL}_{\mathbb{Z}}(\mathcal{U}^\kappa) = \{{}^*f \mid f \in \text{PWL}_{\mathbb{Z}}(\mathbb{R}^\kappa)\}.$$

If $X \subseteq \mathcal{U}^\kappa$, we can consider ${}^*\text{PWL}_{\mathbb{R}}(X)$ and ${}^*\text{PWL}_{\mathbb{Z}}(X)$.

Proposition

Let C be a Zariski closed subset of \mathcal{U}^κ .

- $\mathcal{F}_\kappa / \mathbb{I}_{\mathcal{U}}(C) \cong {}^*\text{PWL}_{\mathbb{R}}(C)$ (vector lattices).
- $\mathcal{F}_\kappa / \mathbb{I}_{\mathcal{U}}(C) \cong {}^*\text{PWL}_{\mathbb{Z}}(C)$ (abelian ℓ -groups).

The Zariski topology on \mathcal{U}^n

We want to understand what these Zariski topologies look like in the finite-dimensional case.

Irreducible closed subsets

We want to understand what these Zariski topologies look like in the finite-dimensional case.

Definition

A closed subset of a topological space is said to be **irreducible** if it is not the union of two proper closed subsets.

Irreducible closed subsets

We want to understand what these Zariski topologies look like in the finite-dimensional case.

Definition

A closed subset of a topological space is said to be **irreducible** if it is not the union of two proper closed subsets.

Irreducible closed in \mathcal{U}^n are exactly the closure of points. They are the subsets $\mathbb{V}_{\mathcal{U}}(I)$ with I prime or $I = \mathcal{F}_n$.

Irreducible closed subsets

We want to understand what these Zariski topologies look like in the finite-dimensional case.

Definition

A closed subset of a topological space is said to be **irreducible** if it is not the union of two proper closed subsets.

Irreducible closed in \mathcal{U}^n are exactly the closure of points. They are the subsets $\mathbb{V}_{\mathcal{U}}(I)$ with I prime or $I = \mathcal{F}_n$.

The irreducible Zariski-closed subsets of \mathbb{R}^n are the semilines starting from the origin ($\mathbb{V}_{\mathbb{R}}(I)$ with I maximal) and the origin ($\mathbb{V}_{\mathbb{R}}(I)$ with $I = \mathcal{F}_n$).

Orthogonal decomposition theorem (Goze 1995)

If $x \in \mathcal{U}^n$, then x can be written in a unique way as $\alpha_1 v_1 + \cdots + \alpha_k v_k$ with v_1, \dots, v_k orthonormal vectors of \mathbb{R}^n and $0 < \alpha_1, \dots, \alpha_n \in \mathcal{U}$ such that α_{i+1}/α_i is infinitesimal.

Orthogonal decomposition theorem (Goze 1995)

If $x \in \mathcal{U}^n$, then x can be written in a unique way as $\alpha_1 v_1 + \dots + \alpha_k v_k$ with v_1, \dots, v_k orthonormal vectors of \mathbb{R}^n and $0 < \alpha_1, \dots, \alpha_n \in \mathcal{U}$ such that α_{i+1}/α_i is infinitesimal.

Thus, we can associate to each $x \in \mathcal{U}^n$ the sequence

$\mathbf{v} = (v_1, \dots, v_k)$ of orthonormal vectors.

We call such sequences **indices**.

Orthogonal decomposition theorem (Goze 1995)

If $x \in \mathcal{U}^n$, then x can be written in a unique way as $\alpha_1 v_1 + \dots + \alpha_k v_k$ with v_1, \dots, v_k orthonormal vectors of \mathbb{R}^n and $0 < \alpha_1, \dots, \alpha_n \in \mathcal{U}$ such that α_{i+1}/α_i is infinitesimal.

Thus, we can associate to each $x \in \mathcal{U}^n$ the sequence $\mathbf{v} = (v_1, \dots, v_k)$ of orthonormal vectors.

We call such sequences **indices**.

Let $\text{Cone}(\mathbf{v})$ be the set of points of \mathcal{U}^n whose index is a truncation of \mathbf{v} .

Orthogonal decomposition theorem (Goze 1995)

If $x \in \mathcal{U}^n$, then x can be written in a unique way as $\alpha_1 v_1 + \dots + \alpha_k v_k$ with v_1, \dots, v_k orthonormal vectors of \mathbb{R}^n and $0 < \alpha_1, \dots, \alpha_n \in \mathcal{U}$ such that α_{i+1}/α_i is infinitesimal.

Thus, we can associate to each $x \in \mathcal{U}^n$ the sequence $\mathbf{v} = (v_1, \dots, v_k)$ of orthonormal vectors.

We call such sequences **indices**.

Let $\text{Cone}(\mathbf{v})$ be the set of points of \mathcal{U}^n whose index is a truncation of \mathbf{v} .

Theorem (C., Lapenta, Spada)

In the Zariski topology of \mathcal{U}^n relative to vector lattices each irreducible closed of \mathcal{U}^n is $\text{Cone}(\mathbf{v})$ for some index \mathbf{v} .

Indices and cones

Every subset $X \subseteq \mathbb{R}^n$ can be associated with a subset *X of \mathcal{U}^n called the **enlargement** of X . Every predicate $P \subseteq \mathbb{R}^n$ and function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ can be enlarged to ${}^*P \subseteq \mathcal{U}^n$ and ${}^*f : \mathcal{U}^n \rightarrow \mathcal{U}$.

Indices and cones

Every subset $X \subseteq \mathbb{R}^n$ can be associated with a subset *X of \mathcal{U}^n called the **enlargement** of X . Every predicate $P \subseteq \mathbb{R}^n$ and function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ can be enlarged to ${}^*P \subseteq \mathcal{U}^n$ and ${}^*f : \mathcal{U}^n \rightarrow \mathcal{U}$.

Transfer principle (Łoś Theorem)

Let φ be a first order formula and ${}^*\varphi$ the formula obtained by replacing every predicate symbol P and every function symbol f with *P and *f . Then φ is true in \mathbb{R} iff ${}^*\varphi$ is true in \mathcal{U} .

Indices and cones

Every subset $X \subseteq \mathbb{R}^n$ can be associated with a subset *X of \mathcal{U}^n called the **enlargement** of X . Every predicate $P \subseteq \mathbb{R}^n$ and function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ can be enlarged to ${}^*P \subseteq \mathcal{U}^n$ and ${}^*f : \mathcal{U}^n \rightarrow \mathcal{U}$.

Transfer principle (Łoś Theorem)

Let φ be a first order formula and ${}^*\varphi$ the formula obtained by replacing every predicate symbol P and every function symbol f with *P and *f . Then φ is true in \mathbb{R} iff ${}^*\varphi$ is true in \mathcal{U} .

If \mathbf{v} is an index, we say that a closed cone of \mathbb{R}^n is a **v-cone** if there exist real numbers $r_2, \dots, r_k > 0$ such that the cone is generated by $\{v_1, v_1 + r_2 v_2, \dots, v_1 + r_2 v_2 + \dots + r_k v_k\}$.

Indices and cones

Every subset $X \subseteq \mathbb{R}^n$ can be associated with a subset *X of \mathcal{U}^n called the **enlargement** of X . Every predicate $P \subseteq \mathbb{R}^n$ and function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ can be enlarged to ${}^*P \subseteq \mathcal{U}^n$ and ${}^*f : \mathcal{U}^n \rightarrow \mathcal{U}$.

Transfer principle (Łoś Theorem)


Let φ be a first order formula and ${}^*\varphi$ the formula obtained by replacing every predicate symbol P and every function symbol f with *P and *f . Then φ is true in \mathbb{R} iff ${}^*\varphi$ is true in \mathcal{U} .

If \mathbf{v} is an index, we say that a closed cone of \mathbb{R}^n is a **v-cone** if there exist real numbers $r_2, \dots, r_k > 0$ such that the cone is generated by $\{v_1, v_1 + r_2 v_2, \dots, v_1 + r_2 v_2 + \dots + r_k v_k\}$.


Proposition

$\text{Cone}(\mathbf{v})$ is the intersection of the enlargements of all the **v-cones**.


$$\mathbf{v} = ((1, 0), (0, 1)).$$




$$\mathbf{v} = ((1, 0), (0, 1)).$$




$$\mathbf{v} = ((1, 0), (0, 1)).$$



$$\mathbf{v} = ((1, 0), (0, 1)).$$



$$\mathbf{v} = ((1, 0), (0, 1)).$$



Theorem (C., Lapenta, and Spada)

*If $f \in \text{PWL}_{\mathbb{R}}(\mathbb{R}^n)$, then $*f$ vanishes on $\text{Cone}(\mathbf{v})$ iff f vanishes on some \mathbf{v} -cone.*

Theorem (C., Lapenta, and Spada)

*If $f \in \text{PWL}_{\mathbb{R}}(\mathbb{R}^n)$, then $*f$ vanishes on $\text{Cone}(\mathbf{v})$ iff f vanishes on some \mathbf{v} -cone.*

As a corollary, we obtain the description of prime ℓ -ideals in finitely generated vector lattices due to Panti.

Theorem (Panti 1999)

Each prime ℓ -ideal of the vector lattice \mathcal{F}_n is of the form $\{f \in \text{PWL}_{\mathbb{R}}(\mathbb{R}^n) \mid f \text{ vanishes on a } \mathbf{v}\text{-cone}\}$ for some index \mathbf{v} .

Theorem (C., Lapenta, and Spada)

*If $f \in \text{PWL}_{\mathbb{R}}(\mathbb{R}^n)$, then $*f$ vanishes on $\text{Cone}(\mathbf{v})$ iff f vanishes on some \mathbf{v} -cone.*

As a corollary, we obtain the description of prime ℓ -ideals in finitely generated vector lattices due to Panti.

Theorem (Panti 1999)

Each prime ℓ -ideal of the vector lattice \mathcal{F}_n is of the form $\{f \in \text{PWL}_{\mathbb{R}}(\mathbb{R}^n) \mid f \text{ vanishes on a } \mathbf{v}\text{-cone}\}$ for some index \mathbf{v} .

Fix a positive infinitesimal $\varepsilon \in \mathcal{U}$. If I is the prime ℓ -ideal of the vector lattice \mathcal{F}_n associated with the index $\mathbf{v} = (v_1, \dots, v_k)$, then $\mathbb{V}_{\mathcal{U}}(I) = \text{cl}\{v_1 + \varepsilon v_2 + \dots + \varepsilon^{k-1} v_k\}$.

Theorem (C., Lapenta, and Spada)

*If $f \in \text{PWL}_{\mathbb{R}}(\mathbb{R}^n)$, then *f vanishes on $\text{Cone}(\mathbf{v})$ iff f vanishes on some \mathbf{v} -cone.*

As a corollary, we obtain the description of prime ℓ -ideals in finitely generated vector lattices due to Panti.

Theorem (Panti 1999)

Each prime ℓ -ideal of the vector lattice \mathcal{F}_n is of the form $\{f \in \text{PWL}_{\mathbb{R}}(\mathbb{R}^n) \mid f \text{ vanishes on a } \mathbf{v}\text{-cone}\}$ for some index \mathbf{v} .

Fix a positive infinitesimal $\varepsilon \in \mathcal{U}$. If I is the prime ℓ -ideal of the vector lattice \mathcal{F}_n associated with the index $\mathbf{v} = (v_1, \dots, v_k)$, then $\mathbb{V}_{\mathcal{U}}(I) = \text{cl}\{v_1 + \varepsilon v_2 + \dots + \varepsilon^{k-1} v_k\}$.

This allows to embed the spectrum of a finitely generated vector lattice V into its dual cone so that $V \cong {}^*\text{PWL}_{\mathbb{R}}(\text{Spec}(V))$.

Abelian ℓ -groups and \mathbb{Z} -reduced indices

Definition

If $w \in \mathbb{R}^n$, let $\langle w \rangle$ be the smallest subspace containing w that admits a basis in \mathbb{Z}^n .

An index $\mathbf{v} = (v_1, \dots, v_k)$ is \mathbb{Z} -reduced if $\langle v_i \rangle$ and $\langle v_j \rangle$ are orthogonal for each $i \neq j$.

Definition

If $w \in \mathbb{R}^n$, let $\langle w \rangle$ be the smallest subspace containing w that admits a basis in \mathbb{Z}^n .

An index $\mathbf{v} = (v_1, \dots, v_k)$ is \mathbb{Z} -reduced if $\langle v_i \rangle$ and $\langle v_j \rangle$ are orthogonal for each $i \neq j$.

Using a sort of Gram-Schmidt process, we can associate to each index \mathbf{v} a unique \mathbb{Z} -reduced index $\text{red}(\mathbf{v})$.

Abelian ℓ -groups and \mathbb{Z} -reduced indices

Definition

If $w \in \mathbb{R}^n$, let $\langle w \rangle$ be the smallest subspace containing w that admits a basis in \mathbb{Z}^n .

An index $\mathbf{v} = (v_1, \dots, v_k)$ is \mathbb{Z} -reduced if $\langle v_i \rangle$ and $\langle v_j \rangle$ are orthogonal for each $i \neq j$.

Using a sort of Gram-Schmidt process, we can associate to each index \mathbf{v} a unique \mathbb{Z} -reduced index $\text{red}(\mathbf{v})$.

Theorem (C., Lapenta, and Spada)

In the Zariski topology of \mathcal{U}^n relative to abelian ℓ -groups each irreducible closed of \mathcal{U}^n is of the form

$$\bigcup \{ \text{Cone}(\mathbf{w}) \mid \text{red}(\mathbf{w}) = \mathbf{v} \}.$$

for some \mathbb{Z} -reduced index \mathbf{v} .

Theorem (C., Lapenta, and Spada)

Let γ be a cardinal. There exists an ultrapower \mathcal{U} of $[0, 1]$ such that:

- The category of κ -generated MV-algebras for some $\kappa \leq \gamma$ is dually equivalent to the category of Zariski closed subsets of \mathcal{U}^κ for some $\kappa \leq \gamma$.
- The category of κ -generated Riesz MV-algebras for some $\kappa \leq \gamma$ is dually equivalent to the category of Zariski closed subsets of \mathcal{U}^κ for some $\kappa \leq \gamma$.

Theorem (C., Lapenta, and Spada)

Let γ be a cardinal. There exists an ultrapower \mathcal{U} of $[0, 1]$ such that:

- The category of κ -generated MV-algebras for some $\kappa \leq \gamma$ is dually equivalent to the category of Zariski closed subsets of \mathcal{U}^κ for some $\kappa \leq \gamma$.
- The category of κ -generated Riesz MV-algebras for some $\kappa \leq \gamma$ is dually equivalent to the category of Zariski closed subsets of \mathcal{U}^κ for some $\kappa \leq \gamma$.

The irreducible closed in \mathcal{U}^n correspond to “infinitesimal simplices”.

Theorem (C., Lapenta, and Spada)

Let γ be a cardinal. There exists an ultrapower \mathcal{U} of $[0, 1]$ such that:

- The category of κ -generated MV-algebras for some $\kappa \leq \gamma$ is dually equivalent to the category of Zariski closed subsets of \mathcal{U}^κ for some $\kappa \leq \gamma$.
- The category of κ -generated Riesz MV-algebras for some $\kappa \leq \gamma$ is dually equivalent to the category of Zariski closed subsets of \mathcal{U}^κ for some $\kappa \leq \gamma$.

The irreducible closed in \mathcal{U}^n correspond to “infinitesimal simplices”.

This is an affine version of the dualities for abelian ℓ -groups and vector lattices.

THANK YOU!