

Math C5.4, Networks, University of Oxford, MT 2022

Problem Sheet 2

Peter Grindrod, Mathematical Institute

1. *Erdos-Renyi*. The simplest and most famous type of random graph is an Erdős–Rényi (ER) graph (which was first studied by Solomonoff and Rapoport). In particular, consider the random-graph ensemble $\mathcal{G}(N, p)$, which is defined as follows: Suppose that there are N nodes. Between each pair of distinct nodes, a single edge exists with uniform and independent probability p . There are no self-edges. A single graph $G \in \mathcal{G}(N, p)$ is generated using this process, and it is interesting to study the properties of collections (“ensembles”) of graphs that are generated in this way. The probability in which each simple graph G with m edges appears in a graph $G \in \mathcal{G}(N, p)$ is

$$P(G) = p^m (1 - p)^{n C_2 - m} . \quad (1)$$

- (a) Write down the total probability of drawing a graph G with exactly m edges from the ensemble $\mathcal{G}(N, p)$, and use it to find the mean value of edges $\langle m \rangle$.
- (b) Calculate the expected mean degree of an ER graph.
- (c) Show, under an appropriate assumption (which you should state), that the degree distribution for an ER graph (in expectation over the ensemble) satisfies

$$p_k \sim e^{-c} \frac{c^k}{k!}, \quad N \rightarrow \infty, \quad (2)$$

where $c = (N - 1)p$.

- (d) Reproduce numerically the results of Figure (14) in the lecture notes.
 - (e) Calculate the global clustering coefficient C of an ER graph (in expectation over the ensemble), and verify your prediction numerically.
 - (f) Show that the diameter of an ER graph is $B + \frac{\ln N}{\ln c}$ as $N \rightarrow \infty$, where B is a constant and $c = (N - 1)p$.
2. *Configuration model*.
 - (a) One way to create an undirected network having a desired vertex degree distribution, is to create new undirected networks from an existing one (with the required distribution). Suppose that we have such a undirected network: select two separate existing edges at random, independently, say (v_a, v_b) and (v_c, v_d) . We will delete the edge (v_a, v_b) and (v_c, v_d) and replace them with new edges (v_a, v_d) and (v_c, v_b) providing that neither of the latter already exists. This is easy to check if we hold the present network by keeping a list of edges. Consider a sparse graph on n vertices with a number $m \ll n(n - 1)/2$ of edges in total, and doing this “swap” independently a total of K consecutive times. What is the probability, q , that any particular edge has NOT been replaced after exactly K edge-pair swaps? Show that if we wish to have $q < 1/10m$ (i.e. that we expect to have all edges replaced) we should have $K > m \ln(10m)/2$.
 - (b) Check this with a numerical implementation.

3. Range Dependent networks.

- (a) Consider an undirected network on an infinite number of vertices indexed by both the negative and positive integers $i = \dots - 2, -1, 0, 1, 2, \dots$. Consider the random network where an edge exist between vertex v_i and vertex v_j ($i \neq j$) with independent probability equal to $f(|i - j|) \equiv \alpha \lambda^{|i - j| - 1}$. Here $0 < \alpha, \lambda \leq 1$, are fixed parameters. For any pair of vertices, v_i and vertex v_j , $|i - j|$ is called the range of the possible connecting edge. How many edges is a vertex expected to have to the right? What is the expected degree of each vertex?
- (b) Generate instances of such a graph on a large number n of vertices, with $\alpha = 1$, so that neighbours are always connected. Calculate the Watts-Strogatz clustering coefficient. How does that vary with $\lambda \in (0, 1]$? Can you produce a figure like Figure 1, below?




FIG. 1: Clustering coefficient versus $\lambda = 0.05, 0.1, 0.15, \dots$ (for $\alpha = 1$). Experiments are all for $n = 500$ plotted as points; solid curve is $3\lambda/(1+\lambda)(1+3\lambda)$ from <https://tinyurl.com/yc5rxnjc>.