

Linear Algebraic Quantifiers

Anuj Dawar

Department of Computer Science and Technology, University of Cambridge

RiC 2022, UCL, 22 September 2022

Descriptive Complexity

Descriptive Complexity

A central question in the field of *Descriptive Complexity* is the question of whether there is a *logic for P*.

Descriptive Complexity

A central question in the field of *Descriptive Complexity* is the question of whether there is a *logic for P*.

On *ordered structures* FP —the extension of first-order logic with a fixed-point operator—suffices. (Immerman-Vardi)

Descriptive Complexity

A central question in the field of *Descriptive Complexity* is the question of whether there is a *logic for P*.

On *ordered structures* **FP**—the extension of first-order logic with a fixed-point operator—suffices. **(Immerman-Vardi)**

FP is not sufficient in the absence of order.

Descriptive Complexity

A central question in the field of *Descriptive Complexity* is the question of whether there is a *logic for P*.

On *ordered structures* FP —the extension of first-order logic with a fixed-point operator—suffices. (Immerman-Vardi)

FP is not sufficient in the absence of order. This can be shown by constructing properties in P not definable in $L_{\infty\omega}^\omega$ —finite variable infinitary logic.

Descriptive Complexity

A central question in the field of *Descriptive Complexity* is the question of whether there is a *logic for P*.

On *ordered structures* FP —the extension of first-order logic with a fixed-point operator—suffices. (Immerman-Vardi)

FP is not sufficient in the absence of order. This can be shown by constructing properties in P not definable in $L_{\infty\omega}^\omega$ —finite variable infinitary logic.

Many extensions of FP with additional operators have been studied.

Descriptive Complexity

A central question in the field of *Descriptive Complexity* is the question of whether there is a *logic for P*.

On *ordered structures* FP —the extension of first-order logic with a fixed-point operator—suffices. (Immerman-Vardi)

FP is not sufficient in the absence of order. This can be shown by constructing properties in P not definable in $L_{\infty\omega}^\omega$ —finite variable infinitary logic.

Many extensions of FP with additional operators have been studied. These are studied through the expressive power of $L_{\infty\omega}^\omega(Q)$, the extension of $L_{\infty\omega}^\omega$ with a set Q of *quantifiers*.

Generalized Quantifiers

Generalized Quantifiers

A *Lindström quantifier* of relational vocabulary σ is given by:
 K —a class of σ -structures, closed under isomorphisms.

Generalized Quantifiers

A *Lindström quantifier* of relational vocabulary σ is given by:

K —a class of σ -structures, closed under isomorphisms.

In this talk, we only consider *finite structures*.

Generalized Quantifiers

A *Lindström quantifier* of relational vocabulary σ is given by:

K —a class of σ -structures, closed under isomorphisms.

In this talk, we only consider *finite structures*.

$L(K)$ is the extension of a *logic* L with the quantifier for K .

Generalized Quantifiers

A *Lindström quantifier* of relational vocabulary σ is given by:

K —a class of σ -structures, closed under isomorphisms.

In this talk, we only consider *finite structures*.

$L(K)$ is the extension of a *logic* L with the quantifier for K .

More generally, for a collection Q of quantifiers, $L(Q)$ is the extension of L with *all* quantifiers in Q .

Logics

Logics

Logics L we are interested in for the purpose of this talk are

Logics

Logics L we are interested in for the purpose of this talk are

- *first-order logic*— $L_{\omega\omega}$ or FO.

Logics

Logics L we are interested in for the purpose of this talk are

- *first-order logic*— $L_{\omega\omega}$ or FO.
- *infinitary logic*— $L_{\infty\omega}$ or $L_{\omega_1\omega}$.

Logics

Logics L we are interested in for the purpose of this talk are

- *first-order logic*— $L_{\omega\omega}$ or FO.
- *infinitary logic*— $L_{\infty\omega}$ or $L_{\omega_1\omega}$.

*The closure of FO under *infinitary* (or *countable*) conjunctions.*

- *k-variable infinitary logic*— $L_{\infty\omega}^k$.

Logics

Logics L we are interested in for the purpose of this talk are

- *first-order logic*— $L_{\omega\omega}$ or FO.
- *infinitary logic*— $L_{\infty\omega}$ or $L_{\omega_1\omega}$.

The closure of FO under infinitary (or countable) conjunctions.

- *k -variable infinitary logic*— $L_{\infty\omega}^k$.
- *finite-variable infinitary logic*— $L_{\infty\omega}^\omega = \bigcup_{k < \omega} L_{\infty\omega}^k$.

Logics

Logics L we are interested in for the purpose of this talk are

- *first-order logic*— $L_{\omega\omega}$ or FO.
- *infinitary logic*— $L_{\infty\omega}$ or $L_{\omega_1\omega}$.

The closure of FO under *infinitary* (or *countable*) conjunctions.

- *k-variable infinitary logic*— $L_{\infty\omega}^k$.
- *finite-variable infinitary logic*— $L_{\infty\omega}^\omega = \bigcup_{k < \omega} L_{\infty\omega}^k$.

Note that the expressive power of $L_{\omega_1\omega}$ on finite structures is *complete*. That is to say, it can define every *isomorphism-closed* class of structures.

Logics with Generalized Quantifiers

Logics with Generalized Quantifiers

If $\sigma = (R_1, \dots, R_r)$ we get a formula

$$K\mathbf{x}(\varphi_1(\mathbf{x}_1), \dots, \varphi_r(\mathbf{x}_r)).$$

Logics with Generalized Quantifiers

If $\sigma = (R_1, \dots, R_r)$ we get a formula

$$K\mathbf{x}(\varphi_1(\mathbf{x}_1), \dots, \varphi_r(\mathbf{x}_r)).$$

$$|\mathbf{x}_i| = \text{ar}(R_i)$$

Logics with Generalized Quantifiers

If $\sigma = (R_1, \dots, R_r)$ we get a formula

$$K\mathbf{x}(\varphi_1(\mathbf{x}_1), \dots, \varphi_r(\mathbf{x}_r)).$$

$$|\mathbf{x}_i| = \text{ar}(R_i)$$

The *arity* of the quantifier K is $\max_i \text{ar}(R_i)$.

Logics with Generalized Quantifiers

If $\sigma = (R_1, \dots, R_r)$ we get a formula

$$K\mathbf{x}(\varphi_1(\mathbf{x}_1), \dots, \varphi_r(\mathbf{x}_r)).$$

$$|\mathbf{x}_i| = \text{ar}(R_i)$$

The *arity* of the quantifier K is $\max_i \text{ar}(R_i)$.

$L(K)$ is the *minimal* extension of L that can express K and is closed under the operations of L , such as

Logics with Generalized Quantifiers

If $\sigma = (R_1, \dots, R_r)$ we get a formula

$$K\mathbf{x}(\varphi_1(\mathbf{x}_1), \dots, \varphi_r(\mathbf{x}_r)).$$

$$|\mathbf{x}_i| = \text{ar}(R_i)$$

The *arity* of the quantifier K is $\max_i \text{ar}(R_i)$.

$L(K)$ is the *minimal* extension of L that can express K and is closed under the operations of L , such as

- *Boolean operations*
- *particularization* (i.e. existential quantification)

Equivalences

Equivalences

For a set of quantifiers Q , write

$$\mathbb{A} \equiv_Q^k \mathbb{B}$$

to denote that \mathbb{A} and \mathbb{B} are not distinguishable in $L_{\infty\omega}^k(Q)$.

Equivalences

For a set of quantifiers Q , write

$$\mathbb{A} \equiv_Q^k \mathbb{B}$$

to denote that \mathbb{A} and \mathbb{B} are not distinguishable in $L_{\infty\omega}^k(Q)$.

For a relational vocabulary τ , we say that \equiv_Q^k is *discrete* if for any pair \mathbb{A}, \mathbb{B} of τ -structures

$$\mathbb{A} \equiv_Q^k \mathbb{B} \quad \text{if, and only if,} \quad \mathbb{A} \cong \mathbb{B}$$

Equivalences

For a set of quantifiers Q , write

$$\mathbb{A} \equiv_Q^k \mathbb{B}$$

to denote that \mathbb{A} and \mathbb{B} are not distinguishable in $L_{\infty\omega}^k(Q)$.

For a relational vocabulary τ , we say that \equiv_Q^k is *discrete* if for any pair \mathbb{A}, \mathbb{B} of τ -structures

$$\mathbb{A} \equiv_Q^k \mathbb{B} \quad \text{if, and only if,} \quad \mathbb{A} \cong \mathbb{B}$$

The following are equivalent:

Equivalences

For a set of quantifiers Q , write

$$\mathbb{A} \equiv_Q^k \mathbb{B}$$

to denote that \mathbb{A} and \mathbb{B} are not distinguishable in $L_{\infty\omega}^k(Q)$.

For a relational vocabulary τ , we say that \equiv_Q^k is *discrete* if for any pair \mathbb{A}, \mathbb{B} of τ -structures

$$\mathbb{A} \equiv_Q^k \mathbb{B} \quad \text{if, and only if,} \quad \mathbb{A} \cong \mathbb{B}$$

The following are equivalent:

- There is some k such that \equiv_Q^k is discrete on τ -structures.
- The expressive power of $L_{\infty\omega}^\omega(Q)$ is complete on τ -structures.

Arity Hierarchy

Arity Hierarchy

Let Q_n denote the collection of *all* n -ary quantifiers.

Arity Hierarchy

Let Q_n denote the collection of *all* n -ary quantifiers.

Theorem (Hella)

For every n , there is a vocabulary τ such that $\equiv_{Q_n}^k$ is not discrete on τ -structures for any k .

Arity Hierarchy

Let Q_n denote the collection of *all* n -ary quantifiers.

Theorem (Hella)

For every n , there is a vocabulary τ such that $\equiv_{Q_n}^k$ is not discrete on τ -structures for any k .

The class of structures not definable in $L_{\infty\omega}^\omega(Q_n)$ can be constructed to be decidable in P .

Arity Hierarchy

Let Q_n denote the collection of *all* n -ary quantifiers.

Theorem (Hella)

For every n , there is a vocabulary τ such that $\equiv_{Q_n}^k$ is not discrete on τ -structures for any k .

The class of structures not definable in $L_{\infty\omega}^\omega(Q_n)$ can be constructed to be decidable in P .

Note: τ necessarily contains relations of arity $\geq n + 1$.

Unary and Binary Quantifiers

Unary and Binary Quantifiers

$L_{\infty\omega}^\omega(Q_1)$ has the same expressive power as $L_{\infty\omega}^\omega(C)$ —

Unary and Binary Quantifiers

$L_{\infty\omega}^\omega(Q_1)$ has the same expressive power as $L_{\infty\omega}^\omega(C)$ —where C is the collection of all *unary counting quantifiers*.

Unary and Binary Quantifiers

$L_{\infty\omega}^\omega(Q_1)$ has the same expressive power as $L_{\infty\omega}^\omega(C)$ —where C is the collection of all *unary counting quantifiers*.

$$\exists \geq n, \exists \leq n$$

Unary and Binary Quantifiers

$L_{\infty\omega}^\omega(Q_1)$ has the same expressive power as $L_{\infty\omega}^\omega(C)$ —where C is the collection of all *unary counting quantifiers*.

$\exists \geq n, \exists \leq n$

Graph properties in P not definable in $L_{\infty\omega}^\omega(C)$ were constructed by **(Cai-Fürer-Immerman)**.

Unary and Binary Quantifiers

$L_{\infty\omega}^\omega(Q_1)$ has the same expressive power as $L_{\infty\omega}^\omega(C)$ —where C is the collection of all *unary counting quantifiers*.

$$\exists \geq n, \exists \leq n$$

Graph properties in P not definable in $L_{\infty\omega}^\omega(C)$ were constructed by **(Cai-Fürer-Immerman)**.

$L_{\infty\omega}^\omega(Q_2)$ can express *all* properties of graphs.

Unary and Binary Quantifiers

$L_{\infty\omega}^{\omega}(Q_1)$ has the same expressive power as $L_{\infty\omega}^{\omega}(C)$ —where C is the collection of all *unary counting quantifiers*.

$$\exists^{\geq n}, \exists^{\leq n}$$

Graph properties in P not definable in $L_{\infty\omega}^{\omega}(C)$ were constructed by **(Cai-Fürer-Immerman)**.

$L_{\infty\omega}^{\omega}(Q_2)$ can express *all* properties of graphs.

These logics are not closed under *first-order interpretations*.

Unary and Binary Quantifiers

$L_{\infty\omega}^{\omega}(Q_1)$ has the same expressive power as $L_{\infty\omega}^{\omega}(C)$ —where C is the collection of all *unary counting quantifiers*.

$$\exists^{\geq n}, \exists^{\leq n}$$

Graph properties in P not definable in $L_{\infty\omega}^{\omega}(C)$ were constructed by **(Cai-Fürer-Immerman)**.

$L_{\infty\omega}^{\omega}(Q_2)$ can express *all* properties of graphs.

These logics are not closed under *first-order interpretations*.

Closure under first-order reductions is a desirable property in *descriptive complexity*, as most interesting complexity classes have it.

First-Order Interpretations

An FO *interpretation* θ of a τ -structure \mathbb{B} in a σ -structure \mathbb{A} is a family of first-order formulas which define the *universe* and *relations* of \mathbb{B} when interpreted in \mathbb{A} .

First-Order Interpretations

An FO *interpretation* θ of a τ -structure \mathbb{B} in a σ -structure \mathbb{A} is a family of first-order formulas which define the *universe* and *relations* of \mathbb{B} when interpreted in \mathbb{A} .

This defines a map from σ -structures to τ -structures, so we write $\mathbb{B} = \theta(\mathbb{A})$.

First-Order Interpretations

An FO *interpretation* θ of a τ -structure \mathbb{B} in a σ -structure \mathbb{A} is a family of first-order formulas which define the *universe* and *relations* of \mathbb{B} when interpreted in \mathbb{A} .

This defines a map from σ -structures to τ -structures, so we write $\mathbb{B} = \theta(\mathbb{A})$.

An FO *reduction* of a class of structures \mathcal{C} to a class \mathcal{D} is a single FO interpretation θ such that $\mathbb{A} \in \mathcal{C}$ if, and only if, $\theta(\mathbb{A}) \in \mathcal{D}$.

First-Order Interpretations

An FO *interpretation* θ of a τ -structure \mathbb{B} in a σ -structure \mathbb{A} is a family of first-order formulas which define the *universe* and *relations* of \mathbb{B} when interpreted in \mathbb{A} .

This defines a map from σ -structures to τ -structures, so we write $\mathbb{B} = \theta(\mathbb{A})$.

An FO *reduction* of a class of structures \mathcal{C} to a class \mathcal{D} is a single FO interpretation θ such that $\mathbb{A} \in \mathcal{C}$ if, and only if, $\theta(\mathbb{A}) \in \mathcal{D}$.

We write $\mathcal{C} \leq_{\text{FO}} \mathcal{D}$.

Vectorized Quantifiers

Vectorized Quantifiers

Let $\sigma = (R_1, \dots, R_r)$ be a relational vocabulary.

Vectorized Quantifiers

Let $\sigma = (R_1, \dots, R_r)$ be a relational vocabulary.

A minimal logic extending L , able to express a property K of σ -structures, and *closed* under first-order interpretations is given by $L(\overline{K})$, where

Vectorized Quantifiers

Let $\sigma = (R_1, \dots, R_r)$ be a relational vocabulary.

A minimal logic extending L , able to express a property K of σ -structures, and *closed* under first-order interpretations is given by $L(\overline{K})$, where

\overline{K} is the collection $\{K_d \mid d \in \omega\}$ of Lindström quantifiers in the vocabularies

$$\sigma_d = (U_d, \sim_d, (R_{i,d})_{i \in [r]})$$

with $\text{ar}(U_d) = d$, $\text{ar}(\sim_d) = 2d$ and $\text{ar}(R_{i,d}) = d \cdot \text{ar}(R_i)$,

Vectorized Quantifiers

Let $\sigma = (R_1, \dots, R_r)$ be a relational vocabulary.

A minimal logic extending L , able to express a property K of σ -structures, and *closed* under first-order interpretations is given by $L(\overline{K})$, where

\overline{K} is the collection $\{K_d \mid d \in \omega\}$ of Lindström quantifiers in the vocabularies

$$\sigma_d = (U_d, \sim_d, (R_{i,d})_{i \in [r]})$$

with $\text{ar}(U_d) = d$, $\text{ar}(\sim_d) = 2d$ and $\text{ar}(R_{i,d}) = d \cdot \text{ar}(R_i)$,
and

$$\mathbb{A} \in K_d \quad \text{iff} \quad (U_d^{\mathbb{A}} / \sim_d^{\mathbb{A}}, (R_{i,d}^{\mathbb{A}})_{i \in [r]}) \in K.$$

Vectorizations of Unary Quantifiers

Vectorizations of Unary Quantifiers

Note that $\overline{K} \not\subseteq Q_n$ for any $n \in \omega$.

Vectorizations of Unary Quantifiers

Note that $\overline{K} \not\subseteq Q_n$ for any $n \in \omega$.

Let

$$\overline{Q_n} = \bigcup_{K \in Q_n} \overline{K}$$

Vectorizations of Unary Quantifiers

Note that $\overline{K} \not\subseteq Q_n$ for any $n \in \omega$.

Let

$$\overline{Q_n} = \bigcup_{K \in Q_n} \overline{K}$$

More generally, for any collection S of quantifiers, let \overline{S} denote the collection of *vectorizations* of quantifiers in S .

Vectorizations of Unary Quantifiers

Note that $\overline{K} \not\subseteq Q_n$ for any $n \in \omega$.

Let

$$\overline{Q_n} = \bigcup_{K \in Q_n} \overline{K}$$

More generally, for any collection S of quantifiers, let \overline{S} denote the collection of *vectorizations* of quantifiers in S .

Theorem

$$L_{\infty\omega}^\omega(\overline{Q_1}) \leq L_{\infty\omega}^\omega(C).$$

Vectorizations of Unary Quantifiers

Note that $\overline{K} \not\subseteq Q_n$ for any $n \in \omega$.

Let

$$\overline{Q_n} = \bigcup_{K \in Q_n} \overline{K}$$

More generally, for any collection S of quantifiers, let \overline{S} denote the collection of *vectorizations* of quantifiers in S .

Theorem

$$L_{\infty\omega}^\omega(\overline{Q_1}) \leq L_{\infty\omega}^\omega(C).$$

In short, vectorization adds nothing to *unary* quantifiers.

Vectorizations of Unary Quantifiers

Note that $\overline{K} \not\subseteq Q_n$ for any $n \in \omega$.

Let

$$\overline{Q_n} = \bigcup_{K \in Q_n} \overline{K}$$

More generally, for any collection S of quantifiers, let \overline{S} denote the collection of *vectorizations* of quantifiers in S .

Theorem

$$L_{\infty\omega}^\omega(\overline{Q_1}) \leq L_{\infty\omega}^\omega(C).$$

In short, vectorization adds nothing to *unary* quantifiers.

Counting tuples can always be replaced by counting elements.

Vectorizations of Binary Quantifiers

Vectorizations of Binary Quantifiers

Theorem

$$L_{\infty\omega} \leq L_{\omega\omega}(\overline{Q_2})$$

Vectorizations of Binary Quantifiers

Theorem

$$L_{\infty\omega} \leq L_{\omega\omega}(\overline{Q_2})$$

In short, with vectorized binary quantifiers, we can express *everything*.

Vectorizations of Binary Quantifiers

Theorem

$$L_{\infty\omega} \leq L_{\omega\omega}(\overline{Q_2})$$

In short, with vectorized binary quantifiers, we can express *everything*.

This follows from the fact that for any vocabulary τ , there is a *first-order definable bi-interpretation* to the vocabulary with one binary relation.

Vectorizations of Binary Quantifiers

Theorem

$$L_{\infty\omega} \leq L_{\omega\omega}(\overline{Q_2})$$

In short, with vectorized binary quantifiers, we can express *everything*.

This follows from the fact that for any vocabulary τ , there is a *first-order definable bi-interpretation* to the vocabulary with one binary relation.

So, for any class K of σ -structures, there is a *first-order interpretation* Φ and a class of graphs G such that

$$\Phi(\mathbb{A}) \in G \quad \text{iff} \quad \mathbb{A} \in K.$$

Restricted Classes of Binary Quantifiers

Restricted Classes of Binary Quantifiers

Thus, when it comes to vectorized quantifiers, the *arity hierarchy* has just two levels.

Restricted Classes of Binary Quantifiers

Thus, when it comes to vectorized quantifiers, the *arity hierarchy* has just two levels.

To get *interesting* classes of vectorized quantifiers beyond the unary, we consider *proper subclasses* of $\overline{Q_2}$.

Restricted Classes of Binary Quantifiers

Thus, when it comes to vectorized quantifiers, the *arity hierarchy* has just two levels.

To get *interesting* classes of vectorized quantifiers beyond the unary, we consider *proper subclasses* of $\overline{Q_2}$.

One way to get interesting classes is to *strengthen* the requirement of *isomorphism invariance*.

Restricted Classes of Binary Quantifiers

Thus, when it comes to vectorized quantifiers, the *arity hierarchy* has just two levels.

To get *interesting* classes of vectorized quantifiers beyond the unary, we consider *proper subclasses* of $\overline{Q_2}$.

One way to get interesting classes is to *strengthen* the requirement of *isomorphism invariance*.

One such strengthening gives us the *linear algebraic quantifiers*.

Isomorphism Closure

Isomorphism Closure

Fix a vocabulary $\sigma = (R_1, \dots, R_r)$ where all relation symbols are *binary*.

Isomorphism Closure

Fix a vocabulary $\sigma = (R_1, \dots, R_r)$ where all relation symbols are *binary*.

Two σ -structures $\mathbb{A} = (A, R_1^A, \dots, R_r^A)$ and $\mathbb{B} = (B, R_1^B, \dots, R_r^B)$ are *isomorphic* if there is a bijection $\beta : A \rightarrow B$ with $\beta(R_i^A) = R_i^B$, for all $i \in [r]$.

Isomorphism Closure

Fix a vocabulary $\sigma = (R_1, \dots, R_r)$ where all relation symbols are *binary*.

Two σ -structures $\mathbb{A} = (A, R_1^A, \dots, R_r^A)$ and $\mathbb{B} = (B, R_1^B, \dots, R_r^B)$ are *isomorphic* if there is a bijection $\beta : A \rightarrow B$ with $\beta(R_i^A) = R_i^B$, for all $i \in [r]$.

Equivalently, if we fix bijections between A and $\{1, \dots, n\}$ on the one hand and B and $\{1, \dots, n\}$ on the other, then we can view each R_i^A or R_i^B as a $n \times n$ matrix with entries in $\{0, 1\}$.

Isomorphism Closure

Fix a vocabulary $\sigma = (R_1, \dots, R_r)$ where all relation symbols are *binary*.

Two σ -structures $\mathbb{A} = (A, R_1^A, \dots, R_r^A)$ and $\mathbb{B} = (B, R_1^B, \dots, R_r^B)$ are *isomorphic* if there is a bijection $\beta : A \rightarrow B$ with $\beta(R_i^A) = R_i^B$, for all $i \in [r]$.

Equivalently, if we fix bijections between A and $\{1, \dots, n\}$ on the one hand and B and $\{1, \dots, n\}$ on the other, then we can view each R_i^A or R_i^B as a $n \times n$ matrix with entries in $\{0, 1\}$.

An *isomorphism* is then an $n \times n$ *permutation matrix* P such that

$$PR_i^A P^{-1} = R_i^B \quad \text{for all } i.$$

Linear Algebraic Equivalence

Linear Algebraic Equivalence

For a field \mathbb{F} , say that $\mathbb{A} = (A, R_1^A, \dots, R_r^A)$ and $\mathbb{B} = (B, R_1^B, \dots, R_r^B)$ are \mathbb{F} -linear algebraically equivalent if

there is an invertible matrix $I \in GL_n(\mathbb{F})$ such that

$$IR_i^A I^{-1} = R_i^B \quad \text{for all } i.$$

Linear Algebraic Equivalence

For a field \mathbb{F} , say that $\mathbb{A} = (A, R_1^A, \dots, R_r^A)$ and $\mathbb{B} = (B, R_1^B, \dots, R_r^B)$ are \mathbb{F} -linear algebraically equivalent if

there is an invertible matrix $I \in GL_n(\mathbb{F})$ such that

$$IR_i^A I^{-1} = R_i^B \quad \text{for all } i.$$

Since all the R_i are $\{0, 1\}$ -matrices, the existence of such an I only depends on the characteristic of \mathbb{F} .

Linear Algebraic Equivalence

For a field \mathbb{F} , say that $\mathbb{A} = (A, R_1^A, \dots, R_r^A)$ and $\mathbb{B} = (B, R_1^B, \dots, R_r^B)$ are \mathbb{F} -linear algebraically equivalent if

there is an invertible matrix $I \in GL_n(\mathbb{F})$ such that

$$IR_i^A I^{-1} = R_i^B \quad \text{for all } i.$$

Since all the R_i are $\{0, 1\}$ -matrices, the existence of such an I only depends on the *characteristic* of \mathbb{F} .

Write $\mathbb{A} \cong_p \mathbb{B}$ to denote that the two structures are \mathbb{F}_p -linear algebraically equivalent, where

Linear Algebraic Equivalence

For a field \mathbb{F} , say that $\mathbb{A} = (A, R_1^A, \dots, R_r^A)$ and $\mathbb{B} = (B, R_1^B, \dots, R_r^B)$ are \mathbb{F} -linear algebraically equivalent if

there is an invertible matrix $I \in GL_n(\mathbb{F})$ such that

$$IR_i^A I^{-1} = R_i^B \quad \text{for all } i.$$

Since all the R_i are $\{0, 1\}$ -matrices, the existence of such an I only depends on the *characteristic* of \mathbb{F} .

Write $\mathbb{A} \cong_p \mathbb{B}$ to denote that the two structures are \mathbb{F}_p -linear algebraically equivalent, where

$p \in \{0\} \cup \text{Primes}$ and \mathbb{F}_p is the *prime field* of characteristic p .

Module Isomorphism

Module Isomorphism

There is a way to see the \mathbb{F}_p -linear algebraic equivalence of $\mathbb{A} = (A, R_1^A, \dots, R_r^A)$ and $\mathbb{B} = (B, R_1^B, \dots, R_r^B)$ as the *isomorphism* of a pair of *modules* over the *polynomial ring*

$$\mathbb{F}_p[x_1, \dots, x_r].$$

Module Isomorphism

There is a way to see the \mathbb{F}_p -linear algebraic equivalence of $\mathbb{A} = (A, R_1^A, \dots, R_r^A)$ and $\mathbb{B} = (B, R_1^B, \dots, R_r^B)$ as the *isomorphism* of a pair of *modules* over the *polynomial ring*

$$\mathbb{F}_p[x_1, \dots, x_r].$$

This is useful in establishing that the problem of deciding $\mathbb{A} \cong_p \mathbb{B}$ is in polynomial time.

Linear Algebraic Quantifiers

Linear Algebraic Quantifiers

Write L_p for the collection of all quantifiers over vocabularies of *binary* relations which are invariant under \cong_p .

Linear Algebraic Quantifiers

Write L_p for the collection of all quantifiers over vocabularies of *binary* relations which are invariant under \cong_p .

For $\Omega \subseteq \{0\} \cup \text{Primes}$, let

$$L_\Omega = \bigcup_{p \in \Omega} L_p.$$

Rank Quantifiers

Rank Quantifiers

For any $p \in \{0\} \cup \text{Primes}$, and $t \in \omega$, let rk_p^t be the quantifier consisting of structures (A, M) where $M \subseteq A \times A$ and

M seen as a matrix in $\mathbb{F}_p^{A \times A}$ has *rank* at least t .

Rank Quantifiers

For any $p \in \{0\} \cup \text{Primes}$, and $t \in \omega$, let rk_p^t be the quantifier consisting of structures (A, M) where $M \subseteq A \times A$ and

M seen as a matrix in $\mathbb{F}_p^{A \times A}$ has *rank* at least t .

Rk_p is the collection of quantifiers $\{\text{rk}_p^t \mid t \in \omega\}$.

Rank Quantifiers

For any $p \in \{0\} \cup \text{Primes}$, and $t \in \omega$, let rk_p^t be the quantifier consisting of structures (A, M) where $M \subseteq A \times A$ and

M seen as a matrix in $\mathbb{F}_p^{A \times A}$ has rank at least t .

Rk_p is the collection of quantifiers $\{\text{rk}_p^t \mid t \in \omega\}$.

Rk is the collection of quantifiers $\bigcup_p \text{Rk}_p$.

Rank Quantifiers

For any $p \in \{0\} \cup \text{Primes}$, and $t \in \omega$, let rk_p^t be the quantifier consisting of structures (A, M) where $M \subseteq A \times A$ and

M seen as a matrix in $\mathbb{F}_p^{A \times A}$ has *rank* at least t .

Rk_p is the collection of quantifiers $\{\text{rk}_p^t \mid t \in \omega\}$.

Rk is the collection of quantifiers $\bigcup_p \text{Rk}_p$.

$L_{\infty\omega}^\omega(\overline{\text{Rk}})$ subsumes *rank logic*, the extension of fixed-point logic with *rank operators* which has been studied in descriptive complexity as a candidate logic for P .

Linear Algebraic Logic

Linear Algebraic Logic

For any $\Omega \subseteq \{0\} \cup \text{Primes}$, we define the Ω -linear algebraic logics.

Linear Algebraic Logic

For any $\Omega \subseteq \{0\} \cup \text{Primes}$, we define the Ω -linear algebraic logics.

$$\text{LA}^k(\Omega) = L_{\infty\omega}^k(\overline{L_\Omega})$$

$$\text{LA}^\omega(\Omega) = L_{\infty\omega}^\omega(\overline{L_\Omega})$$

Linear Algebraic Logic

For any $\Omega \subseteq \{0\} \cup \text{Primes}$, we define the Ω -linear algebraic logics.

$$\text{LA}^k(\Omega) = L_{\infty\omega}^k(\overline{L_\Omega})$$

$$\text{LA}^\omega(\Omega) = L_{\infty\omega}^\omega(\overline{L_\Omega})$$

Also, write $\equiv^{\text{LA}^k(\Omega)}$ to denote indistinguishability in $\text{LA}^k(\Omega)$.

Linear Algebraic Logic

For any $\Omega \subseteq \{0\} \cup \text{Primes}$, we define the Ω -linear algebraic logics.

$$\text{LA}^k(\Omega) = L_{\infty\omega}^k(\overline{L_\Omega})$$

$$\text{LA}^\omega(\Omega) = L_{\infty\omega}^\omega(\overline{L_\Omega})$$

Also, write $\equiv^{\text{LA}^k(\Omega)}$ to denote indistinguishability in $\text{LA}^k(\Omega)$. That is, it is another name for $\equiv_{L_\Omega}^k$.

Linear Algebraic Logic

For any $\Omega \subseteq \{0\} \cup \text{Primes}$, we define the Ω -linear algebraic logics.

$$\text{LA}^k(\Omega) = L_{\infty\omega}^k(\overline{L_\Omega})$$

$$\text{LA}^\omega(\Omega) = L_{\infty\omega}^\omega(\overline{L_\Omega})$$

Also, write $\equiv^{\text{LA}^k(\Omega)}$ to denote indistinguishability in $\text{LA}^k(\Omega)$. That is, it is another name for $\equiv_{L_\Omega}^k$.

This relation is decidable in *polynomial time* (for fixed k) using the module isomorphism algorithm of **Chistov et al.**

Invertible Map Game

Invertible Map Game

The game is played between *Spoiler* and *Duplicator* on \mathbb{A} and \mathbb{B} .

Invertible Map Game

The game is played between *Spoiler* and *Duplicator* on \mathbb{A} and \mathbb{B} .
We have (as usual) k pebbles each on elements of \mathbb{A} and \mathbb{B} .

Invertible Map Game

The game is played between *Spoiler* and *Duplicator* on \mathbb{A} and \mathbb{B} .
We have (as usual) k pebbles each on elements of \mathbb{A} and \mathbb{B} .

Play proceeds in the following steps:

Invertible Map Game

The game is played between *Spoiler* and *Duplicator* on \mathbb{A} and \mathbb{B} . We have (as usual) k pebbles each on elements of \mathbb{A} and \mathbb{B} .

Play proceeds in the following steps:

1. *Spoiler* announces $p_1, \dots, p_{2m} \in [k]$ to move.

Invertible Map Game

The game is played between *Spoiler* and *Duplicator* on \mathbb{A} and \mathbb{B} . We have (as usual) k pebbles each on elements of \mathbb{A} and \mathbb{B} .

Play proceeds in the following steps:

1. *Spoiler* announces $p_1, \dots, p_{2m} \in [k]$ to move.
2. *Spoiler* chooses a *characteristic* p .

Invertible Map Game

The game is played between *Spoiler* and *Duplicator* on \mathbb{A} and \mathbb{B} . We have (as usual) k pebbles each on elements of \mathbb{A} and \mathbb{B} .

Play proceeds in the following steps:

1. *Spoiler* announces $p_1, \dots, p_{2m} \in [k]$ to move.
2. *Spoiler* chooses a *characteristic* p .
3. *Duplicator* gives a *partition* of \mathbb{A}^{2m} into parts P_1, \dots, P_t and of \mathbb{B}^{2m} into parts Q_1, \dots, Q_t .

Invertible Map Game

The game is played between *Spoiler* and *Duplicator* on \mathbb{A} and \mathbb{B} . We have (as usual) k pebbles each on elements of \mathbb{A} and \mathbb{B} .

Play proceeds in the following steps:

1. *Spoiler* announces $p_1, \dots, p_{2m} \in [k]$ to move.
2. *Spoiler* chooses a *characteristic* p .
3. *Duplicator* gives a *partition* of \mathbb{A}^{2m} into parts P_1, \dots, P_t and of \mathbb{B}^{2m} into parts Q_1, \dots, Q_t .

Note: P_i can be thought of as a $\mathbb{A}^m \times \mathbb{A}^m$ 0-1 matrix M_i with $(M_i)_{\bar{a}\bar{b}} = 1$ iff $\bar{a}\bar{b} \in P_i$.

Invertible Map Game

The game is played between *Spoiler* and *Duplicator* on \mathbb{A} and \mathbb{B} . We have (as usual) k pebbles each on elements of \mathbb{A} and \mathbb{B} .

Play proceeds in the following steps:

1. *Spoiler* announces $p_1, \dots, p_{2m} \in [k]$ to move.
2. *Spoiler* chooses a *characteristic* p .
3. *Duplicator* gives a *partition* of \mathbb{A}^{2m} into parts P_1, \dots, P_t and of \mathbb{B}^{2m} into parts Q_1, \dots, Q_t .

Note: P_i can be thought of as a $\mathbb{A}^m \times \mathbb{A}^m$ 0-1 matrix M_i with $(M_i)_{\bar{a}\bar{b}} = 1$ iff $\bar{a}\bar{b} \in P_i$. Similarly, Q_i is a $\mathbb{B}^m \times \mathbb{B}^m$ matrix N_i .

Invertible Map Game

The game is played between *Spoiler* and *Duplicator* on \mathbb{A} and \mathbb{B} . We have (as usual) k pebbles each on elements of \mathbb{A} and \mathbb{B} .

Play proceeds in the following steps:

1. *Spoiler* announces $p_1, \dots, p_{2m} \in [k]$ to move.
2. *Spoiler* chooses a *characteristic* p .
3. *Duplicator* gives a *partition* of \mathbb{A}^{2m} into parts P_1, \dots, P_t and of \mathbb{B}^{2m} into parts Q_1, \dots, Q_t .

Note: P_i can be thought of as a $\mathbb{A}^m \times \mathbb{A}^m$ 0-1 matrix M_i with $(M_i)_{\bar{a}\bar{b}} = 1$ iff $\bar{a}\bar{b} \in P_i$. Similarly, Q_i is a $\mathbb{B}^m \times \mathbb{B}^m$ matrix N_i .

The partitions must satisfy the condition that there is an *invertible* $I \in \mathbb{F}_p^{\mathbb{B}^m \times \mathbb{A}^m}$ such that $M_i = I^{-1}N_iI$ for all i .

Invertible Map Game

The game is played between *Spoiler* and *Duplicator* on \mathbb{A} and \mathbb{B} . We have (as usual) k pebbles each on elements of \mathbb{A} and \mathbb{B} .

Play proceeds in the following steps:

1. *Spoiler* announces $p_1, \dots, p_{2m} \in [k]$ to move.
2. *Spoiler* chooses a *characteristic* p .
3. *Duplicator* gives a *partition* of \mathbb{A}^{2m} into parts P_1, \dots, P_t and of \mathbb{B}^{2m} into parts Q_1, \dots, Q_t .

Note: P_i can be thought of as a $\mathbb{A}^m \times \mathbb{A}^m$ 0-1 matrix M_i with $(M_i)_{\bar{a}\bar{b}} = 1$ iff $\bar{a}\bar{b} \in P_i$. Similarly, Q_i is a $\mathbb{B}^m \times \mathbb{B}^m$ matrix N_i .

The partitions must satisfy the condition that there is an *invertible* $I \in \mathbb{F}_p^{\mathbb{B}^m \times \mathbb{A}^m}$ such that $M_i = I^{-1}N_iI$ for all i .

4. *Spoiler* chooses some $i \in \{1, \dots, t\}$ and an $\bar{a} \in P_i$ and $\bar{b} \in Q_i$ on which the $2m$ pebbles are placed.

Characteristic Zero

Characteristic Zero

Theorem (Holm; D. Vagnozzi)

$$LA^\omega(\{0\}) \leq L^\omega_{\infty\omega}(C).$$

Characteristic Zero

Theorem (Holm; D. Vagnozzi)

$$\text{LA}^\omega(\{0\}) \leq L_{\infty\omega}^\omega(C).$$

Linear algebra over fields of *characteristic zero* can be *simulated by counting*.

Characteristic Zero

Theorem (Holm; D. Vagnozzi)

$$\text{LA}^\omega(\{0\}) \leq L_{\infty\omega}^\omega(C).$$

Linear algebra over fields of *characteristic zero* can be *simulated by counting*.

This essentially follows from the following observation.

Characteristic Zero

Theorem (Holm; D. Vagnozzi)

$$\text{LA}^\omega(\{0\}) \leq L_{\infty\omega}^\omega(C).$$

Linear algebra over fields of *characteristic zero* can be *simulated by counting*.

This essentially follows from the following observation.

For any vocabulary σ of binary relations, and two σ -structures \mathbb{A} and \mathbb{B} ,

$$\mathbb{A} \equiv_C^3 \mathbb{B} \quad \Rightarrow \quad \mathbb{A} \cong_0 \mathbb{B}.$$

Characteristic Zero

Theorem (Holm; D. Vagnozzi)

$$\text{LA}^\omega(\{0\}) \leq L_{\infty\omega}^\omega(C).$$

Linear algebra over fields of *characteristic zero* can be *simulated by counting*.

This essentially follows from the following observation.

For any vocabulary σ of binary relations, and two σ -structures \mathbb{A} and \mathbb{B} ,

$$\mathbb{A} \equiv_C^3 \mathbb{B} \quad \Rightarrow \quad \mathbb{A} \cong_0 \mathbb{B}.$$

\equiv_C^3 can be characterized in terms of *coherent algebras*, and isomorphism of such algebras is witnessed by invertible matrices.

Charecteristic Two

Characteristic Two

Theorem (D., Grohe, Holm, Laubner 2009)

$$L^3_{\omega\omega}(L_2) \not\preceq L^{\omega}_{\infty\omega}(C)$$

Characteristic Two

Theorem (D., Grohe, Holm, Laubner 2009)

$$L^3_{\omega\omega}(L_2) \not\leq L^{\omega}_{\infty\omega}(C)$$

Cai, Fürer and Immerman give a construction of pairs of graphs $G_k, H_k (k \in \omega)$ such that

Characteristic Two

Theorem (D., Grohe, Holm, Laubner 2009)

$$L^3_{\omega\omega}(L_2) \not\leq L^{\omega}_{\infty\omega}(C)$$

Cai, Fürer and Immerman give a construction of pairs of graphs $G_k, H_k (k \in \omega)$ such that

- $G_k \equiv_C^k H_k$; and

Characteristic Two

Theorem (D., Grohe, Holm, Laubner 2009)

$$L_{\omega\omega}^3(L_2) \not\leq L_{\infty\omega}^\omega(C)$$

Cai, Fürer and Immerman give a construction of pairs of graphs $G_k, H_k (k \in \omega)$ such that

- $G_k \equiv_C^k H_k$; and
- $G_k \not\equiv H_k$.

Characteristic Two

Theorem (D., Grohe, Holm, Laubner 2009)

$$L_{\omega\omega}^3(L_2) \not\leq L_{\infty\omega}^\omega(C)$$

Cai, Fürer and Immerman give a construction of pairs of graphs $G_k, H_k (k \in \omega)$ such that

- $G_k \equiv_C^k H_k$; and
- $G_k \not\equiv H_k$.

We can show that there is a single formula φ of $L_{\omega\omega}^3(L_2)$ (indeed of $L_{\omega\omega}^3(\text{Rk}_2)$) such that

$$G_k \models \varphi; \quad H_k \not\models \varphi \quad \text{for all } k.$$

Distinct Characteristics

Distinct Characteristics

Theorem (D., Holm 2012)

For $p, q \in \text{Primes}$ with $p \neq q$,

$$L_{\omega\omega}^{\omega}(L_p) \not\leq L_{\infty\omega}^{\omega}(L_q).$$

Distinct Characteristics

Theorem (D., Holm 2012)

For $p, q \in \text{Primes}$ with $p \neq q$,

$$L_{\omega\omega}^\omega(L_p) \not\preceq L_{\infty\omega}^\omega(L_q).$$

For any prime p , we can construct a class of structures $\text{CFI}(p)$ which codes solvable systems of equations over \mathbb{F}_p .

Distinct Characteristics

Theorem (D., Holm 2012)

For $p, q \in \text{Primes}$ with $p \neq q$,

$$L_{\omega\omega}^\omega(L_p) \not\preceq L_{\infty\omega}^\omega(L_q).$$

For any prime p , we can construct a class of structures $\text{CFI}(p)$ which codes solvable systems of equations over \mathbb{F}_p .

We use a simple version of the *invertible map game* to show that this is not expressible in $L_{\infty\omega}^\omega(L_q)$.

Distinct Characteristics

Theorem (D., Holm 2012)

For $p, q \in \text{Primes}$ with $p \neq q$,

$$L_{\omega\omega}^\omega(L_p) \not\leq L_{\infty\omega}^\omega(L_q).$$

For any prime p , we can construct a class of structures $\text{CFI}(p)$ which codes solvable systems of equations over \mathbb{F}_p .

We use a simple version of the *invertible map game* to show that this is not expressible in $L_{\infty\omega}^\omega(L_q)$.

Note: We do not consider vectorizations here.

Rank Logics

Rank Logics

Let $p \in \text{Primes}$ and $P = \text{Primes} \setminus \{p\}$.

Rank Logics

Let $p \in \text{Primes}$ and $P = \text{Primes} \setminus \{p\}$.

Theorem (Grädel, Pakusa 2017)

$$L_{\omega\omega}^{\omega}(\text{Rk}_p) \not\preceq L_{\infty\omega}^{\omega}(\bigcup_{q \in P} \overline{\text{Rk}_q})$$

Rank Logics

Let $p \in \text{Primes}$ and $P = \text{Primes} \setminus \{p\}$.

Theorem (Grädel, Pakusa 2017)

$$L_{\omega\omega}^{\omega}(\text{Rk}_p) \not\preceq L_{\omega\omega}^{\omega}(\bigcup_{q \in P} \overline{\text{Rk}_q})$$

This is proved by showing that the structures in $\text{CFI}(p)$ can be constructed to be *homogeneous* in a way that guarantees that the quantifiers Rk_q , even vectorized, can be defined in $L_{\omega\omega}^{\omega}(C)$.

Vectorizations

Vectorizations

Let $p \in \text{Primes}$ and $P = \text{Primes} \setminus \{p\}$.

Vectorizations

Let $p \in \text{Primes}$ and $P = \text{Primes} \setminus \{p\}$.

Theorem (D. Grädel, Pakusa 2019)

$$\text{LA}^\omega(\{p\}) \not\preceq \text{LA}^\omega(P).$$

Vectorizations

Let $p \in \text{Primes}$ and $P = \text{Primes} \setminus \{p\}$.

Theorem (D. Grädel, Pakusa 2019)

$$\text{LA}^\omega(\{p\}) \not\leq \text{LA}^\omega(P).$$

In short, as long as Ω does not contain *all primes*, $\text{LA}^\omega(\Omega)$ is *not complete*.

Vectorizations

Let $p \in \text{Primes}$ and $P = \text{Primes} \setminus \{p\}$.

Theorem (D. Grädel, Pakusa 2019)

$$\text{LA}^\omega(\{p\}) \not\leq \text{LA}^\omega(P).$$

In short, as long as Ω does not contain *all primes*, $\text{LA}^\omega(\Omega)$ is *not complete*.

This is established by showing that on the structures in $\text{CFI}(p)$, the equivalence relation $\equiv^{\text{LA}^k(P)}$ can itself be defined in $L_{\infty\omega}^\omega(C)$.

Vectorizations

Let $p \in \text{Primes}$ and $P = \text{Primes} \setminus \{p\}$.

Theorem (D. Grädel, Pakusa 2019)

$$\text{LA}^\omega(\{p\}) \not\leq \text{LA}^\omega(P).$$

In short, as long as Ω does not contain *all primes*, $\text{LA}^\omega(\Omega)$ is *not complete*.

This is established by showing that on the structures in $\text{CFI}(p)$, the equivalence relation $\equiv^{\text{LA}^k(P)}$ can itself be defined in $L_{\infty\omega}^\omega(C)$.

This uses the homogeneity of structures in $\text{CFI}(p)$, along with the fact that the automorphism groups of the structures are Abelian p -groups.

Vectorizations

Let $p \in \text{Primes}$ and $P = \text{Primes} \setminus \{p\}$.

Theorem (D. Grädel, Pakusa 2019)

$$\text{LA}^\omega(\{p\}) \not\leq \text{LA}^\omega(P).$$

In short, as long as Ω does not contain *all primes*, $\text{LA}^\omega(\Omega)$ is *not complete*.

This is established by showing that on the structures in $\text{CFI}(p)$, the equivalence relation $\equiv^{\text{LA}^k(P)}$ can itself be defined in $L_{\infty\omega}^\omega(C)$.

This uses the homogeneity of structures in $\text{CFI}(p)$, along with the fact that the automorphism groups of the structures are Abelian p -groups. This enables us to represent them as *semisimple* \mathbb{F}_q -algebras and apply *Maschke's theorem*.

Rank Logic Again

Rank Logic Again

Theorem (Lichter 2021)

There is a polynomial-time decidable property that is not definable in $L_{\infty\omega}^{\omega}(\overline{\text{Rk}})$.

Rank Logic Again

Theorem (Lichter 2021)

There is a polynomial-time decidable property that is not definable in $L_{\infty\omega}^\omega(\overline{\text{Rk}})$.

The construction is a CFI-like collection of structures encoding systems of linear equations over the *ring* $\mathbb{Z}/\mathbb{Z}_{2^m}$ for growing values of m .

Rank Logic Again

Theorem (Lichter 2021)

There is a polynomial-time decidable property that is not definable in $L_{\infty\omega}^\omega(\overline{\text{Rk}})$.

The construction is a CFI-like collection of structures encoding systems of linear equations over the *ring* $\mathbb{Z}/\mathbb{Z}_{2^m}$ for growing values of m .

The proof uses the **Grädel-Pakusa** argument to show that the quantifiers Rk_p for $p \neq 2$ are useless on these structures.

Rank Logic Again

Theorem (Lichter 2021)

There is a polynomial-time decidable property that is not definable in $L_{\infty\omega}^\omega(\overline{\text{Rk}})$.

The construction is a CFI-like collection of structures encoding systems of linear equations over the *ring* $\mathbb{Z}/\mathbb{Z}_{2^m}$ for growing values of m .

The proof uses the **Grädel-Pakusa** argument to show that the quantifiers Rk_p for $p \neq 2$ are useless on these structures.

It then uses the *invertible map game* to show that $\text{LA}^\omega(\{2\})$ does not distinguish them.

All Characteristics

All Characteristics

Theorem (D., Grädel, Lichter 2022)

Taking Ω to be the set of all characteristics,

There is a polynomial-time decidable property that is not definable in $\text{LA}^\omega(\Omega)$.

All Characteristics

Theorem (D., Grädel, Lichter 2022)

Taking Ω to be the set of all characteristics,

There is a polynomial-time decidable property that is not definable in $\text{LA}^\omega(\Omega)$.

The proof combines the construction of (Lichter 2021) with the algebraic machinery of (D., Grädel, Pakusa 2019).

All Characteristics

Theorem (D., Grädel, Lichter 2022)

Taking Ω to be the set of all characteristics,

There is a polynomial-time decidable property that is not definable in $\text{LA}^\omega(\Omega)$.

The proof combines the construction of (Lichter 2021) with the algebraic machinery of (D., Grädel, Pakusa 2019).

In particular, this shows that for each the expressive power of LA^ω is not complete, and

All Characteristics

Theorem (D., Grädel, Lichter 2022)

Taking Ω to be the set of all characteristics,

There is a polynomial-time decidable property that is not definable in $\text{LA}^\omega(\Omega)$.

The proof combines the construction of (Lichter 2021) with the algebraic machinery of (D., Grädel, Pakusa 2019).

In particular, this shows that for each the expressive power of LA^ω is not complete, and for each k , the equivalence relation \equiv^{LA^k} is not *discrete*.

Conclusions

Conclusions

Linear Algebraic quantifiers are a *natural* class of generalized quantifiers obtained by replacing *isomorphism invariance* by a stronger condition.

Conclusions

Linear Algebraic quantifiers are a *natural* class of generalized quantifiers obtained by replacing *isomorphism invariance* by a stronger condition.

They extend the expressive power of counting quantifiers, but still have nice *algorithmic* properties, like polynomial-time decidable equivalence.

Conclusions

Linear Algebraic quantifiers are a *natural* class of generalized quantifiers obtained by replacing *isomorphism invariance* by a stronger condition.

They extend the expressive power of counting quantifiers, but still have nice *algorithmic* properties, like polynomial-time decidable equivalence.

We have developed sophisticated algebraic machinery for analysing their expressive power, and show it is not complete.