

Window 100/308: samples 990-1009, centre 1000 | target bins: 0, corrected: 0 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 101/308: samples 1000-1019, centre 1010 | target bins: 0, corrected: 0 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 102/308: samples 1010-1029, centre 1020 | target bins: 0, corrected: 0 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 103/308: samples 1020-1039, centre 1030 | target bins: 0, corrected: 0 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 104/308: samples 1030-1049, centre 1040 | target bins: 128, corrected: 52 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 105/308: samples 1040-1059, centre 1050 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 106/308: samples 1050-1069, centre 1060 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 107/308: samples 1060-1079, centre 1070 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 108/308: samples 1070-1089, centre 1080 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 109/308: samples 1080-1099, centre 1090 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 110/308: samples 1090-1109, centre 1100 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 111/308: samples 1100-1119, centre 1110 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 112/308: samples 1110-1129, centre 1120 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 113/308: samples 1120-1139, centre 1130 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 114/308: samples 1130-1149, centre 1140 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 115/308: samples 1140-1159, centre 1150 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 116/308: samples 1150-1169, centre 1160 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 117/308: samples 1160-1179, centre 1170 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 118/308: samples 1170-1189, centre 1180 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 119/308: samples 1180-1199, centre 1190 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 120/308: samples 1190-1209, centre 1200 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 121/308: samples 1200-1219, centre 1210 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 122/308: samples 1210-1229, centre 1220 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 123/308: samples 1220-1239, centre 1230 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 124/308: samples 1230-1249, centre 1240 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 125/308: samples 1240-1259, centre 1250 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 126/308: samples 1250-1269, centre 1260 | target bins: 36, corrected: 36 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 127/308: samples 1260-1279, centre 1270 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 128/308: samples 1270-1289, centre 1280 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 129/308: samples 1280-1299, centre 1290 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 130/308: samples 1290-1309, centre 1300 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 131/308: samples 1300-1319, centre 1310 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 132/308: samples 1310-1329, centre 1320 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 133/308: samples 1320-1339, centre 1330 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 134/308: samples 1330-1349, centre 1340 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 135/308: samples 1340-1359, centre 1350 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 136/308: samples 1350-1369, centre 1360 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 137/308: samples 1360-1379, centre 1370 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 138/308: samples 1370-1389, centre 1380 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 139/308: samples 1380-1399, centre 1390 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 140/308: samples 1390-1409, centre 1400 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 141/308: samples 1400-1419, centre 1410 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 142/308: samples 1410-1429, centre 1420 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 143/308: samples 1420-1439, centre 1430 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 144/308: samples 1430-1449, centre 1440 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 145/308: samples 1440-1459, centre 1450 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 146/308: samples 1450-1469, centre 1460 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 147/308: samples 1460-1479, centre 1470 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 148/308: samples 1470-1489, centre 1480 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 149/308: samples 1480-1499, centre 1490 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 150/308: samples 1490-1509, centre 1500 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 151/308: samples 1500-1519, centre 1510 | target bins: 36, corrected: 36 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 152/308: samples 1510-1529, centre 1520 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 153/308: samples 1520-1539, centre 1530 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 154/308: samples 1530-1549, centre 1540 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 155/308: samples 1540-1559, centre 1550 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 156/308: samples 1550-1569, centre 1560 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 157/308: samples 1560-1579, centre 1570 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 158/308: samples 1570-1589, centre 1580 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 159/308: samples 1580-1599, centre 1590 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 160/308: samples 1590-1609, centre 1600 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 161/308: samples 1600-1619, centre 1610 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 162/308: samples 1610-1629, centre 1620 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 163/308: samples 1620-1639, centre 1630 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 164/308: samples 1630-1649, centre 1640 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 165/308: samples 1640-1659, centre 1650 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 166/308: samples 1650-1669, centre 1660 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 167/308: samples 1660-1679, centre 1670 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 168/308: samples 1670-1689, centre 1680 | target bins: 36, corrected: 36 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 169/308: samples 1680-1699, centre 1690 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 170/308: samples 1690-1709, centre 1700 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 171/308: samples 1700-1719, centre 1710 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 172/308: samples 1710-1729, centre 1720 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 173/308: samples 1720-1739, centre 1730 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 174/308: samples 1730-1749, centre 1740 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 175/308: samples 1740-1759, centre 1750 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 176/308: samples 1750-1769, centre 1760 | target bins: 36, corrected: 36 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 177/308: samples 1760-1779, centre 1770 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 178/308: samples 1770-1789, centre 1780 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 179/308: samples 1780-1799, centre 1790 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 180/308: samples 1790-1809, centre 1800 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 181/308: samples 1800-1819, centre 1810 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 182/308: samples 1810-1829, centre 1820 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 183/308: samples 1820-1839, centre 1830 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 184/308: samples 1830-1849, centre 1840 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 185/308: samples 1840-1859, centre 1850 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 186/308: samples 1850-1869, centre 1860 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 187/308: samples 1860-1879, centre 1870 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 188/308: samples 1870-1889, centre 1880 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 189/308: samples 1880-1899, centre 1890 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 190/308: samples 1890-1909, centre 1900 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 191/308: samples 1900-1919, centre 1910 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 192/308: samples 1910-1929, centre 1920 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 193/308: samples 1920-1939, centre 1930 | target bins: 36, corrected: 36 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 194/308: samples 1930-1949, centre 1940 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 195/308: samples 1940-1959, centre 1950 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 196/308: samples 1950-1969, centre 1960 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 197/308: samples 1960-1979, centre 1970 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 198/308: samples 1970-1989, centre 1980 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 199/308: samples 1980-1999, centre 1990 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 200/308: samples 1990-2009, centre 2000 | target bins: 37, corrected: 37 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 201/308: samples 2000-2019, centre 2010 | target bins: 36, corrected: 36 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 202/308: samples 2010-2029, centre 2020 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 203/308: samples 2020-2039, centre 2030 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 204/308: samples 2030-2049, centre 2040 | target bins: 38, corrected: 38 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 205/308: samples 2040-2059, centre 2050 | target bins: 46, corrected: 44 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 206/308: samples 2050-2069, centre 2060 | target bins: 128, corrected: 52 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 207/308: samples 2060-2079, centre 2070 | target bins: 0, corrected: 0 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 208/308: samples 2070-2089, centre 2080 | target bins: 0, corrected: 0 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$




4) rotation that happened on the target bins (dotted = 174.4 deg)




Window 209/308: samples 2080-2099, centre 2090 | target bins: 0, corrected: 0 | ch 3, ΔC_0 = -12.89 cm


1) where the target appears in the STFT (dots: > -30 dB)




2) ΔC_0 each bin reads (shaded: map ≠ 0 = target footprint + alias images)



3) phase computed for each bin, ph $= -2\pi/\lambda\,\Delta C_0$



4) rotation that happened on the target bins (dotted = 174.4 deg)



delta_C0 map, ch 3: target at pixel 1547, main lobe +-82 px, alias images at +-405 px

ΔC_0 map [cm]

legacy (hold): constant over the line
new (footprint): only where the target appears

Phase SATA added to every sample of the line (err = ||out - ideal|| / ||ideal||)

arg(out) - arg(in) [deg]

legacy map (err 0.000)
new map (err 0.031)
ideal 174.4 deg

Amplitude

amplitude

|input|
|output|, new map

azimuth sample