band k = 2 (f_k = +250 Hz) | window 98/308, centre 980 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +688 Hz → label +188 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg

|STFT| [dB]

target, corrected
target, not corrected

ΔC_0[posaux] [cm]

ph [rad]

rotation [deg]

look angle θ [deg]

absolute Doppler $\tilde{f}$ in band k [Hz]

cell x = posaux

band k = 2 (f_k = +250 Hz) | window 99/308, centre 990 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +676 Hz → label +176 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg

|STFT| [dB]

target, corrected
target, not corrected

ΔC_0[posaux] [cm]

ph [rad]


rotation [deg]

look angle θ [deg]


absolute Doppler $\tilde{f}$ in band k [Hz]

cell x = posaux


band k = 2 (f_k = +250 Hz) | window 100/308, centre 1000 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +663 Hz → label +163 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 101/308, centre 1010 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +651 Hz → label +151 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 102/308, centre 1020 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +639 Hz → label +139 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 103/308, centre 1030 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +626 Hz → label +126 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 104/308, centre 1040 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +614 Hz → label +114 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 105/308, centre 1050 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +602 Hz → label +102 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 106/308, centre 1060 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +589 Hz → label +89 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 107/308, centre 1070 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +577 Hz → label +77 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 108/308, centre 1080 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +564 Hz → label +64 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 109/308, centre 1090 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +552 Hz → label +52 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 110/308, centre 1100 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +540 Hz → label +40 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 111/308, centre 1110 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +527 Hz → label +27 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 112/308, centre 1120 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +515 Hz → label +15 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 113/308, centre 1130 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +503 Hz → label +3 Hz (orange) | other band: folded, points at $x_t - 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 114/308, centre 1140 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +490 Hz → label +490 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]

target, corrected
target, not corrected

ΔC_0[posaux] [cm]

ph [rad]

rotation [deg]

look angle θ [deg]


absolute Doppler $\tilde{f}$ in band k [Hz]

cell x = posaux


band k = 2 (f_k = +250 Hz) | window 115/308, centre 1150 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +478 Hz → label +478 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 116/308, centre 1160 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +466 Hz → label +466 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 117/308, centre 1170 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +453 Hz → label +453 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 118/308, centre 1180 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +441 Hz → label +441 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 119/308, centre 1190 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +429 Hz → label +429 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 120/308, centre 1200 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +416 Hz → label +416 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 121/308, centre 1210 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +404 Hz → label +404 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 122/308, centre 1220 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +392 Hz → label +392 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 123/308, centre 1230 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +379 Hz → label +379 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 124/308, centre 1240 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +367 Hz → label +367 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 125/308, centre 1250 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +355 Hz → label +355 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 126/308, centre 1260 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +342 Hz → label +342 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 127/308, centre 1270 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +330 Hz → label +330 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 128/308, centre 1280 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +318 Hz → label +318 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 129/308, centre 1290 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +305 Hz → label +305 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 130/308, centre 1300 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +293 Hz → label +293 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 131/308, centre 1310 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +281 Hz → label +281 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 132/308, centre 1320 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +268 Hz → label +268 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 133/308, centre 1330 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +256 Hz → label +256 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 134/308, centre 1340 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +244 Hz → label +244 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 135/308, centre 1350 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +231 Hz → label +231 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 136/308, centre 1360 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +219 Hz → label +219 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 137/308, centre 1370 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +207 Hz → label +207 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 138/308, centre 1380 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +194 Hz → label +194 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 139/308, centre 1390 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +182 Hz → label +182 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 140/308, centre 1400 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +169 Hz → label +169 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 141/308, centre 1410 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +157 Hz → label +157 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 142/308, centre 1420 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +145 Hz → label +145 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 143/308, centre 1430 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +132 Hz → label +132 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 144/308, centre 1440 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +120 Hz → label +120 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 145/308, centre 1450 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +108 Hz → label +108 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 146/308, centre 1460 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +95 Hz → label +95 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 147/308, centre 1470 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +83 Hz → label +83 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 148/308, centre 1480 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +71 Hz → label +71 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 149/308, centre 1490 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +58 Hz → label +58 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 150/308, centre 1500 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +46 Hz → label +46 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 151/308, centre 1510 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +34 Hz → label +34 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]

target, corrected
target, not corrected

ΔC_0[posaux] [cm]

ph [rad]

rotation [deg]

look angle θ [deg]


absolute Doppler $\tilde{f}$ in band k [Hz]

cell x = posaux

band k = 2 (f_k = +250 Hz) | window 152/308, centre 1520 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +21 Hz → label +21 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 153/308, centre 1530 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target +9 Hz → label +9 Hz (orange) | own band: points at x_t | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 154/308, centre 1540 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -3 Hz → label +497 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]

target, corrected
target, not corrected

ΔC_0[posaux] [cm]

ph [rad]

rotation [deg]

look angle θ [deg]

absolute Doppler $\tilde{f}$ in band k [Hz]

cell x = posaux

band k = 2 (f_k = +250 Hz) | window 155/308, centre 1550 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -16 Hz → label +484 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 156/308, centre 1560 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -28 Hz → label +472 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 157/308, centre 1570 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -40 Hz → label +460 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 158/308, centre 1580 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -53 Hz → label +447 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 159/308, centre 1590 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -65 Hz → label +435 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 160/308, centre 1600 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -77 Hz → label +423 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 161/308, centre 1610 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -90 Hz → label +410 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 162/308, centre 1620 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -102 Hz → label +398 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 163/308, centre 1630 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -114 Hz → label +386 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 164/308, centre 1640 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -127 Hz → label +373 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 165/308, centre 1650 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -139 Hz → label +361 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 166/308, centre 1660 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -151 Hz → label +349 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 167/308, centre 1670 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -164 Hz → label +336 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 168/308, centre 1680 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -176 Hz → label +324 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 169/308, centre 1690 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -188 Hz → label +312 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 170/308, centre 1700 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -201 Hz → label +299 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 171/308, centre 1710 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -213 Hz → label +287 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 172/308, centre 1720 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -226 Hz → label +274 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 173/308, centre 1730 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -238 Hz → label +262 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 174/308, centre 1740 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -250 Hz → label +250 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 175/308, centre 1750 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -263 Hz → label +237 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 176/308, centre 1760 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -275 Hz → label +225 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 177/308, centre 1770 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -287 Hz → label +213 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 178/308, centre 1780 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -300 Hz → label +200 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 179/308, centre 1790 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -312 Hz → label +188 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 180/308, centre 1800 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -324 Hz → label +176 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 181/308, centre 1810 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -337 Hz → label +163 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 182/308, centre 1820 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -349 Hz → label +151 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 183/308, centre 1830 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -361 Hz → label +139 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 184/308, centre 1840 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -374 Hz → label +126 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 185/308, centre 1850 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -386 Hz → label +114 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 186/308, centre 1860 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -398 Hz → label +102 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 187/308, centre 1870 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -411 Hz → label +89 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 188/308, centre 1880 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -423 Hz → label +77 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]

target, corrected
target, not corrected

ΔC_0[posaux] [cm]

ph [rad]

rotation [deg]

look angle θ [deg]

absolute Doppler $\tilde{f}$ in band k [Hz]

cell x = posaux

band k = 2 (f_k = +250 Hz) | window 189/308, centre 1890 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -435 Hz → label +65 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]

target, corrected
target, not corrected

ΔC_0[posaux] [cm]

ph [rad]

rotation [deg]

look angle θ [deg]

absolute Doppler $\tilde{f}$ in band k [Hz]

cell x = posaux

band k = 2 (f_k = +250 Hz) | window 190/308, centre 1900 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -448 Hz → label +52 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 191/308, centre 1910 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -460 Hz → label +40 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 192/308, centre 1920 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -472 Hz → label +28 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux

band k = 2 (f_k = +250 Hz) | window 193/308, centre 1930 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -485 Hz → label +15 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 194/308, centre 1940 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -497 Hz → label +3 Hz (orange) | other band: folded, points at $x_t + 1X$ | ideal rotation -78.8 deg

|STFT| [dB]

target, corrected
target, not corrected

ΔC_0[posaux] [cm]

ph [rad]


rotation [deg]

look angle θ [deg]


absolute Doppler $\tilde{f}$ in band k [Hz]

cell x = posaux


band k = 2 (f_k = +250 Hz) | window 195/308, centre 1950 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -509 Hz → label +491 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 196/308, centre 1960 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -522 Hz → label +478 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 197/308, centre 1970 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -534 Hz → label +466 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 198/308, centre 1980 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -546 Hz → label +454 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 199/308, centre 1990 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -559 Hz → label +441 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 200/308, centre 2000 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -571 Hz → label +429 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg

|STFT| [dB]
target, corrected
target, not corrected
ΔC_0[posaux] [cm]
ph [rad]
rotation [deg]
look angle θ [deg]
absolute Doppler $\tilde{f}$ in band k [Hz]
cell x = posaux


band k = 2 (f_k = +250 Hz) | window 201/308, centre 2010 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -583 Hz → label +417 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 202/308, centre 2020 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -596 Hz → label +404 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 203/308, centre 2030 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -608 Hz → label +392 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg




band k = 2 (f_k = +250 Hz) | window 204/308, centre 2040 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -621 Hz → label +379 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 205/308, centre 2050 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -633 Hz → label +367 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 206/308, centre 2060 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -645 Hz → label +355 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 207/308, centre 2070 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -658 Hz → label +342 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg

|STFT| [dB]

target, corrected
target, not corrected

ΔC_0[posaux] [cm]

ph [rad]


rotation [deg]

look angle θ [deg]


absolute Doppler $\tilde{f}$ in band k [Hz]

cell x = posaux

band k = 2 (f_k = +250 Hz) | window 208/308, centre 2080 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -670 Hz → label +330 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg



band k = 2 (f_k = +250 Hz) | window 209/308, centre 2090 | channel 0, ΔC_0 = -5.47 cm
true Doppler of the target -682 Hz → label +318 Hz (orange) | other band: folded, points at $x_t + 2X$ | ideal rotation -78.8 deg



band k = 2: the f_a map, window by window (peak of each window)

Doppler [Hz]

true Doppler
peak label, own band
peak label, folded


$\theta(\tilde{f})$ of the peak [deg]

cell the peak points to

own band
folded

rotation on the peak [deg]

window centre [cell]


which part of the illumination belongs to band k (own band points at x_t, the rest at $x_t + mX$)
Doppler [Hz]
600
400
200
0
−200
−400
−600
true Doppler of the target
band k = 2
delta_C0 map of band 2 (direct block and fold images)
ΔC_0 map [cm]
0
−1
−2
−3
−4
−5
arg(out) - arg(in) [deg]
150
100
50
0
−50
−100
−150
SATA band 2 output (error vs ideal 0.020)
ideal -78.8 deg
0
500
1000
1500
2000
2500
3000
azimuth sample