Parametrizations of curves



Def: $\vec{r}:[a,b]\longrightarrow\mathbb{R}^2$ is a parametrization of C if

$$\vec{r}(t)=(x(t),y(t))$$

is in C for all $t\in[a,b]$, and traverses C once as t increases from a to b.

Examples: 1) Half circle



$$\left.\begin{array}{l} x=\cos t \\ y=\sin t \end{array}\right\} \quad \vec{r}(t)=\langle\cos t,\ \sin t\rangle \quad (0\leq t\leq\pi)$$

2) Ellipse

$$\frac{1}{4}(x-5)^2+(y-3)^2=1$$



$$\begin{cases} x=5+2\cos t \\ y=3+\sin t \end{cases} \quad 0\leq t\leq 2\pi$$

---

3) Cusp.

$y^2=x^3$



$C=C_1\cup(-C_2)$

$$C_1:\begin{cases} x=x \\ y=+\sqrt{x^3} \end{cases} \quad (0\leq x\leq 1) \qquad C_2:\begin{cases} x=x \\ y=-\sqrt{x^3} \end{cases} \quad (0\leq x\leq 1)$$

3') A better parametrization:



$$\begin{cases} x=t^2 \\ y=t^3 \end{cases} \quad (-1\leq t\leq 1)$$

Line integrals (16.2)

↳ of functions

↳ of vector fields (work)



Parametrization:

$\vec{r}(t) = (x(t), y(t))$, $a \leq t \leq b$.

$$\text{length}(C) = \lim_{\Delta s_i \to 0} \sum_i \Delta s_i$$

$$= \int_a^b \underbrace{\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}\, dt}_{ds \text{ "arc length element"}}$$

Why:

$$\Delta s_i = \sqrt{(\Delta x_i)^2 + (\Delta y_i)^2}$$

$$= \sqrt{\left(\frac{\Delta x_i}{\Delta t}\right)^2 + \left(\frac{\Delta y_i}{\Delta t}\right)^2}\, \Delta t$$

Definition:

$$\boxed{\begin{aligned} \int_C f(x,y)\, ds &= \lim_{\Delta s_i \to 0} \sum_i f(x_i, y_i)\, \Delta s_i \\ &= \int_a^b f(x(t), y(t)) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}\, dt \\ &= \int_a^b f(\vec{r}(t))\, |\vec{r}\,'(t)|\, dt \end{aligned}}$$

Example: $f(x,y) = 2 + x^2 y$



$x = \cos(t)$

$y = \sin(t)$

$0 \leq t \leq \pi$

$$\int_C f(x,y)\, ds =$$

$$= \int_0^\pi (2 + \cos^2 t \sin t) \underbrace{\sqrt{(\sin t)^2 + (-\cos t)^2}}_{1}\, dt$$

$$= 2t - \frac{\cos^3 t}{3} \Big|_0^\pi = 2\pi + \frac{2}{3}$$

Applications of $\int_C f\,ds$:

1) $\int_C ds$ = length of $C =: L$

2) Average value of f on C $= \frac{1}{L}\int_C f\,ds$

3) $\rho(x,y)$ = density of C (mass / unit length) at (x,y)

$$M = \int_C \rho\,ds$$

$$\bar{x} = \frac{1}{M}\int_C x\rho\,ds$$

$$I_y = \frac{1}{M}\int_C (x^2+z^2)\rho\,ds$$

4) In 2D: area of "fence"



Line integrals of vector fields

$\vec{F} = \langle P, Q, R \rangle$



$\vec{r}(t) = (x(t), y(t), z(t))$, $a \le t \le b$

Definition (Work): $\vec{F} = \langle P, Q, R \rangle$

$$\underbrace{\int_C \vec{F}\cdot d\vec{r}}_{\text{"Work"}} := \lim_{\Delta r_i \to 0} \sum_i \vec{F}(x_i,y_i)\cdot \Delta\vec{r}_i$$

$$= \int_a^b F(\vec{r}(t))\cdot \frac{d\vec{r}}{dt}\,dt$$

Example: $\vec{F} = \langle y, z, x \rangle$



$$\vec{r}(t) = (2,0,0) + t\big((3,4,5)-(2,0,0)\big)$$
$$= (2+t,\ 4t,\ 5t)$$
$$0 \le t \le 1$$

$$\int_C \overbrace{\vec{F}}^{\langle y,z,x\rangle} \cdot \overbrace{d\vec{r}}^{\langle dx,dy,dz\rangle} =$$

$$= \int_0^1 \langle 4t, 5t, 2+t\rangle \cdot \langle 1,4,5\rangle \, dt$$

$$= \int_0^1 4t + 20t + (10+5t) \, dt$$

$$= \cdots = \frac{49}{2} \quad //$$

Another way to write the same thing:

$$\int_C \overbrace{\vec{F}}^{\langle P,Q\rangle} \cdot \overbrace{d\vec{r}}^{\langle dx,dy\rangle} =$$

$$= \int_a^b \langle P, Q\rangle \cdot \overbrace{\left\langle \frac{dx}{dt}, \frac{dy}{dt} \right\rangle}^{\frac{d\vec{r}}{dt}} dt$$

$$= \int_a^b \left(P \frac{dx}{dt} + Q \frac{dy}{dt} \right) dt$$

$$=: \int_C P\,dx + Q\,dy$$

Example:

C = twisted cubic curve

$$\begin{cases} x = t \quad y = t^2 \quad z = t^3 \\ 0 \le t \le 1 \end{cases}$$



$$\vec{F} = \langle \overbrace{xy}^{P}, \overbrace{yz}^{Q}, \overbrace{zx}^{R} \rangle$$

$$\int_C \vec{F} \cdot d\vec{r} =$$

$$= \int_C xy\,dx + yz\,dy + zx\,dz$$

$$= \int_0^1 t^3 \cdot dt + t^5 \cdot 2t\,dt + t^4 \cdot 3t^2\,dt$$

$$= \int_0^1 t^3 + 5t^6 \, dt$$

$$= \frac{t^4}{4} + 5\frac{t^7}{7} \Big|_0^1$$

$$= \frac{1}{4} + \frac{5}{7} = \frac{7+20}{28} = \frac{27}{28} \quad //$$

Work = Change in kinetic energy

$\vec{r}(t)$ = position of particle at time t.

$\vec{F}(t)$ = net force on particle at time t.

$$= m \cdot \vec{a}$$
$$= m \cdot \vec{r}''$$

$$K = \frac{1}{2} m v^2 \qquad \text{(Kinetic energy)}$$
$$= \frac{1}{2} m \vec{r}' \cdot \vec{r}'$$

$$\frac{dK}{dt} = \frac{1}{2} m \left(\vec{r}'' \cdot \vec{r}' + \vec{r}' \cdot \vec{r}'' \right)$$
$$= m \vec{r}'' \cdot \vec{r}'$$
$$= \vec{F}(t) \cdot \vec{r}'(t)$$

Conclusion:

$$\Delta K = \int_a^b \frac{dK}{dt} \, dt$$
$$= \int_a^b \vec{F} \cdot \frac{d\vec{r}}{dt} \, dt$$
$$= \int_C \vec{F} \cdot d\vec{r} = \text{Work}$$

Orientations:



1) $\int_{-C} \vec{F} \cdot d\vec{r} = -\int_C \vec{F} \cdot d\vec{r}$

2) $\int_{-C} f \, ds = +\int_C f \, ds$