



Sexual Dimorphism




- Sexual dimorphism shows extensive interspecific variation
- The evolution of sexual dimorphism is likely multifactorial, but the role of evolvability remains unclear

Evolvability and Modularity

- Low evolvability may slow the pace of evolutionary change
- High evolvability may promote greater morphological diversification and faster evolutionary change




Hypothesis




Data

- 50 species from two simian clades: Platyrrhini and Catarrhini
- 38 linear cranial measurements collected from simian skulls
- ~ 2.520 simian skulls measured
- P matrices estimated for each species
- Sexual dimorphism was measured as the **norm** of the vector difference between males and females




Analysis





$$\text{evolvability}_{(SD)} = \frac{(SD + \epsilon)' P SD}{|SD + \epsilon|^2}$$
$$\text{integration} = 1 - \frac{1}{SD^T P^{-1} SD}$$

$$\text{conditional evolvability}_{(SD)} = \text{evolvability}_{(SD)} \cdot \text{integration}$$



Measurement Error Origin (Primate Crania)




Results



- No relationship was found between evolvability and morphological integration



- Excluding P_max changes the relationship, making it nonlinear and inverse



Conclusions and Next Steps

- Sexual dimorphism tends to evolve along highly evolvable directions
- Integration alone does not explain variation in evolvability among species
- Measurement error does not substantially alter the observed patterns

- Identify additional predictors of evolvability in the direction of sexual dimorphism.
- Test whether these patterns are phylogenetically widespread
- Examine the consequences of removing Pmax on evolvability estimates.