

# Introduction to Bayesian statistics with R

## 4. Priors

---

Olivier Gimenez

last updated: 2025-03-09


## A detour to explore priors

---


## Influence of the prior

---


**Prior  $Beta(0.5, 0.5)$  and posterior survival  $Beta(19.5, 38.5)$**



## Prior $Beta(2, 2)$ and posterior survival $Beta(21, 40)$



## Prior $Beta(20, 1)$ and posterior survival $Beta(39, 49)$



## The role of the prior

- In biological applications, the prior is a convenient means of incorporating expert opinion or information from previous or related studies that would otherwise need to be ignored. We'll get back to that.

# The role of the prior

- In biological applications, the prior is a convenient means of incorporating expert opinion or information from previous or related studies that would otherwise need to be ignored. We'll get back to that.
- With sparse data, the role of the prior can be to enable inference on key parameters that would otherwise be impossible.



## The role of the prior

- In biological applications, the prior is a convenient means of incorporating expert opinion or information from previous or related studies that would otherwise need to be ignored. We'll get back to that.
- With sparse data, the role of the prior can be to enable inference on key parameters that would otherwise be impossible.
- With sufficiently large and informative datasets the prior typically has little effect on the results.

## The role of the prior

- In biological applications, the prior is a convenient means of incorporating expert opinion or information from previous or related studies that would otherwise need to be ignored. We'll get back to that.
- With sparse data, the role of the prior can be to enable inference on key parameters that would otherwise be impossible.
- With sufficiently large and informative datasets the prior typically has little effect on the results.
- Always perform a sensitivity analysis.

## Informative priors vs. no information

- Informative priors aim to reflect information available to the analyst that is gained independently of the data being studied.

## Informative priors vs. no information

- Informative priors aim to reflect information available to the analyst that is gained independently of the data being studied.
- In the absence of any prior information on one or more model parameters we wish to ensure that this lack of knowledge is properly reflected in the prior.

## Informative priors vs. no information

- Informative priors aim to reflect information available to the analyst that is gained independently of the data being studied.
- In the absence of any prior information on one or more model parameters we wish to ensure that this lack of knowledge is properly reflected in the prior.
- Always perform a sensitivity analysis.



## How to incorporate prior information?

---

## Estimating survival using capture-recapture data

- A bird might be captured, missed and recaptured; this is coded 101.



## Estimating survival using capture-recapture data

- A bird might be captured, missed and recaptured; this is coded 101.
- Simplest model relies on constant survival  $\phi$  and detection  $p$  probabilities.

## Estimating survival using capture-recapture data

- A bird might be captured, missed and recaptured; this is coded 101.
- Simplest model relies on constant survival  $\phi$  and detection  $p$  probabilities.
- Likelihood for that particular bird:

$$\Pr(101) = \phi(1 - p)\phi p$$

## Estimating survival using capture-recapture data

- A bird might be captured, missed and recaptured; this is coded 101.
- Simplest model relies on constant survival  $\phi$  and detection  $p$  probabilities.
- Likelihood for that particular bird:

$$\Pr(101) = \phi(1 - p)\phi p$$

- We assume a vague prior:

$$\phi_{prior} \sim \text{Beta}(1, 1) = \text{Uniform}(0, 1)$$

# Notation

- $y_{i,t} = 1$  if individual  $i$  detected at occasion  $t$  and 0 otherwise
- $z_{i,t} = 1$  if individual  $i$  alive between occasions  $t$  and  $t + 1$  and 0 otherwise

$$y_{i,t} \mid z_{i,t} \sim \text{Bernoulli}(p \mid z_{i,t}) \quad [\text{likelihood (observation eq.)}]$$

$$z_{i,t+1} \mid z_{i,t} \sim \text{Bernoulli}(\phi \mid z_{i,t}) \quad [\text{likelihood (state eq.)}]$$

$$\phi \sim \text{Beta}(1, 1) \quad [\text{prior for } \phi]$$

$$p \sim \text{Beta}(1, 1) \quad [\text{prior for } p]$$

## European dippers in Eastern France (1981-1987)



## How to incorporate prior information?

- If no information, mean posterior survival is  $\phi_{posterior} = 0.56$  with credible interval  $[0.51, 0.61]$ .

## How to incorporate prior information?

- If no information, mean posterior survival is  $\phi_{posterior} = 0.56$  with credible interval  $[0.51, 0.61]$ .
- Using information on body mass and annual survival of 27 European passerines, we can predict survival of European dippers using only body mass.

## How to incorporate prior information?

- If no information, mean posterior survival is  $\phi_{posterior} = 0.56$  with credible interval  $[0.51, 0.61]$ .
- Using information on body mass and annual survival of 27 European passerines, we can predict survival of European dippers using only body mass.
- For dippers, body mass is 59.8g, therefore  $\phi = 0.57$  with  $sd = 0.073$ .



## How to incorporate prior information?

- If no information, mean posterior survival is  $\phi_{posterior} = 0.56$  with credible interval  $[0.51, 0.61]$ .
- Using information on body mass and annual survival of 27 European passerines, we can predict survival of European dippers using only body mass.
- For dippers, body mass is 59.8g, therefore  $\phi = 0.57$  with  $sd = 0.073$ .
- Assuming an informative prior  $\phi_{prior} \sim \text{Normal}(0.57, 0.073^2)$ .

## How to incorporate prior information?

- If no information, mean posterior survival is  $\phi_{posterior} = 0.56$  with credible interval  $[0.51, 0.61]$ .
- Using information on body mass and annual survival of 27 European passerines, we can predict survival of European dippers using only body mass.
- For dippers, body mass is 59.8g, therefore  $\phi = 0.57$  with  $sd = 0.073$ .
- Assuming an informative prior  $\phi_{prior} \sim \text{Normal}(0.57, 0.073^2)$ .
- Mean posterior  $\phi_{posterior} = 0.56$  with credible interval  $[0.52, 0.60]$ .

## How to incorporate prior information?

- If no information, mean posterior survival is  $\phi_{posterior} = 0.56$  with credible interval  $[0.51, 0.61]$ .
- Using information on body mass and annual survival of 27 European passerines, we can predict survival of European dippers using only body mass.
- For dippers, body mass is 59.8g, therefore  $\phi = 0.57$  with  $sd = 0.073$ .
- Assuming an informative prior  $\phi_{prior} \sim \text{Normal}(0.57, 0.073^2)$ .
- Mean posterior  $\phi_{posterior} = 0.56$  with credible interval  $[0.52, 0.60]$ .
- No increase of precision in posterior inference.

## How to incorporate prior information?

- Now if you had only the three first years of data, what would have happened?

## How to incorporate prior information?

- Now if you had only the three first years of data, what would have happened?
- Width of credible interval is 0.47 (vague prior) vs. 0.30 (informative prior).

## How to incorporate prior information?

- Now if you had only the three first years of data, what would have happened?
- Width of credible interval is 0.47 (vague prior) vs. 0.30 (informative prior).
- Huge increase of precision in posterior inference (40% gain)!

## Compare **vague** vs. **informative** prior

## Prior elicitation via moment matching

---



## Remember the Beta distribution

- Recall that the Beta distribution is a continuous distribution with values between 0 and 1. Useful for modelling survival or detection probabilities.

## Remember the Beta distribution

- Recall that the Beta distribution is a continuous distribution with values between 0 and 1. Useful for modelling survival or detection probabilities.
- If  $X \sim \text{Beta}(\alpha, \beta)$ , then the first and second moments of  $X$  are:

$$\mu = E(X) = \frac{\alpha}{\alpha + \beta}$$

$$\sigma^2 = \text{Var}(X) = \frac{\alpha\beta}{(\alpha + \beta)^2(\alpha + \beta + 1)}$$

## Moment matching

- In the capture-recapture example, we know a priori that the mean of the probability we're interested in is  $\mu = 0.57$  and its variance is  $\sigma^2 = 0.073^2$ .

## Moment matching

- In the capture-recapture example, we know a priori that the mean of the probability we're interested in is  $\mu = 0.57$  and its variance is  $\sigma^2 = 0.073^2$ .
- Parameters  $\mu$  and  $\sigma^2$  are seen as the moments of a  $Beta(\alpha, \beta)$  distribution.

## Moment matching

- In the capture-recapture example, we know a priori that the mean of the probability we're interested in is  $\mu = 0.57$  and its variance is  $\sigma^2 = 0.073^2$ .
- Parameters  $\mu$  and  $\sigma^2$  are seen as the moments of a  $Beta(\alpha, \beta)$  distribution.
- Now we look for values of  $\alpha$  and  $\beta$  that match the observed moments of the Beta distribution ( $\mu$  and  $\sigma^2$ ).

## Moment matching

- In the capture-recapture example, we know a priori that the mean of the probability we're interested in is  $\mu = 0.57$  and its variance is  $\sigma^2 = 0.073^2$ .
- Parameters  $\mu$  and  $\sigma^2$  are seen as the moments of a  $Beta(\alpha, \beta)$  distribution.
- Now we look for values of  $\alpha$  and  $\beta$  that match the observed moments of the Beta distribution ( $\mu$  and  $\sigma^2$ ).
- We need another set of equations:

$$\alpha = \left( \frac{1 - \mu}{\sigma^2} - \frac{1}{\mu} \right) \mu^2$$

$$\beta = \alpha \left( \frac{1}{\mu} - 1 \right)$$

- For our model, that means:

```
(alpha <- ( (1 - 0.57)/(0.073*0.073) - (1/0.57) )*0.57^2)
#> [1] 25.64636
(beta <- alpha * ( (1/0.57) - 1))
#> [1] 19.34726
```

- For our model, that means:

```
(alpha <- ( (1 - 0.57)/(0.073*0.073) - (1/0.57) )*0.57^2)
#> [1] 25.64636
(beta <- alpha * ( (1/0.57) - 1))
#> [1] 19.34726
```

- Now use  $\phi_{prior} \sim \text{Beta}(\alpha = 25.6, \beta = 19.3)$  instead of  
 $\phi_{prior} \sim \text{Normal}(0.57, 0.073^2)$



## **Your turn: Practical 3**


---

## Prior predictive checks


---

# Linear regression

Unreasonable prior  $\beta \sim N(0, 1000^2)$




Reasonable prior  $\beta \sim N(2, 0.5^2)$




# Logistic regression

Unreasonable  $\text{logit}(\phi) = \beta \sim N(0, 10^2)$



Reasonable  $\text{logit}(\phi) = \beta \sim N(0, 1.5^2)$



## Your turn: Practical 4

---

## Your turn: Practical 4