Prediction of ozone level in Boston

Lucas Emanuel Resck Domingues Lucas Machado Moscheb*

Predicting O3 in Boston

Load and visualize



Data treatment

We noticed that some days do not exist in the dataset, for example, the day August 31, 2001 does not have information in the dataset.

```
##       X   City         State Site.Num Date.Local  O3.Mean
## 148 148 Boston Massachusetts       42 2001-08-28 0.024583
## 149 149 Boston Massachusetts       42 2001-08-29 0.015000
## 150 150 Boston Massachusetts       42 2001-08-30 0.022333
## 151 151 Boston Massachusetts       42 2001-09-01 0.021958
## 152 152 Boston Massachusetts       42 2001-09-02 0.018750
## 153 153 Boston Massachusetts       42 2001-09-03 0.028708
```

Also, there is duplicated days, as June 9, 2002:

*Escola de Matemática Aplicada

```
##       X   City         State Site.Num Date.Local  O3.Mean
## 412 412 Boston Massachusetts       42 2002-06-08 0.022917
## 413 413 Boston Massachusetts       42 2002-06-09 0.036190
## 414 414 Boston Massachusetts       42 2002-06-09 0.037000
## 415 415 Boston Massachusetts       42 2002-06-10 0.023389
```

The duplicated one is easier to deal, but the nan values are harder. First we calculate the mean value between the duplicated.

The rate of NA values is almost 5% of the dataset.

```
## [1] 0.04453367
```

So as to solve that problem, we make a knn imputation using the month ($k = 30$)

```
o3.clean <- knn.impute(as.matrix(o3.ts), k = 30)
o3.clean <- as.ts(o3.clean)
```

Models

Now we develop some models using the train data.

The metric to compare is the Mean Absolute Error (MAE) in the predictions:

```
mae <- function(error)
{
    mean(abs(error))
}
```

Decompose

Regression

Holt-Winters

ARMA