

## Exam 2 (A2)

Class: Bayesian Statistics  
Instructor: Luiz Max Carvalho

02/06/2021

**Turn in date: until 09/06/2021 at 23:59h Brasilia Time.**

- Please read through the whole exam before starting to answer;
- State and prove all non-trivial mathematical results necessary to substantiate your arguments;
- Do not forget to add appropriate scholarly references *at the end* of the document;
- Mathematical expressions also receive punctuation;
- You can write your answer to a question as a point-by-point response or in “essay” form, your call;
- Please hand in a single, **typeset** (  $\LaTeX$ ) PDF file as your final main document. Code appendices are welcome, *in addition* to the main PDF document.
- You may consult any sources, provided you cite **ALL** of your sources (books, papers, blog posts, videos);
- You may use symbolic algebra programs such as Sympy or Wolfram Alpha to help you get through the hairier calculations, provided you cite the tools you have used.
- The exam is worth 100 marks.

## Background

This exam covers applications, namely estimation, prior sensitivity and prediction. You will need a working knowledge of basic computing tools, and knowledge of MCMC is highly valuable. Chapter 6 in [Robert \(2007\)](#) gives an overview of computational techniques for Bayesian statistics.

## Inferring population sizes – theory

Consider the model

$$x_i \sim \text{Binomial}(N, \theta),$$

with **both**  $N$  and  $\theta$  unknown and suppose one observes  $\mathbf{x} = \{x_1, x_2, \dots, x_K\}$ . Here, we will write  $\xi = (N, \theta)$ .

- a) (10 marks) Formulate a hierarchical prior  $(\pi_1)$  for  $N$ , i.e., elicit  $F$  such that  $N \mid \alpha \sim F(\alpha)$  and  $\alpha \sim \Pi_A$ . Justify your choice;

**Solution.** The only restriction prior to the data we have is that  $N \in \mathbb{Z}^+$ . Therefore we are looking for a distribution in this space. We can follow the assumption made by [Raftery \(1988\)](#) where  $N \sim \text{Poisson}(\mu)$ . Other approach I will use is to consider  $N \sim \text{Geometric}(\nu)$ .

The first approach has the advantage in calculations. When

$$x_i \mid N, \theta \sim \text{Binomial}(N, \theta)$$

and

$$N \sim \text{Poisson}(\mu),$$

we have that  $x_i \sim \text{Poisson}(\mu \cdot \theta)$ , as proved in [Appendix A](#). And we can transform prior information about the mean of  $x_i$  in probability distributions for  $\mu \cdot \theta$ .

The second approach serves as a comparative. The geometric distribution is a simple distribution used for discrete survival distribution, but it is interesting because of its simplicity and the compact domain of the parameter since  $\nu \in [0, 1]$ . From that, we can build a model that carries the correlation between  $\nu$  and  $\theta$  whenever necessary in a direct way: we transform the variables to logistic space and defines a bivariate normal distribution.

### Priors to the hyperparameters

Now we shall define the priors to the hyperparameters. Define  $\lambda = \mu \cdot \theta$ . Given that it is more likely that previous research made statements about  $x_i$ , we use the same idea as Raftery considering the prior over  $(\lambda, \theta)$ . I will assume they are independent from now on with

$$\lambda \sim \text{Gamma}(\alpha, \beta),$$

and

$$\theta \sim \text{Beta}(a, b).$$

This simplification may be carried, given that the data brings information to the correlation, and it simplifies the math a lot. The other approach will have two settings. First I suppose  $\nu$  and  $\theta$  independents with

$$\nu \sim \text{Beta}(\alpha_1, \beta_1)$$

and

$$\theta \sim \text{Beta}(\alpha_2, \beta_2).$$

I choose the Beta distribution because it has a flexible shape with a good intuition behind it. Other point is that the Beta distribution forms a conjugate family for the Geometric distribution. Another set up is to consider the correlated case. We do it in the following way:

$$\begin{pmatrix} \text{logit}(\nu) \\ \text{logit}(\theta) \end{pmatrix} \sim \text{Normal}(\eta, \Sigma).$$

This choice is intrinsically linked to the fact the the normal distribution is a good approximation to a series of events, and it has a very good interpretation of the parameters. The problem with this approach is that it is harder to codify prior information. We necessarily need information about  $N$ ,  $\theta$ , and how they relate.

From these three approaches, I will call these approaches in the text *Raftery approach*, *Geometric and independent approach*, and *Geometric and correlated approach*, respectively.

- b) (5 marks) Using the prior from the previous item, write out the full joint posterior kernel for all unknown quantities in the model,  $p(\xi | \mathbf{x})$ . *Hint*: do not forget to include the appropriate indicator functions!;

**Solution.** For the Geometric and correlated approach, it may be impossible to find a simple expression.

Generally speaking, by Bayes' Theorem,

$$\begin{aligned} p(\xi | \mathbf{x}) &\propto l(\xi | \mathbf{x}) \cdot \pi(\xi) \\ &= \left( \prod_{i=1}^n \binom{N}{x_i} \theta^{x_i} (1-\theta)^{N-x_i} \mathbb{1}(N \geq x_i) \right) \cdot \pi(\xi) \\ &= \left( \prod_{i=1}^n \binom{N}{x_i} \right) \theta^S (1-\theta)^{nN-S} \cdot \pi(\xi) \cdot \mathbb{1}(N \geq x_{\max}), \end{aligned} \quad (1)$$

where  $S = \sum_{i=1}^n x_i$ . We shall derive for each case the prior  $\pi(\xi)$ .

(1) **Raftery**:

$$\begin{aligned} \pi(\xi) &= \int_0^\infty \pi(\xi, \lambda) d\lambda \\ &= \int_0^\infty \pi(N | \theta, \lambda) \pi(\theta, \lambda) d\lambda \\ &= \int_0^\infty \frac{e^{-\lambda/\theta} (\lambda/\theta)^N}{N!} \pi(\lambda) \pi(\theta) d\lambda \\ &\propto \int_0^\infty \frac{e^{-\lambda/\theta} (\lambda/\theta)^N}{N!} \lambda^{\alpha-1} e^{-\beta\lambda} \theta^{a-1} (1-\theta)^{b-1} \mathbb{1}(0 < \theta < 1) d\lambda \\ &= \frac{\theta^{a-1-N} (1-\theta)^{b-1}}{N!} \mathbb{1}(0 < \theta < 1) \int_0^\infty \lambda^{\alpha+N-1} e^{-(\beta+1/\theta)\lambda} d\lambda \\ &= \frac{\theta^{a-1-N} (1-\theta)^{b-1}}{N!} \mathbb{1}(0 < \theta < 1) \cdot \frac{\Gamma(\alpha+N)}{(\beta+1/\theta)^{\alpha+N}}, \end{aligned}$$

since the integrand is the kernel of a gamma distribution. Therefore, rewriting,

$$\pi(\xi) \propto \frac{\Gamma(\alpha+N) \theta^{a-1-N} (1-\theta)^{b-1}}{(\beta+1/\theta)^{\alpha+N} N!} \mathbb{1}(0 < \theta < 1) \quad (2)$$

and

$$p(\xi | \mathbf{x}) \propto \left( \prod_{i=1}^n \binom{N}{x_i} \right) \frac{\Gamma(\alpha+N) \theta^{a+S-1-N} (1-\theta)^{b+nN-S-1}}{(\beta+1/\theta)^{\alpha+N} N!} \mathbb{1}(0 < \theta < 1, N \geq x_{\max}) \quad (3)$$

(2) **Geometric and independent:**

$$\begin{aligned}
\pi(\xi) &= \int_0^1 \pi(\xi, \nu) d\nu \\
&= \int_0^1 \pi(N|\nu, \theta) \pi(\nu) \pi(\theta) d\nu \\
&\propto \theta^{\alpha_2-1} (1-\theta)^{\beta_2-1} \mathbb{1}(0 < \theta < 1) \int_0^1 (1-\nu)^N \nu \cdot \nu^{\alpha_1-1} (1-\nu)^{\beta_1-1} d\nu \\
&= \theta^{\alpha_2-1} (1-\theta)^{\beta_2-1} \mathbb{1}(0 < \theta < 1) \int_0^1 \nu^{\alpha_1} (1-\nu)^{N+\beta_1-1} d\nu \\
&= \theta^{\alpha_2-1} (1-\theta)^{\beta_2-1} \mathbb{1}(0 < \theta < 1) B(\alpha_1 + 1, N + \beta_1),
\end{aligned}$$

since the integrand is the kernel of a Beta distribution. Rewriting,

$$\pi(\xi) \propto B(\alpha_1 + 1, N + \beta_1) \theta^{\alpha_2-1} (1-\theta)^{\beta_2-1} \mathbb{1}(0 < \theta < 1) \quad (4)$$

and

$$p(\xi|\mathbf{x}) \propto \left( \prod_{i=1}^n \binom{N}{x_i} \right) B(\alpha_1 + 1, N + \beta_1) \theta^{S+\alpha_2-1} (1-\theta)^{nN+\beta_2-S-1} \mathbb{1}(0 < \theta < 1, N \geq x_{\max}) \quad (5)$$

(3) **Geometric and correlated:** This is the harder case. I have to derive  $\pi(\nu, \theta)$  from  $\pi(\text{logit}(\nu), \text{logit}(\theta))$ . Define  $f(x_1, x_2) = (\text{logit}^{-1}(x_1), \text{logit}^{-1}(x_2))$ . This is an invertible function with  $f^{-1}(y_1, y_2) = (\text{logit}(y_1), \text{logit}(y_2))$ . By the change of variables,

$$\pi(\nu, \theta) = \pi(f^{-1}(\nu, \theta)) \cdot \left| \det \left[ \frac{df^{-1}(z)}{dz} \right]_{z=(\nu, \theta)} \right|.$$

Observe that

$$\det \left[ \frac{df^{-1}(z)}{dz} \right]_{z=(\nu, \theta)} = \det \begin{bmatrix} \frac{d}{d\nu} \text{logit}(\nu) & 0 \\ 0 & \frac{d}{d\theta} \text{logit}(\theta) \end{bmatrix} = \frac{d}{d\nu} \text{logit}(\nu) \cdot \frac{d}{d\theta} \text{logit}(\theta).$$

By the calculation of the Jacobian, we can join everything

$$\begin{aligned}
\pi(\xi) &= \int_0^1 \pi(N|\nu, \theta) \pi(\nu, \theta) d\nu \\
&= \int_0^1 (1-\nu)^N \nu \pi(\text{logit}(\nu), \text{logit}(\theta)) |\text{logit}'(\nu) \cdot \text{logit}'(\theta)| d\nu \\
&= \int_0^1 (1-\nu)^N \nu \pi(\text{logit}(\nu), \text{logit}(\theta)) \frac{1}{\nu(1-\nu)\theta(1-\theta)} d\nu \\
&= \frac{1}{\theta(1-\theta)} \int_0^1 (1-\nu)^{N-1} \pi(\text{logit}(\nu), \text{logit}(\theta)) d\nu.
\end{aligned}$$

Let  $z = (\text{logit}(\nu), \text{logit}(\theta))$ . Therefore,

$$\pi(\xi) \propto \frac{1}{\theta(1-\theta)} \int_0^1 (1-\nu)^{N-1} \exp \left\{ -\frac{1}{2} (z - \eta)^T \Sigma^{-1} (z - \eta) \right\} d\nu \quad (6)$$

and, for  $N \geq x_{\max}$ ,

$$p(\xi|\mathbf{x}) \propto \left( \prod_{i=1}^n \binom{N}{x_i} \right) \theta^{S-1} (1-\theta)^{nN-S-1} \int_0^1 (1-\nu)^{N-1} \exp \left\{ -\frac{1}{2} (z - \eta)^T \Sigma^{-1} (z - \eta) \right\} d\nu. \quad (7)$$

c) (5 marks) Is your model identifiable?

**Solution.** I shall verify the likelihood of the model is identifiable and it brings the information from the data to the parameter. I say a model is identifiable if the implication

$$P_{\theta_1} = P_{\theta_2} \implies \theta_1 = \theta_2$$

is true for the model  $P_\theta$ . Suppose that

$$\binom{N_1}{x} \theta_1^x (1 - \theta_1)^{N_1 - x} = \binom{N_2}{x} \theta_2^x (1 - \theta_2)^{N_2 - x}$$

for every  $x$  between 0 and  $\min(N_1, N_2)$ . Suppose  $N_1 < N_2$ . Therefore the equality is false for  $x = N_1 + 1$ . If  $N_1 > N_2$ , the equality is also false. We conclude  $N_1 = N_2$ , and, therefore,

$$\frac{\theta_1}{1 - \theta_1} = \frac{\theta_2}{1 - \theta_2} \implies \theta_1 = \theta_2.$$

Hence, the binomial model is identifiable. For  $n$  observations, take  $x_1 = \dots = x_n = x$ . We will have

$$f(x|\theta_1, N_1)^n = f(x|\theta_2, N_2)^n \implies f(x|\theta_1, N_1) = f(x|\theta_2, N_2)$$

and the prove follows as previously.

Strictly speaking, given that the prior is proper, and hence the posterior is proper, all the parameters are identifiable. The choice of a suitable prior can resolve the non-identifiability [Xie and Carlin \(2006\)](#). We can also see in the three models, the posterior of  $\xi$  has additional information of the data, coming from the productory and  $S$ .

d) (5 marks) Exhibit the marginal posterior density for  $N$ ,  $p_1(N | \mathbf{x})$ ;

**Solution.** We separate the marginalized posterior:

(1) **Raftery:** Consider the equation 3 and integrate over  $\theta$ . For  $N \geq x_{\max}$ ,

$$p_1(N|\mathbf{x}) \propto \left( \prod_{i=1}^n \binom{N}{x_i} \right) \frac{\Gamma(\alpha + N)}{N!} \int_0^1 \frac{\theta^{\alpha+S-1-N} (1-\theta)^{b+nN-S-1}}{(\beta + 1/\theta)^{\alpha+N}} d\theta. \quad (8)$$

(2) **Geometric independent:** Consider the equation 5 and integrate over  $\theta$ . For  $N \geq x_{\max}$ ,

$$\begin{aligned} p_1(N|\mathbf{x}) &\propto \left( \prod_{i=1}^n \binom{N}{x_i} \right) B(\alpha_1 + 1, N + \beta_1) \int_0^1 \theta^{S+\alpha_2-1} (1-\theta)^{nN+\beta_2-S-1} d\theta \\ &= \left( \prod_{i=1}^n \binom{N}{x_i} \right) B(\alpha_1 + 1, N + \beta_1) B(S + \alpha_2, nN - S + \beta_2). \end{aligned} \quad (9)$$

(3) **Geometric correlated:** Consider the equation 7 and integrate over  $\theta$ . For  $N \geq x_{\max}$ ,

$$\begin{aligned} p_1(N|\mathbf{x}) &\propto \left( \prod_{i=1}^n \binom{N}{x_i} \right) \int_0^1 \int_0^1 \theta^{S-1} (1-\theta)^{nN-S-1} (1-\nu)^{N-1} \\ &\quad \times \exp \left\{ -\frac{1}{2} (z - \eta)^T \Sigma^{-1} (z - \eta) \right\} d\nu d\theta. \end{aligned} \quad (10)$$

e) (5 marks) Return to point (a) above and consider an alternative, uninformative prior structure for  $\xi$ ,  $\pi_2$ . Then, derive  $p_2(N | \mathbf{x})$ ;

**Solution.** For an uninformative prior, for instance, we could consider in the Raftery approach  $\alpha, \beta \rightarrow 0$ . When we do that, we obtain

$$\pi(\lambda) \propto \lambda^{-1},$$

an improper prior and  $a = b = 1$ , what implies  $\theta$  uniformly distributed on the interval  $(0, 1)$ . In that regard,

$$p_2(N|\mathbf{x}) \propto \left( \prod_{i=1}^n \binom{N}{x_i} \right) \frac{\Gamma(N)}{N!} \int_0^1 \frac{\theta^{S-N}(1-\theta)^{nN-S}}{\theta^{-N}} d\theta = \left( \prod_{i=1}^n \binom{N}{x_i} \right) \frac{B(S+1, nN-S+1)}{N}.$$

Simplifying, we have

$$p_2(N|\mathbf{x}) \propto \left( \prod_{i=1}^n \binom{N}{x_i} \right) \frac{(nN-S)!}{(nN+1)!N}. \quad (11)$$

We need to prove  $p_2$  defines a probability distribution with a correct constant. Observe that,

$$\begin{aligned} \left( \prod_{i=1}^n \binom{N}{x_i} \right) \frac{(nN-S)!}{(nN+1)!N} &= \frac{1}{N \prod_{i=1}^n x_i!} \frac{\prod_{i=1}^n \prod_{j=1}^{x_i} (N-x_i+j)}{\prod_{k=1}^{S+1} (nN-S+k)} \frac{N^{-(S+1)}}{N^{-(S+1)}} \\ &\propto \frac{1}{N^2} \frac{\prod_{i=1}^n \prod_{j=1}^{x_i} (1-(x_i-j)/N)}{\prod_{k=1}^{S+1} (n-(S-k)/N)} \end{aligned}$$

given that the numerator has  $x_1 + \dots + x_n = S$  factors, while the denominator has  $S+1$ . Observe that, taking  $n \geq 2$ , for every  $i, j$ , and  $k$  if  $N \geq S$ ,

$$1 - (x_i - j)/N < 1$$

and  $n - (S - k)/N \geq n - 1 + 1/N > 1$ . Therefore, if  $N \geq S$

$$\frac{1}{N^2} \frac{\prod_{i=1}^n \prod_{j=1}^{x_i} (1-(x_i-j)/N)}{\prod_{k=1}^{S+1} (n-(S-k)/N)} \leq \frac{1}{N^2}.$$

and

$$\sum_{N=x_{\max}}^{\infty} \left( \prod_{i=1}^n \binom{N}{x_i} \right) \frac{(nN-S)!}{(nN+1)!N} \leq \sum_{x_{\max}}^{S-1} \left( \prod_{i=1}^n \binom{N}{x_i} \right) \frac{(nN-S)!}{(nN+1)!N} + \sum_{N=S}^{\infty} \frac{1}{N^2} < +\infty,$$

what proves  $p_2(N|\mathbf{x})$  is a probability distribution given the right constant. If  $n = 1$ , we have that

$$\binom{N}{x} \frac{(N-x)!}{(N+1)!N} \propto \frac{1}{N(N+1)} \leq \frac{1}{N^2}$$

also convergent.

- f) (10 marks) Formulate a third prior structure on  $\xi, \pi_3$ , that allows for the closed-form marginalization over the hyperparameters  $\alpha$  – see (a) – and write out  $p_3(N|\mathbf{x})$ ;

**Solution.** The structure of the Geometric independent approach allowed the closed-form marginalization over  $\nu$  and  $\theta$ , as expressed in equation 9. I will denote from now on  $p_4$  the marginalized distribution of Geometric correlated approach, given by expression 10.

- g) (10 marks) Show whether each of the marginal posteriors considered is proper. Then, derive the posterior predictive distribution,  $g_i(\tilde{x}|\mathbf{x})$ , for each of the posteriors considered ( $i = 1, 2, 3, 4$ ).

**Solution.**  $p_1$  (Raftery),  $p_3$  (Geometric independent), and  $p_4$  (Geometric correlated) are proper since the prior is proper, by Bayes' theorem. For a prove, consult the second question from A1 exam [Moschen \(2021\)](#). We proved that the posterior  $p_2$  is proper in exercise (e). Now we derive the posterior predictive distribution

$$g(\tilde{x}|\mathbf{x}) = \sum_{N=x_{\max}}^{\infty} \int_0^1 f(\tilde{x}, \mathbf{x}|\xi) \pi(\xi) d\theta$$

for each of the four priors. Let  $x_{n+1} = \tilde{x}$  for notation simplification.

(1) **Raftery:** Consider the prior from expression 2.

Let  $S' = S + x_{n+1}$  and  $x'_{\max} = \max(x_{\max}, x_{n+1})$ .

$$\begin{aligned} g_1(x_{n+1}|\mathbf{x}) &\propto \sum_{N=x'_{\max}}^{\infty} \int_0^1 \left\{ \prod_{i=1}^{n+1} \binom{N}{x_i} \right\} \theta^{S'} (1-\theta)^{(n+1)N-S'} \frac{\Gamma(\alpha+N) \theta^{\alpha-1-N} (1-\theta)^{b-1}}{(\beta+1/\theta)^{\alpha+N} N!} d\theta \\ &= \sum_{N=x'_{\max}}^{\infty} \left\{ \prod_{i=1}^{n+1} \binom{N}{x_i} \right\} \frac{\Gamma(\alpha+N)}{N!} \int_0^1 \frac{\theta^{a+S'-1-N} (1-\theta)^{(n+1)N-S'+b-1}}{(\beta+1/\theta)^{\alpha+N}} d\theta. \end{aligned}$$

(2) **Uninformative:** This case is the same as Raftery, with  $a = b = 1$  and  $\alpha = \beta = 0$ . Then,

$$\begin{aligned} g_2(x_{n+1}|\mathbf{x}) &\propto \sum_{N=x'_{\max}}^{\infty} \left\{ \prod_{i=1}^{n+1} \binom{N}{x_i} \right\} \frac{\Gamma(N)}{N!} \int_0^1 \frac{\theta^{S'-N} (1-\theta)^{(n+1)N-S'}}{(1/\theta)^N} d\theta \\ &= \sum_{N=x'_{\max}}^{\infty} \left\{ \prod_{i=1}^{n+1} \binom{N}{x_i} \right\} \frac{1}{N} \int_0^1 \theta^{S'} (1-\theta)^{(n+1)N-S'} d\theta \\ &= \sum_{N=x'_{\max}}^{\infty} \left\{ \prod_{i=1}^{n+1} \binom{N}{x_i} \right\} \frac{1}{N} B(S'+1, (n+1)N-S'+1) \\ &= \sum_{N=x'_{\max}}^{\infty} \left\{ \prod_{i=1}^{n+1} \binom{N}{x_i} \right\} \frac{S'![(n+1)N-S']!}{[(n+1)N+1]!N}. \end{aligned}$$

(3) **Geometric independent:** Consider the prior from expression 4.

Let  $B_N = B(\alpha_1 + 1, N + \beta_1)$  to reduce the size of the expression. Then

$$\begin{aligned} g_3(x_{n+1}|\mathbf{x}) &\propto \sum_{N=x'_{\max}}^{\infty} \int_0^1 \left\{ \prod_{i=1}^{n+1} \binom{N}{x_i} \right\} \theta^{S'} (1-\theta)^{(n+1)N-S'} B_N \theta^{\alpha_2-1} (1-\theta)^{\beta_2-1} d\theta \\ &= \sum_{N=x'_{\max}}^{\infty} \left\{ \prod_{i=1}^{n+1} \binom{N}{x_i} \right\} B_N \int_0^1 \theta^{S'+\alpha_2-1} (1-\theta)^{(n+1)N+\beta_2-S'-1} d\theta \\ &= \sum_{N=x'_{\max}}^{\infty} \left\{ \prod_{i=1}^{n+1} \binom{N}{x_i} \right\} B(\alpha_1 + 1, N + \beta_1) B(S' + \alpha_2, (n+1)N + \beta_2 - S'). \end{aligned}$$

(4) **Geometric correlated:** Consider the prior expression 6.

Denote  $I_{\theta,N} = \int_0^1 (1-\nu)^{N-1} \exp\left\{-\frac{1}{2}(z-\eta)^T \Sigma^{-1}(z-\eta)\right\} d\nu$ .

$$\begin{aligned} g_4(x_{n+1}|\mathbf{x}) &\propto \sum_{N=x'_{\max}}^{\infty} \int_0^1 \left\{ \prod_{i=1}^{n+1} \binom{N}{x_i} \right\} \theta^{S'} (1-\theta)^{(n+1)N-S'} \frac{1}{\theta(1-\theta)} \cdot I_{\theta,N} d\theta \\ &= \sum_{N=x'_{\max}}^{\infty} \left\{ \prod_{i=1}^{n+1} \binom{N}{x_i} \right\} \int_0^1 \theta^{S'-1} (1-\theta)^{(n+1)N-S'-1} \cdot I_{\theta,N} d\theta \end{aligned}$$

h) (5 marks) Consider the loss function

$$L(\delta(\mathbf{x}), N) = \left( \frac{\delta(\mathbf{x}) - N}{N} \right)^2. \quad (12)$$

Derive the Bayes estimator under this loss.

**Solution.** The Bayes estimator is

$$\delta^\pi(\mathbf{x}) = \arg \min_d \mathbb{E}^\pi [L(d, N) | \mathbf{x}] = \arg \min_d \mathbb{E}^\pi \left[ \left( \frac{d - N}{N} \right)^2 | \mathbf{x} \right].$$

In general terms,

$$\mathbb{E}^\pi \left[ \left( \frac{d - N}{N} \right)^2 | \mathbf{x} \right] = \mathbb{E}^\pi \left[ \left( \frac{d^2}{N^2} - 2d \frac{1}{N} + 1 \right) | \mathbf{x} \right] = d^2 \mathbb{E}^\pi [N^{-2} | \mathbf{x}] - 2d \mathbb{E}^\pi [N^{-1} | \mathbf{x}] + 1. \quad (13)$$

The minimum argument of a quadratic expression is given By

$$\delta^\pi(\mathbf{x}) = \frac{\mathbb{E}^\pi [N^{-1} | \mathbf{x}]}{\mathbb{E}^\pi [N^{-2} | \mathbf{x}]} = \frac{\sum_{N=x_{\max}}^{\infty} N^{-1} p_i(N | \mathbf{x})}{\sum_{N=x_{\max}}^{\infty} N^{-2} p_i(N | \mathbf{x})},$$

em que  $i = 1, \dots, 4$ .

## Inferring population sizes – practice

Consider the problem of inferring the population sizes of major herbivores ([Carroll and Lombard, 1985](#)). In the first case, one is interested in estimating the number of impala (*Aepyceros melampus*) herds in the Kruger National Park, in northeastern South Africa. In an initial survey collected the following numbers of herds:  $\mathbf{x}_{\text{impala}} = \{15, 20, 21, 23, 26\}$ . Another scientific question is the number of individual waterbuck (*Kobus ellipsiprymnus*) in the same park. The observed numbers of waterbuck in separate sightings were  $\mathbf{x}_{\text{waterbuck}} = \{53, 57, 66, 67, 72\}$  and may be regarded (for simplicity) as independent and identically distributed.



(a) Impala



(b) Waterbuck

Figure 1: Two antelope species whose population sizes we want to estimate.

- i) (20 marks) For each data set, sketch the marginal posterior distributions  $p_1(N | \mathbf{x})$ ,  $p_2(N | \mathbf{x})$  and  $p_3(N | \mathbf{x})$ . Moreover, under each posterior, provide (i) the Bayes estimator under quadratic loss and under the loss in (12) and (ii) a 95% credibility interval for  $N$ . Discuss the differences and similarities between these distributions and estimates: do the prior modelling choices substantially impact the final inferences? If so, how?



**Solution.** I use *Stan* platform with *PyStan* interface for *Python programming language*. The problem is that it does not handle unbounded discrete parameters. Therefore we use the approach suggested in the documentation [Stan Development Team \(2021\)](#) considering the population size a continuous variable. I tried to use the package *PyMC3* for this model (this package handles discrete parameters), but given convergence problems, I do not present the results.

As I will present posteriorly, this problem may be related to the sampler Metropolis. For the Geometric and independent approach, it is possible to compare the analytic curve with Stan result (considering  $N$  a real variable), and Stan got a well approximate result, while PyMC3 had convergence problems.

In Figures XXX and YYY, I sketch the marginal posterior distributions for each approach and each data set. In table XXX, it is shown the Bayes estimator under quadratic loss (the expected value ([Robert, 2007](#), Section 4)), under loss [12](#), and 95% credibility interval.

- j) (25 marks) Let  $\bar{x} = K^{-1} \sum_{k=1}^K x_k$  and  $s^2 = K^{-1} \sum_{k=1}^K (x_k - \bar{x})^2$ . For this problem, a sample is said to be *stable* if  $\bar{x}/s^2 \geq (\sqrt{2} + 1)/\sqrt{2}$  and *unstable* otherwise. Devise a simple method of moments estimator (MME) for  $N$ . Then, using a Monte Carlo simulation, compare the MME to the three Bayes estimators under quadratic loss in terms of relative mean squared error. How do the Bayes estimators compare to MME in terms of the stability of the generated samples? *Hint*: You may want to follow the simulation setup of [Carroll and Lombard \(1985\)](#).

**Solution.** Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

## Appendix

### A Binomial and Poisson

Suppose  $X|N \sim \text{Binomial}(N, p)$  and  $N \sim \text{Poisson}(\mu)$ . We shall derive the distribution of  $X$ . By the Law of total probability,

$$\begin{aligned}
\Pr(X = k) &= \sum_{n=0}^{\infty} \Pr(X = k|N = n) \Pr(N = n) \quad [\Pr(X = k|N = n) = 0 \text{ if } k > n] \\
&= \sum_{n=k}^{\infty} \binom{n}{k} p^k (1-p)^{n-k} \frac{e^{-\mu} \mu^n}{n!} \\
&= \frac{e^{-\mu} p^k}{k!} \sum_{n=k}^{\infty} \frac{(1-p)^{n-k}}{(n-k)!} \mu^{n-k} \mu^k \\
&= \frac{e^{-\mu} (p\mu)^k}{k!} \sum_{n=k}^{\infty} \frac{((1-p)\mu)^{n-k}}{(n-k)!} \\
&= \frac{e^{-\mu} (p\mu)^k}{k!} \sum_{m=0}^{\infty} \frac{((1-p)\mu)^m}{m!} \quad [m = n - k] \\
&= \frac{e^{-\mu} (p\mu)^k}{k!} e^{(1-p)\mu} \\
&= \frac{e^{-p\mu} (p\mu)^k}{k!},
\end{aligned} \tag{14}$$

what implies  $X \sim \text{Poisson}(p \cdot \mu)$ .

## Bibliography

- Carroll, R. J. and Lombard, F. (1985). A note on N estimators for the binomial distribution. *Journal of the American Statistical Association*, 80(390):423–426.
- Moschen, L. M. (2021). Exam a1. Available at <https://github.com/lucamoschen/phd-bayesian-statistics/blob/main/notes/exam/A1.pdf>.
- Raftery, A. E. (1988). Inference for the binomial n parameter: A hierarchical bayes approach. *Biometrika*, 75(2):223–228.
- Robert, C. (2007). *The Bayesian choice: from decision-theoretic foundations to computational implementation*. Springer Science & Business Media.
- Stan Development Team (2021). Mark-recapture models. Available at [https://mc-stan.org/docs/2\\_27/stan-users-guide/mark-recapture-models.html](https://mc-stan.org/docs/2_27/stan-users-guide/mark-recapture-models.html).
- Xie, Y. and Carlin, B. P. (2006). Measures of bayesian learning and identifiability in hierarchical models. *Journal of Statistical Planning and Inference*, 136(10):3458–3477.