

Prevalence estimation

Lucas Machado Moschen

*School of Applied Mathematics,
Fundação Getulio Vargas*

July 26, 2021

1 Introduction

A key question for epidemiologists and public health authorities is about the proportion of individuals exposed to the disease at time t . This quantity can be measured periodically, and the evolution shows how the transmission is going on. For instance, if after a year the proportion grew 50% it would be worrisome. We call it *prevalence*. High prevalence of a disease within a population might mean that there is a high incidence of it or prolonged survival without cure.

This report presents the initial models for my bachelor dissertation entitled “Bayesian analysis of respondent-driven surveys with outcome uncertainty”, which proposes to study prevalence when the diagnostic tests are imperfect and the population is hidden, that is, there is no sampling frame for it [Heckathorn \(1997\)](#).

1.1 Respondent-driven sampling

RDS is commonly used to survey hidden or hard-to-reach populations when no sampling frame exists [Heckathorn \(1997\)](#). In this approach, the researchers select some individuals, called *seeds* from the target population, and give them a fixed amount of *recruitment coupons* to recruit their peers. Each recipient of the coupons reclaims it in the study site, is interviewed, and receives more coupons to continue the recruitment. This process occurs until it reaches some criteria. The sampling is without replacement, so the participants cannot be recruited more than once. Moreover, the respondents inform their *network degree*.

The subjects receive a reward for being interviewed and for each recruitment of their peers which establishes a dual system incentive. The *primary incentive* is the *individual-sanction-based control*, so there is a reward for participating. The second one is the *group-mediated social control* that influences the participants to induce others to comply to get the reward for the recruitment. When social approval is important, recruitment can be even more efficient and cheaper, since material incentive can be converted into symbolic by the individuals. In summary, accepting to be recruited will have a material incentive for both and a symbolic incentive for the recruited, since their peers also participated.

Let $G = (V, E)$ be an undirected graph representing the hidden population. The *recruitment graph* $G_R = (V_R, E_R)$ represents the recruited individuals and the recruitment edge. Given that each individual can be sampled only once, it is not possible to observe the *recruitment-induced subgraph*, that is the induced subgraph generated by V_R . Moreover, the *coupon matrix* C defined by $C_{ij} = 1$ if the i^{th} subject has at least one coupon before the j^{th} recruitment event, is also observed with the recruitment times. Assuming an exponential and independent distribution of the times, the likelihood can be written explicitly, and the distribution interpreted as an exponential random graph model.

These models allowed several applications in social sciences, epidemiology, and statistics, including hidden populations size estimation [Crawford et al. \(2018\)](#), regression [Bastos et al. \(2012\)](#), communicable disease prevalence estimation [Albuquerque et al. \(2009\)](#), among others.

2 Preliminary definitions

Suppose we have a sample indexed by i . Let Y_i be the indicator function of the i^{th} individual's exposure to the disease, and T_i indicating whether the test in the i^{th} individual is positive at time t . Suppose that $\{Y_i\}$ and $\{T_i\}$ are two independent and identically distributed random variables with $\Pr(X = 1) = \theta$ and $\Pr(T = 1) = p$. We say that θ is the prevalence and p is the apparent prevalence in the population.

If the test is perfect, then for every i , $T_i = Y_i$, and $\theta = p$ (with probability one when they are random variables). Unfortunately, this is not true in the real world, what makes important to regard the evaluation of the diagnostic, and the following definitions are used:

Definition 2.1 (Specificity). Probability of a negative test correctly identified. In mathematical terms, conditioned on $Y = 0$, the *specificity* γ_e is the probability of $T = 0$:

$$\gamma_e = \Pr(T = 0 | Y = 0). \quad (1)$$

Definition 2.2 (Sensitivity). Probability of a positive test correctly identified. In mathematical terms, conditioned on $Y = 1$, the *sensitivity* γ_s is the probability of $T = 1$:

$$\gamma_s = \Pr(T = 1 | Y = 1). \quad (2)$$

Theorem 1 (Relation between prevalence and apparent prevalence). *These quantities are related by the following equation:*

$$p = \gamma_s \theta + (1 - \gamma_e)(1 - \theta). \quad (3)$$

Proof. This is a direct application of the definition of conditional probability and the countable additivity axiom of Probability:

$$\begin{aligned} p &= \Pr(T = 1) = \Pr(T = 1, Y = 1) + \Pr(T = 1, Y = 0) \\ &= \Pr(T = 1 | Y = 1) \Pr(Y = 1) + \Pr(T = 1 | Y = 0) \Pr(Y = 0) \\ &= \Pr(T = 1 | Y = 1) \Pr(Y = 1) + (1 - \Pr(T = 0 | Y = 0))(1 - \Pr(Y = 1)) \\ &= \gamma_s \theta + (1 - \gamma_e)(1 - \theta). \end{aligned}$$

□

The intuition behind this equation is pretty simple: the proportion of positive test counts the correct identified exposed individuals and the incorrect identified not exposed. Observe that if $\gamma_s = \gamma_e = 1$, we have the trivial case $p = \theta$. Moreover, if $\gamma_s = \gamma_e = 0.5$, we have that $p = 0.5$ and there is no information about θ .

Remark. Actually, we are interested in the pontual prevalence at time t . Being impossible to test every individual at the same time, we assume that all individuals remain exposed to the disease at time of the last tested individual.

Definition 2.3 (Link function). A class of functions which maps a non-linear relationship to a linear one. Here we consider functions with domain $[0, 1]$. Examples include the logit and probit functions.

3 Prevalence model

Firstly, we make some assumptions to simplify the modeling:

Assumption 1. For a Bayesian modeling, we assume each model's parameter has a probability distribution that incorporates the researcher's uncertainty about it.

Assumption 2. For each individual, we observe k features that are possible risk factors represented by the vector $\mathbf{x}_i \in \mathbb{R}^k$ of the i^{th} individual.

Assumption 3. Suppose that each individual has a probability θ_i of having been exposed to the disease that depends on the prevalence θ and \mathbf{x}_i , not necessarily linearly. We therefore will have the probability of positive test p_i in the i^{th} individual. Therefore, the sequences $\{Y_i\}$ and $\{T_i\}$ are not identically distributed anymore.

Assumption 4. Sensitivity and specificity have the same distribution for all individuals and it only depends on the test used to diagnose.

From above, we develop three different models.

3.1 Perfect tests

The first model supposes the samples are independent and the test is perfect, which means that $\theta_i = p_i$ for all i . Therefore it only considers the risk factors \mathbf{x}_i .

$$\begin{aligned} T_i &\sim \text{Bernoulli}(\theta_i), \\ g(\theta_i) &= g(\theta) + \mathbf{x}_i\beta, \end{aligned} \tag{4}$$

where $g(\cdot)$ is a link function. The parameter $\beta \in \mathbb{R}^k$ is the risk effects. For Bayesian inference, priors on β and θ must be included. We use $\beta \sim \text{Normal}(\mu, \Sigma)$ and $\theta \sim \text{Beta}(\alpha^p, \beta^p)$, where $\mu \in \mathbb{R}^k$, $\Sigma \in \mathbb{R}^{k \times k}$ symmetric positive-definite matrix, $\alpha^p \in \mathbb{R}_{++}$, and $\beta^p \in \mathbb{R}_{++}$ are fixed hyperparameters.

Remark. If the risk effects are zero, i.e $\mathbf{x}_i = 0$, the probability of the i^{th} having been exposed is the prevalence θ , which means that in a population with no risk effects, the probability of a person has the disease is exactly the proportion in this population.

3.1.1 Identifiability

3.1.2 Experiments

https://github.com/lucamoschen/rds-bayesian-analysis/blob/main/exercises/primary_model/model_experiments.ipynb

3.2 Imperfect tests

This model includes the sensitivity and specificity of the diagnostic test.

$$\begin{aligned} T_i &\sim \text{Bernoulli}(p_i) \\ p_i &= \gamma_s \theta_i + (1 - \gamma_e)(1 - \theta_i), \\ g(\theta_i) &= g(\theta) + \mathbf{x}_i\beta, \\ \beta &\sim \text{Normal}(\mu, \Sigma), \\ \theta &\sim \text{Beta}(\alpha^p, \beta^p) \\ \gamma_s &\sim \text{Beta}(\alpha^s, \beta^s), \\ \gamma_e &\sim \text{Beta}(\alpha^e, \beta^e), \end{aligned} \tag{5}$$

where $\alpha^p, \alpha^s, \alpha^e, \beta^p, \beta^s, \beta^e \in \mathbb{R}_{++}$ are fixed hyperparameters. This model does not include prior knowledge about the correlation between specificity and sensitivity.

3.3 Imperfect test and respondent-driven sampling

For now, we consider the network dependence induced by the RDS. Ideally, we should build a model for it and include the associated uncertainty. Since it is not possible (for a while), we

treat it as a random effect for each individual. Conditionally autoregressive (CAR) models in the Gaussian case are used. Therefore:

$$\begin{aligned}
T_i &\sim \text{Bernoulli}(p_i) \\
p_i &= \gamma_s \theta_i + (1 - \gamma_e)(1 - \theta_i), \\
g(\theta_i) &= g(\theta) + \mathbf{x}_i \beta + \omega_i, \\
\omega_i | \{\omega_j\}_{j \neq i}, \tau &\sim \text{Normal} \left(\rho \sum_j b_{ij} \omega_j / b_{i+}, \tau^2 / b_{i+} \right) \\
\beta &\sim \text{Normal}(\mu, \Sigma), \\
\theta &\sim \text{Beta}(\alpha^p, \beta^p) \\
\gamma_s &\sim \text{Beta}(\alpha^s, \beta^s), \\
\gamma_e &\sim \text{Beta}(\alpha^e, \beta^e), \\
\tau &\sim \text{Normal}^+(0, \sigma_\tau^2),
\end{aligned} \tag{6}$$

such that ρ is a fixed parameter in order to the distribution be proper, $[B]_{ij} = b_{ij}$ is a fixed matrix to measure the distance between i and j , and $b_{i+} = \sum_j b_{ij}$. For instance,

$$b_{ij} = \begin{cases} 1, & \text{if } i \text{ recruited } j \text{ or the contrary} \\ 0, & \text{otherwise.} \end{cases}$$

3.3.1 Exponential Random Graph Model (ERGM)

RDS has the constraint of being without replacement. For that reason, we do not observe all links among the samples [Crawford \(2016\)](#). Considering the model developed by Crawford, we can model the matrix B as *Exponential Random Graph Model* (ERGM). Define the following

1. $\mathbf{s} = \text{tril}(BC)^T \mathbf{1} + C^T \mathbf{u}$, such that B is the adjacency matrix of the recruited subjects, C is the *Coupon Matrix*, \mathbf{u} the vector of the number of edge ends belonging to each vertex (in the order of recruitment) that are not connected to any other sampled vertex, and $\text{tril}(M)$ the lower triangle of M .
2. $T(B) = -\lambda \mathbf{s}$, such that λ is the rate of the recruitment time.
3. $V(B) = \sum_{k \text{ is not seed}} \log(\lambda \mathbf{s}_k)$
4. $w = (0, t_2 - t_1, \dots, t_n - t_{n-1})$ is the waiting times between recruitments.

Therefore $\Pr(B|w) \propto \exp[T(B)^T w + V(B)]$. With that, the model becomes

$$\begin{aligned}
T_i &\sim \text{Bernoulli}(p_i) \\
p_i &= \gamma_s \theta_i + (1 - \gamma_e)(1 - \theta_i), \\
g(\theta_i) &= g(\theta) + \mathbf{x}_i \beta + \omega_i, \\
\omega_i | \{\omega_j\}_{j \neq i}, \tau &\sim \text{Normal} \left(\rho \sum_j b_{ij} \omega_j / b_{i+}, \tau^2 / b_{i+} \right) \\
B|w &\propto \exp[T(B)^T w + V(B)] \\
\lambda &\sim \Gamma(\alpha^\lambda, \beta^\lambda), \\
\beta &\sim \text{Normal}(\mu, \Sigma), \\
\theta &\sim \text{Beta}(\alpha^p, \beta^p) \\
\gamma_s &\sim \text{Beta}(\alpha^s, \beta^s), \\
\gamma_e &\sim \text{Beta}(\alpha^e, \beta^e), \\
\tau &\sim \text{Normal}^+(0, \sigma_\tau^2).
\end{aligned} \tag{7}$$

The problem with this model is that we are assigning a posterior distribution for B .

4 Correlation analysis between specificity and sensitivity

In this section, we shall describe how to use the Bivariate Beta [Olkin and Trikalinos \(2015\)](#) to model the correlation between specificity and sensitivity.

References

- Albuquerque, E. M. d. et al. (2009). *Avaliação da técnica de amostragem respondent-driven sampling na estimação de prevalências de doenças transmissíveis em populações organizadas em redes complexas*. PhD thesis, ENSP.
- Bastos, L. S., Pinho, A. A., Codeço, C., and Bastos, F. I. (2012). Binary regression analysis with network structure of respondent-driven sampling data.
- Crawford, F. W. (2016). The graphical structure of respondent-driven sampling. *Sociological Methodology*, 46(1):187–211.
- Crawford, F. W., Wu, J., and Heimer, R. (2018). Hidden population size estimation from respondent-driven sampling: a network approach. *Journal of the American Statistical Association*, 113(522):755–766.
- Heckathorn, D. D. (1997). Respondent-driven sampling: A new approach to the study of hidden populations. *Social Problems*, 44(2):174–199.
- Olkin, I. and Trikalinos, T. A. (2015). Constructions for a bivariate beta distribution. *Statistics & Probability Letters*, 96:54–60.