Debugging time evolution

Lucas Myers

November 18, 2022

1 Checking periodically perturbed free energy

To check that the configuration force matches with the energy, we check the energy for a uniform configuration first, and then for a small perturbation. The latter check gives an estimate for the configuration force by the functional Taylor series expansion:

$$F(Q+\delta Q) = F(Q) + \int_\Omega \left(\frac{\delta F}{\delta Q}\Big|_Q : \delta Q \right) dV + \mathcal{O}\left(|\delta Q|^2\right) \tag{1}$$

with

$$\frac{\delta F}{\delta Q} = \frac{\partial f}{\partial Q} - \nabla \cdot \frac{\partial f}{\partial (\nabla Q)} \tag{2}$$

Since this holds for an arbitrary functional F, and also for an arbitrary (small) perturbation δQ, it's probably true that:

$$f(Q+\delta Q) = f(Q) + \left(\frac{\partial f}{\partial Q} - \nabla \cdot \frac{\partial f}{\partial (\nabla Q)} \right) : \delta Q + \mathcal{O}\left(|\delta Q|^2\right) \tag{3}$$

for each component of the free energy f. For the example we test, we take Q to be:

$$Q = S_0 \begin{bmatrix} \frac{2}{3} & 0 & 0 \\ 0 & -\frac{1}{3} & 0 \\ 0 & 0 & -\frac{1}{3} \end{bmatrix} \tag{4}$$

with

$$\delta Q = S_0 \begin{bmatrix} 0 & \epsilon \sin kx & 0 \\ \epsilon \sin kx & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \tag{5}$$

This corresponds (to first order) to a perturbation of the director so that the total configuration $(Q+\delta Q)$ is a uniform-S configuration with director $\mathbf{n} = (1, \epsilon \sin kx, 0)$.

Here, the nondimensional free energy is given by:

$$F = \int_\Omega \left(-\tfrac{1}{2}\alpha Q : Q + \left(\log 4\pi - \log Z + \Lambda : \left(Q + \tfrac{1}{3} I \right) \right) + \tfrac{1}{2} |\nabla Q|^2 \right) dV \tag{6}$$

Then we can calculate $F(Q)$ partially analytically and partially numerically. Of course the elastic part is zero, and we calculate:

$$-\tfrac{1}{2}\alpha Q : Q = -S_0^2 \frac{\alpha}{3} = -1.2153600266666669 \tag{7}$$

with $S_0 = 0.6751$ and $\alpha = 8.0$. Further, we can calculate numerically (see `calc_lambda` program with `generate_periodic_Q_tensors.py`) Λ and Z for $S_0 = 0.6751$:

$$\begin{aligned} Z &= 3.87017170996747 \\ \Lambda &= \begin{bmatrix} 3.6005952163635766 & 0 & 0 \\ 0 & -1.8002976081817883 & 0 \\ 0 & 0 & -1.8002976081817883 \end{bmatrix} \end{aligned} \tag{8}$$

Then we may calculate F by multiplying this energy density by the size of the domain. The uniform configuration check is then just to compare that number with the energy output from the simulation for a uniform configuration with the same S_0 and α values and domain size. Calculating the energy density from the mean-field interaction gives:

$$f_{\text{entropy}} = 3.60848720197831 \tag{9}$$

Running the simulation and calculating the free energy over the domain gives:

$$F_{\text{mean field}} = -11.99497833 \tag{10}$$

and calculating out the energy density from the entropy term gives:

$$F_{\text{entropy}} = 35.61419603 \tag{11}$$

Comparing these values to the energy density multiplied by $(3.1415926 \times 3.1415926)$ (the domain size) we get similar results.

We may also consider a configuration with a different S_0-value, just to be sure. Running a similar calculation with $S_0 = 0.5$, we get the following values for Λ and Z:

$$\begin{aligned} Z &= 1.8088523960817302 \\ \Lambda &= \begin{bmatrix} 2.323990879382531 & 0 & 0 \\ 0 & -1.1619954396912653 & 0 \\ 0 & 0 & 0 \end{bmatrix} \end{aligned} \tag{12}$$

Then the entropy and mean field part of the free energy density is given by:

$$\begin{aligned} f_{\text{entropy}} &= 3.100327077782174 \\ f_{\text{mean field}} &= -\frac{2}{3} \end{aligned} \tag{13}$$

Both of these corresopnd with the energy densities written to the vtu file.

The configuration force, which is exactly the functional derivative of the free energy, is given by:

$$\frac{\delta F}{\delta Q} = -\alpha Q + \Lambda - \nabla^2 Q \tag{14}$$

We need to evaluate this with the unperturbed Q-tensor – since this is just a uniform configuration, the elastic term will be zero. However, if we explicitly calculate it we find that:

$$\left.\frac{\delta F}{\delta Q}\right|_Q : \delta Q \tag{15}$$

Hence, we actually can't get any information out of this particular perturbation.