


# Colouring Digraphs with Bounded Maximum Degree

Lucas Picasarri-Arrieta

National Institute of Informatics


# Digraph dicolouring

- **$k$ -dicolouring** of  $D$ : partition of  $V(D)$  into  $k$  acyclic subdigraphs (*i.e.* no monochromatic directed cycle).



# Digraph colouring


- **$k$ -dicolouring** of  $D$ : partition of  $V(D)$  into  $k$  acyclic subdigraphs (*i.e.* no monochromatic directed cycle).
- **Dichromatic number**  $\vec{\chi}(D)$ : minimum  $k$  s.t.  $D$  admits a  $k$ -dicolouring.



$$\vec{\chi}(D) = 3$$



# Digraph dicolouring

- **$k$ -dicolouring** of  $D$ : partition of  $V(D)$  into  $k$  acyclic subdigraphs (i.e. no monochromatic directed cycle).
- **Dichromatic number**  $\vec{\chi}(D)$ : minimum  $k$  s.t.  $D$  admits a  $k$ -dicolouring.
- Generalizations of **proper colouring** and **chromatic number**.



# Digraph dicolouring

- **$k$ -dicolouring** of  $D$ : partition of  $V(D)$  into  $k$  acyclic subdigraphs (i.e. no monochromatic directed cycle).
- **Dichromatic number**  $\vec{\chi}(D)$ : minimum  $k$  s.t.  $D$  admits a  $k$ -dicolouring.
- Generalizations of **proper colouring** and **chromatic number**.





$$\chi(G) = \vec{\chi}(\overleftrightarrow{G})$$

# From graphs to digraphs: main questions

Given any result on graph colouring, two questions arise:

- **Question 1:** Does it generalize to all digraphs?
- **Question 2:** If it does, can we strengthen it on oriented graphs?



# Maximum Degrees of a digraph

**Max-max-degree:**  $\Delta_{\max}(D) = \max \left\{ \max(d^-(v), d^+(v)) \mid v \in V(D) \right\}.$

**Max-min-degree:**  $\Delta_{\min}(D) = \max \left\{ \min(d^-(v), d^+(v)) \mid v \in V(D) \right\}.$

**Max-geometric-degree:**  $\tilde{\Delta}(D) = \max \left\{ \sqrt{d^-(v) \cdot d^+(v)} \mid v \in V(D) \right\}.$

$$\Delta(G) = \Delta_{\min}(\overleftrightarrow{G}) = \tilde{\Delta}(\overleftrightarrow{G}) = \Delta_{\max}(\overleftrightarrow{G})$$

**First easy bound:**

$$\vec{\chi}(D) \leq \Delta_{\min}(D) + 1 \leq \tilde{\Delta}(D) + 1 \leq \Delta_{\max}(D) + 1$$

# Maximum Degrees of a digraph

**Max-max-degree:**  $\Delta_{\max}(D) = \max \left\{ \max(d^-(v), d^+(v)) \mid v \in V(D) \right\}.$

**Max-min-degree:**  $\Delta_{\min}(D) = \max \left\{ \min(d^-(v), d^+(v)) \mid v \in V(D) \right\}.$

**Max-geometric-degree:**  $\tilde{\Delta}(D) = \max \left\{ \sqrt{d^-(v) \cdot d^+(v)} \mid v \in V(D) \right\}.$

$$\Delta(G) = \Delta_{\min}(\overleftrightarrow{G}) = \tilde{\Delta}(\overleftrightarrow{G}) = \Delta_{\max}(\overleftrightarrow{G})$$

**First easy bound:**

$$\chi(D) \leq \Delta_{\min}(D) + 1 \leq \tilde{\Delta}(D) + 1 \leq \Delta_{\max}(D) + 1$$



# Maximum Degrees of a digraph

**Max-max-degree:**  $\Delta_{\max}(D) = \max \left\{ \max(d^-(v), d^+(v)) \mid v \in V(D) \right\}.$

**Max-min-degree:**  $\Delta_{\min}(D) = \max \left\{ \min(d^-(v), d^+(v)) \mid v \in V(D) \right\}.$

**Max-geometric-degree:**  $\tilde{\Delta}(D) = \max \left\{ \sqrt{d^-(v) \cdot d^+(v)} \mid v \in V(D) \right\}.$

$$\Delta(G) = \Delta_{\min}(\overleftrightarrow{G}) = \tilde{\Delta}(\overleftrightarrow{G}) = \Delta_{\max}(\overleftrightarrow{G})$$

First easy bound:

$$\vec{\chi}(D) \leq \Delta_{\min}(D) + 1 \leq \tilde{\Delta}(D) + 1 \leq \Delta_{\max}(D) + 1$$

# Maximum Degrees of a digraph

**Max-max-degree:**  $\Delta_{\max}(D) = \max \left\{ \max(d^-(v), d^+(v)) \mid v \in V(D) \right\}.$

**Max-min-degree:**  $\Delta_{\min}(D) = \max \left\{ \min(d^-(v), d^+(v)) \mid v \in V(D) \right\}.$

**Max-geometric-degree:**  $\tilde{\Delta}(D) = \max \left\{ \sqrt{d^-(v) \cdot d^+(v)} \mid v \in V(D) \right\}.$

$$\Delta(G) = \Delta_{\min}(\overleftrightarrow{G}) = \tilde{\Delta}(\overleftrightarrow{G}) = \Delta_{\max}(\overleftrightarrow{G})$$

First easy bound:

$$\vec{\chi}(D) \leq \Delta_{\min}(D) + 1 \leq \tilde{\Delta}(D) + 1 \leq \Delta_{\max}(D) + 1$$

# Maximum Degrees of a digraph

**Max-max-degree:**  $\Delta_{\max}(D) = \max \left\{ \max(d^-(v), d^+(v)) \mid v \in V(D) \right\}.$

**Max-min-degree:**  $\Delta_{\min}(D) = \max \left\{ \min(d^-(v), d^+(v)) \mid v \in V(D) \right\}.$

**Max-geometric-degree:**  $\tilde{\Delta}(D) = \max \left\{ \sqrt{d^-(v) \cdot d^+(v)} \mid v \in V(D) \right\}.$

$$\Delta(G) = \Delta_{\min}(\overleftrightarrow{G}) = \tilde{\Delta}(\overleftrightarrow{G}) = \Delta_{\max}(\overleftrightarrow{G})$$



**First easy bound:**

$$\vec{\chi}(D) \leq \Delta_{\min}(D) + 1 \leq \tilde{\Delta}(D) + 1 \leq \Delta_{\max}(D) + 1$$

# Directed Brooks' Theorem

## Theorem (Brooks 1941)

Let  $G$  be a connected graph, then  $\chi(G) \leq \Delta(G)$  unless  $G$  is an **odd cycle** or a **complete graph**.






# Directed Brooks' Theorem

## Theorem (Brooks 1941)

Let  $G$  be a connected graph, then  $\chi(G) \leq \Delta(G)$  unless  $G$  is an **odd cycle** or a **complete graph**.

## Theorem (Mohar 2010)

Let  $D$  be a connected digraph, then  $\vec{\chi}(D) \leq \Delta_{\max}(D)$  unless  $D$  is a **bidirected odd cycle**, a **bidirected complete graph**, or a **directed cycle**.



## Analogues for $\Delta_{\min}$

### Theorem (Aboulker and Aubian 2022)

Deciding  $\vec{\chi}(D) \leq \Delta_{\min}(D)$  is an **NP-complete** problem.

$\Rightarrow$  **No easy characterisation** (unless  $P=NP$ ) in general.

### Theorem (P. 2023)

Let  $\vec{G}$  be an **oriented graph** with  $\Delta_{\min}(\vec{G}) \geq 2$ , then  $\vec{\chi}(\vec{G}) \leq \Delta_{\min}(\vec{G})$ .

### Corollary

Let  $\vec{G}$  be an **orientation** of  $G$  with  $\Delta(G) \leq 5$ , then  $\vec{\chi}(\vec{G}) \leq 2$ .

# Analogues for $\Delta_{\min}$

## Theorem (Aboulker and Aubian 2022)

Deciding  $\vec{\chi}(D) \leq \Delta_{\min}(D)$  is an **NP-complete** problem.

$\Rightarrow$  **No easy characterisation** (unless  $P=NP$ ) in general.

## Theorem (P. 2023)

Let  $\vec{G}$  be an **oriented graph** with  $\Delta_{\min}(\vec{G}) \geq 2$ , then  $\vec{\chi}(\vec{G}) \leq \Delta_{\min}(\vec{G})$ .

## Corollary

Let  $\vec{G}$  be an **orientation** of  $G$  with  $\Delta(G) \leq 5$ , then  $\vec{\chi}(\vec{G}) \leq 2$ .

# Analogues for $\Delta_{\min}$

## Theorem (Aboulker and Aubian 2022)

Deciding  $\vec{\chi}(D) \leq \Delta_{\min}(D)$  is an **NP-complete** problem.

$\implies$  **No easy characterisation** (unless  $P=NP$ ) in general.

## Theorem (P. 2023)

Let  $\vec{G}$  be an **oriented graph** with  $\Delta_{\min}(\vec{G}) \geq 2$ , then  $\vec{\chi}(\vec{G}) \leq \Delta_{\min}(\vec{G})$ .

## Corollary

Let  $\vec{G}$  be an **orientation** of  $G$  with  $\Delta(G) \leq 5$ , then  $\vec{\chi}(\vec{G}) \leq 2$ .



# Asymptotic bounds

## Conjecture (Erdős and Neumann-Lara 1979)

Let  $\vec{G}$  be an oriented graph, then  $\vec{\chi}(\vec{G}) = O\left(\frac{\Delta_{\max}(\vec{G})}{\log \Delta_{\max}(\vec{G})}\right)$ .

**Remark:** Best possible for **random tournaments**.

## Theorem (Harutyunyan and Mohar 2011)

Every oriented graph  $\vec{G}$  with  $\tilde{\Delta}(\vec{G})$  **large enough** satisfies  $\vec{\chi}(\vec{G}) \leq (1 - e^{-13})\tilde{\Delta}(\vec{G})$ .

- $\vec{\chi}(\vec{G}) \leq 0.816\tilde{\Delta}(\vec{G}) + O(1)$ . [Golowich 2016]
- $\vec{\chi}(\vec{G}) \leq 0.707\tilde{\Delta}(\vec{G}) + O(1)$ . (consequence of) [P. 2023]

**Question:** What about digraphs with bounded **biclique number**?

# Asymptotic bounds

## Conjecture (Erdős and Neumann-Lara 1979)

Let  $\vec{G}$  be an oriented graph, then  $\vec{\chi}(\vec{G}) = O\left(\frac{\Delta_{\max}(\vec{G})}{\log \Delta_{\max}(\vec{G})}\right)$ .

**Remark:** Best possible for **random tournaments**.

## Theorem (Harutyunyan and Mohar 2011)

Every oriented graph  $\vec{G}$  with  $\tilde{\Delta}(\vec{G})$  **large enough** satisfies  $\vec{\chi}(\vec{G}) \leq (1 - e^{-13})\tilde{\Delta}(\vec{G})$ .

- $\vec{\chi}(\vec{G}) \leq 0.816\tilde{\Delta}(\vec{G}) + O(1)$ . [Golowich 2016]
- $\vec{\chi}(\vec{G}) \leq 0.707\tilde{\Delta}(\vec{G}) + O(1)$ . (consequence of) [P. 2023]

**Question:** What about digraphs with bounded **biclique number**?

# Asymptotic bounds

## Conjecture (Erdős and Neumann-Lara 1979)

Let  $\vec{G}$  be an oriented graph, then  $\vec{\chi}(\vec{G}) = O\left(\frac{\Delta_{\max}(\vec{G})}{\log \Delta_{\max}(\vec{G})}\right)$ .

**Remark:** Best possible for **random tournaments**.

## Theorem (Harutyunyan and Mohar 2011)

Every oriented graph  $\vec{G}$  with  $\tilde{\Delta}(\vec{G})$  **large enough** satisfies  $\vec{\chi}(\vec{G}) \leq (1 - e^{-13})\tilde{\Delta}(\vec{G})$ .

- $\vec{\chi}(\vec{G}) \leq 0.816\tilde{\Delta}(\vec{G}) + O(1)$ . [Golowich 2016]
- $\vec{\chi}(\vec{G}) \leq 0.707\tilde{\Delta}(\vec{G}) + O(1)$ . (consequence of) [P. 2023]

**Question:** What about digraphs with bounded **biclique number**?

# Asymptotic bounds

## Conjecture (Erdős and Neumann-Lara 1979)

Let  $\vec{G}$  be an oriented graph, then  $\vec{\chi}(\vec{G}) = O\left(\frac{\Delta_{\max}(\vec{G})}{\log \Delta_{\max}(\vec{G})}\right)$ .

**Remark:** Best possible for **random tournaments**.

## Theorem (Harutyunyan and Mohar 2011)

Every oriented graph  $\vec{G}$  with  $\tilde{\Delta}(\vec{G})$  **large enough** satisfies  $\vec{\chi}(\vec{G}) \leq (1 - e^{-13})\tilde{\Delta}(\vec{G})$ .

- $\vec{\chi}(\vec{G}) \leq 0.816\tilde{\Delta}(\vec{G}) + O(1)$ . [Golowich 2016]
- $\vec{\chi}(\vec{G}) \leq 0.707\tilde{\Delta}(\vec{G}) + O(1)$ . (consequence of) [P. 2023]

**Question:** What about digraphs with bounded **biclique number**?

# Asymptotic bounds

## Conjecture (Erdős and Neumann-Lara 1979)

Let  $\vec{G}$  be an oriented graph, then  $\vec{\chi}(\vec{G}) = O\left(\frac{\Delta_{\max}(\vec{G})}{\log \Delta_{\max}(\vec{G})}\right)$ .

**Remark:** Best possible for **random tournaments**.

## Theorem (Harutyunyan and Mohar 2011)

Every oriented graph  $\vec{G}$  with  $\tilde{\Delta}(\vec{G})$  **large enough** satisfies  $\vec{\chi}(\vec{G}) \leq (1 - e^{-13})\tilde{\Delta}(\vec{G})$ .

- $\vec{\chi}(\vec{G}) \leq 0.816\tilde{\Delta}(\vec{G}) + O(1)$ . [Golowich 2016]
- $\vec{\chi}(\vec{G}) \leq 0.707\tilde{\Delta}(\vec{G}) + O(1)$ . (consequence of) [P. 2023]

**Question:** What about digraphs with bounded **biclique number**?

# Reed's conjecture

$\omega$  = size of a largest clique



## Conjecture (Reed 1998)

Every graph  $G$  satisfies  $\chi(G) \leq \left\lceil \frac{\Delta(G)+1+\omega(G)}{2} \right\rceil$ .

# Reed's conjecture


$\omega$  = size of a largest clique



## Conjecture (Reed 1998)

Every graph  $G$  satisfies  $\chi(G) \leq \left\lceil \frac{\Delta(G)+1+\omega(G)}{2} \right\rceil$ .

**Remark:** If true, it is best possible (Gallai's construction).



$$\Delta = 4, \omega = 2, \chi = 4$$

# Reed's conjecture

$\omega$  = size of a largest clique



## Conjecture (Reed 1998)


Every graph  $G$  satisfies  $\chi(G) \leq \left\lceil \frac{\Delta(G)+1+\omega(G)}{2} \right\rceil$ .

**Remark:** If true, it is best possible (Gallai's construction).

## Theorem (Reed 1998)

There exists  $\varepsilon > 0$  s.t. every graph  $G$  satisfies

$$\chi(G) \leq \lceil (1 - \varepsilon)(\Delta(G) + 1) + \varepsilon\omega(G) \rceil.$$



$$\Delta = 4, \omega = 2, \chi = 4$$



# Reed's conjecture

$\omega$  = size of a largest clique



## Conjecture (Reed 1998)

Every graph  $G$  satisfies  $\chi(G) \leq \left\lceil \frac{\Delta(G)+1+\omega(G)}{2} \right\rceil$ .

**Remark:** If true, it is best possible (Gallai's construction).

## Theorem (Reed 1998)

There exists  $\varepsilon > 0$  s.t. every graph  $G$  satisfies

$$\chi(G) \leq \lceil (1 - \varepsilon)(\Delta(G) + 1) + \varepsilon\omega(G) \rceil.$$

- $\varepsilon = 0.038$


[Bonamy et al. 2015]

- $\varepsilon = 0.077$

[Delcourt and Postle 2017]

- $\varepsilon = 0.119$

[Hurley et al. 2022]



$$\Delta = 4, \omega = 2, \chi = 4$$

# An analogue of Reed's conjecture for digraphs

$\vec{\omega}$  = size of a largest biclique



Conjecture (Kawarabayashi and P. 2024)

Every digraph  $D$  satisfies  $\vec{\chi}(D) \leq \left\lceil \frac{\vec{\Delta}(D) + 1 + \vec{\omega}(D)}{2} \right\rceil$ .

# An analogue of Reed's conjecture for digraphs


$\vec{\omega}$  = size of a largest biclique



Conjecture (Kawarabayashi and P. 2024)

Every digraph  $D$  satisfies  $\vec{\chi}(D) \leq \left\lceil \frac{\vec{\Delta}(D) + 1 + \vec{\omega}(D)}{2} \right\rceil$ .

**Remark:** New tight constructions.



$$\vec{\omega} = k, \vec{\Delta} = 3k - 2, \vec{\chi} = 2k$$

# An analogue of Reed's conjecture for digraphs

$\vec{\omega}$  = size of a largest biclique



## Conjecture (Kawarabayashi and P. 2024)


Every digraph  $D$  satisfies  $\vec{\chi}(D) \leq \left\lceil \frac{\tilde{\Delta}(D) + 1 + \vec{\omega}(D)}{2} \right\rceil$ .

**Remark:** New tight constructions.

## Theorem (Kawarabayashi and P. 2024)

There exists  $\varepsilon > 0$  s.t. every digraph  $D$  satisfies

$$\vec{\chi}(D) \leq \left\lceil (1 - \varepsilon)(\tilde{\Delta}(D) + 1) + \varepsilon \vec{\omega}(D) \right\rceil.$$



$$\vec{\omega} = k, \tilde{\Delta} = 3k - 2, \vec{\chi} = 2k$$

## Sketch of the proof – Lemma 1/3 – Exclude large bicliques

### Lemma

Let  $D$  be a digraph with  $\vec{\omega}(D) > \frac{2}{3}(\Delta_{\max}(D) + 1)$ . Then  $D$  has an **acyclic set** of vertices  $I$  such that  $\vec{\omega}(D - I) = \vec{\omega}(D) - 1$ .


# Sketch of the proof – Lemma 1/3 – Exclude large bicliques

## Lemma

Let  $D$  be a digraph with  $\vec{\omega}(D) > \frac{2}{3}(\Delta_{\max}(D) + 1)$ . Then  $D$  has an **acyclic set** of vertices  $I$  such that  $\vec{\omega}(D - I) = \vec{\omega}(D) - 1$ .

## Sketch of the proof

- 1 The intersection graph of the maximum bicliques is a disjoint union of cliques.




# Sketch of the proof – Lemma 1/3 – Exclude large bicliques

## Lemma

Let  $D$  be a digraph with  $\vec{\omega}(D) > \frac{2}{3}(\Delta_{\max}(D) + 1)$ . Then  $D$  has an **acyclic set** of vertices  $I$  such that  $\vec{\omega}(D - I) = \vec{\omega}(D) - 1$ .

### Sketch of the proof

- 1 The intersection graph of the maximum bicliques is a disjoint union of cliques.
- 2 Take the digraph induced by the intersections of bicliques.




# Sketch of the proof – Lemma 1/3 – Exclude large bicliques

## Lemma

Let  $D$  be a digraph with  $\vec{\omega}(D) > \frac{2}{3}(\Delta_{\max}(D) + 1)$ . Then  $D$  has an **acyclic set** of vertices  $I$  such that  $\vec{\omega}(D - I) = \vec{\omega}(D) - 1$ .

## Sketch of the proof

- 1 The intersection graph of the maximum bicliques is a disjoint union of cliques.
- 2 Take the digraph induced by the intersections of bicliques.
- 3 Use the low density to find an acyclic transversal.






# Sketch of the proof – Lemma 1/3 – Exclude large bicliques

## Lemma

Let  $D$  be a digraph with  $\vec{\omega}(D) > \frac{2}{3}(\Delta_{\max}(D) + 1)$ . Then  $D$  has an **acyclic set** of vertices  $I$  such that  $\vec{\omega}(D - I) = \vec{\omega}(D) - 1$ .

## Sketch of the proof

- 1 The intersection graph of the maximum bicliques is a disjoint union of cliques.
- 2 Take the digraph induced by the intersections of bicliques.
- 3 Use the low density to find an acyclic transversal.



## Corollary

If  $D$  is a minimal counter example to  $\vec{\chi}(D) \leq \left\lceil (1 - \varepsilon)(\tilde{\Delta}(D) + 1) + \varepsilon \vec{\omega}(D) \right\rceil$ , then  $\vec{\omega}(D) \leq \frac{2}{3}(\Delta_{\max}(D) + 1)$ .

## Sketch of the proof – Lemma 2/3 – Exclude dense out-neighbourhoods

### Lemma

If  $D$  is a minimal counterexample to  $\vec{\chi}(D) \leq \left\lceil (1 - \varepsilon)(\tilde{\Delta}(D) + 1) + \varepsilon \vec{\omega}(D) \right\rceil$ , for every  $v \in V(D)$ ,  $|A(D[N^+(v)])| \leq (1 - a)\Delta_{\max}^2(D)$ .


# Sketch of the proof – Lemma 2/3 – Exclude dense out-neighbourhoods

## Lemma

If  $D$  is a minimal counterexample to  $\bar{\chi}(D) \leq \left\lceil (1 - \varepsilon)(\tilde{\Delta}(D) + 1) + \varepsilon \tilde{\omega}(D) \right\rceil$ , for every  $v \in V(D)$ ,  $|A(D[N^+(v)])| \leq (1 - a)\Delta_{\max}^2(D)$ .

## Sketch of the proof

- 1 Remove **a few vertices** from  $N^+(v)$  so the remaining vertices have **almost no out-neighbours** out-side.




# Sketch of the proof – Lemma 2/3 – Exclude dense out-neighbourhoods

## Lemma

If  $D$  is a minimal counterexample to  $\bar{\chi}(D) \leq \left\lceil (1 - \varepsilon)(\tilde{\Delta}(D) + 1) + \varepsilon \bar{\omega}(D) \right\rceil$ , for every  $v \in V(D)$ ,  $|A(D[N^+(v)])| \leq (1 - a)\Delta_{\max}^2(D)$ .

## Sketch of the proof

- 1 Remove **a few vertices** from  $N^+(v)$  so the remaining vertices have **almost no out-neighbours** out-side.
- 2 As  $\bar{\omega} \leq \frac{2}{3}\Delta_{\max}$ , find a large matching in  $\bar{D}[N]$ .




# Sketch of the proof – Lemma 2/3 – Exclude dense out-neighbourhoods

## Lemma

If  $D$  is a minimal counterexample to  $\bar{\chi}(D) \leq \left\lceil (1 - \varepsilon)(\tilde{\Delta}(D) + 1) + \varepsilon \bar{\omega}(D) \right\rceil$ , for every  $v \in V(D)$ ,  $|A(D[N^+(v)])| \leq (1 - a)\Delta_{\max}^2(D)$ .

## Sketch of the proof

- 1 Remove **a few vertices** from  $N^+(v)$  so the remaining vertices have **almost no out-neighbours** out-side.
- 2 As  $\bar{\omega} \leq \frac{2}{3}\Delta_{\max}$ , find a large matching in  $\bar{D}[N]$ .
- 3 Use this matching to extend a dicolouring of  $D - N$  to  $D$ , and contradict the minimality of  $D$ .



## Sketch of the proof – Lemma 3/3 – Coloring sparse digraphs

### Lemma

Let  $D$  be a digraph with  $\Delta_{\max}(D) = \Delta$  *large enough*. If for every vertex  $v \in V(D)$ ,  $|A(D[N^+(v)])| \leq (1 - a)\Delta^2$ , then  $\vec{\chi}(D) \leq (1 - \varepsilon)\tilde{\Delta}(D)$ .


# Sketch of the proof – Lemma 3/3 – Coloring sparse digraphs

## Lemma

Let  $D$  be a digraph with  $\Delta_{\max}(D) = \Delta$  *large enough*. If for every vertex  $v \in V(D)$ ,  $|A(D[N^+(v)])| \leq (1 - a)\Delta^2$ , then  $\vec{\chi}(D) \leq (1 - \varepsilon)\tilde{\Delta}(D)$ .

## Sketch of the proof

- 1 Colour every vertex *uniformly at random* with  $[1, \Delta/2]$ .




# Sketch of the proof – Lemma 3/3 – Coloring sparse digraphs

## Lemma

Let  $D$  be a digraph with  $\Delta_{\max}(D) = \Delta$  *large enough*. If for every vertex  $v \in V(D)$ ,  $|A(D[N^+(v)])| \leq (1 - a)\Delta^2$ , then  $\vec{\chi}(D) \leq (1 - \varepsilon)\tilde{\Delta}(D)$ .

## Sketch of the proof

- 1 Colour every vertex **uniformly at random** with  $[1, \Delta/2]$ .
- 2 **Uncolour** every vertex with both an in-neighbour and an out-neighbour of the same colour.






# Sketch of the proof – Lemma 3/3 – Coloring sparse digraphs

## Lemma

Let  $D$  be a digraph with  $\Delta_{\max}(D) = \Delta$  *large enough*. If for every vertex  $v \in V(D)$ ,  $|A(D[N^+(v)])| \leq (1 - a)\Delta^2$ , then  $\vec{\chi}(D) \leq (1 - \varepsilon)\tilde{\Delta}(D)$ .

## Sketch of the proof

- 1 Colour every vertex **uniformly at random** with  $[1, \Delta/2]$ .
- 2 **Uncolour** every vertex with both an in-neighbour and an out-neighbour of the same colour.
- 3 For fixed  $v \in V(D)$ , with high probability, many colours are **repeated** in  $N^+(v)$ .




# Sketch of the proof – Lemma 3/3 – Coloring sparse digraphs

## Lemma

Let  $D$  be a digraph with  $\Delta_{\max}(D) = \Delta$  *large enough*. If for every vertex  $v \in V(D)$ ,  $|A(D[N^+(v)])| \leq (1 - a)\Delta^2$ , then  $\vec{\chi}(D) \leq (1 - \varepsilon)\tilde{\Delta}(D)$ .

## Sketch of the proof

- 1 Colour every vertex **uniformly at random** with  $[1, \Delta/2]$ .
- 2 **Uncolour** every vertex with both an in-neighbour and an out-neighbour of the same colour.
- 3 For fixed  $v \in V(D)$ , with high probability, many colours are **repeated** in  $N^+(v)$ .
- 4 This holds for **all vertices** with positive probability (Lovasz Local Lemma).




# Sketch of the proof – Lemma 3/3 – Coloring sparse digraphs

## Lemma

Let  $D$  be a digraph with  $\Delta_{\max}(D) = \Delta$  *large enough*. If for every vertex  $v \in V(D)$ ,  $|A(D[N^+(v)])| \leq (1 - a)\Delta^2$ , then  $\bar{\chi}(D) \leq (1 - \varepsilon)\tilde{\Delta}(D)$ .

## Sketch of the proof

- 1 Colour every vertex **uniformly at random** with  $[1, \Delta/2]$ .
- 2 **Uncolour** every vertex with both an in-neighbour and an out-neighbour of the same colour.
- 3 For fixed  $v \in V(D)$ , with high probability, many colours are **repeated** in  $N^+(v)$ .
- 4 This holds for **all vertices** with positive probability (Lovasz Local Lemma).
- 5 **Greedy** colour the uncoloured vertices.



## Further Research

- **Problem:** Prove the existence of  $\varepsilon > 0$  such that every oriented graph  $\vec{G}$  satisfies  $\vec{\chi}(\vec{G}) \leq \lceil (1 - \varepsilon)\Delta^+(\vec{G}) \rceil$ .
- **Problem:** Find the values of  $k \in [1, \Delta + 1]$  for which  $k$ -DICOLOURABILITY is solvable in polynomial time on digraphs  $D$  with  $\Delta_{\max}(D) = \Delta$ .  
Undirected case (Molloy and Reed 2014):  $k \in [\Delta - \sqrt{\Delta}, \Delta + 1]$ .
- **Conjecture (Neumann-Lara 1982):** Every oriented planar graph is 2-colourable.

Thank you!

## Further Research

- **Problem:** Prove the existence of  $\varepsilon > 0$  such that every oriented graph  $\vec{G}$  satisfies  $\vec{\chi}(\vec{G}) \leq \lceil (1 - \varepsilon)\Delta^+(\vec{G}) \rceil$ .
- **Problem:** Find the values of  $k \in [1, \Delta + 1]$  for which  $k$ -DICOLOURABILITY is solvable in polynomial time on digraphs  $D$  with  $\Delta_{\max}(D) = \Delta$ .

Undirected case (Molloy and Reed 2014):  $k \in [\Delta - \sqrt{\Delta}, \Delta + 1]$ .

- **Conjecture (Neumann-Lara 1982):** Every oriented planar graph is 2-colourable.

Thank you!

## Further Research

- **Problem:** Prove the existence of  $\varepsilon > 0$  such that every oriented graph  $\vec{G}$  satisfies  $\vec{\chi}(\vec{G}) \leq \lceil (1 - \varepsilon)\Delta^+(\vec{G}) \rceil$ .
- **Problem:** Find the values of  $k \in [1, \Delta + 1]$  for which  $k$ -DICOLOURABILITY is solvable in polynomial time on digraphs  $D$  with  $\Delta_{\max}(D) = \Delta$ .  
Undirected case (Molloy and Reed 2014):  $k \in [\Delta - \sqrt{\Delta}, \Delta + 1]$ .
- **Conjecture (Neumann-Lara 1982):** Every oriented planar graph is 2-colourable.

Thank you!

## Further Research

- **Problem:** Prove the existence of  $\varepsilon > 0$  such that every oriented graph  $\vec{G}$  satisfies  $\vec{\chi}(\vec{G}) \leq \lceil (1 - \varepsilon)\Delta^+(\vec{G}) \rceil$ .
- **Problem:** Find the values of  $k \in [1, \Delta + 1]$  for which  $k$ -DICOLOURABILITY is solvable in polynomial time on digraphs  $D$  with  $\Delta_{\max}(D) = \Delta$ .  
Undirected case (Molloy and Reed 2014):  $k \in [\Delta - \sqrt{\Delta}, \Delta + 1]$ .
- **Conjecture (Neumann-Lara 1982):** Every oriented planar graph is 2-colourable.

Thank you!